

2020 Edition with All New Updates/Qs Covered up to Jan 2020

Includes Recent Updates of WHO 2015 Lung, WHO 2016 Male Genital System  
and WHO 2017 CNS/Hematopoietic Classifications

Compiled by Leading Faculties and Subject Experts of Pathology

# Complete Review of Pathology & Hematology for NBE

6<sup>th</sup>  
Edition

Covering 2500+ Qs with Explanations, 100+ IBQs & 1000+ Colored Illustrations/Images

Fully Updated from Robbin's 10/e (Basic Edition), Robbin's 9/e,  
Wintrobe's 13/e, Sternberg's 6/e, Ackerman's 11/e, WHO 2015 Lung,  
WHO 2016 Male Genital System and WHO 2017 CNS/Hematopoietic Classifications

Includes  
Chapter-wise  
**NEXT**  
Pattern  
Questions

Special Feature  
First Book Covering  
Sample Video Questions



Recent Qs (Jan) 2020 – 2010

AIIMS Nov 2019 – 2010

JIPMER Dec 2019 – 2010

PGI May 2019 – 2010

AIPG 2012 – 2010



CBS Exam Books



*Vandana Puri*

*My dear students, whenever you find yourself doubting how far you can go, just remember, how far you have already come. Life is just 10% what happens to you and 90% how you respond to it so give it your best shot. I just hope, this book, provides you the best confidence, by which you can face this tough journey of life and reach the most beautiful destination of your life. Wish you loads of success...*



*Praveen Kr Gupta*

*There are no secrets to success. It is the result of right guidance, hard work, and learning from the best. I am sure, this book will serve all your purposes.*

Praveen Kr Gupta • Vandana Puri





# JOIN CBS EXAM BOOKS

## ON FACEBOOK & INSTAGRAM

### & GET A CHANCE TO WIN CBS EXAM BOOKS (New Editions)

#### About CBS Exam Books FB Page

Total Books given as Complimentary: **10,000+** / Amount of PrepCash Distributed: **2,00,000/-**  
Year of Establishment: **2015** / Total likes: **32500+** / Total Posts Shared: **5000+** / Total Contests Conducted: **500+**

#### Join us to Get

- Sample pages of the Books
- Sample Chapter of the Books
- IBQs with Explanations to Clear the Concepts
- Synopses of Important Topics
- Live Session with Authors
- Book Reviews
- Last Minute Revision Tips
- Success Stories of the Toppers
- FAQs from Examinations

& many more...

#### Join us to Participate in

- Books Giveaway
- Online Test Series
- Special Offers
- Bid2Win
- Share, Refer, Like & Win
- Fastest Finger First

& other contests...

#### Join us to win

- Complimentary Books
- Customized Gifts
- PrepCash
- Discount Coupons
- Exciting Offers
- & many more...





CBS FB QR Code

Scan the QR Codes  
to Join CBS Exam Books



CBS Instagram QR Code





# Complete Review of **Pathology** & **Hematology** for NBE

Covering 2500+ Qs with Explanations, 100+ IBQs & 1000+ Colored Illustrations/Images

Fully Updated from Robbin's 10/e (Basic Edition), Robbin's 9/e,  
Wintrobe's 13/e, Sternberg's 6/e, Ackerman's 11/e, WHO 2015 Lung,  
WHO 2016 Male Genital System and WHO 2017 CNS/Hematopoietic Classifications

Includes  
Chapter-wise  
NEXT Pattern  
Questions

— ■ **Sixth Edition** ■ —

**Praveen Kumar Gupta** MBBS, MD

MBBS (Honours), Medical College and Hospital, Kolkata, West Bengal  
MD, Ex Senior Resident (Laboratory Medicine), AIIMS, New Delhi

**Consultant Pathologist & Lab Head**

Primus Super Speciality Hospital, New Delhi

**Vandana Puri** MBBS, MD (Pathology), DNB (Pathology), MNAMS

MBBS, CMC, Coimbatore, Tamil Nadu  
MD & Ex SR Lady Hardinge Medical College, New Delhi  
Ex SRA, Hematology, AIIMS, New Delhi  
Ex Assistant Professor, UCMS, New Delhi

**Associate Professor**

Department of Pathology  
Lady Hardinge Medical College, New Delhi



**CBS Publishers & Distributors Pvt Ltd**

• New Delhi • Bengaluru • Chennai • Kochi • Kolkata • Mumbai  
• Hyderabad • Nagpur • Patna • Pune • Vijayawada





ISBN: 978-81-945783-3-8

Copyright © Authors & Publishers

**Sixth Edition: 2020**

**Fifth Edition: 2019-20**

**Fourth Edition: 2018**

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system without permission, in writing, from the authors and the publishers.

Published by **Satish Kumar Jain** and produced by **Varun Jain** for

**CBS Publishers and Distributors Pvt Ltd**

4819/XI Prahlad Street, 24 Ansari Road, Daryaganj, New Delhi 110 002, India.

Ph: 23289259, 23266861, 23266867 Website: [www.cbspd.com](http://www.cbspd.com)

Fax: 011-23243014

e-mail: [delhi@cbspd.com](mailto:delhi@cbspd.com); [cbspubs@airtelmail.in](mailto:cbspubs@airtelmail.in).

**Corporate Office:** 204 FIE, Industrial Area, Patparganj, Delhi 110 092

Ph: +91-11-4934 4934 Fax: 4934 4935

e-mail: [feedback@cbspd.com](mailto:feedback@cbspd.com); [bhupesharora@cbspd.com](mailto:bhupesharora@cbspd.com)

#### DISCLAIMER

This book contains questions based on important topics frequently asked in previous years National Level PG Entrance Examinations & State Level Examinations in India. Often repeated topics and sub-topics have been included for students' benefit. We do not claim that these questions are exact or similar to questions asked in any recent examinations in India. If any such similarity is found, it is purely coincidental and by chance.

#### Branches

---

- **Bengaluru:** Seema House 2975, 17th Cross, K.R. Road, Banasankari 2nd Stage, Bengaluru 560 070, Karnataka  
Ph: +91-80-26771678/79 Fax: +91-80-26771680 e-mail: [bangalore@cbspd.com](mailto:bangalore@cbspd.com)
- **Chennai:** No. 7, Subbaraya Street, Shenoy Nagar, Chennai 600 030, Tamil Nadu  
Ph: +91-44-26680620, 26681266 Fax: +91-44-42032115 e-mail: [chennai@cbspd.com](mailto:chennai@cbspd.com)
- **Kochi:** 68/1534, 35, 36-Power House Road, Opp. KSEB, Cochin-682018, Kochi, Kerala  
Ph: +91-484-4059061-65 Fax: +91-484-4059065 e-mail: [kochi@cbspd.com](mailto:kochi@cbspd.com)
- **Kolkata:** 6/B, Ground Floor, Rameswar Shaw Road, Kolkata-700014 (West Bengal)  
Ph: +91-33-22891126, 22891127, 22891128 e-mail: [kolkata@cbspd.com](mailto:kolkata@cbspd.com)
- **Mumbai:** 83-C, Dr E Moses Road, Worli, Mumbai-400018, Maharashtra  
Ph: +91-22-24902340/41 Fax: +91-22-24902342 e-mail: [mumbai@cbspd.com](mailto:mumbai@cbspd.com)

#### Representatives

---

- |                                   |                                    |
|-----------------------------------|------------------------------------|
| • <b>Hyderabad</b> +91-9885175004 | • <b>Patna</b> +91-9334159340      |
| • <b>Pune</b> +91-9623451994      | • <b>Vijayawada</b> +91-9000660880 |

Printed at :



---

## *Dedicated to*

*My parents for their love and support, my loving wife Dr Meenakshi and cute little angel Myra, it is a privilege to share my life and love with you.*

**Dr Praveen Kumar Gupta**

*My soulmate Dr Shakti Tiwari, my kids Arsh and Amaira, my parents for their unconditional support and my teachers and students for their inspiration, love and guidance.*

**Dr Vandana Puri**

---



# About the Authors

**Dr Praveen Kr Gupta, MBBS, MD**, is presently working as Consultant Pathologist and Lab Head at Primus Super Speciality Hospital, New Delhi. He did his MBBS from Medical College and Hospital, Kolkata, West Bengal and MD as well as Senior Residency from the prestigious All India Institute of Medical Sciences, New Delhi. He has an excellent academic record and has been the recipient of many awards in various states and national level competitions in Clinical Pathology and Hematology. He has authored several books, including the popular AIIMS PGMEET series and has many National and International publications to his name. Teaching is his passion and he is well known for his emphasis on concept building and focused learning. He is the co-founder of "AIM4AIIMS-PG" group, which has been mentoring and guiding over 50,000 PG aspirants all over the country. It gives us immense pleasure to announce the release of Pathology App by Dr Praveen Kr Gupta, which will aid many students all over the country to understand the basic concepts of pathology. This is available on Android and IOS and can be bought for a nominal cost.



Join Author's **Facebook Discussion Group** – [www.facebook.com/Pathology by Dr Praveen](https://www.facebook.com/Pathology%20by%20Dr%20Praveen%20Gupta/) or <https://www.facebook.com/groups/390769121404969/> or


Scan this QR Code

**Dr Vandana Puri, MBBS, MD (Pathology), DNB (Pathology), MNAMS**, is presently working as Associate Professor Pathology at Lady Hardinge Medical College and Associated Hospitals, New Delhi. She did her MBBS from Coimbatore Medical College, Coimbatore, Tamil Nadu, India. Following this, she pursued her MD and Senior Residency in Pathology from Lady Hardinge Medical College, New Delhi. She did her Senior Research Associateship in Hematology from AIIMS, New Delhi. Before joining Lady Hardinge Medical College, she had worked as Assistant Professor in Department of Pathology, University College of Medical Sciences and GTB Hospital, Delhi. She has more than 40 International and National Publications to her name. Besides, she has been co-author of important chapters in many National and International books. She has vast teaching experience and she orients the students in concept building in advanced pathology by making them understand the basics of pathology. She has more than 54,000 student followers at her Facebook group "Dr Vandana Pathology Study Circle" who are regularly benefitted and updated by discussions. She is also author of book-PICS (*coming soon*) and has both android and iOS apps with DocTutorials where in you can subscribe for pathology. You can sign up both on apps and website too ([www.doctutorials.com](http://www.doctutorials.com))



Follow authors you tube channel and Facebook group

<https://www.youtube.com/channel/UCrYYtBBSscRAhAaUyMJTKpA> or



Scan this QR Code

&

<https://www.Facebook.com/groups/1444287919221567/> or



Scan this QR Code

Join Author's **Facebook Discussion Group** – [www.facebook.com/Dr Vandana Pathology study circle](https://www.facebook.com/Dr%20Vandana%20Pathology%20study%20circle/)



# Preface

“The best way to predict your future is to create it.”

Abraham Lincoln

**A**chieving success in Postgraduate Medical Entrance Exams (PGMEE) and to be able to pursue a specialty of choice in the premier medical institutes of the country has been the ultimate dream of every medical student. To make this dream a reality—a belief in self, setting realistic goals, ‘intelligent hard work’, ‘smart and productive studying’ and selection of the right study material are some of the very important factors.

*To cover more than 15 subjects of medical curriculum in a limited time span is definitely not an easy task, however it's not impossible.*

As, impossible itself says I'm possible!

It is very important to **strengthen your fundamentals and get your basics right! Pathology is the backbone of medical science.** One cannot understand clinical subjects, like **Medicine, Surgery, Pediatrics** and **Gynecology** without being well-versed with the basic pathology of diseases. By the time a student appears for any PGMEE, usually during or after internship, preclinical subjects, like **Pathology tend to fade away from the memory** as the clinical subjects largely dominate the mind.

In the recent NBE pattern examinations, **25–30 MCQs out of 300** are being asked from core pathology. Most of the MCQs asked from Medicine and Surgery also have one or two options related to Pathology, which makes contribution of **50–60 MCQs from pathology; which is almost 15–20% of any examination.** In AIIMS Nov 2015 exam, there were over 60 Image-based Questions which re-emphasized the importance of conceptual learning with figures rather than nearly solving repeats.

Solving recent NBE based, AIIMS, PGI and other state entrance exam questions not only requires **mastery over the repeated MCQs** but also **thorough knowledge** of topics with **special emphasis on concepts and high-yielding facts.**

For the first time AIIMS New Pattern 2019 Model Questions have been added to ace your preparation practice before upcoming AIIMS examination. Chapter-wise NEXT Pattern Qs have been added as per the changing scenario of upcoming examination.

With so many MCQ-based guidebooks flooding the market, there was a felt need for a book on Pathology which is *student friendly, lucid, interesting, not full of big paragraphs loaded with heavy information, but which covers all topics and relevant information that can be revised in limited time and is updated with all recent facts and advances.*

This book is useful **not only** for the students preparing for various PGME exams, but also for the **undergraduate students and pathology students pursuing postgraduation**, it will also help in concept building as well as **quick revision.**

Praveen Kumar Gupta

Vandana Puri

# Highlights of 6<sup>th</sup> edition

The overwhelming response by the students and 100% Pathology (including many Medicines, Pediatrics) MCQs from AIIMS, PGI, JIPMER and NEET entrance exams from this book made it a roaring sensation in the market. It gives us immense pleasure to bring to you the sixth edition of Complete Review of Pathology & Hematology with fully revised content, most recent updates, recent questions, high-yield points and annexures.

- **Pretexts:** Detailed yet concise, pointwise overview of the entire topic. High-yielding MCQs and repeated MCQs have been highlighted.
- **Flow charts:** Flow diagrams have been added wherever necessary for easy understanding including most updated points from recent editions of books. **New chapters on “Diseases of Muscle and Tumors of Bones and Joints” have been added. Over 80 new labelled and explained images have been added for proper explanation of texts and better understanding of Image-Based Questions.**
- **Recent updates:** All additions and changes are according to latest Robbin’s 10th edition and are highlighted separately, so that students are updated with recent advances in relevant topics. These are potential MCQs in upcoming PGMEEs.
- **MCQs:** All MCQs of NBE pattern up to January 2020, AIIMS up to November 2019, PGI up to December 2019 and JIPMER up to November 2019 have been included, and the recent Qs have been highlighted separately. MCQs have been arranged chronologically with recent ones coming earlier, so that more stress is put on recent pattern questions. Repeated MCQs have been clubbed together to avoid unnecessary duplication and time wastage.
- **AIIMS new pattern 2019 model questions:** Since you all must be aware that AIIMS PG exam has announced new patterns of MCQs from this session onwards, we have added many new pattern Qs including “match the following, arrange in sequence, reason and assertion and concept-based multiple answers” in the current edition to benefit the aspiring students.
- **Image-based MCQs:** As PGMEEs have shifted to online CBT exams, there has been an increase in image-based questions. We have included potential image-based MCQs at the end of every topic to familiarize the students with the same. **Students can learn identification points of images from the pretext and then quickly answer Image-based Questions for strong hold in subject.**
- **Video-based questions have been added to cater recent AIIMS pattern.**
- **Authentic explanations:** Explanations from standard and recent edition textbooks have been provided for each answer. Difficult and controversial MCQs have been explained in detail discussing each option and excluding the incorrect ones. This will help a student to develop his/her analytical skills.
- **Annexures:** A new Annexure on Autoantibodies in Autoimmune Disease has been added for quick revision.
- **NEXT pattern questions** have been added.

Although utmost care has been taken to avoid all possible errors, some minor errors might have crept in inadvertently. We request the readers to kindly point out the same and give their valuable suggestions or feedback on the address provided in About the Authors page.

*We wish you all the very best for your upcoming exams and for your bright future!*



# Acknowledgements

Firstly, we would like to express our eternal gratitude to the blessings of Almighty GOD. We thank our parents for their blessings, everlasting love and support.

**We wish to sincerely thank our teachers and faculty for being the source of our inspiration and knowledge, and who helped us to achieve a lot in our personal and professional lines:**

- Prof Manjula Jain-ex. Director Professor, Department of Pathology, LHMC
- Prof AK Mukhopadhyay-Professor & Head, Department of Laboratory Medicine, AIIMS
- Prof Sunita Sharma-Director Professor and Head, Department of Pathology, LHMC
- Prof Renu Saxena-Head of Department, Department of Hematology, AIIMS
- Prof Usha Rusia-Previous Head of Department, Department of Pathology, UCMS
- Prof Meera Sikka -Head of Department, Department of Pathology, UCMS
- Prof Sarman Singh-Professor, Department of Laboratory Medicine, AIIMS
- Prof M Irshad-Professor, Department of Laboratory Medicine, AIIMS
- Prof Kiran Agarwal-Director Professor, Department of Pathology, LHMC
- Prof Shilpi Agarwal-Director Professor, Department of Pathology, LHMC
- Prof Anita Nangia-Director Professor, Department of Pathology, LHMC
- Prof Shailja Shukla-Director Professor, Department of Pathology, LHMC
- Dr Sonal Sharma & Dr Mrinalini Kotru-Professor, Department of Pathology, UCMS
- Dr Subhadra Sharma, Purva Mathur & Dr Arulselvi S-Associate Professor, Department of Laboratory Medicine, AIIMS
- Dr Sangeeta Pahuja & Dr Smita Singh-Associate Professor, Department of Pathology, LHMC
- Dr Mukta Pujani-Associate Professor, Department of Pathology, ESI Medical College, Faridabad
- Dr Venkateswaran Iyer (Associate Professor), Dr Prasenjit Das, Dr Sudip Arava, Dr Soumya (Assistant Professor) Department of Pathology, AIIMS, Prateek Bhatia (Asst Professor, PGI, Chandigarh), Dr Manupriya Nain (Assistant Professor, Hindu Rao Hospital)
- Dr Gaurav Chhabra, Dr Sudip Datta, Dr Raghavendra L, Dr Shyam Prakash-Assistant Professor, Lab Medicine, AIIMS

**A special thanks to Dr Meenakshi (Dr Praveen's wife) for her immense motivation and support. She has been instrumental in conceptualization, formatting and designing of our book.**

***We thank our friends/students/well-wishers, for their invaluable contribution & support in writing this book***

- Dr Shakti Tiwari-Consultant Anesthetist, Sant Parmanand Hospital (Dr Vandana's husband)
- Dr Sawan Kumar-Consultant Pathologist, Primus Hospital, New Delhi, Dr Apoorv Singh (MBBS, AIIMS)
- Dr Zainab Vora, Dr Ravi Sharma, Dr Amit Gupta, Dr Anil Shekhawat (AIIMS, New Delhi)
- Dr Param Prakash, Postgraduate Resident, LHMC for helping in framing of NEXT Pattern Questions

***We are also grateful to our following well-wishers, whose ideas and suggestions have helped us immensely***

- Dr Akhilesh Raj Jhamad, Dr Rajat Jain, Dr Vivek Jain, Dr Thameem Saif, Dr Mukesh Bhatia, Dr Vineet, Dr Nachiketa Bhatia, Dr Apurv Mehra, Dr Sourav Bhatia, Dr Shashwat Ray, Dr Manish Soni, Dr Ashish, Dr Sonu Panwar, Dr Surendra Nath Reddy, Dr Rajeswar Gudadhe, Dr Ashwini, Dr Saurabh Bhatia, Dr Sidharth Sekhar Mishra, Mr Dhruv, Mr Amit Bhatia, Mr Rajiv, Mr Roshan, Dr T Piyush, Dr Kumar Sarvottam, Dr Pritesh Singh, Dr Manjunatha A.

We are extending our special thanks to **Mr Satish Kumar Jain** (Chairman) and **Mr Varun Jain** (Managing Director), M/s CBS Publishers and Distributors Pvt Ltd for their wholehearted support in publication of this book. We have no words to describe the role, efforts, inputs and initiatives undertaken by **Mr Bhupesh Arora** (Vice President - Publishing & Marketing, PGME&N and Nursing Division) for helping and motivating us.

We sincerely thank the entire CBS team for bringing out the book with utmost care and attractive presentation. We would like to thank Dr Mrinalini Bakshi (Editorial Head & Content Strategist) for her editorial support and Ms Nitasha Arora (Production Head & Content Strategist), Dr Anju Dhir (Project Manager & Senior Scientific Coordinator), Mr Shivendu Bhushan Pandey (Senior Editor), Mr Ashutosh Pathak (Senior Proof Reader) and all the production team members Mr Chaman Lal, Mr Prakash Gaur, Mr Phool Kumar, Mr Bunty Kashyap, Ms Tahira Parveen, Ms Manorama Gupta, Ms Babita Verma, Mr Chander Mani, Mr Raju Sharma, Mr Manoj Chaudhary, Mr Vikram Chaudhary, Mr Manoj Malakar, Mr Arun Kumar and Mr Rahul Negi for devoting laborious hours in designing and typesetting of the book.





# Contents

Preface.....	v
Acknowledgements .....	vii
Recent Pattern Questions 2020 at a Glance .....	xiv
AIIMS New Pattern 2019 Model Questions.....	xvii
Sample Video Questions.....	xxxi

## ANNEXURES


1. Important Special Stains and Fixatives	xxxii
2. Important Fixatives	xxxiii
3. Cytokines and Cytokine Receptors	xxxiv
4. Type of Modified Macrophages	xxxv
5. Types of Giant Cells	xxxv
6. Important Translocations	xxxvi
7. Important Genes & Chromosome	xxxvi
8. Chromosomal Abnormality	xxxvii
9. Cancer Predisposition Syndromes and Associated Gene	xxxviii
10. CD Markers used for Hematolymphoid Neoplasms, Location/Characteristics	xxxix
11. Features of the Peripheral Blood Smear	xli
12. Blood-Clotting Factors	xlii
13. Crystals in Urine	xlii
14. Different Types of Urinary Casts	xliii
15. Rosettes	xliv
16. Familial Cancer Syndromes with Cutaneous Manifestations	xlvi
17. Relationship between Proteins and Neurodegenerative Diseases	xlvi
18. Pattern of Injury and Associated Agents	xlvi
19. Morphological Differentiation of Malaria Parasites	xlvi
20. Autoantibodies in Autoimmune Disease	xlvi

## 1. CELL AS A UNIT OF HEALTH AND DISEASE

Theory.....	1
NEXT Pattern Questions.....	10
Multiple Choice Questions.....	11
Answers with Explanations .....	13

## 2. CELL ADAPTATION, INJURY AND DEATH

Theory.....	17
NEXT Pattern Questions.....	34
Image-Based Questions .....	35
Answers of Image-Based Questions .....	36
Multiple Choice Questions .....	37
Answers with Explanations .....	42



### 3. INFLAMMATION AND REPAIR

Theory.....	49
NEXT Pattern Questions.....	67
Image-Based Questions.....	68
Answers of Image-Based Questions.....	69
Multiple Choice Questions.....	70
Answers with Explanations.....	76

### 4. HEMODYNAMICS

Theory.....	85
Image-Based Questions.....	93
Answers of Image-Based Questions.....	94
Multiple Choice Questions.....	95
Answers with Explanations.....	97

### 5. GENETIC DISORDERS

Theory.....	99
NEXT Pattern Questions.....	112
Image-Based Questions.....	114
Answers of Image-Based Questions.....	115
Multiple Choice Questions.....	116
Answers with Explanations.....	124

### 6. DISEASES OF THE IMMUNE SYSTEM

Theory.....	133
NEXT Pattern Questions.....	155
Image-Based Question.....	156
Answer of Image-Based Question.....	156
Multiple Choice Questions.....	157
Answers with Explanations.....	166


### 7. NEOPLASIA

Theory.....	179
NEXT Pattern Questions.....	193
Image-Based Questions.....	194
Answers of Image-Based Questions.....	196
Multiple Choice Questions.....	197
Answers with Explanations.....	204

### 8. DISEASES OF INFANCY AND CHILDHOOD

Theory.....	215
Image-Based Questions.....	220
Answers of Image-Based Questions.....	220
Multiple Choice Questions.....	221
Answers with Explanations.....	223





## 9. WHITE BLOOD CELLS AND ITS DISORDERS

Theory.....	225
Image-Based Questions .....	246
Answers of Image-Based Questions .....	248
Multiple Choice Questions .....	249
Answers with Explanations .....	260

## 10. RED BLOOD CELLS AND ITS DISORDERS

Theory.....	273
Image-Based Questions .....	288
Answers of Image-Based Questions .....	289
Multiple Choice Questions .....	290
Answers with Explanations .....	300

## 11. BLEEDING AND COAGULATION DISORDERS

Theory.....	313
Image-Based Question .....	324
Answer of Image-Based Question.....	324
Multiple Choice Questions .....	325
Answers with Explanations .....	331

## 12. CARDIOVASCULAR SYSTEM AND ITS DISORDERS


Theory.....	339
NEXT Pattern Questions .....	363
Image-Based Questions .....	365
Answers of Image-Based Questions .....	366
Multiple Choice Questions .....	367
Answers with Explanations .....	374

## 13. RESPIRATORY SYSTEM AND ITS DISORDERS

Theory.....	383
NEXT Pattern Questions .....	399
Image-Based Questions .....	400
Answers of Image-Based Questions .....	402
Multiple Choice Questions .....	403
Answers with Explanations .....	409

## 14. GASTROINTESTINAL TRACT AND ITS DISORDERS

Theory.....	417
NEXT Pattern Questions .....	440
Image-Based Questions .....	441
Answers of Image-Based Questions .....	442
Multiple Choice Questions .....	443
Answers with Explanations .....	448



## 15. LIVER, GALLBLADDER, PANCREAS AND ITS DISORDERS

Theory.....	455
NEXT Pattern Questions.....	472
Image-Based Questions.....	473
Answers of Image-Based Questions.....	475
Multiple Choice Questions.....	476
Answers with Explanations.....	483

## 16. RENAL SYSTEM AND ITS DISORDERS

Theory.....	493
NEXT Pattern Questions.....	511
Image-Based Questions.....	512
Answers of Image-Based Questions.....	513
Multiple Choice Questions.....	514
Answers with Explanations.....	522

## 17. MALE AND FEMALE GENITOURINARY TRACT

Theory.....	529
Image-Based Questions.....	547
Answers of Image-Based Questions.....	547
Multiple Choice Questions.....	548
Answers with Explanations.....	552

## 18. BREAST

Theory.....	557
NEXT Pattern Question.....	565
Multiple Choice Questions.....	566
Answers with Explanations.....	568


## 19. ENDOCRINE SYSTEM AND ITS DISORDERS

Theory.....	571
NEXT Pattern Questions.....	583
Image-Based Questions.....	584
Answers of Image-Based Questions.....	586
Multiple Choice Questions.....	587
Answers with Explanations.....	590

## 20. SKIN AND ITS DISORDERS

Theory.....	597
Image-Based Questions.....	607
Answers of Image-Based Questions.....	608
Multiple Choice Questions.....	609
Answers with Explanations.....	610





## 21. CENTRAL NERVOUS SYSTEM AND ITS DISORDERS

Theory.....	613
NEXT Pattern Questions.....	629
Image-Based Questions.....	630
Answers of Image-Based Questions.....	631
Multiple Choice Questions.....	632
Answers with Explanations.....	636

## 22. BLOOD BANKING AND TRANSFUSION MEDICINE

Theory.....	641
Image-Based Questions.....	646
Answers of Image-Based Questions.....	647
Multiple Choice Questions.....	648
Answers with Explanations.....	650

## 23. TUMORS OF SOFT TISSUE & HEAD & NECK

Theory.....	653
Multiple Choice Questions.....	658
Answers with Explanations.....	660

## 24. DISEASES OF MUSCLES

Theory.....	663
Image-Based Questions.....	666
Answers of Image-Based Questions.....	666
Multiple Choice Questions.....	667
Answers with Explanations.....	667

## 25. TUMORS OF BONE AND JOINTS


Theory.....	669
NEXT Pattern Question.....	673
Image-Based Questions.....	674
Answers of Image-Based Questions.....	674
Multiple Choice Questions.....	674
Answers with Explanations.....	674

## 26. RECENT TECHNIQUES IN PATHOLOGY

Theory.....	676
Image-Based Questions.....	680
Answers of Image-Based Questions.....	680


## 27. STAINS AND FIXATIVES

Multiple Choice Questions.....	682
Answers with Explanations.....	685



## Recent Pattern Questions 2020 at a Glance

1. Histological picture of a lesion excised from the right cervical region is shown below. What is your diagnosis?



- Necrotizing granulomatous inflammation
- Neurofibroma
- Schwannoma
- Hodgkin lymphoma


2. Which of the following has a major role in thrombus formation?

- Endothelial injury
- Vasoconstriction
- Platelet activation
- Coagulation cascade

3. Mutation in DNA Helicase causes

- Wermer syndrome
- Werner syndrome
- Sipple syndrome
- Autoimmune lymphoproliferative syndrome

4. Which of the following mode of inheritance is shown below?



- AD
- AR
- X linked dominant
- X linked recessive

5. An 8-year-old child presented with history of recurrent infections. The child had rashes. Investigations revealed low platelets. What could be the probable cause?

- Job syndrome
- Wiskott-Aldrich syndrome
- Henoch-Schonlein purpura
- Hyper IgM syndrome


6. B cells are induced to produce IgE by which of the following?

- IL 2
- IL 4
- IL 1
- IL 6

7. Graft between Identical twins is:


- Allograft
- Xenograft
- Isograft
- Autograft

8. Identify the parasite in the intestinal biopsy of a HIV positive patient.



- Giardia
- CMV
- Amoebic colitis
- Cryptosporidium

9. A 75-year-old male, known smoker presented to pulmonology department with history of cough. Biopsy was taken which showed the following. What is the change shown?




- Dysplasia
- Metaplasia
- Hyperplasia
- Atrophy

Ans.


- a
- a
- b
- a
- b
- b
- c
- d
- b




10. Cardiac biopsy of a patient who died following myocardial infarction is shown below. What is the finding is a feature of reperfusion injury?



- a. Waviness of fibers  
b. Neutrophils in cardiac muscle  
c. Eosinophilic contraction bands  
d. Swelling of cells
11. Bulky friable vegetations are seen in:  
a. Rheumatic carditis  
b. Infective endocarditis  
c. Libman sack's endocarditis  
d. Non-bacterial thrombotic endocarditis
12. Histological picture of a lesion excised from the right cervical region is shown below. What is your diagnosis?




- a. Necrotizing granulomatous inflammation  
b. Neurofibroma  
c. Schwannoma  
d. Hodgkin lymphoma
13. A 30-year old male presented with history of dyspnoea, cough and sputum production. The patient died of respiratory failure. Gross image of lung is shown below. What is the likely etiology?



- a. Cystic fibrosis  
b. Mutation in dynein arms  
c. Alpha 1 antitrypsin deficiency  
d. Antibodies against type IV collagen


14. A 75-year-old male, known smoker presented to pulmonology department with history of cough. Biopsy was taken which showed the following. What is the change shown?



- a. Dysplasia  
b. Metaplasia  
c. Hyperplasia  
d. Atrophy
15. A 23-year-old lady presented with diarrhea, vomiting and poor appetite. Biopsy showed crypt hyperplasia, villous atrophy and CD8+ cells in the lamina propria. Skin manifestations have been shown. What could be the diagnosis?



- a. Whipple's disease  
b. Chronic pancreatitis  
c. Environmental enteropathy  
d. Celiac disease
16. Identify the parasite in the intestinal biopsy of a HIV positive patient.




- a. Giardia  
b. CMV  
c. Amoebic colitis  
d. Cryptosporidium


Ans.

10. c  
11. b  
12. a  
13. c  
14. b  
15. d  
16. d






17. A 5 year-old boy presented with bleeding per rectum. PR showed rectal polyp, biopsy showed the following. What is your diagnosis?




- a. Villous adenoma      b. Peutz-Jeghers polyp  
c. Juvenile polyp      d. Serrated adenoma

18. A 50-year-old male presented with hematuria. Investigations revealed normal glucose levels, proteinuria and creatinine of 9 mg%. Electron microscopic image is shown below. What other investigations could help in the diagnosis?




- a. ANA      b. HIV serology  
c. Electrophoresis      d. Anti GBM antibodies

19. A 30-year-old female presented with 4 cm mass in the right breast. Biopsy showed densely packed cells with bland nuclei and mucin infiltrating the stroma. What is your diagnosis?



- a. Invasive papillary carcinoma  
b. Medullary carcinoma  
c. Apocrine carcinoma  
d. Colloid carcinoma

20. 25-year-old female presented with swelling in front of neck. TSH levels were elevated. Biopsy showed lymphocytic infiltration and Hurthle cells. Which of the following is the possible diagnosis?



- a. Graves' disease  
b. Hashimoto's thyroiditis  
c. Medullary carcinoma thyroid  
d. Papillary carcinoma thyroid


21. Patient came with swelling in midline of neck measuring 2 cm in size. Histopathological examination showed Orphan Annie eye nuclei. What is the most likely diagnosis?

- a. Medullary carcinoma      b. Papillary carcinoma thyroid  
c. Toxic nodular goitre      d. Follicular thyroid carcinoma

22. Most common site of gastrinoma in MEN 1 is:

- a. Stomach      b. Jejunum  
c. Duodenum      d. Appendix

23. A 25-year-old male presented with swelling in the wrist joint. Histopathological examination showed spindle cells and Verocay bodies. What is the most likely diagnosis?



- a. Neurofibroma      b. Schwannoma  
c. Lipoma      d. Squamous cell carcinoma

24. Why is CPDA better than ACD for storage of blood?


- a. Improves oxygen transport  
b. More citrate ions  
c. It is less acidic  
d. Hypertonicity of blood

25. Which of the following translocation is seen in Myxoid liposarcoma?

- a. t(11:22)      b. t(14:18)  
c. t(x:18)      d. t(12:16)

Ans.

17. c  
18. d  
19. d  
20. b  
21. b  
22. c  
23. b  
24. a  
25. d



## AIIMS New Pattern 2019 Model Questions

### GENERAL PATHOLOGY

#### PATTERN 1: MULTIPLE TRUE /FALSE TYPE

1. In an experiment, a cell line derived from a human malignant neoplasm is grown in culture. A human IgG antibody is added to the culture, and the tumor cells become coated by the antibody, but they do not undergo lysis. Next, human cells are added that are negative for CD3, CD19, and surface immunoglobulin, but are positive for CD16 and CD56. The tumor cells are observed to undergo lysis. Which of the following statement denotes correctly about the cell types most likely to have killed the tumor cells?

1. B cells have surface immunoglobulin and can lyse the tumor cells
  2. CD8+ cell are positive for CD16 and can lyse the tumor cells
  3. Dendritic cells can lyse the tumor cells
  4. Macrophage express MHC II and can lyse the tumor cells
  5. Natural killer cells show ADCC and can lyse the tumor cells
- a. Option 1, 3, 4 are true
  - b. Option 2 and 5 are true
  - c. Option 5 is true, all others are false
  - d. All options are false

**Ans. (c) Option 5 is true, all others are false**

CD8 + CELL is not positive for CD16 ans c

2. Which of the following are true about deficiencies in complement pathway

1. Deficiency of decay-accelerating factor → paroxysmal nocturnal hemoglobinuria
  2. Deficiency of C6 and C7 → recurrent pyogenic bacterial infections
  3. Deficiency of C1 esterase inhibitor → hereditary angioedema
  4. Deficiency of C3 and C5 → recurrent pyogenic bacterial infections
  5. Deficiency of C6, C7, and C8 → recurrent infections with Neisseria species
- a. Option 1, 3, 4 are true
  - b. Option 2 and 5 are true
  - c. Option 5 is true, all others are false
  - d. All options are false

**Ans. (a) Option 1, 3, 4 are true**

#### PATTERN 2 : MATCH THE FOLLOWING

- 3.
- |                         |                           |
|-------------------------|---------------------------|
| A. Mast cells           | 1. Granulomatous response |
| B. Langhans giant cells | 2. Cytotoxic lymphocytes  |
| C. CD8 + cells          | 3. Humoral immunity       |

Contd...

D. Dendritic cells	4. Elaborate type I interferons
	5. Surface-bound IgE
	6. Express MHC I

- a. A-5, B-1, C-2, D-4
- b. A-5, B-1, C-2, D-6
- c. A-5, B-1, C-6, D-4
- d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

4.

A. c-abl	1. Small cell carcinoma of the lung
B. L-myc	2. Neuroblastoma
C. N-myc	3. Breast cancer
D. c-myc	4. Burkitt's lymphoma
	5. Chronic myelocytic leukemia (CML)
	6. Squamous cell carcinoma of the lung

- a. A-5, B-1, C-2, D-4
- b. A-5, B-1, C-2, D-6
- c. A-5, B-1, C-6, D-4
- d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

5.

A. CD8	1. Helper T cells,
B. CD4	2. Pan T cell marker
C. CD3	3. T cell receptor
D. CD2	4. Receptor for sheep erythrocyte (E rosette)
	5. Cytotoxic T cells
	6. Receptor for Fc portion of IgG

- a. A-5, B-1, C-2, D-4
- b. A-5, B-1, C-2, D-6
- c. A-5, B-1, C-6, D-4
- d. A-5, B-6, C-2, D-4


**Ans. (a) A-5, B-1, C-2, D-4**

6.

A. Centromere	1. Smith (SLE),
B. Speckled (non-DNA extractable nuclear proteins)	2. Double-stranded DNA (SLE)
C. Rim (peripheral)	3. T cell receptor
D. Nucleolar (RNA)	4. Progressive systemic sclerosis
	5. CREST syndrome
	6. Histones

- a. A-5, B-1, C-2, D-4
- b. A-5, B-1, C-2, D-6
- c. A-5, B-1, C-6, D-4
- d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**



7. Based on tumor and their respective IHC marker match the following


A. Carcinoma	1. CD 45
B. Lymphoma	2. NSE
C. Sarcoma	3. Vimentin
D. Melanoma	4. CK
E. Neuroendocrine Tumor	5. HMB45

- a. A-5, B-1, C-2, D-3, E-4      b. A-4, B-1, C-3, D-5, E-2  
c. A-3, B-2, C-4, D-1, E-5      d. A-1, B-2, C-3, D-4, E-5

Ans. (b) A-4, B-1, C-3, D-5, E-2

### PATTERN 3: SEQUENCE Q

8. A 63-year-old man has increasing malaise and back pain for the past 4 months. A radiograph of his spine shows rounded lucent lesions. The microscopic appearance of his liver with Congo red stain and with polarized light is shown in the figure. What is the correct sequence to diagnose this condition



- a. Biopsy -h & e -congo red -polarising light under congo red  
b. Biopsy - polarising light under congo red - mahogany brown on gross  
c. Biopsy-masson trichrome-electron microscopy  
d. Biopsy -congo red -h & e -polarising light under congo red

Ans. (a) Biopsy -H & E -congo red -polarising light under congo red

Now this is sequence biopsy fb h and e which shows pink colour deposits and then we do special stain like congo red fb seeing it under polarizing mic so ans is A.

9. A 48-year-old woman has fingers that are tapered and claw-like, with decreased motion at the small joints. There are no wrinkle lines on her facial skin. The microscopic appearance of the skin is shown in the figure. The patient also has diffuse interstitial fibrosis of the lungs, with pulmonary hypertension and cor pulmonale. Which of the following dermal inflammatory cells is the most likely initiator of the process that is the cause of her skin disease?

10. An epidemiologic study is conducted to determine risk factors for HIV infection. The study documents that individuals with coexisting sexually transmitted diseases such as chancroid are more likely to become HIV-positive. It is postulated that an inflamed mucosal surface is an ideal location for the transmission of HIV during sexual intercourse. Which of the following cells in these mucosal surfaces is most instrumental in transmitting HIV to CD4+ T lymphocytes?

- a. CD8+ cells      b. Dendritic cells  
c. Natural killer cells      d. Neutrophils  
e. Plasma cells      f. CD4+ lymphocyte  
g. Mast cell

Ans. (b) Dendritic cells; (f) CD4+ lymphocyte

11. Which of the following are the characteristics of Exudative pleural effusion?

1. Pleural fluid protein/serum protein > 0.5  
2. Pleural fluid protein/serum protein < 0.5  
3. Pleural fluid LDH/serum LDH > 0.6  
4. Pleural fluid LDH/serum LDH < 0.6

Select the correct answer using the code given below:

- a. 1 and 3      b. 1 and 4  
c. 2 and 3      d. 2 and 4

Ans. (a) 1 and 3

Light's criteria (Sensitivity 99%, Specificity 98%)


Criteria	Transudate	Exudate
Pleural fluid protein: Serum: protein ratio	<0.5	>0.5
Pleural fluid LDH: Serum LDH	<0.6	>0.6
Pleural fluid LDH	<200	>200

### PATTERN 4: MULTIPLE CORRECT OPTIONS

12. A 37-year-old man who is HIV-positive has noticed an increasing number and size of skin lesions on his face, trunk, and extremities, as shown in the figure, over the past 18 months. Some of the larger lesions appear to be nodular. Which among the following is/ are correct options?

- A. HIV is not detected in the spindle cells that proliferate in Kaposi sarcoma.  
B. Cytomegalovirus is implicated cause of above lesion  
C. Epstein-Barr virus is usually not associated with skin lesions  
D. Kaposi sarcoma is associated with Kaposi sarcoma herpes virus.  
E. Human herpesvirus-8 is the implicated cause  
a. Option A, C, D, E are true  
b. Option A, C, D are true, B and E are false  
c. Option E is true, all others are false  
d. All options are false





**Ans. (a) Option A,C,D,E are true**

This is Kaposi sarcoma. Kaposi sarcoma is associated with Kaposi sarcoma herpes virus (KSHV). KSHV is also called HHV8. Human herpesvirus-8 is the implicated cause of KS – true. HIV is not detected in the spindle cells that proliferate in Kaposi sarcoma. Remember Epstein-Barr virus is usually not associated with skin lesions. So ans is A.

**13. A 23-year-old girl is infected with HPV. Which among below are correct lesions that can be associated with above lesion**

- A. Verruca vulgaris
- B. Condyloma
- C. Cervical neoplasia
- D. Carcinoma of the nasopharynx
- E. African Burkitt's lymphoma
- a. Option A,B,C are true
- b. Option A,C,D are true, B and E are false
- c. Option E is true, all others are false
- d. All options are false

**Ans. (a) Option a,b,c are true**

#### PATTERN 5: BEST ONE ANSWER

**14. Lymphocyte phenotype test is done for?**

- a. Agammaglobulinemia
- b. SCID
- c. Sepsis
- d. Acute leukemia

**Ans. (c) Sepsis**

Clinical signs and symptoms of sepsis are nonspecific and often indistinguishable from those of nonseptic critical illness.

This ambiguity frequently delays the diagnosis of sepsis until culture results can confirm the presence or absence of an infectious organism. Lymphocyte phenotyping can be conducted rapidly and may provide information on the presence of infection before culture results are available. Hence answer is sepsis c.

**15. True about repeats in Fragile X syndrome?**

- a. CGG repeats
- b. CAG repeats
- c. CTG repeats
- d. GCT repeats

**Ans. (c) CTG repeats**

Carcinoid heart disease: PV stenosis, TV regurgitation

#### PATTERN 6: ASSERTION REASONING

**16. Assertion: majority of people with HIV have a symptomatic disease.**

**Reason: HIV causes progressive destruction of CD4 cells.**

- a. If both **Assertion** and **Reason** are true and the **Reason** is the correct explanation of the **Assertion**.
- b. If both **Assertion** and **Reason** are true but the **Reason** is not the correct explanation of the **Assertion**.
- c. If **Assertion** is true but **Reason** is false.
- d. If **Assertion** is false and **Reason** is true.

**Ans. (d) If Assertion is false and Reason is true.**

**17. Assertion: Natural Killer Cells are a part of innate immunity.**

**Reason: Natural Killer Cells do not need previous sensitization**

- a. If both **Assertion** and **Reason** are true and the **Reason** is the correct explanation of the **Assertion**.
- b. If both **Assertion** and **Reason** are true but the **Reason** is not the correct explanation of the **Assertion**.
- c. If **Assertion** is true but **Reason** is false.
- d. If **Assertion** is false and **Reason** is true.

**Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.**

#### PATTERN 6: SCENARIO TYPE Q

**18. A 45-year-old female patient presented with vaginal discharge. Her pap smear revealed HSV infection. What inclusions do u suspect in this case**

- a. Cowdry A bodies
- b. koilocytosis
- c. Guarnieri bodies
- d. Warthin-Finkeldey giant cells
- e. Ground-Glass Change
- f. Atypical lymphocytes

**Ans. (a) Cowdry A bodies**

**19. A 4-year-old child presented with measles. He showed many giant cells in his lymph node biopsy. Which of the following is the correct about inclusions seen in this entity**

- a. Cowdry A bodies
- b. koilocytosis
- c. Guarnieri bodies
- d. Warthin-Finkeldey giant cells
- e. Ground-Glass Change
- f. Atypical lymphocytes

**Ans. (d) Warthin-Finkeldey giant cells**

**20. Deficiency of which immune function results in tubercular infections?**

- 1. Defective phagocytic function
- 2. Defect in T cell function
- 3. Defect in B cell function
- 4. Defect in antibody productivity
- a. Only 1
- b. Only 1 and 3
- c. Only 2
- d. None of above


**Ans. (c) Only 2**

T-lymphocytes play a major role in conferring immunity against M. tuberculosis.

**21. Syndrome X includes:**

- a. Hyperlipidemia, Obesity, Type 2 DM
- b. Obesity, CAD, COPD
- c. Hyponatremia, Hyperlipidemia with type 2 DM
- d. Hyponatremia, Hyperlipidemia with type 2 DM





**Ans. (a) Hyperlipidemia, Obesity, Type 2 DM**

Metabolic syndrome, sometimes known as Syndrome X, is a clustering of at least three of the five following medical conditions: central obesity, high blood pressure, high blood sugar (type 2 diabetes mellitus), high serum triglycerides, and low serum high density lipoprotein (HDL).

**22. Cutaneous manifestations of tuberculosis include:**

- Erythema nodosum and lupus vulgaris
- Erythema marginatum and lupus vulgaris
- Phlyctenular conjunctivitis and erythema multiform
- Pyoderma gangrenosum and Dactylitis

**Ans. (a) Erythema nodosum and lupus vulgaris**

**Lesion associated with TB**

- Erythema nodosum (EN), also known as subacute migratory panniculitis of Villanova an inflammatory condition characterized by inflammation of the fat cells under the skin resulting in tender red nodules or lumps that are usually seen on both shins.
- Lupus vulgaris (also known as tuberculosis luposa) are painful cutaneous tuberculosis skin lesion with nodular appearance, most often on the face around the nose, eyelids, lips, cheeks, ear and neck. It is the most common Mycobacterium tuberculosis skin infection.

**23. Killer T cells which are responsible for defence against intracellular pathogen are expressed by which of the following CD phenotypes?**

- CD 2
- CD 8
- CD 5
- CD 10
- CD 4
- CD 16
- 2, 5, 6
- 3, 4, 5
- 1, 5, 6
- None

**Ans. (a) 2, 5, 6**

Natural killer T-cells/NKT cells include both NK 1.1\* and NK 1.1 as well as CD4, CD4, CD8 cells, CD 16, CD 16

**HEMATOLOGY**

**PATTERN 1: MULTIPLE TRUE FALSE TYPE**

**24. Identify the correct statements regarding Polycythemia**

- Most cases of Polycythemia vera are due to mutation of the JAK2 gene on the short arm of chromosome 9.
- Increase in bone marrow production of RBCs is seen in relative polycythemia
- Ectopic release of EPO may occur in renal cell carcinoma
- Hypoxic stimulus for EPO release can cause increase in bone marrow production of RBCs
- Plasma volume is increased in relative polycythemia
- Option 1, 3, 4 are true
- Option 2 and 5 are true
- Option 5 is true, all others are false
- All options are false

**Ans. (a) Option 1, 3, 4 are true**

Option 2 and 5 is false. No increase in bone marrow production of RBCs in relative polycythemia. plasma volume is decreased.

**25. Safe transfusion practices regarding Patient cross match for a blood transfusion**

- Before blood is transfused into newborns or patients with T-cell deficiencies, it must be irradiated to kill donor lymphocytes
- Blood group AB patients have natural antibodies.
- Major cross match is Patient serum is mixed with a sample of RBCs from a donor unit.
- Blood group O individuals are considered universal donors.
- Direct Coombs test to identify atypical antigens on the persons RBCs
- Option 1, 3, 4 are true
- Option 2 and 5 are true
- Option 5 is true, all others are false
- All options are false

**Ans. (a) Option 1, 3, 4 are true**

Option 2 and 5 is false-Direct Coombs test to identify atypical IgG antibodies on the persons RBCs and Blood group AB patients lack natural antibodies.

**PATTERN 2 : MATCH THE FOLLOWING**

**26.**


A. Spleen in portal hypertension (PH)	1. Macrophages with a soap bubble appearance
B. Niemann-Pick disease	2. Macrophages with a fibrillary appearance
C. Gaucher disease	3. Humoral immunity
D. Red pulp of spleen	4. Fixed macrophages and sinusoids
	5. Thickened ("sugar-coated") capsule
	6. B and T cells

- A-5, B-1, C-2, D-4
- A-5, B-1, C-2, D-6
- A-5, B-1, C-6, D-4
- A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

**Spleen in portal hypertension (PH)**

- Gross findings
  - Spleen is covered by a thickened ("sugar-coated") capsule from perisplenitis
- Microscopic findings
  - Calcium and iron concretions called Gamna-Gandy bodies are deposited in collagen



### PATTERN 3: SEQUENCE Q

27. Identify the correct sequence in primary hemostasis. What is the correct sequence to diagnose this condition
- Vessel injury-Platelet adhesion to Vwf-Platelet release of aggregating agents-platelet gp2b3a binding to fibrinogen
  - Vessel injury-platelet gp2b3a binding to fibrinogen -Platelet adhesion to Vwf-Platelet release of aggregating agents
  - Vessel injury-Platelet adhesion to Vwf-platelet gp2b3a binding to fibrinogen -Platelet release of aggregating agents
  - Platelet adhesion to Vwf-Platelet release of aggregating agents-platelet gp2b3a binding to fibrinogen -injury

Ans. (a) Vessel injury-Platelet adhesion to Vwf-Platelet release of aggregating agents-platelet gp2b3a binding to fibrinogen

### PATTERN 4: MULTIPLE CORRECT OPTIONS

28. A 37-year-old man presents with Thrombocytopenia. What of the following can be causes of decreased platelet count
- Thrombotic thrombocytopenic purpura
  - Hypersplenism
  - Idiopathic thrombocytopenic purpura
  - Thrombotic thrombocytopenic purpura.
  - Decreased production of platelets
- Option A,B,C,D,E are true
  - Option A,C,D are true, Band E are false
  - Option E is true, all others are false
  - All options are false

Ans. (a) Option A,B,C,D,E are true

### PATTERN 5: BEST ONE ANSWER

29. Lymphocytic and histiocytic (L&H) cells are seen in?
- Nodular sclerosis
  - Mixed cellularity classical HL
  - Nodular lymphocyte predominant HL
  - Lymphocyte depleted classical HL

Ans. (c) Nodular lymphocyte predominant HL

### PATTERN 6: ASSERTION REASONING

30. Assertion: antiphospholipid antibodies produces strokes  
Reason: antiphospholipid antibodies are associated with arterial thrombosis specially cerebral vessels
- If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.
  - If both Assertion and Reason are true but the Reason is not the correct explanation of the Assertion.
  - If Assertion is true but Reason is false.
  - If Assertion is false and Reason is true.

Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.

Cerebral vessels thrombosis (most common site of APLA; produces strokes)

31. Assertion: Allergic transfusion reactions are most common transfusion reaction  
Reason: It is due to type I IgE-mediated hypersensitivity reaction (HSR) against proteins (allergens) that are present in the donor blood
- If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.
  - If both Assertion and Reason are true but the Reason is not the correct explanation of the Assertion.
  - If Assertion is true but Reason is false.
  - If Assertion is false and Reason is true .


Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.

### A –understand Allergic transfusion reactions

- Since the patient has been previously sensitized to an allergen that is present in donor blood, IgE antibodies are already present on the patient's mast cells
- Exposure to the allergen from the donor blood leads to cross-linking of IgE antibodies specific for the allergen on the mast cell membranes.
- IgE triggering causes an early phase reaction that is characterized by mast cell release of preformed mediators.
  - Preformed chemicals include histamine, eosinophil chemotactic factor, and serotonin.

### PATTERN 7: SCENERIO TYPE Q

32. A febrile child presented with Lytic lesions are present in the skull, Central diabetes insipidus (CDI) and Exophthalmos




#### Diagnosis

The electron microscopy shown above is typically seen in this child. What is the electron microscopy showing

- Letterer-Siwe disease
- Hand-Schüller-Christian (HSC) disease
- Urticaria pigmentosa
- Lymphocyte depleted classical HL
- Eosinophilic granuloma
- Birbeck granules

Ans. (b) Hand-Schüller-Christian (HSC)



Hand-Schüller-Christian (HSC) disease shows Classic triad due to infiltrative disease

- Lytic lesions are present in the skull.
- Central diabetes insipidus (CDI), due to invasion of the posterior pituitary stalk
- Exophthalmos from infiltration of the orbit 2<sup>nd</sup>– f-Electron micrograph showing racket-shaped Birbeck granules in a histiocyte

**33. Which of the following conditions are associated with prolonged prothrombin time?**

1. Factor VIII deficiency
2. Factor VII deficiency
3. Heparin anticoagulation
4. Warfarin anticoagulation

Select the correct answer using the code given below:

- a. 1 and 4
- b. 1 and 3
- c. 2 and 3
- d. 2 and 4

**Ans. (d) 2 and 4**

The prothrombin time can be prolonged as a result of deficiencies in vitamin K, warfarin therapy, malabsorption, or lack of intestinal colonization by bacteria (such as in newborn). In addition, poor factor VII synthesis (due to liver disease) or increased consumption (in disseminated intravascular coagulation) may prolong the PT.

**34. Match the following vials in column (A) with the color coding in column (B)?**

Column A-Vials	Column B-color coding
A. EDTA vial	a. Light blue
B. Plain vial	b. Purple
C. Coagulation profile vial	c. Yellow
D Heparin vial	d. Red
	e. Gray
	f. Green

**Ans. A-b, B-d, C-a, D-f**

## SYSTEMIC PATHOLOGY

### PATTERN 1 : MULTIPLE TRUE FALSE TYPE

**35. A patient presents with biventricular failure and narrow pulse pressure. His CXR was suggestive of Dilated cardiomyopathy. Which of the following statements tell correctly about etiology**

1. Idiopathic (most common)
2. Genetic (most common)
3. Postpartum state can be a causative
4. Alcohol can cause direct toxicity to cause DCM
5. Most common cause of sudden death in young athletes

- a. Option 1, 3, 4 are true
- b. Option 2 and 5 are true
- c. Option 5 is true, all others are false
- d. All options are false

**Ans. (a) Option 1, 3, 4 are true**

A-Hypertrophic cardiomyopathy (HCM) is most common cause of sudden death in young athletes and is mostly genetic

**36. A 30-year-old man has sudden onset of hematemesis after a weekend in which he consumed large amounts of alcohol. The bleeding stops, but he has another episode under similar circumstances 1 month later. Upper gastroesophageal endoscopy shows longitudinal tears at the gastroesophageal junction. What is the most likely mechanism to cause his hematemesis?**

1. Most cases occur in the context of alcohol abuse
  2. This is a case of cirrhosis from alcohol abuse.
  3. Mallory-Weiss syndrome with esophageal tears results from severe vomiting.
  4. The bleeding due to this condition is usually not very life-threatening
  5. This is case of Herpes simplex virus infection
- a. Option 1, 3, 4 are true
  - b. Option 2 and 5 are true
  - c. Option 5 is true, all others are false
  - d. All options are false

**Ans. (a) Option 1, 3, 4 are true**


Portal hypertension leads to dilation of esophageal submucosal veins, which can bleed profusely; in this case, the patient's age argues against the presence of cirrhosis from alcohol abuse

**37. Bronchiolitis obliterans with organizing pneumonia (BOOP) is characterized histologically in the lung by**

1. Loose fibrous tissue within bronchioles and alveoli
  2. Asteroid bodies in giant cells within bronchioles
  3. It occurs due to smoking
  4. Steroid responsive
  5. Multiple rheumatoid nodules within the interstitial tissue
- a. Option 1,3,4 are true
  - b. Option 2 and 5 are true
  - c. Option 5 is true, all others are false
  - d. All options are false

**Ans. (a) Option 1, 3, 4 are true**

The lungs respond to these agents, causing bronchiolar injury by forming loose, fibrous tissue within the bronchioles (bronchiolitis obliterans) and alveoli (organizing pneumonia).



### PATTERN 2 : MATCH THE FOLLOWING

- 38.**
- |                                   |                                      |
|-----------------------------------|--------------------------------------|
| A. MC site for Cardiac metastasis | 1. Rhabdomyoma                       |
| B. Tuberous sclerosis             | 2. Cardiac myxoma                    |
| C. Ball-valve effect              | 3. Hypertrophic cardiomyopathy (HCM) |
| D. Endocardial fibroelastosis     | 4. Restrictive cardiomyopathy        |
|                                   | 5. Pericardium                       |
|                                   | 6. $\beta$ -Blockers                 |
- a. A-5, B-1, C-2, D-4      b. A-5, B-1, C-2, D-6  
c. A-5, B-1, C-6, D-4      d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

- 39.**
- |                           |          |
|---------------------------|----------|
| A. Turcot syndrome        | 1. SMAD4 |
| B. Juvenile polyposis     | 2. STK11 |
| C. Peutz-Jeghers syndrome | 3. TSC   |
| D. Cowden syndrome        | 4. PTEN  |
|                           | 5. APC   |
|                           | 6. MYH   |
- a. A-5, B-1, C-2, D-4      b. A-5, B-1, C-2, D-6  
c. A-5, B-1, C-6, D-4      d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

- 40.**
- |                      |   |
|----------------------|---|
| A. MEN 2B            | 1. Wermer's Syndrome                                      |
| B. Type 1 MEN        | 2. Medullary carcinoma of thyroid                         |
| C. Sipple's Syndrome | 3. Membranous glomerulonephropathy (MGN)                  |
| D. MEN4              | 4. Heterozygous inactivating mutations in the CDKN1B gene |
|                      | 5. Mucosal neuromas                                       |
|                      | 6. Minimal change disease (lipoid nephrosis)              |
- a. A-5, B-1, C-2, D-4      b. A-5, B-1, C-2, D-6  
c. A-5, B-1, C-6, D-4      d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

A MEN4 is caused by heterozygous inactivating mutations in the CDKN1B gene (12p13.1-p12) encoding p27, a cyclin-dependent kinase inhibitor that acts as a negative regulator of cell cycle progression

#### Type 1 (Wermer's Syndrome)

- Parathyroid
- Pituitary
- Pancreas

#### Type 2 (Sipple's Syndrome)

- Parathyroid
- Medullary carcinoma of thyroid
- Pheochromocytoma

#### Type 3 (MEN 2B)

- Medullary carcinoma of thyroid
- Pheochromocytoma
- Mucosal neuromas

- 41.**
- |                           |                                     |
|---------------------------|-------------------------------------|
| A. Fats                   | 1. PAS-positive, diastase-sensitive |
| B. Glycogen               | 2. Prussian blue                    |
| C. Hemosiderin            | 3. Von Kossa                        |
| D. $\alpha$ 1 antitrypsin | 4. PAS-positive, diastase-resistant |
|                           | 5. Oil red O                        |
|                           | 6. Congo red                        |
- a. A-5, B-1, C-2, D-4      b. A-5, B-1, C-2, D-6  
c. A-5, B-1, C-6, D-4      d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**

- 42.**
- |  |   |
|--|---|
| A. Minimal change disease (lipoid nephrosis) | 1. Thickening of basement membrane ("spikes and domes") |
| B. Membranous glomerulonephropathy (MGN)     | 2. Mesangial deposits                                   |
| C. IgA nephropathy                           | 3. Subendothelial deposits                              |
| D. FSGS                                      | 4. Apol1 mutation                                       |
|  | 5. EM reveals fusion of foot processes of podocytes     |
|  | 6. K w lesions  |
- a. A-5, B-1, C-2, D-4      b. A-5, B-1, C-2, D-6  
c. A-5, B-1, C-6, D-4      d. A-5, B-6, C-2, D-4

**Ans. (a) A-5, B-1, C-2, D-4**


The risk variants G1 (S342G:I384M) and G2 (del.N388/Y389) are two coding variants in the APOL1 gene on chromosome 22q13. The mutant alleles confer protection against trypanosomal infections. Its mutations are associated with FSGS

### PATTERN 3: SEQUENCE Q

- 43. A 63-year-old man died de to mi. Correct sequence of events**
- Coagulation      necrosis-Neutrophils-Macrophages-Granulation tissue and collagen formation
  - Granulation TISSUE -Coagulation necrosis-Neutrophils-Macrophages-collagen formation
  - Neutrophils-Coagulation      necrosis--Macrophages-Granulation tissue and collagen formation
  - Coagulationnecrosis-Neutrophils-Macrophages-collagen formation-Granulation TISSUE

**Ans. (a) Coagulation necrosis- Neutrophils- Macrophages- Granulation tissue and collagen formation**





44. A child presented with VSD, PV stenosis, RVH and cyanosis after 1 year of age. Diagnosis
- Tetralogy of Fallot
  - Coarctation of the aorta
  - TV atresia
  - Truncus arteriosus
  - Complete transposition of the great arteries
  - VSD

**Ans. (a) Tetralogy of Fallot**

Tetralogy of Fallot presents with A-VSD, Infundibular (most common) or PV stenosis, RVH, Dextrorotated aorta with a right-sided aortic arch

45. A 48-year-old woman has leg claudication and hypertension. On examination, increase in the upper extremity blood pressure is observed. Disparity between upper/lower extremity blood pressure >10 mm Hg is seen. Which of the following is the correct diagnosis of this entity?
- Tetralogy of Fallot
  - Coarctation of the aorta
  - TV atresia
  - Truncus arteriosus
  - Complete transposition of the great arteries
  - VSD

**Ans. (b) Coarctation of the aorta**

46. Identify the correct the classic adenoma-carcinoma sequence
- APC -b-catenin -K-RAS -TP53-LOH at 18q21-Telomerase activation
  - APC -TP53 -K-RAS -LOH at 18q21-Telomerase activation
  - APC -K-RAS -TP53-LOH at 18q21-Telomerase activation
  - APC -b-catenin -K-RAS -Telomerase activation -LOH at 18q21

**Ans. (a) APC -b-catenin -K-RAS -TP53-LOH at 18q21-Telomerase activation**


#### PATTERN 4: MULTIPLE CORRECT OPTIONS

47. Many complications are associated with ST ELEVATION MYOCARDIAL INFARCTION (STEMI). WHICH among below are correct options about the above lesion
- Congestive heart failure: Usually occurs within the first 24 hours
  - Mural thrombus: has high danger of embolization
  - Ventricular fibrillation: MCC death in STEMI
  - Fibrinous pericarditis occurs between day 1 to 7 of a STEMI.
  - Myocardial rupture: MC at 3-7 days
- Option A,B,C,D,E are true
  - Option A,C,D are true, B and e are false
  - Option E is true, all others are false
  - All options are false

**Ans. (a) Option A,B,C,D,E are true**

48. A 65-year-old woman is being treated in the hospital for pneumonia complicated by septicemia. She has required multiple antibiotics and was intubated and mechanically ventilated earlier in the course. On day 20 of hospitalization, she has abdominal distention. At laparotomy, a portion of distal ileum and cecum is resected. The gross appearance of the mucosal surface is shown in the figure.



- The opened colon shows pseudomembranes that are patches of fibrinopurulent debris attached to the mucosa.
  - It's a complication of broadspectrum antibiotic therapy
  - It results from overgrowth of Clostridium difficile or other organisms that are capable of inflicting mucosal injury
  - Clostridium septicum is most often associated with malignancy or immunosuppression
  - Its image of toxic megacolon is an uncommon complication of ulcerative colitis.
- Option A,B,C,D are true
  - Option A,C,D are true, B and e are false
  - Option e is true, all others are false
  - All options are false



**Ans. (a) Option a,b,c,d are true**

The opened colon shows pseudomembranes that are patches of fibrinopurulent debris attached to the mucosa. Pseudomembranous enterocolitis is a complication of broad-spectrum antibiotic therapy, which alters gut flora to allow overgrowth of Clostridium difficile or other organisms that are capable of inflicting mucosal injury. Clostridium septicum infection can lead to myonecrosis that is most often associated with malignancy or immunosuppression. This gross pattern also can appear from ischemic injury that is vascular or mechanical, but this patient's history and the time course support an iatrogenic cause. An ischemic colitis resulting from mesenteric artery thrombosis could appear similar, but it is not associated with C. difficile. A dilated, thinned, toxic megacolon is an uncommon complication of ulcerative colitis

49. A 57-year-old male presents with a lesion similar to that seen in this gross photograph of a sagittal section of the lung. Which one of the listed characteristics, if present in this lesion, would favor the diagnosis of mesothelioma?
- Arises from the pleural surfaces
  - Negative staining with CEA and Leu-M1.
  - Long microvilli seen by electron microscopy
  - Lamellar bodies seen by electron microscopy
  - Its image of adenocarcinoma



- a. Option a,b,c are true
- b. Option a,c,d are true, b and e are false
- c. Option e is true, all others are false
- d. All options are false

**Ans. (a) Option a,b,c are true**


#### PATTERN 5: BEST ONE ANSWER

50. Sterile, nondestructive vegetations present on the MITRAL VALVE?
- a. Libman-Sacks endocarditis
  - b. Infective endocarditis (IE)
  - c. Nonbacterial thrombotic endocarditis
  - d. Carcinoid heart disease

**Ans. (c) Nonbacterial thrombotic endocarditis**

Carcinoid heart disease: PV stenosis, TV regurgitation

51. A 53-year-old woman undergoes a routine checkup. The only abnormal finding is a stool specimen that contains Occult blood. Colonoscopy shows a 1.5-cm, solitary, rounded, erythematous polyp on a 0.5-cm stalk at the splenic flexure. Her colonic lesion is most likely associated with which of the following?
- a. Low risk for development of carcinoma
  - b. Inheritance of an abnormal tumor suppressor gene
  - c. Presence of similar lesions in the small intestine
  - d. Risk for development of endometrial carcinoma



**Ans. (a) Low risk for development of carcinoma**

The figure shows a solitary pedunculated adenoma of the colon with no evidence of malignancy

52. Histologic sections from a 3-cm mass found in the mandible of a 55-year-old female reveal a tumor consisting of nests of tumor cells that appear dark and crowded at the periphery of the nests and loose in the center (similar to the stellate reticulum of a developing tooth). Grossly, the lesions consist of multiple cysts filled with a thick, "motor oil"-like fluid. What is the correct diagnosis for this tumor?
- a. Pleomorphic adenoma
  - b. Ameloblastoma
  - c. Mucoepidermoid carcinoma
  - d. Adenoid cystic carcinoma
  - e. Acinic cell carcinoma

**Ans. (b) Ameloblastoma**


Rare tumor of the oral cavity (found most commonly in the mandible) that is similar to the enamel organ of the tooth is the ameloblastoma. This locally aggressive tumor consists of nests of cells that at their periphery are similar to ameloblasts and centrally are similar to the stellate reticulum of the developing tooth

#### PATTERN 6: ASSERTION REASONING


53. Assertion: Pharynx is the only site for infection leading to Rheumatic fever (RF).  
Reason: Nephrogenic strains of group A streptococcus that produce poststreptococcal glomerulonephritis, lack the types of matrix (M) proteins (virulence factors) in their cell wall that are present in pharyngeal strains; hence they never produce RF.
- a. If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.
  - b. If both Assertion and Reason are true but the Reason is not the correct explanation of the Assertion.
  - c. If Assertion is true but Reason is false.
  - d. If Assertion is false and Reason is true.

**Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.**

54. Assertion: patient presents with secretory diarrhea syndrome.  
Reason: The figure shows a large villous adenoma which can cause these manifestations



- a. If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.
- b. If both Assertion and Reason are true but the Reason is not the correct explanation of the Assertion.
- c. If Assertion is true but Reason is false.
- d. If Assertion is false and Reason is true.



**Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.**

**55. Assertion: squamous cell carcinomas of the urinary bladder are quite rare except in Egypt and other areas of the Middle East.**

**Reason: these areas predominantly have schistosomiasis**

- If both **Assertion** and **Reason** are true and the Reason is the correct explanation of the Assertion.
- If both **Assertion** and **Reason** are true but the Reason is not the correct explanation of the **Assertion**.
- If **Assertion** is true but **Reason** is false.
- If **Assertion** is false and **Reason** is true.

**Ans. (a) If both Assertion and Reason are true and the Reason is the correct explanation of the Assertion.**

#### PATTERN 7: EMQ

**56. A 19-year-old man is advised to see his physician because genetic screening has detected a disease in other family members. On physical examination, a stool sample is positive for occult blood. A colonoscopy is performed, followed by a colectomy. The figure shows the gross appearance of the mucosal surface of the colectomy specimen**

- Molecular analysis of this patient's normal fibroblasts is most likely to show a mutation in which of the following genes?
- Patients of chron's disease are associated with which mutation?



- |         |         |
|---------|---------|
| a. APC  | b. MLH1 |
| c. KRAS | d. NOD2 |
| e. p53  | f. MSH2 |

**Ans. 1-a, 2 - d**

This young patient's colon shows hundreds of polyps. This is most likely a case of familial adenomatous polyposis (FAP) syndrome, which results from inheritance of one mutant copy of the APC tumor-suppressor gene (a few FAP cases are associated with DNA mismatch repair genes). Every somatic cell of this patient most likely has

one defective copy of the APC gene. Polyps are formed when the second copy of the APC gene is lost in many colon epithelial cells. Without treatment, colon cancers arise in 100% of these patients because of accumulation of additional mutations in one or more polyps, typically before 30 years of age. Patients with a gene for hereditary nonpolyposis colorectal carcinoma, such as MLH1 and MSH2, also have an inherited susceptibility to develop colon cancer, but in contrast to patients with FAP, they do not develop numerous polyps. Sporadic colon cancers may have CpG island hypermethylation along with KRAS mutations, whereas others have p53 mutations, but the somatic cells of patients with these cancers do not show abnormalities of these genes. NOD2 mutations are linked with Crohn's disease.

**57. A 66-year-old male presents to his family physician, complaining of bloating, which has developed over the course of several weeks. After evaluation, an exploratory laparotomy is performed. When the peritoneum is entered, the surgeon finds a large amount of gelatinous material. Of the following, what is the most likely anatomic site for the condition causing this change?**

- |                    |             |
|--------------------|-------------|
| a. Lung            | b. Liver    |
| c. Stomach         | d. Appendix |
| e. Large intestine | f. Prostate |

**Ans. (d) Appendix**

The description is that of pseudomyxoma peritonei. The gross features are associated with both benign and malignant neoplasms, and the classification and naming of the lesion is controversial; however, in a vast majority of cases, the neoplasm responsible for the changes in the peritoneal cavity is found in the appendix (D). Although a variety of neoplasms could give rise to the finding of pseudomyxoma peritonei, neoplasms in the appendix are by far the most common source, which indicates that the other answers (A-C, E-F) are not the best choice.

**58. Consider the following statements regarding changes in pregnancy:**

- Plasma volume increases up to 30-50%
- Pregnancy is a hypercoagulable state
- Hematocrit is decreased
- Total plasma proteins increases

**Which of the statements given above is/are correct?**

- |                 |                  |
|-----------------|------------------|
| a. 1 only       | b. 1, 2, 3 and 4 |
| c. 1 and 2 only | d. 3 and 4 only  |

**Ans. (b) 1, 2, 3 and 4**

#### Hematological changes in pregnancy

- Plasma volume: Increase by 30-40%
- Fibrinogen: Increase by 50%
- Hematocrit: Decrease by 6%
- Total plasma proteins: Increase by 30%





59. A 50-year old post menopausal woman comes with complaints of bleeding per vaginum. Which one of the following investigations is NOT required

- a. Endometrial biopsy      b. Diagnostic laparoscopy
- c. Hysteroscopy              d. Pap smear

**Ans. (b) Diagnostic laparoscopy**

#### Evaluation of postmenopausal bleeding

- Abnormal endometrium may have to be investigated by a hysteroscopy with a biopsy or a dilation and curettage.
- Pap smear is done to rule out carcinoma cervix

60. Which of the following statements regarding  $\beta$  Human chorionic Gonadotropin are NOT correct

- 1. It is a glycoprotein hormone.
- 2. Serum levels increase in pregnancy, germ cells tumor and gestational trophoblastic disease
- 3. Its levels are same in single and multiple pregnancy
- 4. It has common and Alpha subunit with other hormones FSH, LH and TSH.

Select the correct answer using the code given below:

- a. 1, 2 and 4                      b. 1, 2 and 3
- c. 2, 3 and 4                      d. 1, 3 and 4

**Ans. (a) 1, 2 and 4**

High levels of hCG could be detected in multiple pregnancy (levels are not same in single pregnancy and multiple pregnancy).

61. Which of the following is/are the common infectious syndrome (s) associated with Klebsiella pneumonia?

- 1. Pneumonia
- 2. Intra-abdominal infections
- 3. Hepatitis

Select the correct answer using the code given below:

- a. 1 only                          b. 1 and 2 only
- c. 1, 2 and 3                      d. 2 only

**Ans. (c) 1, 2 and 3**

The range of clinical disease caused by Klebsiella pneumonia includes pneumonia thrombophlebitis, urinary tract infection, cholecystitis, diarrhea, upper respiratory tract infection, wound infection, osteomyelitis, meningitis, and bacteremia and septicemia.

62. Consider the following statements with regard to acute anterior poliomyelitis:

- 1. It is caused by a virus belonging to picornavirus family
- 2. Muscle pain and cramps may be associated with diffuse transient fasciculations
- 3. Tonsillectomy reduces the risk of bulbar poliomyelitis
- 4. Cerebrospinal fluid may show mild pleocytosis with increase polymorphonuclear cells in early course of disease

Which of the above statements are correct?

- a. 1, 2 and 3                      b. 2, 3 and 4
- c. 1, 2 and 4                      d. 1, 3 and 4

**Ans. (c) 1, 2 and 4**

#### Polio

- Poliovirus, the causative agent of poliomyelitis (commonly known as polio), is a human enterovirus and member of the family of picomaviridae.
- Virus invasion cause inflammation of the nerve cells of anterior horn cells of spinal cord, leading to damage or destruction of motor neuron ganglia. With the destruction of nerve cells, the muscles no longer receive signals from the brain or spinal cord; without nerve stimulation, the muscles atrophy, becoming weak, floppy and poorly controlled, and finally completely paralyzed. Maximum paralysis progresses rapidly (two or four days), and usually involves fever and muscle pain.
- Tonsillectomy increases the risk of bulbar poliomyelitis.
- Analysis of the patient's cerebrospinal fluid (CSF), which is collected by a lumbar puncture ("spinal tap"), reveals an increased number of white blood cells and a mildly elevated protein level.

63. Which of the following are the Indications for lung transplantation?

- 1. Emphysema
- 2. Primary pulmonary hypertension
- 3. Obliterative bronchiolitis

Select the correct answer using the code given below:

- a. 1 and 2 only                      b. 2 and 3 only
- c. 1 and 3 only                      d. 1, 2 and 3

**Ans. (a) 1 and 2 only**

As of 2005, the most common reasons for lung transplantation in the United States were:

- 27% chronic obstructive pulmonary disease (COPD), including emphysema;
- 16% idiopathic pulmonary fibrosis;
- 14% cystic fibrosis;
- 12% idiopathic (formerly known as "primary") pulmonary hypertension;
- 5% alpha 1 antitrypsin deficiency;
- 2% replacing previously transplanted lungs that have since failed;
- 24% other causes, including bronchiectasis and sarcoidosis.

64. Which of the following are common risk factors for contrast induced Nephrotoxicity?

- 1. Use of high osmolality, ionic contrast media
- 2. Diabetes Mellitus
- 3. Myeloma


Select the correct answer using the code given below:

- a. 1 and 2 only                      b. 2 and 3 only
- c. 1 and 3 only                      d. 1, 2 and 3

**Ans. (d) 1, 2 and 3**

There are multiple risk factors of contrast induced nephropathy, whereof a review in 2016 emphasized chronic kidney disease, diabetes mellitus, high blood pressure, reduced intravascular volume, use of contrast agents with high osmolality (limited use today) and/or old age.





**65. In a 40-year-old woman, pap smear shows atypical glandular cells, The next step of management should be:**

- Repeat pap smear after three months
- Colposcopic directed cervical biopsy
- Colposcopy: cervical biopsy, endocervical curettage and endometrial biopsy
- Hysteroscopy and directed endometrial biopsy

**Ans. (c) Colposcopy, cervical biopsy, endocervical curettage and endometrial biopsy.**

The Papanicolaou test (abbreviated as Pap test, also known as Pap smear, cervical smear, or smear test) is a method of cervical screening used to detect potentially precancerous and cancerous processes in the cervix. Abnormal findings are often followed up by more sensitive diagnostic procedures, and if warranted, interventions that aim to prevent progression to cervical cancer. The test was invented by and named for doctor Aurel Babe' and doctor Georgios Papanikolaou.

**66. A patient develops skin necrosis 3 days after being started on warfarin for deep vein thrombosis. What is the most likely cause?**

- Antiphospholipid antibody syndrome
  - Protein C deficiency
  - Disseminated intravascular coagulation
  - Thrombotic thrombocytopenic .....
- Only 1
  - Only 3
  - 1 and 3
  - 1, 2, 3

**Ans. (b) Only 3**

Protein C is vitamin K dependent. Patients with Protein C deficiency are at an increased risk of developing skin necrosis while on warfarin. Protein C has a short half-life (8 hours) compared with other vitamin K dependent factors and therefore is rapidly depleted with warfarin initiation, resulting in a transient hypercoagulable state.

**67. Which of the following is associated with hypercoagulable state?**

- Protein C deficiency
- Antiphospholipid syndrome
- Homocysteinemia

**Select the correct answer using the code given below:**

- Only 1
- Only 3
- 1 and 3
- 1, 2, 3

**Ans. (d) 1, 2 and 3**

### Congenital & acquired hypercoagulable states

Congenital	Acquired
1. Protein C deficiency	1. Antiphospholipid antibody syndrome
2. Protein S deficiency	2. Malignancy

Contd...

Congenital	Acquired
3. Antithrombin deficiency	3. Surgery/Trauma
4. Factor V Leiden	4. Pregnancy/Oral contraceptives
5. Prothrombin gene G2010A mutation	5. Prolonged immobilization
6. Hyper-homocysteinemia	6. Older age
7. Dysfibrinolysis	

**68. Which of the following conditions is associated with cigarette smoking?**

- Non-specific interstitial pneumonia
  - Acute interstitial pneumonia
  - Cryptogenic organizing pneumonia
  - Desquamative interstitial pneumonia
- Only 1
  - Only 3
  - Only 4

**Ans. (c) Only 4**

Desquamative interstitial pneumonia is a form of idiopathic interstitial pneumonia featuring elevated levels of macrophages. Its name is derived from the former belief that these macrophages were pneumocytes that had desquamated. It is associated with patients with a history of smoking.

**69. Consider the following statements about infective endocarditis:**

- Modified Duke criteria are used for clinical diagnosis
- Echocardiographic findings form one of the major Duke criteria
- Presence of one major and two minor criteria is considered as diagnostic of endocarditis
- Presence of glomerulonephritis is a minor Duke criterion


**Which of the statements given above are correct?**

- 1, 2 and 3 only
- 1, 2 and 4 only
- 3 and 4 only
- 1, 2, 3 and 4

**Ans. (b) 1, 2 and 4 only**

### Infective endocarditis

- Established in 1994 by the Duke Endocarditis Service and revised in 2000, the Duke criteria are a collection of major and minor criteria used to establish a diagnosis of infective endocarditis. According to the Duke criteria, diagnosis of infective endocarditis can be definite, possible, or rejected.
- Evidence of endocardial involvement with positive echocardiogram is a major criteria
- Immunological problems is a minor criteria: glomerulonephritis, Osler's nodes, Roth's spots, Rheumatoid factor.



### MULTIPLE COMPLETION TYPE

70. A young female presented to medicine OPD with complaints of diarrhoea and fatigue. She had history of significant weight loss. Her physical examination is non significant and stool examination is normal. An endoscopy performed after microscopic examination and modified her diet. She is started on a special diet with no wheat or barely gains products. The dietary substitution causes marked improvement in her symptoms. Which of the following microscopic findings to be seen in the biopsy specimen?
- a. Lymphatic obstructions
  - b. Noncaseating granulomas
  - c. Atrophy of villi with blunting and flattening
  - d. Foamy macrophages within lamina propria
  - e. Increased in intraepithelial lymphocytes.
1. a, c are correct                      2. a, d are correct  
3. c, b are correct                      4. c, d are correct  
5. c, e are correct

---

**Ans. (5) c, e are correct**

---





# Sample Video Questions

1. Identify the technique?



- a. Bone marrow aspiration
- b. CSF aspirate
- c. Blood culture
- d. Pleural tap

2. Identify the technique?



- a. Bone marrow aspirate
- b. FNAC
- c. Blood culture
- d. Pleural tap

3. What is this brush used for



- a. Liquid based cytology
- b. Conventional cytology
- c. CSF needle
- d. Pleural tapping

4. What is this way of taking samples called:



- a. Order of draw
- b. Collection sample order
- c. Universal protocol
- d. None

5. What is the solution used when there is blood spill



- a. Hypochlorite
- b.  $H_2SO_4$
- c. HCl
- d. Formalin

For video, scan this QR Code



Ans.

- 1. a
- 2. b
- 3. a
- 4. a
- 5. a



## Annexure 1

### 1. Important Special Stains and Fixatives


Name of stain	Elements stained
For Microorganisms	
<i>Ziehl-Neelsen stain, Kinyoun stain</i>	Acid-Fast Organism
<i>May - Grünwald Giemsa Stain</i>	Bacteria, blood elements
<i>Gram stain</i>	Bacteria
<i>Toluidine Method, Steiner method</i>	Helicobacter pylori (stained black)
<i>Grocott's methenamine silver method, PAS</i>	Fungi
<i>Macchiavello stain</i>	Rickettsia and viral inclusions
<i>Shikata's orcein stain</i>	Hepatitis B Antigen
<i>Mucicarmine</i>	Cryptococcus
<i>Warthin – Starry method</i>	Spirochetes
<i>Gomori Methenamine silver</i>	Fungus (stained black)
<i>Calcofluor white</i>	Acanthamoeba (stained white)
For Connective tissue and lipids	
<i>Hematoxylin &amp; Eosin stain (H&amp;E)</i>	All tissues (most commonly used stain)
<i>Trichrome Stain</i>	Collagen
<i>Verhoeff - Van Gieson's stain (Best for Elastin)</i>	Elastic fibers
<i>Luna stain</i>	Elastin & Mast cells
<i>Silver Methenamine stain</i>	Reticulin
<i>Oil red O stain (on Fresh specimen/Frozen section)</i>	Fat
<i>Sudan black (on fixed specimen)</i>	
<i>Mallory's PTAH stain</i>	Muscle striations
<i>MSB (martius scarlet blue) (1<sup>st</sup> stain to stain fibrin in various stages)</i>	Fibrin
<i>PAS, Silver Methenamine stain</i>	Basement membrane
<i>Bielschowsky (silver stain)</i>	Neurofibrillary tangles, Senile plaques
<i>Luxol fast blue</i>	Myelin
<i>Papanicolau stain</i>	Cervical Exfoliative cytology
For Carbohydrates	
<i>PAS</i>	Glycogen/neutral mucin or mucoprotein
<i>Alcian blue</i> (at pH 2.5: positive for acid mucopolysaccharides)	Differentiates Acid & neutral mucopolysaccharides
<i>Mucicarmine stain (specific)</i>	Acidic epithelial Mucin
<i>Alcian blue at pH 1</i>	Highly acidic mucins (sulphated mucins)



For Amyloid	
<b>Congo Red stain</b>	Amyloid (Gold standard is Congo red staining with apple green birefringence under polarized light)
<b>Metachromatic stains like crystal violet</b>	
For Minerals, pigments and miscellaneous	
<b>Von Kossa stain (most commonly used)</b>	Calcium
<b>alizarin red S at pH 4.2 (specific for Calcium)</b>	
<b>Prussian blue</b>	Iron
<b>Fontana Masson Silver stain</b>	Melanin
<b>Modified fouchets</b>	Bile pigments
<b>Orcein</b>	Copper
<b>Modified rhodamine (method of choice)</b>	
For Hematological cells	
<b>Romanowsky stain (Giemsa, Leishman, Wright, Jenner)</b>	Routine blood & bone marrow
<b>Myeloperoxidase</b>	Myeloid cells
<b>Sudan Black B</b>	Myeloid cells
<b>Non-specific Esterase</b>	Monocytic cells
<b>Acid Phosphatase</b>	T-lymphocytes
<b>PAS</b>	Lymphoblasts (block positive), Dysplastic Erythroids, Megakaryocytes (granular positive)
<b>TRAP (Tartrate-resistant Acid Phosphatase)</b>	Hairy cell Leukemia
<b>Toluidine blue</b>	Mast cells & Basophils

## 2. Important Fixatives

HISTOPATHOLOGY	
<b>Routine fixative</b>	10% buffered normal Formalin (most commonly used)
<b>Electron Microscopy</b>	Glutaraldehyde (most commonly used), Osmium tetroxide
<b>Special Tissues</b>	<b>Fixatives</b>
<b>GI Biopsies &amp; testicular Bx for infertility diagnosis</b>	Bouin's fluid
<b>Bone Marrow Biopsies</b>	Zenker's fluid, B-5
<b>Brain Tissues</b>	Formalin ammonium Bromide
<b>Adrenal Medulla</b>	Orth's fluid
CYTOPATHOLOGY	
<b>Nuclear fixatives</b>	Carnoy's Fluid
<b>Cytoplasmic Fixatives</b>	Champy's Fluid
<b>PAP smears</b>	95% ethyl alcohol
<b>Cell blocks</b>	Bouin's fluid



# Annexure 2

## 3. Cytokines and Cytokine Receptors

Cytokine	Receptor	Cell Source	Cell Target	Biologic Activity
<b>IL-1,</b>		Monocytes/macrophages, B cells	All cells	<ul style="list-style-type: none"> <li>Explained in detail in text</li> </ul>
<b>IL-2</b>		<b>T cells<sup>q</sup></b>	T cells, B cells	<ul style="list-style-type: none"> <li>T cell activation and proliferation<sup>q</sup>,</li> <li>B cell growth<sup>q</sup></li> </ul>
<b>IL-3</b>		T cells	<b>Bone marrow progenitors<sup>q</sup></b>	<ul style="list-style-type: none"> <li>Hematopoietic progenitor stimulation.<sup>q</sup></li> </ul>
<b>IL-4</b>		T cells		<ul style="list-style-type: none"> <li>T<sub>H</sub>2 differentiation and proliferation.<sup>q</sup></li> <li>B cell Ig class switching<sup>q</sup></li> </ul>
<b>IL-5</b>		<b>T cells, mast cells &amp; eosinophils</b>	<b>Eosinophils,<sup>q</sup> basophils<sup>q</sup></b>	<ul style="list-style-type: none"> <li>Eosinophil activation.<sup>q</sup></li> </ul>
<b>IL-6</b>	<b>Gp130</b>	<b>Monocytes/macrophages, B cells</b>		<ul style="list-style-type: none"> <li>Myeloma cell<sup>q</sup> growth,</li> <li>Osteoclast growth and activation.<sup>q</sup></li> <li>T and B cell differentiation and growth<sup>q</sup></li> </ul>
<b>IL-7</b>		Bone marrow, <b>thymic epithelial cells<sup>q</sup></b>		<ul style="list-style-type: none"> <li>Important for T-cell development<sup>q</sup></li> </ul>
<b>IL-8</b>	<b>CXCR1, CXCR2</b>	Monocytes/macrophages	<b>Neutrophils</b>	<ul style="list-style-type: none"> <li>Neutrophil migration<sup>q</sup> (<i>neutrophil chemotactic factor</i>)</li> <li>Stimulates angiogenesis.<sup>q</sup></li> <li>Suppresses hepatic precursor proliferation.<sup>q</sup></li> </ul>
<b>IL-9</b>		T cells	Bone marrow progenitors	<ul style="list-style-type: none"> <li>Induces mast cell proliferation and function<sup>q</sup></li> </ul>
<b>IL-10</b>		Monocytes/ macrophages, T cells, B cells		<ul style="list-style-type: none"> <li>Anti-inflammatory molecule<sup>q</sup></li> </ul>
<b>IL-11</b>	<b>Gp130</b>	Bone marrow stromal cells	Megakaryocytes,	<ul style="list-style-type: none"> <li>Induces megakaryocyte colony formation<sup>q</sup></li> </ul>
<b>IL-12</b>		Activated macrophages	T cells	<ul style="list-style-type: none"> <li>Induces T<sub>H</sub>1 T helper cell formation &amp; lymphokine-activated killer cell formation.<sup>q</sup></li> </ul>
<b>IL-13</b>		T cells (T <sub>H</sub> 2)		<ul style="list-style-type: none"> <li>Inhibits macrophage proinflammatory cytokine production.<sup>q</sup></li> </ul>
<b>IL-14</b>		T cells	B cells	<ul style="list-style-type: none"> <li>Induces B cell proliferation</li> </ul>
<b>IL-15</b>		Monocytes/macrophages,	T cells	<ul style="list-style-type: none"> <li>T cell activation and proliferation.</li> <li>Promotes angiogenesis<sup>q</sup></li> </ul>
<b>IL-16</b>	<b>CD4</b>	Mast cells,CD8+ T cells	CD4+ T cells,	<ul style="list-style-type: none"> <li>Chemoattraction of CD4+ T cells,<sup>q</sup></li> <li>Inhibits HIV replication.<sup>q</sup></li> </ul>
<b>IL-17</b>		CD4+ T cells		<ul style="list-style-type: none"> <li>Neutrophil recruitment<sup>q</sup></li> </ul>
<b>IL-18</b>		Keratinocytes, macrophages		<ul style="list-style-type: none"> <li>Upregulated IFN production<sup>q</sup></li> </ul>
<b>IL-21</b>		CD4 T cells	NK cells	<ul style="list-style-type: none"> <li>Down-regulates NK cell activating molecules, NKG2D/DAP10<sup>q</sup></li> </ul>
<b>IL-23</b>		Macrophages	T cells	<ul style="list-style-type: none"> <li>Opposite effects of IL-12 (IL-17, -IFN)<sup>q</sup></li> </ul>



# Annexure 3


## 4. Type of Modified Macrophages

Modified Macrophages	Location
<i>Adipose tissue macrophages</i>	Adipose tissue
<i>Monocyte</i>	Bone Marrow/Blood
<i>Kupffer cell</i>	Liver
<i>Sinus histiocytes</i>	Lymph node
<i>Alveolar macrophages (dust cell)</i>	Pulmonary alveolus of lungs
<i>Tissue macrophages (Histiocyte) leading to Giant cells</i>	Connective tissues
<i>Langerhans cell</i>	Skin and mucosa
<i>Microglia</i>	Central Nervous System
<i>Hofbauer cell</i>	Placenta
<i>Intraglomerular mesangial cell</i>	Kidney
<i>Osteoclasts</i>	Bone
<i>Epithelioid cells</i>	Granulomas
<i>Red Pulp Macrophage (Sinusoidal lining cells)</i>	Red pulp of spleen
<i>Peritoneal macrophages</i>	Peritoneal cavity

## 5. Types of Giant Cells

Variety	Characteristics
<i>Physiological giant cells</i>	Osteoclasts, syncytiotrophoblasts and megakaryocytes.
<i>Langhan giant cell</i>	Nuclei present in the periphery, in a <b>horseshoe pattern</b> , seen in is <b>tuberculosis</b> . <sup>Q</sup>
<i>Foreign body giant cell</i>	<ul style="list-style-type: none"> <li>Nuclei are arranged randomly or haphazardly here.</li> <li>Seen in granuloma formed by <b>foreign bodies</b>, like sutures (intravenous drug abuse, talc)</li> <li>Appears <b>refractile</b><sup>Q</sup> when viewed with polarized light</li> </ul>
<i>Touton giant cells</i>	<ul style="list-style-type: none"> <li>Seen in <b>xanthomas</b>, <b>fat necrosis</b>, <b>xanthogranulomatous inflammation</b>, <b>dermatofibroma</b>.<sup>Q</sup></li> <li>Formed by fusion of epithelioid cells</li> <li>Contain a ring of nuclei surrounded by foamy cytoplasm.</li> </ul>
<i>Warthin-Finkeldey giant cells</i>	Measles. <sup>Q</sup>
<i>Reed-Sternberg cells</i>	Hodgkin's lymphoma <sup>Q</sup>
<i>Tumor giant cells</i> <sup>Q</sup>	Tumors, e.g. HCC





# Annexure 4

## 6. Important Translocations

Translocation	Gene (Chromosome)	Malignancy
(9;22)(q34;q11) <sup>Q</sup>	Bcr–Abl	Chronic myeloid leukemia <sup>Q</sup>
(11;14)(q13;q32)	Bcl1–IgH	Mantle cell lymphoma <sup>Q</sup>
(8;21) <sup>Q</sup> (15;17) <sup>Q</sup> (16;16) <sup>Q</sup>	RUNX1–RUNX1T1 PML–RARA CBFB–MYH11	Acute myeloid leukemia <sup>Q</sup>
(14;18)(q32;q21)	BCL2–IgH	Follicular lymphoma <sup>Q</sup>
(11;22)(q24;q12)	FLI1–EWS	Ewing's sarcoma <sup>Q</sup>
(1;7)(p34;q35) <sup>Q</sup>	LCK–TCRB	T cell acute lymphocytic leukemia
(8;14)(q24;q32)	Myc–IgH	Burkitt's lymphoma <sup>Q</sup> B cell acute lymphocytic leukemia <sup>Q</sup>
(2;13)(q35;q14)	PAX3 –FKHR/ALV	Alveolar rhabdomyosarcoma <sup>Q</sup>
(1;13)(p36;q14)	PAX7 –KHR/ALV	
Inv (2p13;p11.2-14)	REL–NRG	Non-Hodgkin's lymphoma
(10;17)(q11.2;q23)	RET–PKAR1A	Thyroid carcinoma
(1;3)(p34;p21) <sup>Q</sup>	TAL1–TCTA <sup>Q</sup>	Acute T cell leukemia <sup>Q</sup>
Inv1(q23;q31)	TRK–TPM3	Colon carcinoma
(12;22)(q13;q12) <sup>Q</sup>	ATF1–EWS	Malignant melanoma of soft parts
(11;22)(p13;q12) <sup>Q</sup>	WT1–EWS <sup>Q</sup>	Desmoplastic small round cell tumor <sup>Q</sup>

## 7. Important Genes & Chromosome


Gene	Chromosome No.
Rb	13q14.3
p53	17q13.1
APC	5q21
NF1 (Hamartin)	17q11
NF2	22p12
WT <sub>1</sub>	11p13
BRCA1	17q21
BRCA2	13q12
VHL	3p25
p16	9p21
RET (MEN2 syndrome)	10q
ABO	9
Rh	1
PRSS1	7q
FXN (Friedrich ataxia)	9q21.1
DMPK (Myotonic dystrophy)	19q13.3
Menin (MEN1 syndrome)	11q



## 8. Chromosomal Abnormality

Chromosomes involved	Neoplasm
t(9;22)	CML <sup>a</sup> , ALL (p190, FAB types L1 and L2)
t(8;14)	Burkitt's <sup>a</sup> lymphoma, ALL (type L3), Immunoblastic B cell lymphoma
t(15;17)	Acute Promyelocytic Leukemia <sup>a</sup>
t(11;14)	Mantle zone Lymphoma, Multiple myeloma, CLL <sup>a</sup>
t(11;22)	Ewing's sarcoma <sup>a</sup>
t(14;18)	Follicular lymphomas, DLBCL (20%) <sup>a</sup>
t(6;14)	Cystadenocarcinoma of ovary <sup>a</sup>
t(3;8)	Renal adenocarcinoma, Mixed parotid tumor (benign)
Trisomy 12	Chronic lymphocytic leukemia
Chr 8 and 17	Blast crisis of CML <sup>a</sup>
5q- and 7q-	MDS, AML
1p-	Neuroblastoma <sup>a</sup>
13q-	Retinoblastoma <sup>a</sup> , Osteosarcoma <sup>a</sup> , Small- cell carcinoma of lung <sup>a</sup>
Chr 22	Meningioma





# Annexure 5

## 9. Cancer Predisposition Syndromes and Associated Gene

Syndrome	Gene	Chr.	Inherit.	Tumors
<i>Ataxia telangiectasia</i>	<i>ATM</i>	11q22	AR	Breast
<i>Bloom syndrome</i>	<i>BLM</i>	15q26	AR	Acute Leukemia
<i>Cowden syndrome</i>	<i>PTEN</i>	10q23	AD	Breast, thyroid
<i>Familial adenomatous polyposis</i>	<i>APC</i>	5q21	AD	Intestinal adenoma, colorectal
<i>Familial melanoma</i>	<i>p16INK4</i>	9p21	AD	Melanoma, pancreatic
<i>Familial Wilms' tumor</i>	<i>WT1</i>	11p13	AD	Kidney (pediatric)
<i>Hereditary breast/ovarian cancer</i>	<i>BRCA1 BRCA2</i>	17q21 13q12	AD	Breast, ovarian, colon, prostate
<i>Her. diffuse gastric Ca</i>	<i>CDH1</i>	16q22	AD	Stomach
<i>Hereditary multiple exostoses</i>	<i>EXT1 EXT2</i>	8q24	AD	Exostoses, chondrosarcoma
<i>Hereditary retinoblastoma</i>	<i>RB1</i>	13q14	AD	Retinoblastoma, osteosarcoma
<i>Hereditary nonpolyposis colon cancer (HNPCC)</i>	<i>MSH2/6</i>	2p16	AD	Colon, endometrial, ovarian, stomach, small bowel, ureter Ca
<i>Hereditary papillary renal carcinoma</i>	<i>MET</i>	7q31	AD	Papillary kidney
<i>Juvenile polyposis</i>	<i>SMAD4</i>	18q21	AD	Gastrointestinal, pancreatic
<i>Li-Fraumeni Synd</i>	<i>TP53</i>	17p13	AD	Sarcoma, breast
<i>Multiple endocrine neoplasia type 1</i>	<i>MEN1</i>	11q13	AD	Parathyroid, endocrine, pancreas, and pituitary
<i>Multiple endocrine neoplasia type 2a</i>	<i>RET</i>	10q11	AD	Medullary thyroid carcinoma, pheochromocytoma
<i>Neurofibromatosis type 1</i>	<i>NF1</i>	17q11	AD	Neurofibroma, neurofibrosarcoma, brain
<i>Neurofibromatosis type 2</i>	<i>NF2</i>	22q12	AD	Vestibular schwannoma, meningioma, spine
<i>NBCCS (Gorlin syndrome)</i>	<i>PTCH</i>	9q22	AD	Basal cell carcinoma, medulloblastoma, jaw cysts
<i>Tuberous sclerosis</i>	<i>TSC</i>	9q34 16p13	AD	Angiofibroma, renal angiomyolipoma
<i>Von Hippel-Lindau</i>	<i>VHL</i>	3p25	AD	Kidney, cerebellum, pheochromocytoma



# Annexure 6


## 10. CD Markers used for Hematolymphoid Neoplasms, Location/Characteristics

Hematological cells	Immunophenotyping Markers (on Flow Cytometry)
<i>Blasts</i>	<b>CD34</b> , <sup>q</sup> tdt (Lymphoblasts), HLADR
<i>RBCs</i>	Glycophorin A
<i>Megakaryocytic marker</i>	<b>CD41</b> , <sup>q</sup> <b>CD61</b>
<i>WBCs</i>	<b>CD 45</b> <sup>q</sup> (Leukocyte Common Antigen) <sup>q</sup>
<i>Myeloid cells</i>	<b>Anti- MPO</b> , <sup>q</sup> <b>CD13</b> , <b>CD33</b> , <b>CD14</b> , <b>CD117</b>
<i>B –cells</i>	<b>CD19</b> , <sup>q</sup> <b>CD20</b> , <b>CD22</b> , <b>FMF7</b> , <b>CD23</b> , <b>CD79 a</b> , <b>CD79 b</b> , S Ig, IgM
<i>T-cells</i>	<b>CD3</b> , <sup>q</sup> <b>CD2</b> , <b>CD5</b> , <b>CD7</b> , <b>CD8</b> , <b>TCR-α/β</b> , <b>TCR-γ/δ</b>
<i>NK cells</i>	<b>CD16</b> , <sup>q</sup> <b>CD56</b> , <sup>q</sup> <b>CD57</b>
<i>Plasma cells</i>	<b>CD38</b> , <sup>q</sup> <b>CD138</b> , <sup>q</sup> kappa and Lambda light chains

CD Marker	Location/ Characteristics
<b>CD1</b>	<b>CD1a:</b> Cortical thymocytes, dendritic cells (DCs), epidermal Langerhans cells <b>CD1b:</b> Langerhans cells <b>CD1c:</b> Thymocytes, B cells, mantle zone, and umbilical cord <b>CD1d:</b> Intestinal epithelium, kidney tubular epithelia, hepatocytes, and thymus
<b>CD2</b>	Thymocytes, T cells
<b>CD3</b>	Signaling component of <b>T cell receptor</b> (TCR) complex
<b>CD4</b>	A co-receptor for MHC Class II; also a receptor <b>used by HIV</b> to enter T cells
<b>CD5</b>	T cells, <b>thymocytes</b> , and <b>B cells</b> , <b>Seen in CLL</b> , Mantle cell lymphoma and T-cell lymphoma
<b>CD7</b>	<b>Thymocytes</b> , some T cells
<b>CD8</b>	Co-receptor for <b>MHC Class I</b>
<b>CD9</b>	<b>Pre B cells</b> , eosinophils, basophils and platelets
<b>CD10</b>	<b>Pre-B cells</b> , <b>germinal-center B cells</b> , ALL, <b>follicular B- cell lymphomas</b>
<b>CD11c</b>	Dendritic cells and <b>hairy cell leukemia</b> cells
<b>CD13</b>	Myelomonocytic cells, AML, lymphoma and lymphocytic leukemia
<b>CD14</b>	<b>Macrophages</b> which binds to bacterial lipopolysaccharide
<b>CD16</b>	Fc receptor for IgG on NK cells
<b>CD19</b>	Component of the <b>B-cell co-receptor</b>
<b>CD20</b>	Transmembrane protein found on <b>B cells</b>
<b>CD21</b>	<b>CR2</b> , mature B cells, <b>Receptor for complement (C3d)</b> and <b>Epstein-Barr virus (EBV)</b>
<b>CD22</b>	Inhibitory receptor for <b>B cell receptor (BCR)</b> signaling
<b>CD23</b>	Mature B cells, monocytes, activated macrophages, <b>Seen in CLL</b>
<b>CD24</b>	<b>B lymphocytes</b>
<b>CD25</b>	High-affinity receptor for <b>IL-2</b> ; <b>Hairy cell leukemias</b>
<b>CD28</b>	T-cells, <b>co-stimulatory effect on the T-cell</b>
<b>CD30</b>	Present on <b>activated T and B cell</b> ; Hodgkin disease, <b>anaplastic large cell lymphomas</b>
<b>CD31</b>	<b>PECAM-1</b> , a cell adhesion molecule on platelets and endothelial cells

Contd....


















CD Marker	Location/ Characteristics
<b>CD33</b>	Immature <b>myeloid cells, AML blasts and mature monocytes</b>
<b>CD34</b>	<b>Stem cell marker</b> , adhesion, found on hematopoietic precursors
<b>CD38</b>	<b>Plasma</b> , and B & T activated cells
<b>CD40</b>	Co-stimulatory protein found on APCs induces antibody <b>isotype switching in B cells</b>
<b>CD41</b>	GpIIb/IIIa causes platelet aggregation, mutation causes <b>Glanzmann thrombasthenia</b>
<b>CD42</b>	GpIb/V/IX causes platelet adhesion, deficiency results in <b>Bernard-Soulier Syndrome</b>
<b>CD44</b>	Matrix adhesion molecules
<b>CD45</b>	Leucocyte common antigen (LCA)
<b>CD54</b>	Intercellular adhesion molecule -1 ( <b>ICAM-1</b> )
<b>CD55</b>	Complement decay-accelerating factor ( <b>DAF</b> )
<b>CD56</b>	<b>NCAM</b> (neural cell <b>adhesion molecule</b> ) on NK Cells some T-lymphocytes
<b>CD59</b>	Membrane attack complex inhibition factor ( <b>MACIF</b> )
<b>CD61</b>	Integrin $\alpha IIb\beta 3$ (gpIIb/IIIa) on platelets; major role is in platelet aggregation
<b>CD62E</b>	E-selectin
<b>CD62L</b>	L-selectin
<b>CD62P</b>	P-selectin
<b>CD64</b>	<b>Fc-gamma receptor 1 (fcyri)</b> on macrophages and monocytes
<b>CD68</b>	Used as immunocytochemical marker for staining of monocytes/macrophages
<b>CD69</b>	An early <b>activation marker on T cells and NK cells</b>
<b>CD71</b>	<b>Transferrin receptor</b> , mediates cellular uptake of iron
<b>CD72</b>	Mediator of B-cell - T-cell interactions
<b>CD 80 (B7-1)</b>	When bound to CD28 on T-cells, can provide the <b>co-stimulatory effect</b> . Causes up-regulation of a <b>high affinity IL-2 receptor</b> allowing T cells to proliferate
<b>CD 86 (B7-2)</b>	
<b>CD91</b>	<b>Low density lipoprotein (LDL)</b> receptor-related protein 1 (LRP1)
<b>CD95</b>	<b>Fas Receptor- receptor for Fas ligand</b> , an extrinsic apoptotic signal
<b>CD103</b>	<b>Hairy cell leukemia</b> (most specific)
<b>CD106</b>	VCAM-1
<b>CD117</b>	<b>c-kit</b> , the receptor for Stem Cell Factor, Myeloid marker
<b>CD122</b>	Beta subunit of IL-2 receptor0
<b>CD133</b>	Hematopoietic and CNS stem cell marker, Astrocytoma proliferation
<b>CD138</b>	A plasma cell-surface glycoprotein, known as syndecan-1
<b>CD141</b>	<b>Thrombomodulin</b>
<b>CD144</b>	VE-Cadherin adhesion molecule on the <b>vascular endothelium</b>
<b>CD209</b>	DC-SIGN, C-type lectin receptor found on dendritic cell subsets
<b>CD235a</b>	<b>Glycophorin</b> , a protein on blood cells



# Annexure 7

## 11. Features of the Peripheral Blood Smear

RBC Morphology	Images	Definition	Associated conditions
<b>Normal RBC</b>		<b>Central pallor</b>	None
<b>Polychromasia</b>		<b>Large, bluish</b> (due to residual RNA) RBCs lacking central pallor	<b>Rapid production &amp; release</b> of RBCs from BM; <b>Elevated retic % (hemolytic anemia)</b>
<b>Basophilic stippling</b>		<b>Small bluish dots</b> in RBCs ( <b>clustered polyribosomes</b> )	<b>Hemolytic anemia &amp; lead poisoning</b> (lead inhibits pyrimidine 5' nucleotidase, which normally digests the residual RNA)
<b>Pappenheimer bodies</b>		<b>Grayish, irregular inclusions</b> in RBCs composed of aggregates of <b>ribosomes, ferritin &amp; mitochondria</b>	Hemoglobinopathies, splenic hypofunction; megaloblastic anemia
<b>Target RBCs</b>		<b>Central round staining</b> in RBCs	<b>Liver disease, Dysbetalipoproteinemia, Hemolytic anemia, Severe IDA</b>
<b>Heinz bodies</b>		Several grayish, round inclusions represent aggregates of denatured hemoglobin	Indicative of oxidative injury <b>G6PD deficiency</b> , or unstable hemoglobin
<b>Howell-jolly bodies</b>		<b>Few purplish inclusions</b> on RBC represent residual fragments of nuclei containing chromatin	<b>Splenic hypofunction, post-splenectomy, severe hemolytic anemia</b>
<b>Schistocytes</b>		<b>Fragmented RBCs</b> helmet-shaped cells; indicative of shearing of the erythrocyte within the circulation	Microangiopathic hemolytic anemias, including <b>DIC, TTP or HUS</b> , mechanical causes of hemolysis, such as prosthetic valves
<b>Spherocytes</b>		RBCs that have <b>lost their central pallor</b> and appear <b>spherical</b> .	<b>H.S, AIHA, Toxins, Burns</b>
<b>Teardrop cells</b>		<b>Pear-shaped RBCs</b> indicative of mechanical stress on the RBC during release from the BM or passage through the spleen	<b>Thalassemia, megaloblastic anemia; myelofibrosis, myelophthisis (BM replacement)</b>
<b>Burr cells (echinocytes)</b>		<b>Smooth undulations</b> present on RBC surface circumferentially	<b>Uremia, drying artifact</b>
<b>Spur cells (acanthocytes)</b>		<b>Spiny points</b> present on the surface circumferentially; reflective of normal lipid composition of RBC membrane	<b>Liver disease, abetalipoproteinemia, RBCs lacking the Kell blood group antigen</b>





## 12. Blood-Clotting Factors

Factor	Names	Plasma Half-Life
I	Fibrinogen <sup>a</sup>	2–4 d <sup>a</sup>
II	Prothrombin <sup>a</sup>	3–4 d <sup>a</sup>
III	Thromboplastin <sup>a</sup>	–
IV	Calcium <sup>a</sup>	–
V	Proaccelerin, labile factor <sup>a</sup> , accelerator globulin	36 hr <sup>a</sup>
VII	Proconvertin, SPCA, stable factor	4–6 hr (min) <sup>a</sup>
VIII	Antihemophilic factor A (AHF) <sup>a</sup>	8–12 hr <sup>a</sup>
IX	Christmas factor <sup>a</sup> , antihemophilic factor B <sup>a</sup>	18–24 hr
X	Stuart–Prower factor <sup>a</sup>	40–60 hr <sup>a</sup>
XI	Plasma thromboplastin antecedent (PTA), antihemophilic factor C	40–70 hr
XII	Hageman factor <sup>a</sup> , glass factor	60 hr
XIII	Fibrin-stabilizing factor, Laki–Lorand factor <sup>a</sup>	11–14 d (max) <sup>a</sup>
HMW-K	High-molecular-weight kininogen, Fitzgerald factor <sup>a</sup>	150 hr
Pre-K <sub>a</sub>	Prekallikrein, Fletcher factor	35 hr
PL	Platelet phospholipid	–

# Annexure 8

## 13. Crystals in Urine


Triple Phosphate		<ul style="list-style-type: none"> <li>Seen in <b>alkaline urine</b></li> <li>Rectangular shape</li> </ul>
Uric acid		<ul style="list-style-type: none"> <li>Seen in <b>acidic urine</b></li> <li>Brown lemon shaped or star shaped</li> <li>Birefringent with polarized light</li> </ul>
Calcium oxalate		<ul style="list-style-type: none"> <li>Envelope shaped</li> <li>Seen in <b>acidic urine</b></li> </ul>



## 14. Different Types of Urinary Casts






Type	Images	Description	Significance
<b>Plain casts (Cast Matrix)</b>			
Hyaline		Consists of Tamm-Horsfall protein	Nonspecific; <b>0-2/lpf is normal<sup>q</sup></b> <b>Most frequently observed cast<sup>q</sup></b>  Increased number in: <b>renal diseases, dehydration<sup>q</sup>, fever<sup>q</sup>, congestive heart failure, diuretic therapy<sup>q</sup></b>
Waxy		Easily visualized due to high refractive index	Observed most frequently in <b>Chronic Renal Failure<sup>q</sup></b> ;  Broad waxy casts are called “ <b>Renal Failure Casts</b> ” <sup>q</sup>  Also seen in <b>acute &amp; chronic Renal Allograft Rejection<sup>q</sup></b>
<b>Cellular casts</b>			
RBC		Appears red-orange	<b>Pathognomonic of glomerulonephritis (RPGN)<sup>q</sup></b>  Indicator of <b>bleeding within the nephron</b>  Also seen in: <b>IgA Nephropathy<sup>q</sup>, Lupus Nephritis, Bacterial endocarditis, Renal infarction<sup>q</sup></b>
WBC		Protein matrix with WBCs	<b>Pyelonephritis<sup>q</sup> &amp; tubulointerstitial disease<sup>q</sup></b> , Also seen in <b>glomerular diseases<sup>q</sup>, Lupus Nephritis, Interstitial Nephritis<sup>q</sup></b>
Epithelial cell		Protein matrix with tubular cells	<b>Acute Tubular Necrosis<sup>q</sup>, Viral infection (CMV), Heavy Metal<sup>q</sup>, Salicylate poisoning<sup>q</sup></b>
<b>Inclusion Casts</b>			
Granular		Glycoprotein matrix with protein or cellular debris	Glomerular, <b>tubular, tubulointerstitial diseases, Renal graft rejection</b> , lead poisoning, after strenuous exercise  Coarse: in <b>Renal Papillary Necrosis<sup>q</sup></b>  Fine: May be <b>physiological<sup>q</sup>, Hyperparathyroidism<sup>q</sup></b>
Fatty		“ <b>Maltese cross pattern</b> ” in polarized light	Commonly seen in heavy proteinuria, characteristic of <b>nephrotic syndrome<sup>q</sup></b>
Crystal	(See Next page)	Urates, <b>Ca oxalate<sup>q</sup></b>	Indicate deposition of <b>crystals</b> in tubules or <b>collecting ducts</b>
<b>Other Casts</b>			
Pigment		Granular casts with pigment stain	Usually occurs in acute kidney injury due to <b>hemolysis<sup>q</sup> or rhabdomyolysis<sup>q</sup></b> or in acute tubular necrosis; <b>Bilirubin cast</b> in <b>obstructive jaundice</b>
Broad		Diameter 2 to 6 times of normal casts	Typically seen in <b>Chronic Renal Failure<sup>q</sup></b>  Indicate poor prognosis <sup>q</sup>





# Annexure 9

## 15. Rosettes

Rosette Type	Schematic Diagram	Definition	Associated Tumors
<b>Homer Wright rosette</b>		<ul style="list-style-type: none"> <li>Ball-like arrangement of cells that enclose meshwork of fibers.</li> <li><b>Fibers represent primitive neuronal processes<sup>Q</sup></b></li> </ul>	<ul style="list-style-type: none"> <li><b>Neuroblastoma,<sup>Q</sup></b></li> <li><b>Medulloblastoma,<sup>Q</sup></b></li> <li>PNET</li> <li>Pineoblastoma</li> </ul>
<b>Flexner-Wintersteiner rosette</b>		<ul style="list-style-type: none"> <li>Tumor cells circumscribe a <b>central lumen</b></li> <li>Lumen contains small <b>cytoplasmic extensions of the encircling cells.</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Retinoblastoma,<sup>Q</sup></b></li> <li>Pineoblastoma,</li> <li>Medulloepithelioma</li> </ul>
<b>True ependymal rosette</b>		<ul style="list-style-type: none"> <li>Rosettes with an <b>empty</b> appearing lumen</li> </ul>	<ul style="list-style-type: none"> <li><b>Ependymoma<sup>Q</sup></b></li> </ul>
<b>Perivascular pseudorosette</b>		<ul style="list-style-type: none"> <li>Spoke-wheel arrangement of cells</li> <li>Tapered cellular processes <b>radiates around a wall of a centrally-placed vessel.</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Ependymoma,<sup>Q</sup></b></li> <li>Medulloblastoma,</li> <li>PNET</li> <li>Central neurocytoma,</li> <li>Glioblastoma,</li> <li>Monomorphous pilomyxoid astrocytomas</li> </ul>
<b>Pineocytomatous &amp; Neurocytic rosette</b>		<ul style="list-style-type: none"> <li><b>Neuropil-rich rosettes</b> similar to Homer-Wright rosettes</li> <li><b>Larger and more irregular in contour</b></li> </ul>	<ul style="list-style-type: none"> <li><b>Pineocytoma<sup>Q</sup></b></li> <li><b>Central neurocytoma</b></li> </ul>




## 16. Familial Cancer Syndromes with Cutaneous Manifestations

Disease	Inheritance	Chromosomal Location	Gene/Protein
<i>Ataxia-telangiectasia</i>	AR <sup>a</sup>	11q22.3	<i>ATM/ATM</i> <sup>a</sup>
<i>Nevoid basal cell carcinoma syndrome</i>	AD	9q22.3	<i>PTCH/PTCH</i> <sup>a</sup>
<i>Cowden syndrome</i>	AD	10q23	<i>PTEN/PTEN</i> <sup>a</sup>
<i>Familial melanoma syndrome</i>	AD	<b>9p21</b> <sup>a</sup>	<i>CDKN2/p16/INK4</i> ; <sup>a</sup> <i>CDKN2/p14/ARF</i> <sup>a</sup>
<i>Muir-Torre syndrome</i>	AD	2p22	<i>MSH2/MSH2</i> ; <sup>a</sup> <i>MLH1/MLH1</i>
<i>Tuberous sclerosis</i>	AD	9q34, 16p13.3	<i>TSC1/hamartin</i> ; <sup>a</sup> <i>TSC2/tuberin</i> <sup>a</sup>
<i>Neurofibromatosis I</i>	AD	<b>17q11.2</b> <sup>a</sup>	<i>NF1/neurofibromin</i> <sup>a</sup>
<i>Neurofibromatosis II</i>	AD	22q12.2	<i>NF2/merlin</i> <sup>a</sup>
<i>Xeroderma pigmentosum</i>	AR <sup>a</sup>	9q22 and others	<i>XPA/XPA</i> and others

## 17. Relationship between Proteins and Neurodegenerative Diseases


Protein	Diseases with Inclusions
A $\beta$	Alzheimer disease <sup>a</sup>
Tau	Alzheimer disease <sup>a</sup> Frontotemporal lobar degeneration <sup>a</sup> Parkinson disease (with LRRK2 mutations) <sup>a</sup> Progressive supranuclear palsy Corticobasal degeneration <sup>a</sup>
TPD-43	Frontotemporal lobar degeneration Amyotrophic lateral sclerosis <sup>a</sup>
FUS	Frontotemporal lobar degeneration <sup>a</sup> Amyotrophic lateral sclerosis
$\alpha$ -synuclein	Parkinson disease <sup>a</sup> Multiple system atrophy <sup>a</sup>
Polyglutamine aggregates (distinct proteins per disease)	Huntington's disease <sup>a</sup> Some forms of spinocerebellar ataxia Spinal bulbar muscular atrophy



# Annexure 10









## 18. Pattern of Injury and Associated Agents

Pattern of Injury	Examples of Associated Agents
Budd-Chiari syndrome	Oral contraceptives
<i>Peliosis hepatis (Dilated sinusoids)</i>	<b>Anabolic steroids, Danazol, OCPs, tamoxifen<sup>q</sup></b>
<i>Hepatic adenoma</i>	Oral contraceptives, anabolic steroids
<i>Hepatocellular carcinoma</i>	<b>Thorotrast<sup>q</sup></b>
<i>Cholangiocarcinoma</i>	Thorotrast
<i>Angiosarcoma</i>	Thorotrast, <b>vinyl chloride, Arsenic, Thorium dioxide<sup>q</sup></b>
<i>Cholestasis</i>	<b>OCPs and anabolic steroids<sup>q</sup></b>
<i>Cholestatic hepatitis</i>	Chlorpromazine, Statins
<i>Spotty hepatocyte necrosis</i>	Methyldopa, phenytoin
<i>Submassive necrosis, zone 3</i>	Paracetamol, <b>halothane<sup>q</sup></b>
<i>Massive necrosis</i>	<b>Isoniazid, phenytoin<sup>q</sup></b>
<i>Macrovesicular Steatosis</i>	Ethanol, methotrexate, corticosteroids, TPN
<i>Microvesicular Steatohepatitis with Mallory bodies</i>	<b>Amiodarone, ethanol<sup>q</sup></b>
<i>Fibrosis and cirrhosis</i>	Methotrexate (Periportal fibrosis), isoniazid, enalapril
<i>Noncaseating Granulomas</i>	<b>Sulfonamides<sup>q</sup></b>




# Annexure 11

## 19. Morphological Differentiation of Malaria Parasites

	<i>P. falciparum</i>	<i>P. vivax</i>	<i>P. ovale</i>	<i>P. malariae</i>
<b>Infected red cells</b>	Normal size, <sup>a</sup> Maurer's clefts <sup>b</sup>	Enlarged; Schuffner's dots <sup>c</sup>	Enlarged; oval and fimbriated; Schuffner's dots <sup>c</sup>	Normal or microcytic; stippling not usually seen
<b>Ring forms (early trophozoites)</b>	Delicate; frequently 2 or more; accolé forms; <sup>d</sup> small chromatin dot  Falciparum Ring forms	Large, thick; usually single (occasionally 2 in cell; large chromatin dot)  Vivax Trophozoite	Thick compact rings 	Very small, compact rings 
<b>Later trophozoites</b>	Compact, vacuolated; sometimes 2 chromatin dots	Amoeboid; central vacuole; light blue cytoplasm	Smaller than <i>P. Vivax</i> , slightly amoeboid	Band across cell; deep blue cytoplasm
<b>Schizonts</b>	18-24 merozoites filling 2/3 of cell	12-24 merozoites, irregularly arranged	8-12 merozoites filling 3/4 of cell	6-12 merozoites in daisy-head around central mass of pigment
<b>Pigment</b>	Dark to black clumped mass	Fine granular; yellow brown	Coarse light brown	Dark, prominent at all stages
<b>Gametocytes</b>	Crescent of sausage-shaped; diffuse chromatin; single nucleus  Falciparum gametocyte	Spherical compact, almost fills cell; single nucleus 	Oval, fills 3/4 of cell; similar to but smaller than <i>P. vivax</i> 	Round; fills 1/2 to 2/3 of cell; similar to <i>P. vivax</i> but smaller, with no Schuffner's dots 





# Annexure 12

## 20. Autoantibodies in Autoimmune Disease

Autoantibodies	Disease	Test sensitivity (%)
Anti-acetylcholine receptor	Myasthenia gravis	>85
Anti-basement membrane	Goodpasture syndrome	>90
Anticentromere	CREST syndrome Diffuse systemic sclerosis	40 <2
Antiendomysial IgA	Celiac disease	95
Antigliadin IgA	Celiac disease	80
Anti histone	Drug-induced lupus	90-95
Anti-insulin	Systemic lupus erythematosus, Type 1 diabetes	50-70 50
Anti-islet cell	Type 1 diabetes	75-80
Anti-intrinsic factor	Pernicious anemia	60
Anti-parietal cell	Pernicious anemia	90
Anti microsomal	Hashimoto thyroiditis	97
Anti-Smith (Sm)	Systemic lupus erythematosus	20-30
Anti-SS-A (Ro)	Sjögren syndrome Systemic lupus erythematosus	70-95 30-50
Anti-SS-B (La)	Sjögren syndrome	60-90
Antithyroglobulin	Systemic lupus erythematosus Hashimoto thyroiditis	10-15 85
Anti-tissue transglutaminase IgA	Celiac disease	98
Anti-DNA topoisomerase	Diffuse systemic sclerosis	30-70
Antimitochondrial	CREST syndrome Primary biliary cirrhosis	10-20 90-100
Anti myeloperoxidase	Microscopic polyangiitis	80 (p-ANCA)
Antinuclear	Systemic lupus erythematosus Systemic sclerosis Dermatomyositis	-100 70-90 <30
Antiproteinase 3	Polymyositis MCTD Primary biliary cirrhosis Wegener granulomatosis	30-60 95-99 50 >90 (c-ANCA)
Anti-ribonucleoprotein	MCTD Systemic lupus erythematosus	85 30-40
Anti-TSH receptor	Graves disease	85



# Cell as a Unit of Health and Disease

## Key Points

- » Human genome contains **3.2 billion DNA base pairs**
- » **The most common forms of DNA variation in the human genome is Single-Nucleotide Polymorphisms**
- » **Epigenetics is heritable changes<sup>o</sup> in gene expression, not caused by alterations in DNA sequence**
- » Proteins involved in tight junctions are **Occludin, Claudin, Zonulin and Catenin**
- » **All growth factors are proto-oncogenes, except  $TGF\beta$ ,<sup>o</sup> which is a tumor suppressor gene**
- » **Type IV collagen is a nonfibrillary collagen which contribute to the structures of planar basement membranes**
- » **Bone Morphogenic Protein (BMP) is both mitogenic and morphogenic**
- » **Collagen I is the most abundant collagen in the body**
- » **Stem cells are characterized by two important properties: Self-renewal and Asymmetric division**
- » **Transdifferentiation<sup>o</sup>: Irreversible conversion of cells from one differentiated cell type to another**
- » **Shinya Yamanaka and Sir John Gurdon were awarded Nobel Prize for IPSC in 2012**

## Key Recent Updates

CRISPR–clustered regularly interspaced short palindromic repeats forms genome editing technology.



## THE HUMAN GENOME

- Human genome contains **3.2 billion DNA base pairs**.<sup>Q</sup>
- 20,000 protein-encoding genes**<sup>Q</sup>, comprising only **1.5% of the genome**.<sup>Q</sup>
- Function of these **protein encoding genes**: **enzymes, structural components, and signaling molecules** and to **assemble and maintain** all cells in the body.

### Noncoding DNA

- It refers to the **98.5%** of human genome that does **not encode proteins**.
- The amount of noncoding DNA varies greatly among species**. E.g., in bacteria, only 2% of genome is noncoding DNA.
- Noncoding DNA is **also transcribed into functional noncoding RNA molecules** (e.g., **transfer RNA, ribosomal RNA, and regulatory RNAs**)

9<sup>th</sup>

### Latest Update

#### ENCODE (ENCyclopedia of DNA Elements) project: 2007

- This project has systematically **mapped regions of transcription, transcription factor association, chromatin structure and histone modification**.
- Striking conclusion: 80% of the human genome, even the noncoding regions either binds proteins or regulate gene expression.**

Major **classes of functional nonprotein-coding sequences**:


Nonprotein coding sequences	Characteristics
<b>Promoter &amp; enhancer regions</b> <sup>Q</sup>	Provide <b>binding sites</b> for transcription factors
<b>Binding sites for factors</b>	<b>Organize and maintain</b> higher order <b>chromatin structures</b>
<b>Noncoding regulatory RNAs</b>	Regulate gene expression: <b>miRNAs</b> and <b>long noncoding RNAs</b>
<b>Mobile genetic elements</b> <sup>Q</sup> ( <b>transposons</b> /" <b>jumping genes</b> ")	Segments <b>that move around the genome</b> , exhibiting <b>wide variation</b> in number and positioning; Role in gene regulation and chromatin organization
<b>Special structural regions of DNA</b>	Chromosome ends ( <b>Telomeres</b> ) and "tethers" ( <b>centromeres</b> )

## POLYMORPHISM

**Any two individuals share greater than 99.5% of their DNA sequences**.<sup>Q</sup> So what is the reason of genetic variations?

The most common forms of DNA variation in the human genome is shown in Flowchart 1.

**Flowchart 1:** DNA variation in human genome



### Role of SNPs

- SNPs located in **noncoding regions** are **regulatory elements**<sup>Q</sup> in the genome
- They **alter gene expression**<sup>Q</sup> and have **direct influence** on **disease susceptibility**
- Even if any SNP has no effect on gene function (**Neutral SNP**)<sup>Q</sup>, it may be coinherited with the "actual disease causing gene," if located close to that gene → "**Linkage disequilibrium**"<sup>Q</sup>
- May act as **markers of multigenic complex diseases**, E.g. Diabetes, Hypertension.

### Epigenetics

- Definition**
  - Heritable changes**<sup>Q</sup> in **gene expression**, not caused by alterations in **DNA sequence**.<sup>Q</sup>
- Epigenetic factors**
  - Histones and histone-modifying factors
    - Histone methylation, Histone acetylation, Histone phosphorylation, DNA methylation, Chromatin organizing factors**



- **Significance**
  - Epigenetic **Dysregulation** → **central role in malignancy**
  - Many other diseases are associated with **inherited or acquired epigenetic alterations**. E.g. **Genomic imprinting in Prader Willi syndrome**
  - Epigenetic alterations like histone acetylation and DNA methylation are **reversible** and are **responsive to drugs**; So, **HDAC inhibitors and DNA methylation inhibitors** are being tested in the treatment of cancer
- **Diagnosis**
  - **Sequencing**
  - Chip on chip (**Microarray technology**)
  - Using **Methylation specific primers** in Polymerase chain reaction (PCR)
  - **Bisulphite method**: Bisulphite **converts unmethylated cytosine to uracil**, which **acts like thymine** in downstream reactions. The unmethylated (modified) DNA is detected by sequence analysis.



### High Yield Facts


- **SNP** is the most common type of **DNA polymorphism**<sup>Q</sup>
- **SNP** can be **detected** by **SNP Cytogenomic Array**.
- **Linkage analysis** can be used to detect **SNPs** with **profound effects and high penetrance**.<sup>Q</sup>
- **Linkage analysis** is used to **identify unknown genes**<sup>Q</sup> associated with a disease
- **GWAS** can be used to **identify unknown genes NOT** in 'Linkage disequilibrium'<sup>Q</sup>
- The 2 types of genetic polymorphisms most useful for linkage analysis are **SNPs and repeat length polymorphisms**<sup>Q</sup>
- Repeat length polymorphisms can be **mini-satellite (1-3 kb)**<sup>Q</sup> or **Micro-satellite repeats (<1 kb)**<sup>Q</sup>
- **Genome-wide association study (GWAS)** refers to 'Genome Wide Association Studies'<sup>Q</sup>
- In **GWAS**, **entire genome of large number of individuals** with and without a disease are examined for common **genetic variants or polymorphisms** that are over represented in patients with the disease.

## NONCODING REGULATORY RNA

**A. Micro-RNA (miRNA):** Small **noncoding RNA molecule (22 nucleotides)** which causes **RNA silencing** and **post-transcriptional regulation** of gene expression.

- Generation and mode of action of miRNA is shown in Flowchart 2.

**Flowchart 2:** Generation of MicroRNA (miRNA) and their mode of action



### Small Interfering RNAs (siRNAs)

- miRNA **introduced experimentally** into cells and inhibit them
- So-called "**knockdown technology**"
- Possible **therapeutic agents** to **silence pathogenic genes**, such as **oncogenes** involved in neoplastic transformation.

**B. Long Noncoding RNA (lncRNA or long ncRNAs): Nonprotein coding transcripts longer than 200 nucleotides**<sup>Q</sup>

**lncRNAs modulate gene expression in following ways:**

Mechanism	Characteristics
<b>Gene activation</b>	Can <b>facilitate transcription factor binding</b> → promote gene activation
<b>Gene suppression</b>	Can preemptively <b>bind transcription factors</b> → prevent gene transcription.
<b>Histone &amp; DNA modification</b>	Bind to and <b>direct acetylases/methylases</b> (or deacetylases/demethylases)
<b>Assembly of protein complexes</b>	Act as scaffolding to stabilize secondary/tertiary structures and/or multi-subunit complexes influencing chromatin architecture or gene activity
<b>Repressive function</b>	Example: <b>XIST</b> → role in physiologic X chromosome inactivation



### High Yield Facts

- **lncRNAs** have been found to have link with diseases like **atherosclerosis & cancer**.
- **XIST** refers to '**X-inactive specific transcript**'<sup>Q</sup>
- **XIST** is a **RNA gene (17 kb)** on the **X chromosome** that acts as **major effector of the X inactivation process**.
- **XIST** is **expressed on the inactive X chromosome** and not on the active one.





## CELLULAR HOUSEKEEPING

### 'Housekeeping Genes'<sup>Q</sup>

- **Constitutive genes** that are **expressed in all cells of an organism<sup>Q</sup>**
- **Required for the maintenance of basic cellular functions** like protection from environment, nutrient acquisition, communication, movement, renewal of senescent molecules, molecular catabolism, and energy generation.
- Many housekeeping functions are **compartmentalized within intracellular organelles**.

### Housekeeping Functions of Different Intracellular Organelles

Organelle	House-keeping function
<i>Rough ER*</i>	<b>Synthesize new proteins<sup>Q</sup></b> for the plasma membrane
<i>Golgi apparatus</i>	<b>Assembles Proteins</b> physically
<i>Smooth ER*</i>	<b>Hormone<sup>Q</sup></b> and <b>lipoprotein</b> synthesis, <b>modification</b> of hydrophobic <sup>Q</sup> compounds into water-soluble molecules
<i>Lysosomes</i>	<b>Digestion<sup>Q</sup></b> of <b>proteins, polysaccharides, lipids, nucleic acids</b>
<i>Proteasomes</i>	Selectively <b>chews up denatured proteins</b> , releasing <b>peptides<sup>Q</sup></b>
<i>Peroxisomes</i>	<b>Breakdown of fatty acids<sup>Q</sup></b> , generating <b>hydrogen peroxide</b>
<i>Endosomal vesicles</i>	<b>Shuttle internalized material<sup>Q</sup></b> to appropriate intracellular sites
<i>Mitochondria</i>	<b>Synthesize ATP<sup>Q</sup></b> through <b>oxidative phosphorylation</b> & synthesize metabolic <b>intermediates</b> needed for anabolic metabolism

\* ER → Endoplasmic Reticulum

## PLASMA MEMBRANE

**Fluid bilayer of amphipathic phospholipids** with **hydrophilic head** and **hydrophobic lipid tails** that forms a **barrier to passive diffusion** of large or charged molecules.

### Phospholipids in Plasma Membrane and their Functions

Phospholipid	Location	Function
<i>Phosphatidylinositol</i>	Inner leaflet of membrane	Acts as <b>electrostatic scaffold</b> for intracellular proteins <b>Hydrolyzed by phospholipase C</b> to generate 2 <sup>nd</sup> messengers like diacylglycerol ( <b>DAG</b> ) and inositol trisphosphate ( <b>IP3</b> ). <sup>Q</sup>
<i>Phosphatidylserine</i>	Inner face	Confers a <b>–ve charge<sup>Q</sup></b> involved in electrostatic protein interactions
	Extracellular face	Acts as <b>"eat me" signal<sup>Q</sup></b> for phagocytes, in cells undergoing <b>apoptosis<sup>Q</sup></b>
	Platelets	Serves as a <b>cofactor in the clotting<sup>Q</sup></b> of blood
<i>Glycolipids and sphingomyelin</i>	Extracellular face	Important in cell-cell and cell-matrix interactions including <b>inflammatory cell recruitment</b> and <b>sperm-egg interaction<sup>Q</sup></b>

R9<sup>th</sup>

### Latest Update

**Mechanisms of movement across plasma membrane**

#### 1. Passive Membrane Diffusion:

- Transfers:
  - ◆ **Small, nonpolar<sup>Q</sup>** molecules (e.g. O<sub>2</sub> and CO<sub>2</sub>)
  - ◆ **Hydrophobic<sup>Q</sup>** molecules (e.g., estradiol or vitamin D)
  - ◆ **Polar molecules <75 daltons<sup>Q</sup>** (e.g., water, ethanol, and urea).
- **Effective barrier to: polar molecules >75 daltons e.g., Glucose & Ions**, due to their charge and high degree of hydration.

Contd...



## 2. Channels and Carrier proteins:

- For low molecular weight molecules (<1000 daltons)
- Each transported molecule (e.g., ion, sugar, nucleotide) requires a specific transporter

Characteristics	Channel proteins	Carrier proteins
<b>Mechanism</b>	Create <b>hydrophilic<sup>Q</sup></b> pores	Bind their specific solute → <b>undergo conformational changes</b> → transfer the ligand across the membrane
<b>Speed of transport</b>	<b>Rapid<sup>Q</sup></b>	<b>Relatively slow<sup>Q</sup></b>
<b>Direction of movement</b>	<b>Along</b> concentration gradient	<b>Against<sup>Q</sup></b> concentration gradient
<b>Active transport</b>	<b>Cannot occur<sup>Q</sup></b>	<b>Can mediate active transport<sup>Q</sup></b>

## Receptor-mediated and Fluid-phase Uptake

R<sup>9th</sup>

### Latest Update

**Endocytosis:** Mechanism of uptake of **fluids or macromolecules** by the cell; Occurs by two major mechanisms:

Characteristics	Caveolae-mediated endocytosis	Receptor-mediated endocytosis
<b>Includes</b>	<b>Potocytosis ("cellular sipping")<sup>Q</sup></b>	<b>Pinocytosis ("cellular drinking")</b>
<b>Protein involved</b>	Caveolin	Clathrin
<b>Mechanism</b>	<p><b>Caveolae ("little caves")</b> or <b>noncoated</b> plasma membrane invaginations are formed</p> <p>↓</p> <p>Internalization of caveolae with any bound molecules and associated extracellular fluid</p>	<p>Begin at <b>Clathrin-coated pit</b> in plasma membrane</p> <p>↓</p> <p><b>Invaginates &amp; pinches off</b> to form a <b>clathrin-coated vesicle</b> containing the macromolecules</p> <p>↓</p> <p><b>Vesicles uncoat and fuse with</b> an acidic intracellular structure (<b>Early endosome</b>)</p> <p>↓</p> <p><b>Discharge their contents</b> for digestion &amp; passage to <b>lysosome</b></p> <p>↓</p> <p>Endocytosed vesicles <b>may recycle</b> back to plasma membrane for another round of ingestion</p>
<b>Significance</b>	Transports <b>vitamins like folate</b> & Regulate <b>transmembrane signaling</b> and/or <b>cellular adhesion</b> via <b>Integrins</b>	Major uptake mechanism for certain macromolecules like <b>transferrin</b> and <b>low-density lipoprotein (LDL)</b>



### High Yield Facts

- Multi Drug Resistance (**MDR**) protein is a **Transporter ATPase<sup>Q</sup>** which pumps polar compounds (Chemotherapy drugs) out of cells & **may render cancer cells resistant** to treatment
- **Hypertonicity** (extracellular salt > in cytosol) causes a **net movement of water out<sup>Q</sup>** of cells.
- **Hypotonicity** causes a net movement of **water into cells.<sup>Q</sup>**
- Most cytosolic enzymes work at pH 7.4 whereas **lysosomal enzymes** function best at **pH ≤ 5.<sup>Q</sup>**
- **Exocytosis:** The process by which **membrane-bound vesicles<sup>Q</sup>** **fuse with the plasma membrane** and discharge their contents to the **extracellular space**.
- **Transcytosis** is the **movement of endocytosed vesicles** between the **apical and basolateral<sup>Q</sup>** compartments of cells, for transferring **intact proteins<sup>Q</sup>** across epithelial barriers.
- **Phagocytosis** involves **non-clathrin-mediated<sup>Q</sup>** membrane invagination of large particles, by specialized phagocytes.

## CYTOSKELETON

**Intracellular scaffolding** of proteins that helps a cell to adopt a **particular shape,<sup>Q</sup>** **maintain polarity,<sup>Q</sup>** **organize the relationship of intracellular organelles,** and **move.<sup>Q</sup>**

### Three Major Classes of Cytoskeletal Proteins

#### 1. Actin Microfilaments

- **5- to 9-nm diameter fibrils** formed from the **globular protein actin (G-actin)**,
- **Most abundant cytosolic protein in cells.<sup>Q</sup>**
- G-actin monomers **noncovalently polymerize** into long **filaments (F-actin)** that intertwine to form **double-stranded helices** with a defined polarity; (positive and negative)
- In **muscle:** Myosin binds to actin with ATP and **cause muscle contraction.<sup>Q</sup>**
- In **nonmuscle cells:** **F-actin** assembles into well-organized **bundles and networks** that control **cell shape and movement.<sup>Q</sup>**



## 2. Intermediate Filaments

- Impart **tensile strength**<sup>Q</sup> to tolerate **mechanical stress**.
- Characteristic **tissue-specific patterns of expression**<sup>Q</sup>
- Can be **useful for assigning a cell of origin**<sup>Q</sup> for **poorly differentiated tumors**.

Intermediate Filaments	Tissue
<b>Lamin A, B, and C</b>	<b>Nuclear lamina</b> of all cells
<b>Vimentin</b>	Mesenchymal cells ( <b>fibroblasts, endothelium</b> ) <sup>Q</sup>
<b>Desmin</b>	<b>Muscle cells</b> , <sup>Q</sup> forming the scaffold on which actin & myosin contract
<b>Neurofilaments</b>	<b>Axons of neurons</b> <sup>Q</sup> , imparting <b>strength and rigidity</b>
<b>Glial fibrillary acidic protein</b>	<b>Glial cells</b> around neurons
<b>Cytokeratins</b>	<b>Acidic (type I)</b> and <b>neutral/basic (type II)</b> ; different types present in different cells: used as cell markers

## 3. Microtubules

- 25-nm thick fibrils composed of non-covalently polymerized dimers of  $\alpha$ - and  $\beta$ -tubulin
- 2 varieties** of these motor proteins:
  - Kinesins: for anterograde (– to +) transport
  - Dyneins: for retrograde (+ to –) transport
- Functions:**
  - Serve as connecting cables for “**molecular motor**” proteins that use **ATP to move vesicles, organelles, or other molecules** around the cells;
  - Participate in **sister chromatid separation during mitosis**.
  - Adapted to **form motile cilia** (bronchial epithelium) or **flagella (in sperm)**.<sup>Q</sup>



## High Yield Facts

- The largest protein molecule in skeletal muscle is **Titin**.
- Communicating junctions (Gap junctions)** consists of **connexons**<sup>Q</sup> formed by **connexins**<sup>Q</sup>

## CELL-CELL INTERACTIONS

Cells **interact** and **communicate with one another** by forming **junctions** that provide **mechanical links** and enable surface receptors to **recognize ligands on other cells**.

### Three Basic Types


#### 1. Occluding junctions (Tight junctions):

- Seal adjacent cells together to create a **continuous barrier** that **restricts**<sup>Q</sup> the **paracellular (between cells) movement of ions and other molecules**.
- Proteins involved are **Occludin, Claudin, Zonulin and Catenin**.<sup>Q</sup>

#### 2. Anchoring junctions (Desmosomes)<sup>Q</sup>

- Mechanically attach cell and their intracellular cytoskeletons—to other cells or to the extracellular matrix (ECM)**
- Types**

<b>Spot desmosome or Macula adherens</b>	When the adhesion focus is <b>small and rivet-like</b> ; Proteins involved: <b>desmogleins and desmocollins</b>
<b>Hemidesmosome</b>	When such a focus attaches the <b>cell to the ECM</b> integrins; E-cadherins
<b>Belt desmosomes</b>	Similar adhesion domains occurring as <b>broad bands between cells</b>
<b>Cell-cell desmosomal junctions</b>	Formed by homotypic association of <b>transmembrane glycoproteins</b> called <b>Cadherins</b>



Interaction of cells with extracellular matrix (ECM)



### 3. Communicating junctions (Gap junctions)

- Mediate the passage of **chemical or electrical signals** from **one cell to another**.
  - Consists of a dense planar array of 1.5–2-nm **pores** called **connexons**<sup>Q</sup>
  - Formed by hexamers of transmembrane **proteins** called **connexins**<sup>Q</sup>

## SIGNAL TRANSDUCTION PATHWAYS

- **First messengers:** The extracellular ligands that bind to receptors
- **Second messengers:** The intracellular mediators of action, following binding of ligand to receptor

### Newer receptors that are must to know

Receptors	Mechanism	Role in
<b>Notch family</b> receptors	<b>Proteolytic cleavage of the receptor</b> ↓ Nuclear translocation of the cytoplasmic piece ↓ Formation of a transcription complex	Development: <ul style="list-style-type: none"> <li>• Neuronal development</li> <li>• Angiogenesis</li> <li>• T cell development</li> </ul> Diseases: <b>T-ALL, Multiple Sclerosis</b>
<b>Frizzled family receptors</b> for <b>Wnt protein ligands</b>	APC-β-catenin degradation complex not formed ↓ β-catenin translocates to the nucleus ↓ Causes <b>transcriptional activation</b> & cell proliferation	<ul style="list-style-type: none"> <li>• Action of <b>APC gene</b></li> <li>• Disease: Adenomatosis Polyposis Coli &amp; <b>Colorectal Carcinoma</b><sup>Q</sup></li> </ul>



### High Yield Facts

- Mutation in **TTN gene** on chromosome 2 causes **dilated Cardiomyopathy**
- **TGF β** family includes **BMP**<sup>Q</sup>, **Activins**, **Inhibins**, **Mullerian Inhibiting substance**<sup>Q</sup>
- **TGF β** is an **anti-inflammatory**<sup>Q</sup> cytokine and causes **fibrosis**<sup>Q</sup>

## GROWTH FACTORS

- Growth factor activity is mediated through binding to specific receptors, ultimately influencing the expression of genes that can:
  - **Promote entry of cells into the cell cycle**
  - **Relieve blocks** on cell cycle progression (thus promoting replication)
  - **Prevent apoptosis**
  - Enhance **biosynthesis of cellular components** (nucleic acids, proteins, lipids, carbohydrates) required for a mother cell to give rise to two daughter cells



### High Yield Facts

- All growth factors are **proto-oncogenes**, except **TGF β**<sup>Q</sup>, which is a **tumor suppressor gene**.
- **Bone Morphogenic Protein (BMP)** is both **mitogenic & morphogenic**<sup>Q</sup>
- **Hypoxia**<sup>Q</sup> is the most important inducer of **VEGF** production
- **EGF receptor family** includes **ERB-B1(EGFR1) & ERB-B2 (HER2)**<sup>Q</sup>
- Mutation of **ERB-B1** causes **Lung Adenocarcinoma**
- Mutation of **ERB-B2** causes **Breast Carcinoma**

## EXTRACELLULAR MATRIX (ECM)

Cell interactions with ECM are critical for **development and healing**, as well as for maintaining **normal tissue architecture**.


### Functions of ECM

- **Mechanical support**<sup>Q</sup> for cell **anchorage**, **migration** and cellular **polarity**
- **Control of cell proliferation**, Scaffolding for **tissue renewal**, Establishment of **tissue microenvironments**

### Two Basic forms of ECM

Interstitial matrix & Basement membrane.


Characteristics	Interstitial matrix	Basement membrane
<b>Location</b>	Present in the spaces <b>between cells</b> in connective tissue, and <b>between parenchymal epithelium and supportive</b> vascular and smooth muscle structures	<b>"Chicken wire" mesh (porous)</b> Present <b>between epithelium &amp; mesenchymal cells</b>
<b>Synthesized by</b>	<b>Mesenchymal cells</b> (e.g., <b>fibroblasts</b> )	Overlying <b>epithelium</b> & underlying <b>mesenchymal cells</b>
<b>Major constituents</b>	<b>Fibrillar &amp; non-fibrillar</b> collagens, <b>fibronectin</b> , <b>elastin</b> , <b>proteoglycans</b> and <b>hyaluronate</b>	<b>Amorphous non-fibrillar type IV collagen</b> and <b>laminin</b>



## Components of the Extracellular Matrix

- **Fibrous structural proteins**
  - **Collagens:** confer tensile strength and recoil.
  - **Elastin**
    - Help tissues to **recoil and recover their shape** after physical deformation
    - Found in: **Cardiac valves**, large blood vessels, uterus, skin and ligaments.
    - Consist of a **central core of elastin** with an associated **mesh-like network** composed of **fibrillin**.
- **Proteoglycans:**
  - **Hydrated compressible gels** that confer **resistance to compressive forces**
  - In joint cartilage: provides a layer of **lubrication between adjacent bony surfaces**.
  - **Structure:** Proteoglycans consists of **long polysaccharides** (glycosaminoglycans) eg: Keratin and chondroitin sulphate
  - Action of proteoglycan is shown in Flowchart 3

**Flowchart 3:** Action of proteoglycan



- **Adhesive Glycoproteins and Adhesion Receptors**
  - Involved in **cell-to-cell adhesion**, linking **cells to the ECM**, and the interactions **between ECM components**.
  - Prototypical adhesive glycoproteins include:
    - ◆ **Fibronectin:** A major component of the **interstitial ECM**
    - ◆ **Laminin:** A major constituent of **basement membrane** (most abundant glycoprotein in basement membrane)
    - ◆ **Integrins:** are representative of the **adhesion receptors**, also known as **cell adhesion molecules (CAMs)**

## High Yield Facts



- **Fibrillin** synthetic defects lead to skeletal abnormalities and weakened aortic walls, as in individuals with **Marfan's syndrome**

### Fibrillar Collagen

- **Types I, II, III, and V collagens** form **linear fibrils** stabilized by **interchain hydrogen bonding**.
- **Found in:** bone, tendon, cartilage, blood vessels, and skin, healing wounds and scars.

### Non-Fibrillar Collagens

- **Type IV collagen:** contribute to the structures of planar basement membranes
- **Type IX collagen** in **cartilage:** help regulate **collagen fibril diameters** or collagen-collagen **interactions** via 'Fibril-Associated Collagen with Interrupted Triple helices' (**FACITs**)
- **Type VII collagen:** provide **anchoring fibrils** to basement membrane beneath stratified squamous epithelium

## High Yield Facts



- **Cell Adhesion** molecules include **immunoglobulins, cadherins<sup>a</sup> and selectins**
- **E-Cadherin** gene mutation is associated with **Gastric Carcinoma<sup>a</sup>**
- **Germ-line loss of function mutation** in **E-Cadherin** gene (**CDH1**) is associated with **Autosomal Dominant familial Gastric Carcinoma**
- **E-Cadherin** binds and sequesters **β-catenin** in the **WNT pathway<sup>a</sup>**.
- **Laminin<sup>a</sup>** is the most abundant glycoprotein in basement membrane
- **Collagen I** is most the most abundant collagen in the body.<sup>a</sup>
- **Collagen I** is the main component of the organic part of bone.<sup>a</sup>
- **Collagen II** is most abundant in **Cartilage<sup>a</sup>**.

## STEM CELLS


### Definition

**Undifferentiated cells<sup>a</sup>** capable of giving rise to one or more different types of **specialized cells** and replace **damaged cells** and maintain tissue populations.

### Stem Cells are Characterized by Two Important Properties

- **Self-renewal<sup>a</sup>:** Which permits stem cells to **maintain their numbers**.
- **Asymmetric division<sup>a</sup>:** In which **one daughter cell enters a differentiation** pathway and gives rise to mature cells, while the **other remains undifferentiated and retains its self-renewal capacity**.






## Types of Stem Cells

### Embryonic Stem Cells (ES Cells)

#### Characteristics

- Most **undifferentiated**<sup>o</sup>
- Present in the **inner cell mass**<sup>o</sup> of the blastocyst
- Limitless cell renewal capacity**<sup>o</sup>
- Totipotent**<sup>o</sup>: Can give rise to every cell in the body



ES cells induced under appropriate culture conditions form specialized cells of **all three germ cell layers**<sup>o</sup>, including neurons, cardiac muscle, liver cells, and pancreatic islet cells.

### Tissue Stem Cells (Also Called Adult Stem Cells)

**Undifferentiated cells**, found throughout the body after development that multiply to **replenish dying cells** and **regenerate damaged tissues**.

**Important types of adult stem cells are:**

#### A. Hematopoietic Stem Cells


- Found in **Bone Marrow**<sup>o</sup>
- Replenish all **cellular elements of the blood**
- Isolated directly from **bone marrow** or from the **peripheral blood** after administration of colony stimulating factors (CSF)<sup>o</sup>
- Stem cells **can be used to repopulate marrows** depleted after chemotherapy (e.g., for leukemia),
- Provide **normal precursors** to correct various blood cell defects<sup>o</sup>

#### B. Mesenchymal Stem Cells


- Found in **Bone marrow**<sup>o</sup>
- Multipotent** cells that can differentiate into a variety of stromal cells including chondrocytes (cartilage), osteocytes (bone), adipocytes (fat), and myocytes (muscle)<sup>o</sup>
- Manufactures **stromal cellular scaffolding** for **tissue regeneration**<sup>o</sup>

**C. Stem cell niches:** Microenvironment where **tissue stem cells** are found, which **interacts with stem cells** to **regulate cell fate normally**

Stem Cell Niches	Location
<b>Skin stem cells</b>	Hair follicle bulge <sup>o</sup> , Sebaceous glands <sup>o</sup> , Interfollicular areas <sup>o</sup> of the surface epidermis
<b>Small intestine stem cells</b> <sup>o</sup>	Base of a crypt <sup>o</sup> , above Paneth cells
<b>Liver stem cells (oval cells)</b> <sup>o</sup>	Canals of Hering <sup>o</sup>
<b>Corneal stem cells</b> <sup>o</sup>	Limbus region <sup>o</sup>



Types of stem cells



- **Transdifferentiation<sup>Q</sup>**: Irreversible conversion of cells from one differentiated cell type to another<sup>Q</sup>
- **Developmental plasticity<sup>Q</sup>**: capacity of a cell to transdifferentiate into diverse lineages<sup>Q</sup>
- **Dedifferentiation<sup>Q</sup>**: Reverse developmental process in which differentiated cells with specialized functions become undifferentiated progenitor cells.<sup>Q</sup>
- **Transit amplifying cells**: rapidly dividing cells<sup>Q</sup> generated by somatic stem cells
- **Progenitor cells**: Cells that lose the capacity of self-perpetuation and have **restricted developmental potential<sup>Q</sup>**
- **Induced pluripotent stem cell (iPSC)<sup>Q</sup>**: Adult stem cells (skin or blood cells) that have been reprogramed back into an embryonic-like pluripotent state<sup>Q</sup>
- iPSC can be used to develop an **unlimited source of any type of human cell** needed for therapeutic purposes.
- Shinya Yamanaka and Sir John Gurdon were awarded **Nobel Prize for iPSC in 2012<sup>Q</sup>**



## NEXT Pattern Questions



1. A study of peripheral blood smears shows that neutrophil nuclei of women have a Barr body, whereas those of men do not. The Barr body is an inactivated X chromosome. Which of the following forms of RNA is most likely to play a role in Barr body formation?
- a. lncRNA                      b. mRNA                      c. miRNA                      d. siRNA

### Ans. (a) lncRNA

- There are forms of noncoding RNA that play a role in gene expression. Long noncoding RNA (lncRNA) segments greater than 200 nucleotides in length can bind to chromatin to restrict access of RNA polymerase to coding segments. The X chromosome transcribes XIST, a lncRNA that binds to and represses X chromosome expression. However, not all genes on the “inactive” X chromosome are switched off. The RNA transcribed from nuclear DNA that directs protein synthesis through translation is mRNA. MicroRNAs (miRNAs) are noncoding RNA sequences that inhibit the translation of mRNAs. Gene-silencing RNAs (small interfering RNAs [siRNAs]) have the same function as miRNAs, but they are produced synthetically for experimental purposes. Transfer RNA (tRNA) participates in the translation of mRNA to proteins by linking to specific amino acids.



2. At the site of a surgical incision, endothelial cells elaborate vascular endothelial growth factor. There is sprouting with migration of endothelial cells into the wound to establish new capillaries. Which of the following intracellular proteins is most important in facilitating movement of endothelial cells?
- a. Actin                      b. Cytokeratin  
c. Desmin                      d. Lamin

### Ans. (a) Actin


- Actin is a microfilament involved with cell movement.
- The other possibilities listed in B to D are intermediate filaments, which are larger than actin but smaller than myosin (a thick filament interdigitating with actin, required for muscle movement). Cytokeratins form cytoskeletal elements of epithelial cells. Desmin forms the scaffold in muscle cells on which actin and myosin contract. Lamin is associated with the nuclear membrane.



3. A 62-year-old man has increasing knee pain with movement for the past 10 years. The knee joint surfaces are eroded and the joint space narrowed. There is loss of compressibility and lubrication of articular cartilaginous surfaces. Loss of which of the following extracellular matrix components has most likely occurred in this man?
- a. Elastin                      b. Fibronectin  
c. Hyaluronan                      d. Integrin

### Ans. (c) Hyaluronan

- He has osteoarthritis, or degenerative joint disease, with loss of articular hyaline cartilage. Hyaluronan (hyaluronic acid) is a large mucopolysaccharide, one form of proteoglycan, which forms a hydrated, compressible gel contributing to the shock-absorbing function of joint surfaces. Elastin is a fibrillar protein that provides recoil in tissues such as skin, arterial walls, and ligaments that need to stretch and return to their original shape. Fibronectin is a form of glycoprotein that serves an adhesive function. Integrins are glycoproteins that serve as cellular receptors for extracellular matrix components; they can link to intracellular actin so that cells can alter their shape and mobility




## Multiple Choice Questions

1. Which of the following tumor(s) is/are related to DICER1 Gene mutation? (PGI May 2019)
- Retinoblastoma
  - Pleuropulmonary blastoma
  - Cystic nephroma
  - Thyroid carcinoma
  - Sertoli-Leydig cell tumor

2. Integrin binds to: (AIIMS Nov 18)
- Fibronectin
  - Vitronectin
  - Collagen
  - Laminin

3. The small inner circles in the given image of EM signifies which of the following? (AIIMS May 18)



- Neuro transmitter
  - Neurosecretory granules
  - Collagen fibril
  - Microtubules
4. False about micro satellites is? (JIPMER 18)
- Repeat size more than 10 to 15 nucleotides
  - More prone to variation
  - Found in colonic carcinoma
  - DNA repeats present
5. Type 1 collagen is present in all except? (AIIMS May 18)
- Bone
  - Cartilage
  - Ligament
  - Aponeurosis
6. Which among the following is not seen in disorder to deficiency in elastin production? (AIIMS May 18)
- Aortic dissection
  - Lens subluxation
  - Ligament hyperlaxity
  - bone fracture
7. Which Vitamin increases iron Absorption? (AIIMS May 18)
- Vitamin C
  - Biotin
  - Vitamin B6
  - Vitamin E
8. Which of the following plays a role in gene editing? (AIIMS May 2017)
- Gene Xper
  - CRISPR
  - Health care apps
  - Big data
9. Cell to cell permeability occurs through: (Recent Question 2016-17)
- Occludin
  - Zona adherens
  - Connexins
  - Zonulin
10. Tensile strength of tendon depends on (Recent Question 2016-17)
- Fibrillin
  - Collagen
  - Fibronectin
  - Elastin

11. Which of the following mechanism is mainly involved in Genomic imprinting? (AIIMS Nov 2015)

- Methylation
- Acetylation
- Deamination
- Phosphorylation

12. The term pathology was coined by? (APPGMEE 2015)

- Robert Kochs
- Rudolf Virchow
- Lois Pasteur
- Gregor Mendal

13. Which Collagen is typical of basement membrane? (AIIMS May 2015)

- Type I
- Type V
- Type IV
- Type III

14. Which of the following function is done by RNAi in a gene? (AIIMS May 2015)

- Knock in
- Knock out
- Knock down
- Knock up

15. All of the following are Intermediate filament except? (Recent Question 2016)

- Lamin
- Cadherin
- Vimentin
- Desmin

16. Tight junction consists of all except? (Recent Question 2016)

- Occludin
- Claudin
- Zonulin
- Cadherin

17. Titin protein mutated in? (Recent Question 2016)

- DCM
- HOCM
- RCM
- Non functional cardiomyopathy

18. Peripheral protein in cell membrane are attached by? (Recent Question 2016)

- GpI
- Desmosome
- Catenins
- Cadherins

19. Bridging fibrosis in large wounds is due to (Recent Question 2015)

- Keratinocyte growth factor
- Epidermal growth factor
- Platelet derived growth factor
- Transforming growth factor-β

20. Major cytokine involved in fibrosis (Recent Question 2015)

- Transforming growth factor-α
- Transforming growth factor-β
- Fibroblast growth factor
- Epidermal growth factor

21. Types of collage playing important role in wound healing (Recent Question 2015)

- I and III
- II and V
- III and IV
- V and IX

22. Oval stem cells are located in (Recent Question 2015)


- Canal of schlemm
- Canal of herring
- Space of disse
- Basallamina of myotubules

23. Stem cells are present in? (Recent Question 2015)

- Cornea
- Base of crypts
- Bile duct of liver
- Mesonephros


24. Human genome contains (Recent Question 2015)

- 3.2 billion DNA base pairs
- 2.3 billion DNA base pairs
- 3.2 million DNA base pairs
- 2.3 million DNA base pairs



25. **Protein-encoding genes comprises what percentage of genome** (Recent Question 2015)
  - a. 98.5%
  - b. 1.5%
  - c. 5%
  - d. 95%
26. **Jumping genes take part in:** (Recent Question 2015)
  - a. Gene regulation
  - b. Chromosomal aberrations
  - c. Gene movement in a species
  - d. Carry out gene amplification
27. **The most-common form of DNA variation is?** (Recent Question 2015)
  - a. Single nucleotide polymorphism
  - b. Copy Number Variations (CNVs)
  - c. Transposons
  - d. Mutations
28. **Linkage disequilibrium refers to?** (Recent Question 2015)
  - a. Never co-inherited gene
  - b. Inherited but not with disease causing gene
  - c. Genetic mutations
  - d. Co-inherited with the disease causing gene
29. **Heritable changes in gene expression not caused by alterations in DNA sequence refers to?** (Recent Question 2015)
  - a. Genetics
  - b. Epigenetics
  - c. Mutations
  - d. Transposons
30. **Epigenetic factors refers to?** (Recent Question 2015)
  - a. Histone methylation
  - b. Histone phosphorylation
  - c. DNA methylation
  - d. All of the above
31. **Function of Peroxisomes is?** (Recent Question 2014)
  - a. Protein synthesis
  - b. Carbohydrate metabolism
  - c. Generating  $H_2O_2$
  - d. DNA replication
32. **Collagen present in cornea?** (Recent Question 2015)
  - a. Type 1
  - b. Type 2
  - c. Type 4
  - d. Type 7
33. **Gene silencing RNA-** (Recent Question 2014)
  - a. rRNA
  - b. tRNA
  - c. miRNA
  - d. None
34. **Epigenetics deals with genetic modification that do not alter the sequence of DNA. All of the following can detect epigenetic modifications except?** (AIIMS Nov 14)
  - a. Chip on chip
  - b. Bisulphite method
  - c. HPLC
  - d. Methylation specific PCR
35. **Coding DNA constitutes what proportion of total DNA?** (AIIMS May 2014)
  - a. 0.02
  - b. 0.25
  - c. 0.4
  - d. 0.1
36. **GWAS stands for:** (PGI Nov 2014)
  - a. Genome wide Association syndrome
  - b. Genome wide Association studies
  - c. Genetic wide array studies
  - d. Genetic wide amplification studies
  - e. Genomic way of association studies
37. **Genetic polymorphisms include -** (PGI Nov 10)
  - a. SNP
  - b. Microsatellites
  - c. Mutations
  - d. Translocation
  - e. Mini satellites
38. **Function of miRNA is/are:** (PGI May 12)
  - a. Gene silencing
  - b. Gene activation
  - c. Transcription inhibition
  - d. Translation repression
  - e. Breaking of messenger RNA
39. **Most abundant collagen in the body?** (Recent Question 2015)
  - a. Type I
  - b. Type II
  - c. Type III
  - d. Type IV
40. **FACIT collagen is?** (Recent Question 2015)
  - a. Type I
  - b. Type III
  - c. Type IX
  - d. Type XI
41. **Gap junctions comprise of?** (Recent Question 2015)
  - a. Catenins
  - b. Cadherins
  - c. Connexins
  - d. Claudin
42. **All of the following are true about TGF- $\beta$  except:** (Recent Question 2014)
  - a. Anti-inflammatory
  - b. Causes fibrosis
  - c. Tumor suppressor gene
  - d. Anti-Angiogenic
43. **Which of the following statement is not true?** (Recent Question 2013)
  - a. Cytokeratin is the marker of muscle cells
  - b. Vimentin is used to stain fibroblasts
  - c. Neurofilaments are present in neurons
  - d. Desmin can be used as a marker for muscle cells
44. **Which of the following is both morphogenic and mitogenic?** (AIIMS May 2014)
  - a. Insulin growth factor
  - b. Bone morphogenic factor
  - c. Fibroblast growth factor
  - d. Epidermal growth factor
45. **The function of Proteoglycans and Hyaluronan is ?**
  - a. Lubrication and reduces resistance
  - b. Cell growth
  - c. Connect ECM components
  - d. Provides tensile strength
46. **Mutation in COL4A5 chain leads to?** (AIIMS May 2013)
  - a. Alport's syndrome
  - b. Good pasture's syndrome
  - c. Hereditary Non-polypsis Colon Cancer
  - d. Xeroderma Pigmentosum
47. **Oncogene tyrosine kinase involves:** (PGI May 2012)
  - a. PML-BRCA 1
  - b. BCR-ABL
  - c. HAAJ
  - d. JUN
  - e. NJKK
48. **Epidermal growth factor is/are formed by:** (PGI May 12)
  - a. Platelet
  - b. Fibroblast
  - c. Mast cell
  - d. Endothelial cell
  - e. Keratinocyte
49. **Which of the following statements is not correct regarding stem cell?** (DPG 11)
  - a. Developmental elasticity
  - b. Transdifferentiation
  - c. Can be harvested from embryo
  - d. Knock out mice made possible because of it
50. **Which of the following is involved in stem cell self-renewal?** (PGI May 2014)
  - a. Oct 3/4
  - b. sox 2
  - c. c-myc
  - d. FLT3 ligand
  - e. c-kit





**51. Which of the following statements about hematopoietic stem cell is false? (AP PGME 2014)**

- Stem cells have self-renewal property
- Subset of stem cells normally circulate in peripheral blood
- Marrow derived stem cells can seed other tissues and develop into non-hematopoietic cells as well
- Stem cells resemble lymphoblasts morphologically

**52. In an ablated animal, If Myeloid Stem Cells are injected, which type of cells are induced after the incubation period? (AIIMS May 12)**

- T-Lymphocyte
- Erythroid
- Fibroblast
- Hematopoietic Stem Cells

**53. Location of small intestine stem cells is? (Recent Question 12)**

- Jejunum epithelial cells
- Terminal ileum mucosa
- Crypt base
- Mucosa throughout

**54. True about stem cell (PGI Nov 12)**

- Undifferentiated
- Pluripotent
- Dedicated to one cell line
- Used for cell repair and tissue regeneration
- Can form any cell



## Answers with Explanations

**1. Ans. (b) Pleuropulmonary blastoma; (c) Cystic nephroma; (d) Thyroid carcinoma; (e) Sertoli-Leydig cell tumor (Ref: Robbins 9th/pg 5; pg 1033)**

Dicer is a type III cytoplasmic endoribonuclease that is involved in the maturation of several classes of small non-coding RNAs, such as microRNAs. Germline loss-of-function mutations in DICER1 are associated with pleuropulmonary blastoma (PPB), ovarian sex cord-stromal tumors, ciliary body medulloepitheliomas, nasal chondromesenchymal hamartomas, multinodular goiter and differentiated thyroid carcinomas, cystic nephroma and, more rarely, anaplastic sarcoma of kidney. DICER1 mutations have also been documented in pediatric tumors including Wilms' tumor and in pituitary blastoma (PitB).

**2. Ans. (d) Laminin > Fibronectin**

Integrins localised in the plasma membrane are the major adhesion receptors connecting cells with components of the extracellular matrix. Integrins interact directly with laminin, fibronectin present in the basal lamina and intracellularly contact actin through intermediate proteins, such as alpha-actinin, vinculin, and talin

**3. Ans. (b) Neurosecretory granules**

Neurosecretion is the storage, synthesis and release of hormones from neurons. These neurohormones, produced by neurosecretory cells, are normally secreted from nerve cells in the brain that then circulate into the blood.

**4. Ans. (a) Repeat size more than 10 to 15 nucleotides**

**5. Ans. (b) Cartilage**

Please Refer to Explanation of Q.21

**6. Ans. (d) Bone fracture**

**7. Ans. (a) Vitamin C**

**8. Ans. (b) CRISPR (Ref: R 9th/p 5-6)**

CRISPR-Cas9 is a genome editing tool essential in adaptive immunity in select bacteria enabling the organisms to respond to and eliminate invading genetic material

**9. Ans. (c) Connexins (Ref: Robbins 9th/ pg 11)**

- Communicating junctions (Gap junctions): mediate the passage of chemical or electrical signals from one cell to another.
- Consists of pores called connexions and formed by hexamers of transmembrane proteins called connexins

**10. Ans. (b) Collagen (Ref: Robbins 9th/ pg 12-13)**

Fibrous structural proteins like Collagens confer tensile strength and & elastins provide recoil to the tension

**11. Ans. (a) Methylation (Ref: Robbins 9th/pg 3-4)**

- Genomic imprinting selectively inactivates either the maternal or paternal allele.
- Occurs in the ovum or the sperm, before fertilization, and then is stably transmitted to all somatic cells through mitosis


### Mechanisms

- Histones and histone modifying factors**
  - Histone methylation
    - Histone acetylation
    - Histone phosphorylation
- DNA methylation**
- Chromatin organizing factors**

**12. Ans. (b) Rudolf Virchow (Ref: Robbins 9th/pg 1)**

Pathology (from the Ancient Greek roots of pathos, meaning "experience" or "suffering", and -logia, "an account of") is a significant component of the causal study of disease and a major field in modern medicine and diagnosis. The term cellular pathology was coined by Rudolf Virchow.





13. Ans. (c) **Type IV** (Ref: Robbins 9th/pg 20-23)

14. Ans. (c) **Knock down** (Ref: Robbins 9th/pg 3-4)

**Small interfering RNAs (siRNAs)**

- Short RNA sequences **introduced experimentally** into cells
- Their action is similar to **endogenous miRNAs**.
- Synthetic **siRNAs targeted against specific mRNA** have become useful laboratory tools to study gene function (so-called “**knockdown technology**”)

15. Ans. (b) **Cadherin** (Ref: Robbins 9th/pg 21-24)

*Intermediate filaments are 10-nm diameter fibrils which provide tensile strength to a cell.*

*Cadherins are cell-cell desmosomal junctions are formed by homotypic association of transmembrane glycoproteins.*

16. Ans. (d) **Cadherin** (Ref: Robbins 9th/pg 21-24)

**Occluding Junctions (Tight Junctions)**

- Seal adjacent cells together to create a **continuous barrier** that **restricts** the **paracellular (between cells) movement of ions and other molecules**.
- Proteins involved are **Occludin, Claudin, Zonulin and Catenin**

17. Ans. (a) **DCM** (Ref: Robbins 9th/pg 21-24)

Mutations in TTN, a gene on chr 2q31 that encodes titin (so-called because it is the largest protein expressed in humans) causes 20% of all cases of Dilated Cardiomyopathy (DCM)

18. Ans. (a) **GPI** (Ref: Robbins 9th/pg 21-24)

*Proteins are linked to cell via glycoposphatidylinositol (GPI) structures.*

19. Ans. (d) **Transforming growth factor-β** (Ref: R 9th/pg 20)

**Functions of TGFβ are:**

- Anti-inflammatory, fibrosis, tumor suppressor gene, angiogenesis

20. Ans. (b) **Transforming growth factor-β** (Ref: R 9th/pg 20)

21. Ans. (a) **I and III** (Ref: Robbins 9th/pg 20-23)

**Major types of Collagens & disorders associated with them:**

Type	Present in	Disorders
I	Bone <sup>q</sup> , Cornea, <sup>q</sup> Scar tissue, tendons, skin, artery walls, endomysium of myofibrils <sup>q</sup> , fibrocartilage, teeth	Osteogenesis-imperfecta <sup>q</sup> , Ehlers-Danlos Syndrome <sup>q</sup> , Caffey's disease <sup>q</sup>
II	Hyaline cartilage <sup>q</sup> , Vitreous humor <sup>q</sup> of the eye.	Collagenopathy

Type	Present in	Disorders
III	Granulation tissue <sup>q</sup> , Reticular fiber, artery walls, skin, intestines and the uterus	Ehlers-Danlos Syndrome <sup>q</sup>
IV	Basement membrane <sup>q</sup> , eye lens <sup>q</sup> , Glomerular Basement membrane	Alport syndrome <sup>q</sup> (COL4α5) Goodpasture's syndrome <sup>q</sup> (COL4α3)

22. Ans. (b) **Canal of herring** (Ref: Robbins 9th/ pg 27-28)

*Liver stem cells (oval cells)<sup>q</sup> are present in canals of Hering<sup>q</sup>*

23. Ans. (b) **Base of crypts** (Ref: Robbins 9th/pg 27-28)

24. Ans. (a) **3.2 billion DNA base pairs** (Ref: Robbins 9th/ pg 1)

25. Ans. (b) **1.5%** (Ref: Robbins 9th/ pg 1)

In humans, there are 20,000 protein-encoding genes<sup>q</sup>, comprising only 1.5% of the genome.<sup>q</sup>

26. Ans. (a) **Gene regulation** (Ref: Robbins 9th/ pg 2)

**Jumping genes**

- They are **Mobile genetic elements**, also called ‘**transposons**’
- Segments that **move around the genome**, exhibiting **wide variation** in number and positioning;
- Role in **gene regulation** and **chromatin organization**

27. Ans. (a) **Single nucleotide polymorphism** (Ref: R 9th/ pg 3)

**Single-Nucleotide Polymorphisms (SNPs)**

- Variations at **single nucleotide positions**<sup>q</sup>
- **Most prevalent**<sup>q</sup> and **important form of genetic variation**.
- Always **bi-allelic**<sup>q</sup>
- Can occur in **exons, introns, intergenic regions and coding regions**
- **1% of SNPs occur in coding regions**

28. Ans. (d) **Co-inherited with the disease causing gene**

(Ref: Robbins 9th/ pg 3)

Even if any SNP has no effect on gene function (Neutral SNP)<sup>q</sup>, it may be co-inherited with the “actual disease causing gene,” if located close to that gene. This is called “Linkage disequilibrium”<sup>q</sup>

29. Ans. (b) **Epigenetics** (Ref: Robbins 9th/ pg 3-4)

Epigenetics refers to heritable changes in gene expression, not caused by alterations in DNA sequence.

Epigenetic factors include:

- Histones and **histone modifying factors** like Histone methylation, acetylation and phosphorylation
- **DNA methylation**
- **Chromatin organizing factors**

30. Ans. (d) **All of the above** (Ref: Robbins 9th/ pg 3-4)

Contd...



31. Ans. (c) **Generating  $H_2O_2$**  (Ref: Robbins 9th/ pg 6-7)

Functions of some important cell organelles:

- **Lysosomes:**
- **Digestion<sup>Q</sup> of proteins, polysaccharides, lipids, nucleic acids**
- **Proteasomes:**
  - Selectively **chews up denatured proteins, releasing peptides<sup>Q</sup>**
- **Peroxisomes:**
  - **Breakdown of fatty acids,<sup>Q</sup> generating hydrogen peroxide**

32. Ans. (a) **Type I** (Ref: Robbins 9th/ pg 20-23)

33. Ans. (c) **miRNA** (Ref: Robbins 9th/ pg 4-5)

34. Ans. (c) **HPLC** (Ref: Robbins 9th/ pg 3-4)

35. Ans. (a) **0.02** (Ref: Robbins 9th/ pg 1)

Coding DNA constitutes 0.02 or 1.5-2% of total DNA; Refer Ans 3 above

36. Ans. (b) **Genome wide association studies**

(Ref: Robbins 9th/ pg 179)

GWAS refers to 'Genome Wide Association Studies'<sup>Q</sup>

In GWAS, entire genome of large number of individuals with and without a disease are examined for common genetic variants or polymorphisms that are overrepresented in patients with the disease.

37. Ans. (a, b, e) **a. SNP; b. Microsatellites; e. Mini satellites**

(Ref: Robbins 9th/ pg 3)

- The 2 types of genetic polymorphisms are **single nucleotide polymorphisms (SNPs) & repeat length polymorphisms**
- Repeat length polymorphisms can be **mini-satellite (1-3 kb)<sup>Q</sup> or Micro-satellite repeats (<1 kb)<sup>Q</sup>**

38. Ans. (a, d, e) **a. Gene silencing; d. Translation repression; e. Breaking of messenger RNA**

(Ref: Robbins 9th/ pg 4-5)

39. Ans. (b) **Type II** (Ref: Robbins 9th/ pg 20-23)

- **Collagen I is most the most abundant collagen in the body.<sup>Q</sup>**
- **Collagen I is the main component of the organic part of bone.<sup>Q</sup>**
- **Collagen II is most abundant in Cartilage.<sup>Q</sup>**

40. Ans. (c) **Type IX** (Ref: Robbins 9th/ pg 20-23)

Type IX collagen in cartilage: help regulate collagen fibril diameters or collagen-collagen interactions via 'Fibril-Associated Collagen with Interrupted Triple helices' (FACITs).

41. Ans. (c) **Connexins** (Ref: Robbins 9th/ pg 11)

Communicating junctions (Gap junctions)

- Mediate the passage of **chemical or electrical signals** from **one cell to another**.
- Consists of a dense planar array of 1.5-2-nm **pores** called **connexons<sup>Q</sup>**
- Formed by hexamers of transmembrane **proteins** called **connexins<sup>Q</sup>**

42. Ans. (d) **Anti-Angiogenic** (Ref: Robbins 9th/ pg 20)

43. Ans. (a) **Cytokeratin is the marker of muscle cells**

(Ref: Robbins 9th/ pg 11)

44. Ans. (b, a) **b. Bone morphogenic factor > a. Insulin like Growth Factor**

(Ref: Robbins 9th/ pg 19-20; International journal of molecular medicine 2007;20: 53-57)

Discussing the options one by one:

<b>A. Insulin growth factor (IGF)</b>	<ul style="list-style-type: none"> <li>• IGF-I is mitogenic for <b>growth plate chondrocytes</b> and stimulates chondrocyte synthesis of matrix macromolecules, including proteoglycan and collagen.</li> <li>• Causes <b>skeletal growth and development</b> by acting on epiphyseal growth plate</li> </ul>
<b>B. Bone morphogenic factor or protein (BMP)</b>	<ul style="list-style-type: none"> <li>• Act as <b>chemotactic, mitogenic</b> and help in <b>differentiating mechanisms</b></li> <li>• During embryogenesis, BMPs <b>regulate dorsal-ventral patterning</b>, establishment of embryonic <b>body plan</b>, <b>cell apoptosis</b>, <b>differentiation of neural cells</b>, <b>patterning of the limb bud</b> and <b>epithelial-mesenchymal interactions</b> during organogenesis (morphogenesis)</li> </ul>
<b>C. Fibroblast growth factor (FGF)</b>	<ul style="list-style-type: none"> <li>• <b>Chemotactic and Mitogenic</b> for and <b>fibroblasts</b>;</li> <li>• Stimulates <b>angiogenesis</b> and ECM protein synthesis</li> </ul>
<b>D. Epidermal growth factor (EGF)</b>	<ul style="list-style-type: none"> <li>• <b>Mitogenic</b> for hepatocytes and fibroblasts;</li> <li>• Stimulates <b>granulation tissue formation</b></li> </ul>

45. Ans. (a) **Lubrication and reduces resistance**

(Ref: Robbins 9th/ pg 20-24)

Function of proteoglycans and hyaluronan is to provide a layer of lubrication between adjacent bony surfaces and reduces friction & resistance.



46. Ans. (a) **Alport's syndrome** (Ref: Robbins 9th/ pg 20-23)

47. Ans. (b, d) **b. BCR-ABL; d. JUN** (Ref: Robbins 9th/pg 16-17)

48. Ans. (e) **Keratinocyte** (Ref: Robbins 9th/pg 19)

EGF and TGF- $\alpha$  are produced by macrophages and a variety of epithelial cells, including keratinocytes and are mitogenic for hepatocytes, fibroblasts, and a host of epithelial cells.

49. Ans. (a) **Developmental elasticity**

(Ref: Robbins 9th/pg 27-28)

Stem cells show developmental plasticity (transdifferentiation) and not developmental elasticity;

**Transdifferentiation** indicates a change in the lineage commitment of a stem cell.

50. Ans. (a, b, d, e) **a. Oct 3/4; b. sox 2; d. FLT3 ligand; e. c-kit**

(Ref: Robbins 9th/ pg 27-28, 580-581)

Discussing the options one by one:

<b>A. Oct 3/ pg4 (octamer-binding transcription factor)</b>	<ul style="list-style-type: none"> <li>Also known as <b>POU5F1</b>;</li> <li>It is critically involved in the <b>self-renewal</b> of undifferentiated embryonic stem cells.</li> <li>It is frequently <b>used as a marker for undifferentiated cells</b>.</li> </ul>
<b>B. Sox 2</b>	SRY (sex determining region Y)-box 2, also known as SOX2, is a transcription factor that is essential for maintaining self-renewal or pluripotency of undifferentiated embryonic stem cells
<b>C. C myc</b>	Proto-oncogene involved in Burkitt lymphoma; Not involved in stem cell self-renewal;
<b>D. FLT3 ligand</b>	Flt3 ligand (FL) is a hematopoietic cytokine. It is structurally homologous to stem cell factor (SCF) and colony stimulating factor 1 (CSF-1). Flt3 ligand <b>stimulates proliferation &amp; differentiation of various blood cell progenitors</b> .
<b>E. C-kit</b>	<ul style="list-style-type: none"> <li>Mast/stem cell growth factor receptor (SCFR), also known as CD117</li> <li>Signalling through CD117 plays a role in <b>cell survival, proliferation, and differentiation</b>.</li> <li>It binds to a substance called stem cell factor (SCF), which causes <b>stem cell proliferation</b></li> </ul>

51. Ans. (b) **Subset of stem cells normally circulate in peripheral blood** (Ref: Robbins 9th/ pg 27-28)

Discussing the options one by one:

A.	True	<b>Self-renewal takes place by asymmetric division</b> , which permits stem cells to <b>maintain their numbers</b>
B.	False	Hematopoietic stem cells <b>can be found in peripheral blood</b> after administration of colony stimulating factors (CSF) <sup>a</sup>
C.	True	<b>Mesenchymal stem cells</b> found in <b>Bone marrow</b> are <b>multipotent</b> cells that can differentiate into a variety of stromal cells including chondrocytes (cartilage), osteocytes (bone), adipocytes (fat), and myocytes (muscle).
D.	True	Stem cells resemble lymphocytes or lymphoblasts morphologically

52. Ans. (b) **Erythroid** (Ref: Robbins 9th/ pg 27-28, 580-581)

Neutrophils, Monocytes, Basophils, Erythroids and Megakaryocytes are myeloid in origin while lymphocytes are Lymphoid in origin. Hematopoietic Stem Cells give rise to myeloid stem cells and not the reverse.

53. Ans. (c) **Crypts base** (Ref: 9th/ pg 27-28)

54. Ans. (a, b, d, e) **a. Undifferentiated; b. Pluripotent; d. Used for cell repair and tissue regeneration; e. Can form any cell** (Ref: Robbins 9th/ pg 27-28)

Discussing the options one by one:

- Undifferentiated → True**
- Pluripotent → True; embryonic stem cells or ES cells, are pluripotent, that is, they can generate all tissues of the body**
- Dedicated to one cell line False, stem cells are pluripotent or multipotent and can give rise to multiple cell lines
- Used for cell repair and tissue regeneration True; Mesenchymal stem cells manufacture stromal cellular scaffolding for tissue regeneration**
- Can form any cell → True; Embryonic (pluripotent) stem cells can generate all tissues of the body. Pluripotent stem cells give rise to multipotent stem cells, which have more restricted developmental potential, and eventually produce differentiated cells from the three embryonic layers.**

# Cell Adaptation, Injury and Death

## Key Points

- » **Adaptations** are **reversible changes** in response to changes in environment
- » The **adaptive responses** may consist of **Hypertrophy, Hyperplasia, Atrophy and Metaplasia**
- » **Hypoxia is the most common cause** of cell injury and cell death
- » **The earliest change of reversible cell injury is cellular swelling**
- » **Membrane damage is a central factor** in pathogenesis of irreversible injury
- » **The most common type of necrosis: Coagulative necrosis**
- » The **mitochondrial pathway** is the **major mechanism** of apoptosis in all mammalian cells
- » Apoptosis in contrast to necrosis **does not elicit inflammation**
- » Calcification **begins** in **mitochondria**
- » **Most common site of metastatic calcification is lungs followed by kidneys**
- » **Sirtuins have a role in longevity, Diabetes and Aging**

## Key Recent Updates

- » Marker of Autophagy–**LC3**
- » **Beclin I** initiates autophagy
- »  $\alpha_1$  antitrypsin deficiency is disease caused by misfolded protein resulting from ER stress induced cell loss and functional deficiency of protein.







## CELL ADAPTATIONS

### Hypertrophy


- **Increase in the size of cells**<sup>Q</sup>
- **Seen in:** Nondividing cells.<sup>Q</sup>
- **Stimulus:** Increased workload<sup>Q</sup> (Most common)
- **Mechanism:** Increased production of cellular proteins.<sup>Q</sup>
- **Examples**
  - Uterus during pregnancy.<sup>Q</sup>
  - Bulging muscles of body builders.
  - Hypertension or faulty valves → heart hypertrophy
  - Breast in lactation<sup>Q</sup>
- **Outcome**
  - Reversible process; Does not predispose to malignancy<sup>Q</sup>




Normal uterus



Hypertrophied uterus



Normal muscle



Hypertrophied smooth muscle

### Hyperplasia

- Increase in **number of cells** in an organ or tissue in response to a stimulus.<sup>Q</sup>
- **Seen in:** Dividing cells.<sup>Q</sup>
- **Mechanism:**
- **Growth factor-driven proliferation**<sup>Q</sup> of mature cells
- Increased output of new cells from tissue stem cells (rare)
- **Types with examples**
  - **Physiologic hyperplasia:**
    - Action of **hormones or growth factors** in hormone responsive organs
  - Breast in pregnancy and during puberty
  - **Bone marrow**<sup>Q</sup> in response to deficiency of terminally differentiated blood cells.


- Compensatory increase after damage or resection, e.g. **Liver regeneration after partial hepatectomy**<sup>Q</sup>

#### ○ Pathologic hyperplasia


Excessive action of hormones or growth factors acting on target cells.

- **Endometrial hyperplasia**
- **Benign prostatic hyperplasia**
- **Skin warts**-by viral infections such as HPV<sup>Q</sup>

- **Outcome:** Constitutes a fertile soil in which **cancerous proliferations** may eventually arise



Normal endometrium



↑number of glands

Endometrial hyperplasia

R10<sup>th</sup>

Latest Update

- Combined hypertrophy + hyperplasia: uterus in pregnancy<sup>Q</sup>
- Hypertrophy: Breast in lactation<sup>Q</sup>
- Atrophy: Breast at menopause<sup>Q</sup>


### Atrophy

- Reduction in the size of an organ or tissue due to a **decrease in cell size and number**<sup>Q</sup>
- **Mechanism:** Decreased protein synthesis<sup>Q</sup> and increased protein degradation in cells.
- **Types with examples**
  - **Physiologic atrophy:**
    - During **normal development** e.g **notochord and thyroglossal duct**<sup>Q</sup>
    - Decrease in size of **uterus after parturition**<sup>Q</sup>
  - **Pathologic atrophy**
    - **Disuse** atrophy: Most common due to decreased workload






- **Denervation atrophy:** Due to loss of innervation
- **Senile atrophy:** Due to diminished **blood supply**; seen in **aging brain**<sup>Q</sup>
- **Pressure atrophy:** Atrophy of surrounding uninvolved tissues due to an enlarging benign tumor
- **Brown atrophy-lipofuscin granules** seen → **brown discoloration of tissue**
- **Inadequate nutrition:** protein-calorie malnutrition
- **Loss of endocrine stimulation: menopause** → physiologic **atrophy** of the **endometrium**, vaginal epithelium & **breast**




## Metaplasia

- **One differentiated cell type (epithelial or mesenchymal) is replaced by another cell type; Reversible**



- **Mechanism: Reprogramming of stem cells or undifferentiated mesenchymal cells**<sup>Q</sup> in normal tissues
- **Outcome:** If persistent, can initiate **malignant transformation** in metaplastic epithelium






## CELL INJURY



- **Hypoxia** is the most common cause of cell injury
- **Ischemia** is the most common cause of hypoxia

## Reversible Injury

- **Cellular swelling (Hydropic Change/vacuolar degeneration):** Microscopic examination may show small, clear vacuoles within the cytoplasm; these represent distended and pinched-off **segments of the endoplasmic reticulum (ER)**
- Features of **reversible cell injury** seen in **light microscopy** are **cellular swelling and fatty change**.<sup>Q</sup>
- **The earliest change of reversible cell injury- cellular swelling**<sup>Q</sup>
- **The earliest morphological change** in reversible cell injury<sup>Q</sup> occur due to **accumulation of water intracellularly**<sup>Q</sup>
- Other features are demonstrated in the flow diagram below
- **Nuclear alterations** in reversible injury include **disaggregation of granular and fibrillar elements**



Light microscope findings showing hydropic change/cellular swelling in PCT of kidney (Reversible cell injury)



Loss of microvilli<sup>Q</sup>  
**Reversible cell injury:** A. Electron microscopy of normal Cell.  
B. electron microscopy of reversible cell injury

\* mv-microvilli



## POINTS TO REMEMBER

- **Myelin figures** are rolled-up or are present in scroll-like arrangement of **lipid bilayer** within a cell
- Myelin figures are derived from damaged membranes of organelles and plasma membrane<sup>Q</sup>
- Myelin figures can be seen both in reversible and irreversible cell injury

## Irreversible Cell Injury

- **Membrane damage is** the central factor for irreversible injury
- Two phenomena that **consistently characterize irreversibility** are **irreversible mitochondrial dysfunction** and **profound disturbances in membrane function**
- Electron microscopy features of irreversible injury are: large amorphous densities in mitochondria<sup>Q</sup> and intracytoplasmic myelin figures<sup>Q</sup>






### Features of Irreversible Injury


- Severe swelling of mitochondria<sup>Q</sup> and large, flocculant, amorphous densities<sup>Q</sup> due to increased Ca<sup>2+</sup> influx<sup>Q</sup>
- Severe swelling of lysosomes<sup>Q</sup>: Injury to lysosomal membrane with release of enzymes leading to:
  - Decreased basophilia, Increased eosinophilia<sup>Q</sup>
  - Nuclear changes (Pyknosis, karyolysis, Karyorrhexis)<sup>Q</sup>
  - Protein digestion<sup>Q</sup>
  - Severe damage to plasma membrane<sup>Q</sup>

### Stains

- Stain used in histopathology is **Hematoxylin and Eosin**.
- Hematoxylin stains—DNA & RNA of nucleus.
- **Eosin**: Stains cytoplasmic process.
- Necrotic/dying cells appear more pink as
  - Eosin binds strongly to denatured proteins
  - Loss of DNA & RNA causes loss of blue color



Mic: Coagulative necrosis kidney



## CELL DEATH









### POINTS TO REMEMBER


- Cell death can occur by apoptosis or necrosis



Cell death	Apoptosis	Necrosis
Cell size	Reduced	Enlarged
Membrane	Intact	Disrupted
Inflammation ( <b>Most Important</b> )	Absent	Present
Energy	Active	Passive

### Necrosis

- Necrosis is a form of cell death in which cellular membranes fall apart, and cellular enzymes leak out and ultimately digest the cell
- Necrotic cells have a **glassy, homogeneous appearance** due to loss of lighter staining glycogen particles
- The cytoplasm of necrotic cells becomes vacuolated and appears **"moth-eaten"** due to enzymatic digestion of cytoplasmic organelles

Types/Patterns	Characteristics	Images	
<b>Coagulative necrosis</b> (E.g. Infarction of all solid organs except CNS)	<ul style="list-style-type: none"> <li><b>Most common type<sup>a</sup></b></li> <li><b>Architecture</b> of dead tissues is <b>preserved</b></li> <li><b>Mechanism:</b> Protein denaturation</li> </ul>	 <p>Gross image of kidney showing coagulative necrosis</p>	 <p>Loss of nuclei pink cytoplasm</p> <p>Microscopic finding showing coagulative necrosis (preserved cell outlines with loss of nuclei)</p>
<b>Liquefactive necrosis</b>	<ul style="list-style-type: none"> <li><b>Digestion</b> of the dead cells, resulting in <b>transformation of tissue into a liquid</b> viscous mass</li> <li>Focal bacterial or, occasionally, fungal <b>infections</b> are seen; E.g. Necrosis in CNS<sup>a</sup></li> <li><b>Mechanism:</b> Enzymatic action.</li> </ul>	 <p>Gross showing liquefactive necrosis</p> <p>Gross: Brain showing liquefactive necrosis</p>	 <p>Liquefactive necrosis in the brain</p> <p>Mic: Image shows dissolution of tissue</p>
<b>Gangrenous necrosis</b>	<ul style="list-style-type: none"> <li>Commonly used term in clinical practice, but <b>not a specific pattern of cell death<sup>a</sup></b></li> <li><b>Wet</b> gangrene (type of Liquefactive necrosis) and <b>Dry</b> gangrene (type of Coagulative necrosis)</li> </ul>	 <p>Well demarcation between normal and abnormal</p> <p>Dry gangrene</p>	 <p>Intestine</p> <p>Wet gangrene</p>
<b>Caseous necrosis (CN)</b>	<ul style="list-style-type: none"> <li>Characterised by <b>granuloma</b>-collection of fragmented or lysed cells &amp; <b>amorphous granular debris<sup>a</sup></b> enclosed within a distinctive inflammatory border.</li> <li>E.g. <b>Tuberculous infection<sup>a</sup></b> fungi-histoplasma<sup>a</sup>, coccidiomycosis<sup>a</sup></li> </ul>	 <p>Gross-lung showing CN (red arrow)</p>	 <p>Microscopy-showing CN (green arrow) along with Langhans giant cell (red arrow)</p>



Types/ Patterns	Characteristics	Images
<b>Fat necrosis</b>	<ul style="list-style-type: none"> <li>• Focal areas of fat destruction; Not a specific pattern of necrosis.<sup>Q</sup></li> <li>• Lipases → split triglyceride esters → fatty acids → combine with calcium to produce grossly visible chalky-white areas (fat saponification)<sup>Q</sup></li> <li>• E.g. Acute pancreatitis, Injury to breast, Abdomen, Buttocks<sup>Q</sup></li> </ul>	 <p>Microscopy-showing chalky white deposits (arrow)</p>
<b>Fibrinoid necrosis</b>	<ul style="list-style-type: none"> <li>• Due to immune reactions involving blood vessels</li> <li>• Deposits of “antigen –antibody complexes”<sup>Q</sup> and fibrin that has leaked out of vessels, result in a bright pink and amorphous appearance in H &amp; E stains.<sup>Q</sup></li> <li>• Seen in PAN, malignant hypertension, acute rheumatic fever, Libman sacks endocarditis<sup>Q</sup></li> </ul>	 <p>Microscopy-wall of artery showing fibrinoid necrosis (arrow)</p>



### High Yield Facts

- Coagulative necrosis: The most common type of necrosis<sup>Q</sup>: occurs due to degenerated cytoplasmic proteins<sup>Q</sup>
- A localized area of coagulative necrosis is called an infarct.<sup>Q</sup>
- Caseous necrosis is most often seen in tuberculous infection<sup>Q</sup>
- Cause of caseous necrosis-mycolic acid<sup>Q</sup>
- Zenker's degeneration is a true necrosis<sup>Q</sup> (coagulative necrosis)<sup>Q</sup> affecting muscles (skeletal > cardiac)<sup>Q</sup>, during acute infections (especially typhoid<sup>Q</sup>).
- In Zenker's degeneration, Rectus and diaphragm are the most common muscles affected<sup>Q</sup>

### Apoptosis


- Pathway of cell death induced by a tightly regulated suicide program<sup>Q</sup> leading to activation of intrinsic enzymes that degrade the cell's own DNA and proteins.<sup>Q</sup>
- Important characteristics
  - No loss of membrane integrity<sup>Q</sup>
  - No leakage of cellular contents<sup>Q</sup>
  - No host reaction<sup>Q</sup>
- Examples:
  - Physiologic Situations
    - Embryogenesis<sup>Q</sup>
    - Hormone-dependent tissues upon hormone withdrawal

- Cell loss in proliferating cell populations<sup>Q</sup>
- Epithelial cells- in intestinal crypts, as to maintain a constant number (homeostasis).
- Elimination of potentially harmful self-reactive lymphocytes<sup>Q</sup>
- Death of host cells that have served their useful purpose- neutrophils in an acute inflammatory response<sup>Q</sup>
- Pathological situations
  - DNA damage- radiational injuries<sup>Q</sup>
  - Accumulation of misfolded proteins<sup>Q</sup>
  - Cell death in certain viral infections- E.g. councilman bodies in hepatitis B virus<sup>Q</sup>
  - Atrophy in parenchymal organs after duct obstruction. E.g. pancreas, parotids and kidneys<sup>Q</sup>
- Mechanisms:
  - Two pathways: Mitochondrial pathway (intrinsic) and the death receptor pathway (extrinsic)
  - Activation of caspases (cysteine protease enzymes that cleave proteins after aspartic residues)<sup>Q</sup>

### Intrinsic (Mitochondrial) Pathway of Apoptosis

- Major mechanism<sup>Q</sup> of apoptosis in all mammalian cells.
- Results from release of pro-apoptotic molecules from mitochondrial intermembrane space into the cytoplasm.<sup>Q</sup>






\***Sensors:** Contain 3<sup>rd</sup> of 4 BH domains (BH3-only proteins)—BAD, BID, BIM, Puma, Noxa

### Extrinsic (Death Receptor-Initiated) Pathway of Apoptosis

- Initiated by **engagement of plasma membrane death receptors**
- Death receptors are members of the **TNF receptor family**<sup>Q</sup> that contain a cytoplasmic domain involved in protein-protein interactions and is called the **death domain**<sup>Q</sup> because it is essential for delivering apoptotic signals.
- Death receptors:** TNFR1 and FasR (CD95)



Regulatory molecules in Apoptosis	
Anti-apoptotic	Pro-apoptotic
<p>BCL2, BCL-XL, and MCL1</p> <p>Possess four BH domains (called BH 1-4)<sup>Q</sup></p> <ul style="list-style-type: none"> <li>Present in <b>outer mitochondrial &amp; ER membranes &amp; cytosol</b></li> <li>Prevent leakage of <b>cytochrome c</b> into the cytosol</li> </ul>	<p>BAX and BAK</p> <p>Possess four BH domains (called BH 1-4)</p> <ul style="list-style-type: none"> <li>Form a <b>channel in outer mitochondrial membrane</b></li> <li>Allows <b>leakage of cyt c</b> from intermembranous space</li> </ul>


### Biochemical Features of Apoptosis

- Protein cleavage**
  - Active caspases**<sup>Q</sup> cleave many vital cellular proteins and break up nuclear scaffold and cytoskeleton
- DNA Breakdown**<sup>Q</sup>
  - Characteristic breakdown of DNA into large 50 to 300 kilobase piece
  - Subsequent **internucleosomal cleavage**<sup>Q</sup> of DNA into oligonucleosomes, in multiples of **180 to 200 base pairs**<sup>Q</sup>, by Ca<sup>2+</sup> and Mg<sup>2+</sup> dependent **endonucleases**<sup>Q</sup>
- Phagocytic recognition:** Expression of **phosphatidyl serine, thrombospondin**<sup>Q</sup> → early recognition of dead cells by macrophages → phagocytosis **without the release of proinflammatory cellular components**<sup>Q</sup>




## Morphology of Apoptosis

- Cell shrinkage<sup>Q</sup>
- Chromatin condensation: most characteristic feature<sup>Q</sup> of apoptosis.
- Formation of cytoplasmic blebs and apoptotic bodies<sup>Q</sup>
- Plasma membrane remains intact<sup>Q</sup> during apoptosis
- Apoptosis in contrast to necrosis **doesn't** elicit inflammation<sup>Q</sup>.




## Diagnosis of Apoptosis

- Chromatin condensation is seen by hematoxylin, Feulgen and acridine orange staining.
- Estimation of **cytochrome 'c'**<sup>Q</sup>
- Estimation of activated caspases
- Estimation of **Annexin V** (apoptotic cells express **phosphatidylserine / thrombospondin**<sup>Q</sup> on the outer layer of plasma membrane because of which these cells are recognized by the dye **Annexin V**<sup>Q</sup>.)
- DNA breakdown (**Internucleosome cleavage by endonuclease into 200bp oligonucleosomes is a characteristic**)<sup>Q</sup> at specific sites → fragments can be detected by '**step ladder pattern**' on agarose gel electrophoresis or TUNEL (TdT mediated d-UTP Nick End Labelling) technique<sup>Q</sup>.



## Electron Microscopy of apoptosis




## High Yield Facts

- Initiator caspase:
  - Intrinsic pathway – caspase 9<sup>Q</sup>
  - Extrinsic pathway – caspase 8, 10<sup>Q</sup>
- Executionary pathway-caspase 3 (most important), 6<sup>Q</sup>
- Necrosis and apoptosis together<sup>Q</sup>-Injurious stimuli like chemotherapy and radiation can induce apoptosis<sup>Q</sup> if the insult is mild but large doses of same stimuli can induce necrotic cell death<sup>Q</sup>


## Necroptosis

Programed necrosis or "caspase-independent" programed cell death<sup>Q</sup>





## Pyroptosis



## High Yield Facts

### Examples of Necroptosis



- Formation of the mammalian bone growth plate<sup>Q</sup>
- Cell death in steatohepatitis and acute pancreatitis<sup>Q</sup>
- Reperfusion injury<sup>Q</sup>
- Neurodegenerative diseases such as Parkinson's disease<sup>Q</sup>

## FREE RADICALS


- Chemical species that have a **single unpaired electron**<sup>Q</sup> in an outer orbit.
- Unpaired electrons are highly reactive and attack inorganic or organic chemicals
- **Oxidative stress**<sup>Q</sup> - excess of these free radicals due to either increased production or decreased scavenging of reactive oxygen species (ROS).
- Implicated in **cell injury, cancer, aging, and some degenerative diseases** such as Alzheimer disease.<sup>Q</sup>

### Generation of Free Radicals<sup>Q</sup>

The generation and role of ROS in cell injury is given below.







## High Yield Facts

- **Superoxide dismutase:** Enzyme that protects the brain from free radical injury<sup>Q</sup>
- **Telltale sign** of free radical injury: **Lipofuscin pigment**<sup>Q</sup>

## CELLULAR ACCUMULATIONS

### Lipids

#### Triglycerides

- **Steatosis** (Fatty Change): accumulations of triglycerides within parenchymal cells
- **Organs** involved-liver, heart, muscle, and kidney
- **Tigered effect:** Fatty change in heart

### Cholesterol and Cholesterol Esters

- **Foam cells:** Cholesterol-laden macrophages
- Accumulation of cholesterol and cholesterol esters occurs in following conditions:
  - **Xanthomas**
  - **Cholesterosis:** In the lamina propria of **gall bladder**<sup>Q</sup>
  - **Niemann-Pick disease, type C:** Mutations affecting an enzyme involved in cholesterol trafficking, resulting in cholesterol accumulation in multiple organs

#### Phospholipids

Components of the **myelin figures**<sup>Q</sup> found in necrotic cells




Image showing foam cells




Image shows macrovascular steatosis




Image showing cholesterol clefts (arrow)

### Proteins

#### Resorption Droplets

- **Appear as rounded, eosinophilic droplets, vacuoles, or aggregates in the cytoplasm.**<sup>Q</sup>
- Examples of Protein Accumulates are given below:






Image showing reabsorption droplets in PCT



Cell showing many Russell bodies





## Hyaline Change

- Any intracellular or extracellular accumulation with pink homogenous appearance
- Descriptive histologic term rather than a specific marker for cell injury


## Glycogen

- Appears as clear vacuoles within the cytoplasm

- Best fixed in absolute alcohol

### Best stain:


- Best carmine or PAS with diastase (Diastase hydrolyzes glycogen)
- Seen in:
  - Diabetes mellitus:** Glycogen accumulates in renal tubular epithelial cells, liver cells,  $\beta$  cells of the islets of Langerhans, and heart muscle cells
  - Glycogen storage diseases<sup>Q</sup>**, or glycogenoses



## Calcification

- Abnormal tissue deposition of **calcium salts**, with small amounts of **iron, magnesium and other mineral salts<sup>Q</sup>**
- It can be of the following two types:

Features	Dystrophic	Metastatic
Seen in	Dead tissues <sup>Q</sup> , damaged & aging tissues	Living tissues
Serum Calcium	Normal <sup>Q</sup>	Elevated
Sites	Seen in cell injury and necrosis <sup>Q</sup>	Mainly affects gastric mucosa <sup>Q</sup> , kidneys <sup>Q</sup> , lungs <sup>Q</sup> , systemic arteries <sup>Q</sup> & pulmonary veins. <sup>Q</sup>
Effect	Organ dysfunction	Usually no clinical dysfunction, but massive involvement may cause organ damage
Etiology	<p><b>"R-A-T-T"</b></p> <p><b>R</b>-Rheumatic heart ds (Cardiac valves)</p> <p><b>A</b>-Atheromatous plaque</p> <p><b>T</b>-Tubercular lymph node</p> <p><b>T-Tumors (MP-PG)</b></p> <ul style="list-style-type: none"> <li>Meningioma, Mesothelioma</li> <li>Papillary carcinoma of thyroid, Ovary (serous carcinoma)</li> <li>Prolactinoma</li> <li>Glucagonoma</li> </ul> <p>Examples of Dystrophic Calcification</p> <ul style="list-style-type: none"> <li>Psammoma bodies<sup>Q</sup></li> <li>Asbestos bodies<sup>Q</sup></li> <li>Mönckeberg's sclerosis<sup>Q</sup></li> </ul>	<p><b>PRDR (Postgraduate Reads Dr.)</b></p> <ul style="list-style-type: none"> <li>HyperParathyroidism-               <ul style="list-style-type: none"> <li>Parathyroid tumors</li> <li>Ectopic PTHRP from malignant tumors</li> </ul> </li> <li>Resorption of bone tissue-               <ul style="list-style-type: none"> <li>Multiple myeloma<sup>Q</sup></li> <li>Diffuse skeletal metastasis</li> <li>Accelerated bone turnover-Pagets' disease</li> <li>Immobilization</li> </ul> </li> <li>Vitamin <b>D</b>-related disorders               <ul style="list-style-type: none"> <li>Vitamin D intoxication,</li> <li>Sarcoidosis (Macrophages activate vit D precursor)</li> <li>Idiopathic hypercalcemia of infancy (<b>Williams syndrome</b>)<sup>Q</sup></li> </ul> </li> <li><b>Renal failure</b>-Due to secondary hyperparathyroidism</li> <li><b>Others:</b> Milk alkali syndrome, Aluminum intoxication</li> <li>Vitamin A toxicity</li> </ul>



Monckeberg sclerosis  
(Calcification of media of BV)






Image showing psammoma bodies




## Stains

Most common stain used for calcium-Von kossa



Von kossa gives black colour to calcium (arrow)

Most specific stain used for calcium Alizarin Red




Alizarin Red Stain For Calcium

## High Yield Facts

- Calcification **begins** in **mitochondria**<sup>q</sup>
- **Mönckeberg's sclerosis**<sup>q</sup> (**medial calcific sclerosis**) i.e it's a type of **dystrophic calcification**.<sup>q</sup> Calcium deposits are found in **tunica media** of arteries is a misnomer.
- **Internal alkaline environment**<sup>q</sup> in tissues favors **metastatic calcification**
- **Most common site of metastatic calcification is lungs > kidneys**<sup>q</sup>



## Pigments



### High Yield Facts

- The pigments **do not usually evoke any inflammatory response**.<sup>o</sup>
- Iron is normally **carried with transferrin in circulation**.<sup>o</sup>
- In cells, it is stored in association with a protein, **apoferritin**, to form **ferritin micelles**<sup>o</sup>
- Whenever there is a local or systemic excess of iron, **ferritin forms hemosiderin granules**<sup>o</sup>
- Hemosiderin is considered as a **degraded and oxidized form** of ferritin.<sup>o</sup>
- **Ferritin is soluble** while **hemosiderin is insoluble**.<sup>o</sup>


### Stains

- **Masson Fontana**-melanin
- Remember **Masson trichrome** is not for pigment, it distinguishes collagen from muscle
- **Long ZN stain** for Lipofuscin
- **Perls Prussian blue** for hemosiderin (Fe<sup>3+</sup>)
- The tissue-bound ferric ions subsequently are visualized by treatment with **potassium ferrocyanide** to form bright blue deposits of ferric ferrocyanide or Prussian blue.
- **Lillie's method** is used for picking up ferrous iron



## CELLULAR AGING

The major causes of cellular aging are:



### Cellular Senescence

- Normal cells have a limited capacity for replication
- After a fixed number of divisions, cells become arrested in a terminally nondividing state, known as “replicative senescence”<sup>Q</sup>

Two mechanisms are responsible for cellular senescence

Telomere Attrition	Activation of Tumor Suppressor Genes
<ul style="list-style-type: none"> <li>Telomeres are short repeated sequences at the ends of DNA which ensure complete replication of chromosome.<sup>Q</sup></li> <li>Telomere length is maintained by enzyme called telomerase.<sup>Q</sup></li> <li>Telomerase is RNA dependent DNA polymerase</li> </ul>	<p>CDKN2A locus encodes a tumor suppressor gene called p16 or INK4a, which controls G1 to S phase progression during the cell cycle<sup>Q</sup></p> <p>↓</p> <p>Protects the cells from uncontrolled mitogenic signals and pushes cells along the senescence pathway.<sup>Q</sup></p>
<ul style="list-style-type: none"> <li>Telomerase is absent in somatic cells, present only in germ cells and present at low levels in stem cells<sup>Q</sup></li> <li>Replicative senescence<sup>Q</sup> - When somatic cells replicate, telomeres become progressively shortened → signals cell cycle arrest</li> </ul>	

### Defective Protein Homeostasis

- Both normal folding and degradation of misfolded proteins (by autophagy and ubiquitin-proteasome system) are impaired with aging
- Rapamycin has multiple effects including promotion of autophagy and thus increases the life span of middle aged mice

### Dysregulated Nutrient Sensing

Caloric restriction increases lon reducing the signaling intensity of the IGF-1 pathway<sup>Q</sup> and by increasing sirtuins<sup>Q</sup>

### High Yield Facts



#### Hayflick limit

- Normal cells can undergo 60–70 cell divisions in their lifetime. This limit is called Hayflick limit.
- Cause of Hayflick limit is progressive telomere shortening with cell division.

### High Yield Facts



Insulin and insulin-like growth factor 1 (IGF-1) signalling pathway

- Growth hormone secretion by the pituitary stimulates IGF-1 (AKA somatomedin C)
- IGF1 promotes anabolic state as well as cell growth, replication and aging
- It has two downstream targets- AKT and mTOR (mammalian target of rapamycin)
- Rapamycin might increase the life span of middle aged mice by attenuation of IGF-1



## Sirtuins

- Family of **NAD-dependent protein deacetylases**.
- Sirtuin-6<sup>Q</sup>** promotes the expression of genes whose products increase longevity.
- These include **proteins that inhibit metabolic activity<sup>Q</sup>, reduce apoptosis<sup>Q</sup>, stimulate protein folding<sup>Q</sup>, inhibit the harmful effects of oxygen free radicals<sup>Q</sup> and activating DNA repair enzymes through deacylation<sup>Q</sup>**
- Red wine may activate sirtuins and thus increase life span**
- Sirtuins also **increase insulin sensitivity and glucose metabolism**, and may be **targets for the treatment of diabetes**.

## High Yield Facts

- Defective DNA helicase causes **Werner's syndrome (premature aging)**
- Average hematopoietic stem cell suffers **14 new mutations per year<sup>Q</sup>**
- Telomerase is **reactivated in immortalized cancer cells<sup>Q</sup>**
- Mutation of CDKN2A is seen in melanomas<sup>Q</sup>**
- Free radical mediated damage shows-Lipofuscin accumulation**
- Sirtuins have a role in aging, diabetes and cancers<sup>Q</sup>**

R<sup>9th</sup>

## Latest Update

### AUTOPHAGY


- Autophagy ("self-eating") refers to lysosomal digestion of the cell's own components
- It is an **adaptation** to nutrient deprivation
- If the stress is too severe for the process to cope with it, it results in cell death by apoptosis
- Physiological** autophagy is seen in aging and exercise
- It is of three types:
  - Chaperone-mediated**
    - Direct translocation** through lysosomal membrane by chaperone proteins.
  - Microautophagy**
    - Inward invagination of lysosomal membrane for delivery.
  - Macroautophagy**
    - Its **major form** of autophagy
    - It involves sequestration and transportation of cytosol content in a double-membrane bound autophagic vacuole (**autophagosome**).

### Autophagy Plays a Role in Human Diseases

- Cancer:** Autophagy can both promote cancer growth and act as a defense against cancers.
- Neurodegenerative disorders:**
  - Alzheimer's disease—autophagosomes is **accelerated**
  - Huntington's disease—mutant huntingtin **impairs** autophagy.
- Infectious diseases:** Macrophage-specific deletion of **Atg5** increases susceptibility to tuberculosis.
- Polymorphisms in a gene involved in autophagy are associated with **inflammatory bowel disease**


## High Yield Facts

- Nutrient starvation induces autophagy in eukaryotic cells through **inhibition of TOR** (target of rapamycin)
- Nucleation is started by **Beclin 1**
- LC3**-useful marker for autophagy





## ENDOPLASMIC RETICULUM STRESS



### Diseases caused by Misfolded proteins


Disease	Affected Protein	Pathogenesis
<b>Diseases caused by mutant protein that are degraded, leading to their deficiency</b>		
Cystic fibrosis	CFTR	Loss of CFTR leads to defects in chloride transport and death of affected cells
Familial hypercholesterolemia	LDL receptor	Loss of LDL receptor leads to hypercholesterolemia
Tay-Sachs disease	Hexosaminidase $\beta$ subunit	Lack of the lysosomal enzyme leads to accumulation of GM2 gangliosides in neurons
<b>Diseases caused by Misfolded proteins that result in ER stress-induced cell loss</b>		
Retinitis pigmentosa	Rhodopsin	Abnormal folding of rhodopsin causes photoreceptor loss and cell death, resulting in blindness
Creutzfeldt-Jakob disease	Prions	Abnormal folding of PrP <sup>Sc</sup> causes neuronal cell death
Alzheimer's disease	A $\beta$	Abnormal folding of A $\beta$ peptide causes aggregation within neurons & apoptosis
<b>Diseases caused by Misfolded protein that result from both ER stress-induced cell loss and functional deficiency of the protein</b>		
Alpha-1-anti-trypsin deficiency	$\alpha$ -1 anti-trypsin	Storage of nonfunctional protein in hepatocytes cause apoptosis; absence of enzymatic activity in lungs causes elastic tissue destruction giving rise to emphysema



## NEXT Pattern Questions



1. A 36-year-old obese woman has experienced heartburn from gastric reflux for the past 5 years after eating large meals. She undergoes upper gastrointestinal endoscopy, and a biopsy specimen of the distal esophagus is obtained. Which of the following microscopic changes, seen in the figure, has most likely occurred?



- a. Columnar metaplasia      b. Goblet cell hyperplasia      c. Lamina propria atrophy      d. Squamous dysplasia


Ans. (a) **Columnar metaplasia**

(Ref: Robins Basic Pathology 10th ed/pg 50)

- Inflammation from reflux of gastric acid has resulted in replacement of normal esophageal squamous epithelium by intestinal-type columnar epithelium with goblet cells. Such conversion of one adult cell type to another cell type is called metaplasia, and it occurs when stimuli reprogram stem cells. Goblet cells are not normal constituents of the esophageal mucosa, and they are a minor part of this metaplastic process. The lamina propria has some inflammatory cells, but it does not atrophy. The squamous epithelium does not become dysplastic from acid reflux, but the columnar metaplasia may progress to dysplasia, not seen here, if the abnormal stimuli continue. These cells are not significantly increased in size (hypertrophic).



2. While doing a screening chest radiograph in an asymptomatic 37-year-old man a 3 cm nodule in the middle lobe of his right lung was found. The nodule on sectioning shows a sharply circumscribed mass with a soft, white center. The microscopic appearance is shown in the figure. The serum interferon gamma release assay is positive. Which of the following pathologic processes has most likely occurred in this nodule?



- a. Apoptosis      b. Caseous necrosis      c. Coagulative necrosis      d. Fat necrosis

Ans. (b) **Caseous necrosis**

(Ref: Robins Basic Pathology 10th ed/pg 35)

- The grossly cheese like appearance gives this form of necrosis its name—caseous necrosis. The figure shows amorphous pink acellular material at the upper right surrounded by epithelioid macrophages, and a Langhans giant cell is visible at the upper left. In the lung, tuberculosis and fungal infections are most likely to produce this pattern of tissue injury. Apoptosis involves individual cells, without grossly apparent extensive or localized areas of tissue necrosis. Coagulative necrosis is more typical of ischemic tissue injury. Fat necrosis most often occurs in the breast and pancreas. Fatty change is most often a feature of hepatocyte injury, and the cell integrity is maintained. Gangrene characterizes extensive necrosis of multiple cell types in a body region or organ. Liquefactive necrosis is seen in neutrophilic abscesses or ischemic cerebral injury.





Q's

3. An experiment introduces a knockout gene mutation into a cell line. The frequency of shrunken cells with chromatin clumping, karyorrhexis, and cytoplasmic blebbing is increased compared with a cell line without the mutation. Overall survival of the mutant cell line is reduced. Which of the following genes is most likely to be affected by this mutation?

- a. BAX      b. BCL2      c. C-MYC      d. FAS

Ans. (b) **BCL2**


(Ref: Robins Basic Pathology 10th ed/pg 37)

- These histologic findings are typical of apoptosis. The BCL2 gene product inhibits cellular apoptosis by binding to Apaf-1. Hence, the knockout removes this inhibition. The BAX gene product promotes apoptosis, and a knockout would protect against apoptosis. The C-MYC gene is involved with oncogenesis. The FAS gene encodes for a cellular receptor for Fas ligand that signals apoptosis. Activity of the p53 (TP53) gene normally stimulates apoptosis, but mutation favors cell survival.




## Image-Based Questions

1. A 45-year-old male had history of hepatitis B infection. On liver biopsy, multiple councilman bodies were seen. Electrophoresis was done to pick up whether he was undergoing apoptosis or necrosis. Identify the pattern in lane 2 and diagnosis.




- a. Step ladder pattern, apoptosis  
b. Smeared pattern, necrosis  
c. Step ladder pattern, necrosis  
d. Smeared pattern, apoptosis

3. A 50-year-old female comes with history of discharge per vaginum. On examination, her squamocolumnar junction of cervix appears erythematous. She underwent cervical biopsy which showed following findings. Describe the change marked by arrow.




- a. Squamous metaplasia  
b. Columnar metaplasia  
c. Transitional metaplasia  
d. None

2. Earliest morphological change seen in reversible cellular injury? Inset shows normal hepatocytes as control.




- a. Hydropic change      b. Fatty change  
c. Necrosis      d. None

4. A 60-year-old asymptomatic female shows following change in tunica media of blood vessels. Diagnosis is:




- a. Medial calcification      b. Medial fibrosis  
c. Amyloidosis      d. None





5. A 50-year-old female underwent cholecystectomy on examination, the lamina propria of gall bladder was seen infiltrated by foam cells. Diagnosis is:



- |                  |                    |
|------------------|--------------------|
| a. Cholesterosis | b. Atherosclerosis |
| c. Steatosis     | d. None            |



## Answers of Image-Based Questions

1. Ans. (a) **Step ladder pattern, apoptosis**

- This is step ladder pattern which is typically seen in apoptosis, its also called DNA laddering characterized by the activation of endogenous endonucleases with subsequent cleavage of chromatin DNA into internucleosomal fragments of roughly 180-200 base pairs (bp).
- Smear pattern is seen in necrosis.

2. Ans. (a) **Hydropic change**

- Cellular swelling is the first manifestation of almost all forms of injury to cells. Cellular swelling appears whenever cells are incapable of maintaining ionic and fluid homeostasis and is the result of failure of energy-dependent ion pumps in the plasma membrane. It is reversible.
- On microscopic examination, small clear vacuoles may be seen within the cytoplasm; these represent distended and pinched-off segments of the ER. This pattern of nonlethal injury is sometimes called hydropic change or vacuolar degeneration.
- This is a case of hydropic change in hepatocytes (control normal hepatocytes are seen in inset).

3. Ans. (a) **Squamous metaplasia**


- Here we are seeing endocervix lined by columnar epithelium. And underlying stroma shows endocervical glands. Here columnar epithelium is being changed to squamous epithelium suggestive of squamous metaplasia. Inciting cause for this metaplasia is chronic cervicitis.

4. Ans. (a) **Medial calcification**

- Medial artery calcification (MAC) is also known as Mönckeberg's arteriosclerosis, is a nonobstructive condition leading to reduced arterial compliance that is commonly considered as a nonsignificant finding.
- With the H&E stain, calcium appear deep blue-purple.

5. Ans. (a) **Cholesterosis of the gallbladder**

- Here we are seeing accumulation of foam cells in lamina propria of gallbladder suggestive of cholesterosis



## Multiple Choice Questions

### ADAPTATIONS

- 1. True about Metaplasia is:** (JIPMER 2016)
  - a. Involves only epithelial cells
  - b. Is irreversible
  - c. Occurs at stem cells level
  - d. Columnar is the most common type
- 2. An example of metaplasia is?** (Recent Question 2015)
  - a. CIN
  - b. Barrets
  - c. Adenoma
  - d. Bronchial carcinoid
- 3. All the following are true regarding hypertrophy except:** (Recent Question 2015)
  - a. Increase in cell size without increase in number
  - b. DNA content same as in normal cells
  - c. Increase in cell size is due to synthesise of more cellular proteins
  - d. Associated with a switch of contractile proteins from adult to fetal or neonatal forms
- 4. Find the false statement about metaplasia:** (Recent Question 2015)
  - a. Reversible
  - b. No loss of polarity
  - c. Reprogramming of stem cells
  - d. Pleomorphism present
- 5. Increase in the number of goblet cells in the non-respiratory terminal bronchiole is an example of:** (Recent Question 2015)
  - a. Anaplasia
  - b. Dysplasia
  - c. Metaplasia
  - d. Hyperplasia
- 6. In respiratory tract metaplasia occurs from:** (Recent Question 2015)
  - a. Squamous to columnar
  - b. Columnar to cuboidal
  - c. Columnar to squamous
  - d. Cuboidal to squamous
- 7. In Vitamin-A deficiency, cancerous lesions occur due to:**
  - a. Metaplasia
  - b. Dysplasia (MH PG 2014)
  - c. Aplasia
  - d. Hyperplasia
- 8. Definition of hyperplasia is:** (Recent Question 2015)
  - a. Increase in number of cells
  - b. Increase in size of cells
  - c. Change in type of cell
  - d. Increase in nuclear:cytoplasmic ratio
- 9. All are cellular adaptations except:** (Recent Question 2014)
  - a. Hypertrophy
  - b. Hyperplasia
  - c. Necrosis
  - d. Metaplasia
- 10. Decrease in cell size refers to:** (DNB 2012)
  - a. Atrophy
  - b. Metaplasia
  - c. Hyperplasia
  - d. Hypertrophy
- 11. Metaplasia arises from reprogramming of** (JIPMER 2012)
  - a. Stem cells
  - b. Stellate cells
  - c. Squamous cells
  - d. Columnar cells

### CELL INJURY

- 12. All of the followings are signs of reversible cell injury; except:** (AIIMS May 19)
  - a. Loss of microvilli
  - b. Cell Swelling
  - c. Bleb formation
  - d. Dense Mitochondrial deposit
- 13. Features of irreversible cell injury is/are?** (PGI Nov 2017)
  - a. Lysosomal rupture
  - b. Pyknosis
  - c. Bleb formation on membrane
  - d. Severe mitochondrial dysfunction
- 14. A Patient have acquired syndrome associated with defective breakdown and disposal of intracellular fatty acids. Which intracellular organelle is concerned with this mechanism?** (Recent Questions 2016-17, JIPMER 2014)
  - a. Mitochondria
  - b. Peroxisomes
  - c. Lysosomes
  - d. Smooth endoplasmic reticulum
- 15. In myocardium reperfusion injury is due to?** (Recent Question 2016)
  - a. Ca
  - b. Mg
  - c. K
  - d. Mn
- 16. Which of the following is not a sign of reversible cell injury?** (Recent Question 2015)
  - a. ATP depletion
  - b. Cell shrinkage
  - c. Fatty acid deposition
  - d. Reduction of phosphorylation
- 17. Earliest feature of reversible cell injury is:** (Recent Question 2015)
  - a. Cellular swelling
  - b. Decreased ATP
  - c. Clumping of chromatin
  - d. Decreased protein synthesis
- 18. First manifestation in cell injury** (Recent Question 2015)
  - a. Pyknosis
  - b. Cell swelling
  - c. Nuclear fragmentation
  - d. Nuclear lysis
- 19. The following is not a reversible cell injury** (Recent Question 2015)
  - a. Loss of microvilli in plasma membrane
  - b. Lipid vacuoles in cytoplasm
  - c. Mitochondrial swelling and small amorphous densities
  - d. Moth eaten appearance of cytoplasm
- 20. All are features of reversible injury of cell, except** (Recent Question 2014-15)
  - a. Blebs
  - b. Amorphous densities in mitochondrial matrix
  - c. Loss of microvilli
  - d. Cellular swelling




- 21. Cells seen in chronic infection of pseudomonas** (Recent Question 2014)  
 a. Neutrophils                      b. Eosinophils  
 c. Lymphocytes                    d. Macrophage
- 22. Cells most sensitive to hypoxia are:** (AIIMS May 2014)  
 a. Myocardial cells  
 b. Neurons  
 c. Hepatocytes  
 d. Renal tubular epithelial cells
- 23. Irreversible cell injury:** (Recent Question 2013)  
 a. Mitochondrial densities  
 b. Cellular swelling  
 c. Blebs  
 d. None
- 24. In cell death, myelin figures are derived from:** (Recent Question 2013)  
 a. Nucleus                            b. Cell membrane  
 c. Cytoplasm                        d. Mitochondria
- 25. Which finding on electron microscopy indicates irreversible cell injury:** (AIIMS May 12, Nov 02)  
 a. Dilatation of endoplasmic reticulum.  
 b. Dissociation of ribosomes from rough endoplasmic reticulum  
 c. Flocculent amorphous densities in the mitochondria  
 d. Myelin figures

#### CELL DEATH

- 26. Which of the following is activated by intrinsic or extrinsic pathways?** (AIIMS Nov 2019)  
 a. Necroptosis  
 b. Apoptosis  
 c. Necrosis  
 d. Ferroptosis
- 27. Acute inflammatory response is seen in:** (AIIMS May 19)  
 a. Pyroptosis                      b. Necroptosis  
 c. Necrosis                        d. Autophagy
- 28. Which of the following is involved in apoptosis pathway?** (JIPMER Dec 19)  
 a. Myc  
 b. p53 and caspases  
 c. APC  
 d. VHL
- 29. Which of the following statements is false about apoptosis?** (AIIMS Nov 18)  
 a. No inflammation  
 b. Plasma membrane intact  
 c. Organelle swelling  
 d. Affected by dedicated genes
- 30. Which of the following can recognize dead material?** (AIIMS Nov 18)  
 a. NET                                b. Inflammasome  
 c. Necrosis                        d. Toll like receptor
- 31. Necrotic cells are recognized by NOD like receptors which then activates inflammasomes. Which of these are anti apoptotic gene?** (PGI Nov 2018)  
 a. BCL                                b. BCL-2  
 c. BAD                                d. BAX  
 e. MCL

- 32. IL-1 is activated by?** (Recent Question 2019)  
 a. Caspase 1                        b. Caspase 3  
 c. Caspase 5                        d. Caspase 8
- 33. Which of the following is an antiapoptotic gene?** (Recent Question 2019)  
 a. Bcl2                                b. Bcl - XL  
 c. BAX                                d. Both a & b
- 34. Which of the following type of necrosis is seen in immune complex deposition in blood vessel?** (Recent Question 2019)  
 a. Coagulative necrosis        b. Liquefactive necrosis  
 c. Fibrinoid necrosis            d. Caseous necrosis
- 35. For programmed cell death type 2 and autophagy, which is apoptotic genes?** (JIPMER 2017)  
 a. BCL                                b. BAX  
 c. BCL-XL                        d. BIM
- 36. True about p53 gene?** (PGI Nov 2016)  
 a. Tumor suppression        b. Proapoptotic  
 c. Antiapoptotic                d. Cell repair
- 37. True about caspases are?** (PGI Nov 2016)  
 a. They are enzymes starting apoptosis  
 b. They inhibit apoptosis  
 c. They are receptors of apoptosis  
 d. They are proteases which cause cellular death in apoptosis found in irreversible cell damage
- 38. About the given image true is?** (AIIMS May 16)



- a. C is showing necrosis  
 b. C is showing apoptosis  
 c. C is showing normal cells  
 d. A is showing apoptosis
- 39. True about necroptosis is all except?** (Recent Question 2016-17)  
 a. Caspase 1 & 11 is involved  
 b. Caspase independent  
 c. Failure of activation of caspase 8  
 d. Lipid peroxidation is seen
- 40. Which of the following is an antiapoptotic gene** (Recent Question 2016-17)  
 a. BAX                                b. BAD  
 c. BCL-XL                        d. BIM
- 41. SMAC/DIAMBLO is a** (Recent Question 2016-17)  
 a. Anti apoptotic protein  
 b. Induces necrosis  
 c. Acts both as anti and pro apoptotic protein  
 d. Pro-apoptotic protein
- 42. Antiapoptotic gene?** (Recent Question 2016-17)  
 a. FLIP                                b. P53  
 c. BAX                                d. BIM
- 43. Necrosis seen in chronic pancreatitis is?** (Recent Question 2016)  
 a. Fatty                                b. Coagulative  
 c. Liquifactive                    d. Casseous



- 44. False about apoptosis?** (Recent Question 2015)  
 a. Inflammation present  
 b. Program cell death  
 c. Normal physiology  
 d. Genetically determined by a cell
- 45. Earliest change in cell death is?** (Recent Question 2015)  
 a. Karyolysis  
 b. Loss of plasma membrane  
 c. Cell swelling  
 d. Karyorrhexis
- 46. Which of the following induces apoptosis** (Recent Question 2015)  
 a. Oleic acid  
 b. Myristic acid  
 c. Glucocorticoid  
 d. Isoprenoid
- 47. Which of the following is an execution caspase** (Recent Question 2015)  
 a. Caspase 3  
 b. Caspase 5  
 c. Caspase 8  
 d. Caspase 9
- 48. Diabetic foot is an example of** (Recent Question 2015)  
 a. Dry gangrene  
 b. Wet gangrene  
 c. Gas gangrene  
 d. Necrotizing inflammation
- 49. Annexin V is a marker of** (Recent Question 2015)  
 a. Necrosis  
 b. Gangrene  
 c. Aging  
 d. Apoptosis
- 50. Cell organelle which plays a pivotal role in apoptosis** (Recent Question 2015)  
 a. Nucleus  
 b. Mitochondria  
 c. Golgi apparatus  
 d. Plasma membrane
- 51. Fibrinoid necrosis is seen in all except** (Recent Question 2015)  
 a. Malignant hypertension  
 b. Polyarteritis nodosa  
 c. Diabetic glomerulosclerosis  
 d. Rheumatic heart disease
- 52. Ladder pattern of DNA electrophoresis is seen in** (Recent Question 2015)  
 a. Necrosis  
 b. Apoptosis  
 c. Cytolysis  
 d. Karyorrhexis
- 53. The following is not a sensor of apoptosis** (Recent Question 2015)  
 a. Puma  
 b. Noxa  
 c. Bax  
 d. BAD
- 54. Intrinsic pathway of apoptosis is initiated by all the following except** (Recent Question 2015)  
 a. Growth factor withdrawal  
 b. DNA damage  
 c. Protein misfolding  
 d. Type 1 TNF receptor
- 55. Defective apoptosis and increased cell survival is seen in** (Recent Question 2015)  
 a. Autoimmune disease  
 b. Neurodegenerative disease  
 c. Viral infections  
 d. Ischemic injury
- 56. Receptor associated kinases 1 (RIP1) and 3 (RIP3) are involved in** (Recent Question 2015)  
 a. Necrosis  
 b. Apoptosis  
 c. Necroptosis  
 d. Pyroptosis
- 57. Inflammasome is formed in** (Recent Question 2015)  
 a. Necrosis  
 b. Apoptosis  
 c. Necroptosis  
 d. Pyroptosis
- 58. Find the true statement regarding fibrinoid necrosis** (Recent Question 2015)  
 a. Denaturation of structural proteins  
 b. Granuloma formation  
 c. Abscess formation  
 d. Fibrin deposition
- 59. All are true regarding apoptosis except** (Recent Question 2015)  
 a. Active process  
 b. Cell size decreases  
 c. Inflammation absent  
 d. Smear pattern in electrophoresis
- 60. The following is a pro-apoptotic factor** (WB PGMEE 2016, Recent Question 2015)  
 a. Bax  
 b. Bcl-2  
 c. Bcl-xL  
 d. Mcl-1
- 61. Find the wrong match** (Recent Question 2015)  
 a. Coagulative necrosis-Tuberculosis  
 b. Fat necrosis -Acute pancreatitis  
 c. Liquefactive necrosis -Brain  
 d. Fibrinoid necrosis -Malignant hypertension
- 62. Fibrinoid necrosis is seen in all the following except** (Recent Question 2015)  
 a. Malignant hypertension  
 b. Aschoff's nodule  
 c. Polyarteritis nodosa  
 d. Diabetic glomerulosclerosis
- 63. Apoptosis is induced by** (JIPMER 2015)  
 a. Caspases  
 b. DNA synthesis  
 c. Activation of caspases  
 d. Kinase pathway
- 64. Hypoxic death of brain tissue results in:** (APPGMEE 2015)  
 a. Gangrenous necrosis  
 b. Liquefactive necrosis  
 c. Coagulative necrosis  
 d. Fibrinoid necrosis
- 65. True about bcl-2:** (PGI May 2015)  
 a. ↑ Apoptosis  
 b. ↓ Apoptosis  
 c. ↑ Resistance of tumour to treatment  
 d. Only associated with follicular lymphoma  
 e. Cause meningioma
- 66. True about Apoptosis are all except** (Recent Question 2014-15)  
 a. Inflammation is present  
 b. Chromosomal breakage  
 c. Clumping of chromatin  
 d. Cell shrinkage
- 67. Not true about apoptosis** (Recent Question 2014-15)  
 a. Increase in lysosomal enzyme  
 b. Increase in caspases  
 c. Phosphatidyl serine has important role  
 d. Internucleosomal cleavage of nucleus
- 68. Apoptotic bodies are** (Recent Question 2014-15)  
 a. Clumped chromatin bodies  
 b. Pyknotic nucleus without organelles  
 c. Cell membrane bound with organelles  
 d. No nucleus with organelles
- 69. "Caspase-independent" programmed cell death** (Recent Question 2015)  
 a. Necrosis  
 b. Necroptosis  
 c. Apoptosis  
 d. None






70. **Fibrinoid necrosis is seen in** (Recent Question 2015)  
a. Malignant HTN      b. Benign hypertension  
c. Diabetes      d. Acute on chronic gangrene
71. **Immune complexes mediated necrosis is of which type?** (Recent Question 2013)  
a. Coagulative necrosis      b. Liquefactive necrosis  
c. Caseous necrosis      d. Fibrinoid necrosis
72. **Which one of the following is an antiapoptotic protein/gene?** (WB PGME 2016, Recent Question 2013)  
a. BAK      b. BCL-2  
c. BAX      d. BIM
73. **Coagulative necrosis is due to:** (Recent Question 2013)  
a. Denaturation of protein      b. Enzymatic digestion  
c. Infection      d. None
74. **Fat necrosis is common in:** (Recent Question 2013)  
a. Omentum      b. Breast  
c. Retroperitoneal fat      d. All of the above
75. **In apoptosis, cytochrome C acts through:** (Recent Question 2013)  
a. Apaf 1      b. Bcl-2  
c. FADD      d. TNF
76. **Which of the following is not seen in apoptosis?** (AIIMS May 2013)  
a. Chromatin condensation  
b. DNA fragmentation  
c. Inflammation  
d. Cell membrane shrinkage
77. **CD 95 is a marker of:** (AIIMS May 2013)  
a. Intrinsic pathway of apoptosis  
b. Extrinsic pathway of apoptosis  
c. Monocyte  
d. Leucocyte
78. **Apoptosis- all are true except?** (JIPMER 2013)  
a. Normal physiological process of programmed cell death  
b. Products removed by phagocytosis  
c. Plasma membrane zeiosis  
d. Causes inflammation that damage surrounding cells
79. **Apoptosis does not occur by normal capsase pathway in?** (DNB Aug. 12 Pattern)  
a. Liver      b. Muscle  
c. Neurons      d. Skin
80. **Type of necrosis in Myocarial Infarcction?** (JIPMER 2012, UP10)  
a. Caseous      b. Coagulative  
c. Liquifactive      d. Fibrinoid
81. **Which of the following organelles plays a pivotal role in apoptosis?** (AIIMS May 10, AI 11, 09)  
a. Mitochondria      b. Endoplasmic reticulum  
c. Nucleus      d. Golgi apparatus
82. **Which of the following has a direct role in apoptosis?** (DNB Dec 11)  
a. Nitric oxide      b. Adenylcyclase  
c. cAMP      d. Cytochrome C
83. **Apoptosis is inhibited by:** (MH 11)  
a. p53      b. nMYC  
c. RAS      d. Bcl-2
84. **Characteristic feature of apoptosis:** (AI 10)  
a. Cell membrane intact      b. Cytoplasmic eosinophilia  
c. Nuclear moulding      d. Cell swelling

85. **The characteristic feature of apoptosis on light microscopy is:** (AI 10)  
a. Cellular swelling      b. Nuclear compaction  
c. Intact cell membrane      d. Cytoplasmic eosinophilia
86. **All of the following are features of apoptosis, except:** (AI 10)  
a. Cellular swelling      b. Nuclear compaction  
c. Intact cell membrane      d. Cytoplasmic eosinophilia
87. **Coagulative necrosis is seen in:** (AIIMS May 10)  
a. T.B.      b. Sarcoidosis  
c. Cryptococcal infection      d. Wet Gangrene
88. **Caspases are associated with:** (AIIMS May 10)  
a. Organogenesis      b. Hydropic degeneration  
c. Collagen hyalinization      d. Morphology
89. **True about apoptosis:** (PGI May 2010)  
a. Increase in lysosomal enzyme  
b. Increase in caspases  
c. Phosphatidyl serine has important role  
d. Internucleosomal cleavage of nucleus
90. **In apoptosis, permeabilization of membrane occur in:** (PGI Nov 2010)  
a. Nuclear membrane      b. Cytoplasmic membrane  
c. Lysosome      d. Ribosome  
e. Mitochondrial membrane

#### CELLULAR ACCUMULATIONS

91. **Dystrophic calcification is seen in:** (AIIMS May 19)  
a. Paget's disease of bone  
b. Lung involvement in sarcoidosis  
c. Immobilized healing fracture  
d. Myositis ossificans
92. **Dystrophic calcification is/are found in?** (PGI Nov 2017)  
a. Monckeberg's medial sclerosis  
b. Papillary carcinoma thyroid  
c. Hyperparathyroidism  
d. Meningioma  
e. Vitamin D intoxication
93. **Given below is the histopathology of liver biopsy of hemochromatosis. Which of the following stain is used?** (AIIMS Nov 16)



- a. Von kossa      b. Alcian blue  
c. Prussian blue      d. Crystal violet
94. **Which of the following stain and the material stained by it is correct?** (PGI Nov 2016)  
a. Perl stain-Iron  
b. Collagen- Von kossa  
c. Elastin-VG  
d. Copper-rhodamine



- 95. Triggered myocardium is deposition of ?**  
(Recent Question 2016-17)  
a. Fat                                      b. Hyaline  
c. Glycogen                              d. Protein
- 96. Psammoma bodies are absent in?**  
(MAHA 2016, Recent Question 2015)  
a. Meningioma                              b. Papillary ca of ovary  
c. Prolactinoma                              d. Seminoma
- 97. Principle of Prussian blue stain:**  
(Recent Question 2015)  
a. Ferrous to ferricyanide  
b. Ferrocyanide to ferroferric cyanide  
c. Ferroferricyanide to ferrocyanide  
d. Ferrocyanide to ferricferrocyanide
- 98. Metastatic calcification seen in all except:**  
(Recent Question 2015)  
a. Multiple myeloma                              b. Breast cancer  
c. Atherosclerosis                              d. Renal failure
- 99. Intracellular calcification begins in**  
(Recent Question 2015)  
a. Nucleus                              b. Cytoplasm  
c. Golgi complex                              d. Mitochondria
- 100. Most commonly involved organ in metastatic calcification**  
(Recent Question 2015)  
a. Kidney                              b. Heart  
c. Lungs                              d. Aorta
- 101. The following is not a disease caused by misfolding of proteins**  
(Recent Question 2015)  
a. Cystic fibrosis  
b. Parkinsons disease  
c. Alzheimer's disease  
d. Creutzfeldt-Jacob disease
- 102. Intracellular calcification begins in**  
(Recent Question 2015)  
a. Nucleus                              b. Cytoplasm  
c. Golgi complex                              d. Mitochondria
- 103. A patient died of alzheimer's disease. At autopsy, heart contains yellow brown finely granular pigment which are/due to:**  
(Recent Question 2015)  
a. Hemosiderin-iron overload  
b. Lipochrome-wear and tear  
c. Glcoge-glycogen storage disorder  
d. Fat-athlerosclerosis
- 104. Stain for fat all except** (Recent Question 2014-15)  
a. Oil red O                              b. Sudan black  
c. Sudan III                              d. Congo red
- 105. Psammoma bodies features are all except:**  
(Recent Question 2015)  
a. Seen in meningioma  
b. Concentric whorled appearance  
c. Seen in papillary thyroid carcinoma  
d. Seen in teratoma
- 106. Dystrophic calcification is seen in:**  
(Recent Question 2014)  
a. Dying tissue                              b. Hypercalcemia  
c. Hyperprathyroidism                              d. Calcific metabolic disease
- 107. Steatosis means:** (Recent Question 2014)  
a. Fatty change  
b. Accumulation of triglyceride  
c. Accumulation of glycogen  
d. Accumulation of pigment

- 108. Dystrophic calcification is seen in all except:**  
(Recent Question 2014)  
a. Lymph node  
b. Lungs  
c. Kidneys  
d. Rheumatic valves
- 109. Oncocytes are found in all except:** (DNB June 11)  
a. Thyroid                              b. Pancreas  
c. Pituitary                              d. Pineal
- 110. Which of the following is a pathological calcification?**  
(DNB June 10)  
a. Suprasellar calcification  
b. Basal ganglia calcification  
c. Pineal body calcification  
d. Choroid calcification

## CELL AGING

- 111. Highest telomerase activity is seen in**  
(Recent Question 2015)  
a. Stem cells                              b. Somatic cells  
c. Germ cells                              d. Benign tumors
- 112. True regarding sirtuin functions** (Recent Question 2015)  
a. Decrease metabolic activity  
b. Reduce apoptosis  
c. Stimulate protein folding  
d. Inhibit harmful effects of oxygen free radicals
- 113. Werner disease is associated with?**  
(Recent Question Aug 13)  
a. Intestinal polyps  
b. Multiple cancer  
c. Lax joints  
d. Premature ageing
- 114. Sirtuin is associated with?** (DNB Nov. 12 Pattern)  
a. Cancer                              b. Diabetes  
c. Ageing                              d. All of the above
- 115. Which of the following is associated with aging** (AIIMS May 10)  
a. Reduced cross linkages in collagen  
b. Increased free radical injury  
c. Decreased Somatic mutations in DNA  
d. Increased superoxide dismutase levels

## FREE RADICALS

- 116. Organelle where  $H_2O_2$  is produced and destroyed is**  
(Recent Question 2016-17)  
a. Peroxisome  
b. Lysosome  
c. Golgi body  
d. Ribosome
- 117. The enzyme that protects brain from free radical injury:**  
(Recent Question 2015)  
a. Superoxide dismutase  
b. Catalase  
c. Glutathione peroxidase  
d. Monoamine oxidase
- 118. Which of the following is true about glutathione & glutathione peroxidase:** (PGI May 2015)  
a. Act as scavenger of free radicle  
b. Glutathione has anti-oxidant property  
c. Reduced glutathione can chemically detoxify  $H_2O_2$   
d. Oxidized glutathione can chemically detoxify  $H_2O_2$



**119. Which of these is not responsible for removal of free radicals?**  
(Recent Question 2015)

- Catalase
- Superoxide Dismutase
- NADPH oxidase
- Glutathione peroxidase

**120. Pathologic Effects of Free Radicals are all except:**  
(Recent Question 2015)

- Lipid peroxidation in membranes.
- Oxidative modification of proteins
- Single- and double-strand breaks in DNA
- Synthesis of new proteins



## Answers with Explanations

**1. Ans. (c) Occurs at stem cells level**

Option a & b are false.

- Cell adaptations are reversible & b both epithelial & mesenchymal

Option c is true.

- Metaplasia is change in phenotype of cells due to stem cell reprogramming

Option d is false.

- Most common type of metaplasia is squamous metaplasia

**2. Ans. (b) Barrets** (Ref: Robbins 9th/pg 37-38; 8th/pg 10)

**3. Ans. (b) DNA content same as in normal cells**

(Ref: Robbins 9th/pg 34; 8th/pg 6, Rev Esp Cardiol. 2006; 59:473-86. - Vol. 59)

**Hypertrophy:** Characterized by an increment in cardiomyocyte size, with increased protein synthesis and changes in the organization of the sarcomeric structure. DNA content also increases so b is false

**4. Ans. (d) Pleomorphism present**

(Ref: Robbins 9th/pg 37-38)

Option a - true. All the adaptations are reversible.

Option b - true. loss of polarity is a f/o dysplasia

Option c - true - **Mechanism of metaplasia: Reprogramming of stem cells<sup>o</sup>** in normal tissues or of **undifferentiated mesenchymal cells present in connective tissue.**

**5. Ans. (c) Metaplasia**

(Ref: Robbins 9th/pg 37-38; 8th/pg 10)

Increase in the number of goblet cells in the non-respiratory terminal bronchiole is an example of intestinal metaplasia.

**6. Ans. (c) Columnar to squamous** (Ref: R 9th/pg 37-38)

In smokers, pseudostratified ciliated columnar epithelium of respiratory tract changes to squamous epithelium

**7. Ans. (a) Metaplasia**

(Ref: Harshmohan Pathology For Dental 4th ed, pg 81)

In Vitamin-A deficiency, squamous metaplasia of conjunctiva leading to xerophthalmia.

**8. Ans. (a) Increase in number of cells**

(Ref: R 9th/pg 35-36)

**9. Ans. (c) Necrosis** (Ref: Robbins 9th/pg 34; 8th/pg 6)

Necrosis is cell death all others are cell adaptations

**10. Ans. (a) Atrophy** (Ref: Robbins 9th/pg 36-37)

**Atrophy: decrease in cell size and number.<sup>o</sup>**

**11. Ans. (a) Stem cells** (Ref: 9th/pg 36; 8th/pg 8)

**Mechanism of metaplasia: Reprogramming of stem cells<sup>o</sup>** in normal tissues or of **undifferentiated mesenchymal cells present in connective tissue.<sup>o</sup>**

**12. Ans. (d) Dense Mitochondrial deposit**

(Ref: Robbins 9th/pg 42)

**13. Ans. (a, b, d); a. Lysosomal rupture; b. Pyknosis; d. Severe mitochondrial dysfunction**

Bleb formation on membrane which are cytoplasmic blebs are a feature of reversible injury.

**14. Ans. (c) Lysosomes**

**15. Ans. (a) Ca**

(Ref: Gross GJ, Kersten JR, Warltier DC. Mechanisms of postischemic contractile dysfunction. Ann Thorac Surg. 1999; 68: 1898-1904.)

**Mediators of reperfusion injury are: Oxygen Free Radicals, Endothelial Dysfunction and Microvascular Injury, Alterations in Calcium Handling**

**16. Ans. (b) Cell shrinkage** (Ref: Robbins 9th/38; 8th/pg 17)

Features of **reversible cell injury** seen in **light microscopy** are **cellular swelling and fatty change**. Cell shrinkage is a feature of apoptosis (cell death)

**17. Ans. (b) Decreased ATP** (Ref: Robbins 9th/38; 8th/pg 17)

**Earlier molecular changes in cellular injury are decreased O<sub>2</sub> to issue leading to decreased oxidative phosphorylation leading to decreased ATP**



## 18. Ans. (b) Cell swelling

(Ref: Robbins 9th/pg 46; 8th/pg 12)

**Cloudy swelling<sup>Q</sup>** - earliest morphological change in reversible cell injury<sup>Q</sup> due to **accumulation of water intracellularly**

## 19. Ans. (d) Moth eaten appearance of cytoplasm

(Ref: Robbins 9th/pg 40; 8th/pg 12)

Necrotic cells have a more glassy, homogeneous appearance, mostly because of the loss of glycogen particles.

When enzymes have digested cytoplasmic organelles, the cytoplasm becomes vacuolated and appears "moth-eaten."

## 20. Ans. (b) Amorphous densities in mitochondrial matrix

(Ref: Robbins 9th/pg 40; 8th/pg 12)

Amorphous densities in mitochondria are features of irreversible cell injury-option b is false

## 21. Ans. (a) Neutrophils

(Ref: <http://www.hopkinsmedicine.org/mcp/Education/300.713%20Lectures/Infectious.pdf>)

- Acute response to infection is characterized by neutrophils
- Chronic response to infection is characterized by infiltrate by macrophages and monocytes
- Pseudomonas is an exception in that its chronic response is also characterized by neutrophils

## 22. Ans. (b) Neurons (Ref: Robbins 9th/pg 130; 8th/pg 129)

- **Tissue vulnerability to hypoxia-**
- **Neurons** undergo irreversible damage **3 to 4 minutes<sup>Q</sup>** of ischemia
- **Myocardial cells** die after only **20 to 30 minutes<sup>Q</sup>** of ischemia
- **Fibroblasts** within myocardium remain viable even after many hours of ischemia

## 23. Ans. (a) Mitochondrial densities

(Ref: R 9th/pg 42; 8th/pg 19)

**Features of irreversible injury include:**

**Severe swelling of mitochondria<sup>Q</sup>:** Large, flocculent, amorphous densities<sup>Q</sup> develop in mitochondrial matrix (**Increased Ca<sup>2+</sup> influx**)<sup>Q</sup>

## 24. Ans. (b) Cell membrane (Ref: R 9th/pg 42; 8th/pg 9,19)

### Myelin Figures<sup>Q</sup>

- Rolled-up or scroll-like arrangement of a **lipid bilayer within a cell<sup>Q</sup>**
- Derived from **damaged cell membranes<sup>Q</sup>**

## 25. Ans. (c) Flocculent amorphous densities in the mitochondria

(Ref: Robbins 9th/pg 42; 8th/pg 19; Refer to Ans 29)

Please note: myelin figures first appear in reversible cellular injury and become more pronounced in irreversible cell

injury. Hence if this q was asked in PGI, answer should be both c and d. but if we have to mark one..mark c.

## 26. Ans. (b) Apoptosis

(Ref: Robbins 9th ed/pg 53)

## 27. Ans. (c) Necrosis

(Ref: Robbins 9th ed/pg 50)

## 28. Ans. (b) p53 and caspases

(Ref: to question number 36 and 37)

## 29. Ans. (c) Organelle swelling

### Morphology of Apoptosis

- **Cell shrinkage<sup>Q</sup>**
- **Chromatin condensation:** most characteristic feature<sup>Q</sup> of apoptosis
- **Formation of cytoplasmic blebs and apoptotic bodies<sup>Q</sup>**
- **Plasma membrane** remains **intact<sup>Q</sup>** during apoptosis
- Apoptosis in contrast to necrosis **doesn't elicit inflammation<sup>Q</sup>**.

## 30. Ans. (b) Inflammasome

## 31. Ans. (a, e) a. BCL; e. MCL

## 32. Ans. (a) Caspase 1

## 33. Ans. (d) Both a & b

## 34. Ans. (c) Fibrinoid necrosis

## 35. Ans. (b) BAK

BAK and BAX are proapoptotic genes.

## 36. Ans. (a, b) a. Tumor suppression; b. Proapoptotic

(Ref: Robbins 9th/pg 53)

P53 is tumor suppressor gene (see neoplasia chapter) and also induces Bax which causes apoptosis

## 37. Ans. (a, d) a. They are enzyme starting apoptosis; d. They are proteases which cause cellular death in apoptosis found in irreversible cell damage

(Ref: Robbins 9th/pg 53)

Caspases are cysteine proteases that cleave after aspartic acid. These proteins have crucial role in apoptosis

## 38. Ans. (a) C is showing necrosis (Ref: Robbins 9th/pg 53)

C is showing smudged pattern S/O necrosis

## 39. Ans. (b, c) b. Caspase independent; c. Failure of activation of caspase 8 (Ref: Robbins 9th/pg 59)

Necroptosis occurs when caspase 8 cannot be activated, in such cases apoptosis is mediated by RIP1 and RIP3 proteins (caspase independent)





40. Ans. (c) **BCL-XL**
41. Ans. (d) **Pro-apoptotic protein**
42. Ans. (a) **FLIP**
43. Ans. (a) **Fatty** (Ref: Robbins 9th/pg 43; 8th/pg 15,16)  
**Fat necrosis:** Seen in **acute pancreatitis, breast, omentum**
44. Ans. (a) **Inflammation present**  
(Ref: 9th/pg 53; 8th/pg 25,26)  
Apoptosis in contrast to necrosis **doesn't elicit inflammation**<sup>Q</sup>
45. Ans. (c) **Cell swelling** (Ref: Robbins 9th/pg 46; 8th/pg 12)  
Earliest change in any form of cell injury is cellular swelling due to accumulation of water intracellularly.
46. Ans. (d) **Isoprenoid** (Ref: Korean J Physiol Pharmacol. 2013 Feb;17(1):43-50.)
- **Palmitic acid** (PAM), one of the most common saturated fatty acid (SFA) in animals and plants, has been shown to induce **apoptosis**
  - Farnesol (FOH) and other **isoprenoid** alcohols induce **apoptosis** in various carcinoma cells and inhibit tumorigenesis in several in vivo models.
47. Ans. (a) **Caspase 3** (Ref: Robbins 9th/pg 56; 8th/pg 26)  
There are two executionary caspases 3 and 6.  
Most important executionary caspase is caspase 3
48. Ans. (a) **Dry gangrene**  
(Ref: A Practical Manual of Diabetic Foot Care By Michael E. Edmonds; pg 144)  
The combined changes of angiopathy and neuropathy give rise to Diabetic Foot which usually leads to **dry gangrene** in patients of diabetes. In patients with uncontrolled diabetes, however wet gangrene can also ensue.
49. Ans. (d) **Apoptosis**  
(Ref: Robbins 9th/pg 53; Robbins 9th/pg 29-30)  
Apoptotic cells express **phosphatidylserine/thrombospondin**<sup>Q</sup> on the outer layer of plasma membrane because of which these cells are recognized by the dye **Annexin V**
50. Ans. (b) **Mitochondria**  
(Ref: Robbins 9th/pg 38; 8th/pg 17)
51. Ans. (c) **Diabetic glomerulosclerosis** (Ref: R 9th/pg 44)  
**Fibrinoid necrosis:** Seen in **PAN, malignant hypertension, acute rheumatic fever**
52. Ans. (b) **Apoptosis**  
(Ref: Robbins 9th/pg 53; R 9th/pg 29-30)

DNA breakdown (**Internucleosome cleavage by endonuclease into 200 bp oligonucleosomes is a characteristic**)<sup>Q</sup> at specific sites can be detected by '**step ladder pattern**' on gel electrophoresis or **TUNEL (TdT mediated d-UTP Nick End Labelling) technique**

53. Ans. (c) **Bax** (Ref: Robbins 9th/pg 55; 8th/pg 28)  
BAD, BIM, BID, Puma, and Noxa are sensors of apoptosis
54. Ans. (d) **Type 1 TNF receptor** (Ref: Robbins 9th/pg 53)  
**Lack of survival signals (option a) and DNA damage (option b and c) trigger intrinsic pathway of apoptosis**  
**Extrinsic pathway is activated by** Death receptors are members of the **TNF receptor family**
55. Ans. (a) **Autoimmune disease**  
(Ref: AGING, Vol 4, No 5, pp 330-349, Del Puerto HL, Martins AS, Milsted A; et al. (2011). "Canine distemper virus induces apoptosis in cervical tumor derived cell lines". Virol. J. 8 (1): 334)  
Inhibition of apoptosis can result in a number of cancers, autoimmune diseases, inflammatory diseases, and viral infections.  
Defective Neuronal apoptosis plays an important role in neurodegenerative disorders like **Parkinson's disease (PD), Alzheimer's disease (AD)**
56. Ans. (c) **Necroptosis** (Ref: Robbins 9th/pg 58-59)  
**Necroptosis:** Genetically programmed signal transduction event **without caspase activation** → "**caspase-independent**" programmed cell death, it is dependent on signaling by the **RIP1 and RIP3 complex**
57. Ans. (d) **Pyroptosis** (Ref: Robbins 9th/pg 58-59)  
**Pyroptosis:** Promotes the activation of **inflammasome** which activates **Caspase-1**<sup>Q</sup> which releases biologically active form of IL-1 from the precursor.
58. Ans. (d) **Fibrin deposition**  
(Ref: R 9th/pg 44; 8th/pg 15,16)  
**Fibrinoid necrosis:** Deposits of "**antigen -antibody complexes**"<sup>Q</sup> & **fibrin** that has leaked out of vessels, result in a bright **pink and amorphous appearance in H & E stains**
59. Ans. (d) **Smear pattern in electrophoresis**  
(Ref: R 9th/pg 53)  
*Smear pattern is seen in necrosis*
60. Ans. (a) **Bax** (Ref: Robbins 9th/pg 55; 8th/pg 28)
- **Proapoptotic:** Apaf1, cytochrome c, Bak, Bax, Bim, AIF, P53, Caspases, CD95 (FAS), TNF R1 and Smac/Diablo
  - **Antiapoptotic:** Bcl2, Bcl<sub>XL</sub>, Bcl<sub>X</sub>, FLIP, McL-1, IAP



61. Ans. (a) **Coagulative necrosis-Tuberculosis**

(Ref: Robbins 9th/pg 43; 8th/pg 15,16)

62. Ans. (d) **Diabetic glomerulosclerosis** (Ref: R 9th/pg 44)

63. Ans. (c) **Activation of caspases** (Ref: R 9th/pg 52, 53, 56)

Activation of caspases > caspases alone

64. Ans. (b) **Liquefactive necrosis** (Ref: Robbins 9th/pg 43)

Necrosis in **central nervous system**: Liquefactive necrosis

65. Ans. (b, c) **b. ↓ Apoptosis; c. ↑ Resistance of tumor to treatment** (Ref: Robbins 9th/pg 55; 8th/pg 28)

bcl-2: antiapoptotic molecule-option b is true  
Option c is true : resistance of tumour cells to therapy can be caused by a failure in the ability to initiate apoptosis  
Option d is false: bcl2 mutation though is hallmark of follicular carcinoma but is seen in many other cancers also  
Option e is false: tumor suppressor NF2 is disrupted in approximately half of all meningiomas

66. Ans. (a) **Inflammation is present**

(Ref: R 9th/pg 29, 30, 53)

67. Ans. (a) **Increase in lysosomal enzyme**

(Ref: Robbins 9th/pg 53; Robbins 9th/pg 29-30)

68. Ans. (c) **Cell membrane bound with organelles**

(Ref: Robbins 9th/pg 52; 8th/pg 25)

Apoptotic cells break up into fragments, called apoptotic bodies, which contain portions of the cytoplasm and nucleus. The plasma membrane of the apoptotic cell and bodies remains intact, but its structure is altered in such a way that these become “tasty” targets for phagocytes.

69. Ans. (b) **Necroptosis** (Ref: Robbins 9th/pg 58-59)

Necroptosis is triggered **by ligation of TNFR1 and Viral proteins of RNA and DNA viruses.**<sup>Q</sup>

70. Ans. (a) **Malignant HTN** (Ref: 9th/pg 44; 8th/pg 15,16)

71. Ans. (d) **Fibrinoid necrosis** (Ref: Robbins 9th/pg 44)

72. Ans. (b) **BCL-2** (Ref: Robbins 9th/pg 55; 8th/pg 28)

Anti-apoptotic.	Pro-apoptotic	Sensors
BCL2, BCL-XL, and MCL1	BAX and BAK	BAD, BIM, BID, Puma, and Noxa

73. Ans. (a) **Denaturation of protein**

(Ref: Robbins 9th/pg 50)

- **Coagulative necrosis: most common type of necrosis**
- **Occurs due to degenerated cytoplasmic proteins**

74. Ans. (d) **All** (Ref: Robbins 9th/pg 43; 8th/pg 15,16)

- Fat necrosis
- **Not a specific pattern of necrosis.**<sup>Q</sup>
- Signifies focal areas of fat destruction
- **Released lipases** split the triglyceride esters contained within fat cells.
- The **fatty acids**, so derived, combine with calcium to produce grossly visible **chalky-white areas (fat saponification)**<sup>Q</sup>
- Seen in **acute pancreatitis, breast, omentum**<sup>Q</sup>

75. Ans. (a) **Apaf 1**

(Ref: Robbins 9th/pg 54-55; 8th/pg 29)

Leakage of cytochrome c, other proteins form the inner mitochondrial membrane to cytosol

↓  
Cytochrome C binds to **APAF-1**<sup>Q</sup>

↓  
Activation of caspase-9-critical initiator caspase of the mitochondrial pathway

76. Ans. (c) **Inflammation**

(Ref: Robbins 9th/pg 53; 8th/pg 25,26)

- Apoptosis in contrast to necrosis **doesn't elicit inflammation**<sup>Q</sup>.

77. Ans. (b) **Extrinsic pathway of apoptosis**

(Ref: R 9th/pg 53)

**The extrinsic pathway of apoptosis** is activated by cross-linking members of the tumor necrosis factor (TNF) receptor superfamily, such as **CD95 (Fas) and death receptors DR4 and DR5**, by their receptors, Fas ligand or TRAIL (TNF-related apoptosis-inducing ligand), respectively.

78. Ans. (d) **Causes inflammation that damage surrounding cells** (Ref: Robbins 9th/pg 53; 8th/pg 25, 26)

**Blebbing** or **zeiosis** is the formation of blebs, it is seen in apoptotosis to form apoptotic bodies

79. Ans. (c) **Neurons** (Ref: The Central Nervous System in Pediatrics Critical Illness page 13)

80. Ans. (b) **Coagulative**

(Ref: Robbins 9th/pg 43; 8th/pg 15, 16)

MC type of necrosis in Myocarial Infacrction- coagulative

81. Ans. (a) **Mitochondria**

(Ref: Essential of apoptosis 2003:245)

**Mitochondria is the most important organelle involved in apoptosis initiation and regulation.**

82. Ans. (d) **Cytochrome C**

(Ref: R 9th/pg 53-54; 8th/pg 28, 29)



83. Ans. (d) **Bcl-2** (Ref: Robbins 9th/pg 55; 8th/pg 28)

84. Ans. (a) **Cell membrane intact** (Ref: Robbins 9th/pg 53)

- Most characteristic feature is condensation of nuclear chromatin
- Intact cell membrane and lack of inflammation differentiates apoptosis from necrosis
- Nuclear molding is seen in small cell carcinoma, lung
- Cytoplasmic eosinophilia- nonspecific finding. Occurs both in necrosis and apoptosis
- If we have to mark one: mark a since that occurs exclusively in apoptosis

85. Ans. (b) **Nuclear compaction** (Ref: Robbins 9th/pg 53)

Here student gets confused in 2 options b and c..

- On light microscopy, most characteristic feature is condensation of nuclear chromatin and not intactness of cell membrane
- Intact cell membrane and lack of inflammation differentiates apoptosis from necrosis.

86. Ans. (a) **Cellular swelling**

(Ref: R 9th/pg 53; 8th/pg 25,26)

**Cellular shrinkage and not cellular swelling is a feature of apoptosis**

87. Ans. (a) **TB** (Ref: Robbins 9th/pg 43'; 8th/pg 15,16)

Type of necrosis in TB: **Caseous necrosis.**

**Caseous necrosis is a variant of coagulative necrosis**

88. Ans. (a) **Organogenesis**

(Ref: <http://onlinelibrary.wiley.com/doi/10.1002/bdra.10090/abstract>)

Caspases are key mediators in the regulation and execution of apoptosis, a crucial part of the morphogenetic process during limb development.

89. Ans. (b, c, d); **b. Increase in caspases; c. Phosphatidyl serine has important role; d. Internucleosomal cleavage of nucleus**

(Ref: Robbins 9th/pg 56; 8th/pg 30)

- In healthy cells, phosphatidylserine is present on the inner leaflet of the plasma membrane, but in apoptotic cells this phospholipid "flips" out and is expressed on the outer layer of the membrane, where it is recognized by several macrophage receptors.
- Apoptosis occurs due to activation of enzymes called **caspases (cysteine proteases that cleave proteins after aspartic residues)**<sup>Q</sup>.
- Apoptotic cell exhibit a characteristic breakdown of DNA into large 50 to 300 kilobase piece. Subsequently there is **internucleosomal cleavage**<sup>Q</sup> of DNA into oligonucleosomes, in multiples of 180 to **200 base pairs**<sup>Q</sup>, by  $\text{Ca}^{2+}$  &  $\text{Mg}^{2+}$  dependent **endonucleases**<sup>Q</sup>. The fragments may be visualized by **agarose gel electrophoresis**<sup>Q</sup> as **DNA ladders**<sup>Q</sup>.

90. Ans. (e) **Mitochondrial membrane**

(Ref: Robbins 9th/pg 56)

Mitochondrial membrane permeabilization is hallmark of apoptosis

91. Ans. (d) **Myositis ossificans**

(Ref: Robbins 9th ed/pg 65)

92. Ans. (a, b, c); **a. Monckeberg's medial sclerosis; b. Papillary carcinoma thyroid; d. Meningioma**

Hyperparathyroidism and Vitamin D intoxication cause hypercalcemia and so are associated with dystrophic calcification.

93. Ans. (c) **Prussian blue**

(Ref: Complete review of pathology and Hematology 2nd ed; Praveen Kumar, Vandana Puri Annexure 4)

This is a straight forward question on the **stain used for Iron in tissue** (Hemochromatosis) which is Prussian blue stain.

94. Ans. (a, c, d) **a. Perl stain-Iron; c. Elastin-VG; d. Copper-rhodamine**

(Ref: Complete Review of Pathology and Hematology 2nd ed; Praveen Kumar, Vandana Puri Annexure 4)

95. Ans. (a) **Fat** (Ref: Robbins 9th/pg 62)

FAT accumulation in myocardium causes fatty change (tigered effect)

96. Ans. (d) **Seminoma** (Ref: Robbins 9th/pg 65; 8th/pg 38)

**Psammoma bodies are type of dystrophic calcification seen in - Meningioma, Papillary carcinoma of thyroid, ovary (serous carcinoma), prolactinoma, Somatostatinoma, mesothelioma**

97. Ans. (b) **Ferrocyanide to ferroferric cyanide**

(Ref: Dacie and Lewis Practical Haematology 11<sup>th</sup>ed, pg 338)

Hemosiderin reacts with potassium ferrocyanide to form a blue compound, ferri ferrocyanide; this reaction is the basis of a positive Prussian-blue (Perls') reaction.

98. Ans. (c) **Atherosclerosis**

(Ref: Robbins 9th/pg 65; 8th/pg 38)

Atheromatous plaque shows **Dystrophic calcification**  
**All others are causes of metastatic calcification**

99. Ans. (d) **Mitochondria**

(Ref: Robbins 9th/pg 62; 8th/pg 33)

Foam cells - cholesterol-laden macrophages

100. Ans. (c) **Lungs** (Ref: Robbins 9th/pg 65; 8th/pg 38)

Metastatic calcification is seen most commonly in lungs followed by kidneys



**101. Ans. (b) Parkinsons disease**

(Ref: <https://en.wikipedia.org/wiki/Proteopathy>)

Answer should be **none**; but if we have to mark one go for b. as a, c, d are well known proteinopathies whereas b is still in phase of research.

The proteopathies (also known as **proteinopathies**, **protein conformational disorders**, or **protein misfolding diseases**) include - Alzheimer's disease, Prion diseases, Parkinson's disease, Huntington's disease, Cystic Fibrosis, AL (light chain) amyloidosis

**102. Ans. (d) Mitochondria**

(Ref: Robbins 9th/pg 65; 8th/pg 38)

**103. Ans. (b) Lipochrome-wear and tear**

(Ref: Robbins 9th/pg 64; 8th/pg 37)

- Lipofuscin: **Perinuclear**<sup>Q</sup> brown-coloured pigment, Seen in **aging**<sup>Q</sup>, **protein energy malnutrition**<sup>Q</sup> and **cancer** **cahcxia**.

**104. Ans. (d) Congo red** (Ref: Bancroft theory and practice of histological stains 7th ed, pg 165)

**105. Ans. (d) Seen in teratoma** (Ref: R 9th/pg 65; 8th/pg 38)

- Psammoma bodies (PBs) are concentric lamellated calcified structures, observed most commonly in papillary thyroid carcinoma (PTC), meningioma, and papillary serous cystadenocarcinoma of ovary
- They represent a process of dystrophic calcification

**106. Ans. (a) Dying tissue** (Ref: Robbins 9th/pg 65; 8th/pg 38)

**107. Ans. (a, b) a. Fatty change; b. Accumulation of triglyceride** (Ref: Robbins 9th/pg 62; 8th/pg 33)

**Steatosis** (Fatty Change): accumulations of triglycerides within parenchymal cells

**Organs** involved-liver, heart, muscle, and kidney

**108. Ans. (b, c) b. Lungs; c. Kidneys** (Ref: Robbins 9th/pg 65)

Dystrophic calcification occurs in dead and damaged tissues and not in normal organs

**109. Ans. (d) Pineal**

(Ref: Rosai and Ackerman's Surgical Pathology 10ed volume 2.. Pathol Annu. 1992; 27(Pt 1):263-304)

- **Oncocyte** is an epithelial cell characterized by an **excessive amount of mitochondria**
- **Characterized by** abundant acidophilic, granular cytoplasm
- **Other names:** *Oxyphilic cell*, *Askanazy cell*, *Hürthle cell* (thyroid gland only), *Apocrine metaplasia* (breast gland only)
- They are present in benign tumors of **salivary glands, thyroid, parathyroid, kidney, lung and pituitary**

**110. Ans. (a) Suprasellar calcification**

(Ref: J Pediatr Neurosci. 2011 Oct; 6 (Suppl1): S46-S55)

Suprasellar calcifications are always pathological  
Other 3 can be physiological as well

**111. Ans. (c) Germ cells** (Ref: Robbins 9th/pg 67; 8th/pg 40)

**112. Ans. (a) Decrease metabolic activity**

(Ref: R 9th/pg 67, 68)

- **Sirtuins:** These include proteins that inhibit metabolic activity<sup>Q</sup>, reduce apoptosis<sup>Q</sup>, stimulate protein folding<sup>Q</sup>, inhibit the harmful effects of oxygen free radicals<sup>Q</sup> and activating DNA repair enzymes through deacetylation<sup>Q</sup>

**113. Ans. (d) Premature ageing**

(Ref: R 9th/pg 66; 8th/pg 40)

Syndromes associated with defective DNA repair	
<b>Defective DNA helicase</b>   <b>Werner syndrome<sup>Q</sup></b> (Premature aging)	<b>Defective DNA repair syndromes<sup>Q</sup></b>   <b>BAX-F</b> Bloom Syndrome <sup>Q</sup> Ataxia Telangiectasia <sup>Q</sup> Xeroderma Pigmentosa <sup>Q</sup> Fanconi Anemia <sup>Q</sup>

**114. Ans. (d) All of the above** (Ref: Robbins 9th/pg 67,68)

**Sirtuins**

- Family of **NAD-dependent protein deacetylases**.
- Sirtuin-6<sup>Q</sup> promotes the expression of genes whose products increase longevity.
- Since these proteins increase longevity, activate DNA repair enzymes so prevent against cancer  
Sirtuins also increase insulin sensitivity and glucose metabolism, and may be targets for the treatment of diabetes.

**115. Ans. (b) Increased free radical injury**

(Ref: Robbins 9th/pg 66-68; 8th/pg 39, 40)

**116. Ans. (a) Peroxisome**

**117. Ans. (a) Superoxide dismutase**

(Ref: Robbins 9th/pg 48)

**Superoxide dismutase is an antioxidant and known to protect brain from free radical injury**

**118. Ans. (a, b, c); a. Act as scavenger of free radicle; b. Glutathione has anti-oxidant property; c. Reduced glutathione can chemically detoxify H<sub>2</sub>O<sub>2</sub>**

(Ref: Robbins 9th/pg 48; 8th/pg 13,14)

- Glutathione peroxidase:  $\text{H}_2\text{O}_2 + 2\text{GSH} \rightarrow \text{GSSG} + 2\text{H}_2\text{O}$
- Reduced glutathione can chemically detoxify  $\text{H}_2\text{O}_2$





- Intracellular ratio of oxidized glutathione (GSSG) to reduced glutathione (GSH) is a reflection of the oxidative state of the cell
- Important indicator of the cell's ability to detoxify ROS.

#### 119. Ans. (c) **NADPH oxidase**

(Ref: R 9th/pg 48; 8th/pg 13,14)

Antioxidants

Enzymes	Non enzymes
<ul style="list-style-type: none"> <li>• <b>Catalase</b>- Present in <b>peroxisomes</b><sup>Q</sup> decomposes <math>H_2O_2</math></li> <li>• <b>Superoxidase dismutases</b>: <math>O_2^-</math> to <math>H_2O_2</math></li> <li>• <b>Manganese-SOD</b>, which is localized in <b>mitochondria</b>,</li> <li>• <b>Copper-zinc-SOD</b>, which is found in the <b>cytosol</b>.</li> </ul>	<p><b>Antioxidants</b>- vitamins E, A and C glutathione in the cytosol.</p> <p><b>Transferrin, ferritin, lactoferrin, and ceruloplasmin</b><sup>Q</sup> - Minimise the reactivity of metals by binding with them</p>
<p><b>Glutathione peroxidase</b>: <math>H_2O_2 + 2GSH \rightarrow GSSG + 2H_2O</math></p> <ul style="list-style-type: none"> <li>• Important indicator of the cell's ability to detoxify ROS<sup>Q</sup></li> </ul>	

#### **NADPH oxidase**

Oxidizes **NADPH**<sup>Q</sup> (reduced nicotinamide-adenine dinucleotide phosphate) and, in the process, **reduces oxygen to superoxide**.<sup>Q</sup>

#### 120. Ans. (d) **Synthesis of new proteins** (Ref: R 9th/pg 49)

Pathologic Effects of Free Radicals are

- Lipid peroxidation in membranes- result in extensive membrane damage.
- Oxidative modification of proteins
- Single- and double-strand breaks in DNA cross-linking of DNA strands, and formation of adducts

# Inflammation and Repair

## Key Points

- » **Hallmark<sup>o</sup>** of acute inflammation: **Increased vascular permeability**
- » Endothelial cell expression of **E-selectin is a hallmark of acute cytokine mediated inflammation**
- » **Tissue destruction: Hallmark of chronic inflammation**
- » **Granulation tissue<sup>o</sup>** is the hallmark of the fibrogenic repair.
- » **Chemotaxis: Unidirectional movement<sup>o</sup>** of the leukocytes **towards site of injury** along a **chemical gradient**
- » **Opsonisation:** Phagocytosis requires **polymerization of actin filaments**
- » Pinocytosis is due to **internalization into clathrin-coated pits**
- » **Catarrhal inflammation: Commonest type of inflammation**
- » Maximum strength gained 70% of strength of normal skin

## Key Recent Updates


- » Albumin is **negative** acute phase reactant
- » Emperipolesis is seen in Rosai Dorfman disease
- » **NETosis** is neutrophil cell death pathway which is protective to body but can sometimes lead to autoimmune diseases.




## OVERVIEW OF INFLAMMATION

- **Definition:** Response of the vascularized tissues to an injurious stimulus is called inflammation.
- **Types:** Acute inflammation & Chronic inflammation

Features	Acute	Chronic
<b>Onset</b>	Early: minutes to hours	Slow: days to months
<b>Cellular infiltrate</b>	<b>Neutrophils<sup>Q</sup></b>	<b>Mononuclear cells</b> -Monocytes/macrophages, lymphocytes, plasma cells
<b>Exceptions</b>	<b>Acute typhoid fever (neutropenia)</b>	Chronic <b>pseudomonas</b> infection (neutrophilia)
<b>Tissue injury</b>	Mild and self-limited	Severe and progressive



Neutrophils (arrow)



Lymphocyte (green arrow) and macrophages (yellow arrow)

## High Yield Facts



### Celsus four cardinal signs of inflammation

- **Rubor:** redness (due to **vasodilation<sup>Q</sup>** of small blood vessels)
- **Tumor:** swelling (due to exudation of fluid)
- **Calor:** heat (due to increased blood flow i.e **hyperemia due to vasodilation<sup>Q</sup>**)
- **Dolor:** pain (due to stretching of tissue due to edema and chemical mediator **bradykinin<sup>Q</sup>**)
- **Function laesa**-loss of function (Added by Virchow)

## ACUTE INFLAMMATION

### Three Major Components

1. **Vasodilation**
2. **Increased vascular permeability- the hallmark<sup>Q</sup>**
3. **Emigration of leucocytes** from microcirculation and their activation


### Vascular Changes

- **Vasoconstriction:** **First<sup>Q</sup>** change in the blood vessels, **transient<sup>Q</sup>** in nature.
- **Vasodilation:** **Second** change in the blood vessels lasting for a **longer duration<sup>Q</sup>**. It first occurs in **arterioles**. Results in **increased blood flow** → **redness (rubor)** & sensation of **warmth (calor)**
- **Increased permeability:** **Hallmark<sup>Q</sup>** of acute inflammation; **Maximally seen in the venules.<sup>Q</sup>**

### Mechanism of Increased Vascular Permeability

Name	Mechanism	Caused by	Affected Vessels
<b>Immediate transient response<sup>Q</sup> (ITR)</b> Rapid short lived (15–30 min) reversible	<i>Formation of endothelial gaps</i>	<ul style="list-style-type: none"> <li>• Histamine<sup>Q</sup>, bradykinin</li> <li>• Contraction of endothelial cell cytoskeleton<sup>Q</sup></li> </ul>	Venules
<b>Immediate sustained response<sup>Q</sup> (ISR)</b> Rapid & long lived	<i>Direct endothelial injury</i>	<ul style="list-style-type: none"> <li>• Burns</li> <li>• Endothelial cell necrosis and detachment<sup>Q</sup></li> </ul>	Venules, capillaries and arterioles
<b>Delayed prolonged leakage<sup>Q</sup> (DPL)</b> (appears after 2–12 hr)	<i>Mild endothelial damage</i> e.g., Late-appearing sunburn <sup>Q</sup>	<ul style="list-style-type: none"> <li>• Thermal and radiation injury<sup>Q</sup></li> </ul>	Venules and capillaries
<b>LMI (Leukocyte-mediated endothelial injury) late &amp; sustained</b>	<i>WBC damage</i>	<ul style="list-style-type: none"> <li>• Activated leukocytes</li> <li>• Endothelial injury or detachment<sup>Q</sup></li> </ul>	<b>Venules (mostly);</b> pulmonary and glomerular capillaries
<b>Increased transcytosis</b>	<i>Formation of vesiculo-vacuolar organelles<sup>Q</sup></i> (Intercellular channels)	<ul style="list-style-type: none"> <li>• Histamine and VEGF</li> </ul>	Venules

Note: Endothelial cell retraction is same as endothelial cell contraction (Robbins 10th ed)



## Cellular Changes


**Leukocyte recruitment** to sites of inflammation from vessel lumen can be divided into the following steps:

- **In the lumen:** Margination, rolling, and adhesion to endothelium
- **Across lumen:** Migration across the **endothelium and vessel wall**<sup>Q</sup>
- **Outside the lumen:** Migration in the **tissues toward a chemotactic stimulus**<sup>Q</sup>



## High Yield Facts

- Formation of endothelial gaps (**Immediate transient response**) is the **most common mechanism**<sup>Q</sup> for increased permeability.
- Most important immediate mediator responsible for **Immediate transient response**<sup>Q</sup> is **histamine**<sup>Q</sup>
- The loss of fluid and increased vessel diameter leads to slower blood flow, concentration of red cells in small vessels, and increased viscosity of the blood.



## In the Lumen

### Margination

- Movement of the leukocytes **principally neutrophils** towards the periphery of the blood vessel
- Occurs due to **increased vascular permeability**<sup>Q</sup>
- **Vasodilatation and selectins** have important role

### Rolling

- **Transient adhesion** of **leukocytes** with the **endothelial cells**
  - Mediated by **selectins**.<sup>Q</sup>

Molecule	Distribution	Activated by	Ligand (interact with)
<b>P-selectin (CD62P)</b>	Platelets <sup>Q</sup> and Endothelium <sup>Q</sup>	(TNF, IL-1), histamine, or thrombin	<b>Sialyl-Lewis X</b> expressed on leucocytes
<b>E-selectin (CD62E)</b>	Endothelium <sup>Q</sup>	(TNF, IL-1)	<b>Sialyl-Lewis X</b> expressed on leucocytes
<b>L-selectin (CD62L)</b>	<b>Leucocytes</b> <sup>Q</sup> : Neutrophils, monocytes and T cells (naive and central memory), B cells (naive)	—	<b>Sialyl-Lewis X/PNAd</b> on Glycoprotein adhesion molecules ( <b>GlyCAM-1</b> ), <b>CD34</b> , <b>MAdCAM-1</b> <sup>Q</sup> expressed on endothelium






## Adhesion

- **Firm attachment** of the **leukocytes to the endothelial cells**
- Mediated by **heterodimeric leukocyte surface proteins called integrins**.

Integrin	Molecule of Integrin	Ligand on endothelium
$\beta 1$ -integrins	VLA molecules (VLA-4 (CD49a/CD29) or $\alpha_4\beta 7$ (CD49D/CD29)	VCAM-1 <sup>Q</sup>
$\beta 2$ -integrins	LFA-1 or Mac-1(CD11a/CD18)	ICAM-1 <sup>Q</sup>



## High Yield Facts

- **Pavementing**: Endothelium appears to be lined by white cells. This is due to margination.
  - Endothelial cell expression of **E-selectin is a hallmark of acute cytokine-mediated inflammation.<sup>Q</sup>**
  - **P Selectin is stored in Endothelium weibel palade bodies<sup>Q</sup> and Platelets  $\alpha$  granules<sup>Q</sup>**
  - **L-selectin helps in Lymphocytes homing to high endothelial venules<sup>Q</sup>**
- Integrins** are transmembrane receptors that are the bridges for cell-cell and cell-extracellular matrix (ECM) interactions. Few ligands for integrins:
- **VCAM & ICAM** – on the endothelium
  - **GPIIb/IIIa**, an integrin on the surface of blood platelets
  - **Fibronectin, vitronectin, collagen, and laminin.**

## Across the Lumen

### Leukocyte Migration through Endothelium: Transmigration or Diapedesis<sup>Q</sup>


- Occurs mainly in **postcapillary venules.<sup>Q</sup>**
- Most important molecule: **PECAM-1 (platelet endothelial cell adhesion molecule) or CD31.<sup>Q</sup>**
- After traversing the endothelium, leukocytes pierce basement membrane, probably by secreting **collagenases,<sup>Q</sup>** and **enter extravascular tissue.**

## Outside the Lumen

### Migration in the Tissues Toward a Chemotactic Stimulus: Chemotaxis

- **Unidirectional movement<sup>Q</sup>** of the leukocytes **towards site of injury along a chemical gradient.**
- **Bind to G-protein coupled receptors<sup>Q</sup>** on the surface of leukocyte
- Causes **actin polymerization** and all movements.

### Chemotactic Agents



## PHAGOCYTOSIS AND CLEARANCE OF THE OFFENDING AGENT

Phagocytosis involves following three sequential steps



## Recognition and Attachment

- Particles to be ingested by leukocytes (microbes and dead cells) are **recognized by receptors present on the leucocyte surface** like:<sup>Q</sup>

<b>Mannose receptors</b>	<ul style="list-style-type: none"> <li>Recognizes <b>microbes and not host cells</b>.<sup>Q</sup></li> <li>Part of molecules found on <b>microbial cell walls</b>.</li> <li>Mammalian glycoproteins and glycolipids contain <b>terminal sialic acid or N-acetylgalactosamine</b><sup>Q</sup></li> </ul>
<b>Scavenger receptors</b>	<ul style="list-style-type: none"> <li>Bind microbes and oxidized LDL particles<sup>Q</sup> that can <b>no longer bind to LDL receptor</b></li> </ul>
<b>Macrophage integrins</b>	<ul style="list-style-type: none"> <li>Mac-1 integrins (<b>CD11b/CD18</b>)- bind microbes for phagocytosis</li> </ul>

## Opsonization

- Coating of the bacteria** so that they are **easily phagocytosed** by the **leucocytes**.
- Chemicals causing opsonization are called **opsonins**
- Opsonization **increases efficiency** of phagocytosis.
- Phagocytosis can occur without opsonization also.<sup>Q</sup>
- Opsonins include:**
  - C3b<sup>Q</sup>, Fc fragment of IgG antibodies<sup>Q</sup>, Plasma lectins-mannose-binding lectin<sup>Q</sup> and Fibrinogen<sup>Q</sup>, C reactive protein<sup>Q</sup>**

## Engulfment

- Occurs after the particle is bound to phagocyte receptors
- Leucocyte **cytoplasm** forms pseudopods and leads to formation of phagosome
- Plasma membrane pinches off to form a vesicle
- Fuses with a lysosomal granule, resulting in discharge of the granule's contents into the phagolysosome




### High Yield Facts


- Phagocytosis requires **polymerization of actin filaments**<sup>Q</sup>
- Pinocytosis is due to **internalization into clathrin-coated pits**
- Pinocytosis is **not dependent on the actin cytoskeleton**.<sup>Q</sup>

## INTRACELLULAR DESTRUCTION OF MICROBES AND DEBRIS

Final step in the elimination of infectious agents and necrotic cells.



## Oxygen-dependent Killing



### High Yield Facts

- Respiratory burst** is also called oxidative burst
- Respiratory burst** results in rapid release of **reactive oxygen species**<sup>Q</sup> from different types of cells
- NADPH oxidase** is also called **phagocyte oxidase** or **phagocyte NADPH oxidase** (PHOX)

## Enzymes Involved in Respiratory Burst

Enzyme	Function
<b>NADPH oxidase</b>	Chiefly <sup>Q</sup> responsible for the formation of <b>hydrogen peroxide</b> <sup>Q</sup>
<b>Catalase (peroxisomes)</b> <sup>Q</sup>	Degrades <b>hydrogen peroxide</b> into water and oxygen. <sup>Q</sup>
<b>Superoxide Dismutase (SOD)</b>	Causes conversion of <b>superoxide ion</b> into <b>hydrogen peroxide</b> . <sup>Q</sup>
<b>Glutathione peroxidase</b>	Causes conversion of <b>reduced glutathione</b> to its homodimer. <sup>Q</sup>

## Oxygen Independent Killing

### Lysosomal Enzymes and Other Lysosomal Proteins

- Neutrophils and monocytes** contain **lysosomal granule**<sup>Q</sup> that contribute to **microbial killing**
- Neutrophils have **two main types of granules**



Primary (azurophilic) granules	Secondary (specific) granules
Secreted at <b>higher concentration</b> of agonists <sup>Q</sup>	Secreted at <b>lower concentration</b> of agonists <sup>Q</sup>
<ul style="list-style-type: none"> <li>• Myeloperoxidase<sup>Q</sup></li> <li>• Bactericidal factors (lysozyme and defensins)</li> <li>• Acid hydrolases<sup>Q</sup></li> <li>• Neutral proteases: <ul style="list-style-type: none"> <li>▪ Elastase</li> <li>▪ Cathepsin G</li> <li>▪ Nonspecific collagenases<sup>Q</sup></li> <li>▪ Proteinase 3</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• Lysozyme</li> <li>• Collagenase<sup>Q</sup> (Type IV)</li> <li>• Gelatinase</li> <li>• Lactoferrin<sup>Q</sup></li> <li>• Plasminogen activator</li> <li>• Histaminase</li> <li>• Alkaline Phosphatase</li> </ul>

**\*Tertiary granules:** Develop during chemotaxis, contain **gelatinases<sup>Q</sup>** and **acid hydrolases<sup>Q</sup>**

### Other Microbicidal Granule Contents

Granule Contents	Characteristics
<b>Defensins</b>	Cationic <b>arginine-rich granule peptides<sup>Q</sup></b> that are toxic to microbes
<b>Cathelicidins</b>	<ul style="list-style-type: none"> <li>• Antimicrobial proteins found in neutrophils</li> <li>• Vitamin D upregulates genetic expression of Human cathelicidin antimicrobial protein (Hcap18)<sup>Q</sup> <small>RG<sup>Q</sup></small></li> </ul>
<b>Lysozyme</b>	<ul style="list-style-type: none"> <li>• Hydrolyzes <b>muramic acid-N-acetylglucosamine bond<sup>Q</sup></b></li> <li>• Found in the glycopeptide coat of all bacteria</li> </ul>
<b>Lactoferrin</b>	<b>Iron-binding<sup>Q</sup></b> protein present in specific granules
<b>Major basic protein</b>	Cationic protein of eosinophils, which has limited bactericidal activity but is cytotoxic to many parasites

## DEFECTS IN LEUCOCYTE FUNCTION

### Inherited Defect in Phagolysosome Function


#### A. Chédiak-Higashi Syndrome

- Autosomal recessive
- Defects in the **lysosomal transport protein LYST<sup>Q</sup>** encoded by the gene **CHS1** at 1q42<sup>Q</sup>
- **Defective fusion of phagosomes and lysosomes in phagocytes<sup>Q</sup>** (causing susceptibility to infections)
- **Abnormalities in melanocytes<sup>Q</sup>** (leading to **albinism**)
- **Abnormalities in cells of the nervous system<sup>Q</sup>** (associated with nerve defects)
- **Abnormalities in platelets<sup>Q</sup>** (causing **bleeding disorders**).

#### Leukocyte Abnormalities

- **Neutropenia (most common)<sup>Q</sup>**
- Leukocytes contain **giant granules<sup>Q</sup>** (characteristic) result from **aberrant phagolysosome fusion<sup>Q</sup>**

- Defective degranulation, impaired chemotaxis, delayed microbial killing and NK cell function is also impaired



#### B. Chronic Granulomatous Disease

- **Inherited Defect in Microbicidal Activity**
- Inherited defect in genes encoding components of **phagocyte oxidase<sup>Q</sup>**
- **X Linked:** Defect in membrane-bound components (**gp91phox<sup>Q</sup>**)
- **Autosomal recessive** defects: Defect in (**p47phox** and **p67phox<sup>Q</sup>**).
- Formation of **widespread granulomas<sup>Q</sup>**

#### Diagnosis

- **Nitroblue-tetrazolium (NBT) test:**
  - Depends upon direct **reduction of NBT by superoxide free radical** to form an insoluble **formazan<sup>Q</sup>**
  - **Negative** in chronic granulomatous disease and **positive in normal individuals**.
  - This test tells whether or not PHOX enzymes are present, **not how much they are affected<sup>Q</sup>**
- **Dihydrorhodamine (DHR) test**
- **Cytochrome C reduction assay:**
  - **Quantitative:** Amount of **superoxide** a patient's phagocytes can produce<sup>Q</sup>

#### C. Myeloperoxidase Deficiency

- Both **catalase positive & catalase negative organisms survive within phagocytes and cause infections<sup>Q</sup>**
- In CGD catalase negative organisms can be killed<sup>Q</sup> -  $H_2O_2$  in not broken due to catalase so  $H_2O_2 + MPO + Cl^-$  forms HOCl. Which kills microbes)




R10<sup>th</sup>

## Latest Update

### Neutrophil Extracellular Traps (NET)

- **Extracellular fibrillar networks<sup>Q</sup>** that provide a high concentration of antimicrobial substances **at sites of infection**
- **Prevents spread of microbes by trapping them in the fibrils.<sup>Q</sup>**
- **Beneficial suicide:** Neutrophils nucleus undergoing apoptosis to make NETs (**Netosis**)




Beneficial suicide Neutrophils

- Consist of a **viscous meshwork of nuclear chromatin which includes histones and associated DNA<sup>Q</sup>**
- **Arginine** is the most important amino acid for NET

- **Nuclear chromatin in the NETs:** Source of nuclear antigens in systemic autoimmune diseases<sup>Q</sup>

### Frustrated Phagocytosis

- Seen when phagocytes encounter materials **that cannot be easily ingested**
- Eg immune complexes deposited on **immovable flat surfaces (e.g., glomerular basement membrane),<sup>Q</sup>**
- Inability of WBCs to ingest these substances → strong activation & **release of large amounts of lysosomal enzymes into the extracellular environment** → **Leukocyte-mediated tissue injury<sup>Q</sup>**



A-NET; B-Cells

### D. Acquired Leucocyte Deficiencies

- **Bone marrow suppression:** Tumors, radiation & chemotherapy: Decreased Production<sup>Q</sup> of leukocytes
- **Diabetes,<sup>Q</sup> malignancy, sepsis, chronic dialysis:** Affects **Adhesion and chemotaxis<sup>Q</sup>**
- **Leukemia,<sup>Q</sup> anemia, sepsis, diabetes, malnutrition:** Affects Phagocytosis and microbicidal activity

- **TGF- $\beta$ <sup>Q</sup>**
- **IL-10<sup>Q</sup>**
- Resolvin and proteins (Derived from **polyunsaturated fatty acids**)<sup>Q</sup>
- **Cholinergic discharge** that **inhibit the production of TNF** in macrophages<sup>Q</sup>

## TERMINATION OF THE ACUTE INFLAMMATORY RESPONSE

### Anti-inflammatory Mediators


- **Lipoxins<sup>Q</sup>**

### Mnemonic

#### Anti inflammatory mediators

T – TGF- $\beta$   
I – IL-10  
L – Lipoxins

## MEDIATORS OF INFLAMMATION







## Cellular Mediators

Mediator	Characteristics
<b>Histamine</b>	<ul style="list-style-type: none"> <li>Formed from the amino acid '<b>histidine</b>'</li> <li><b>Sources: Mast cells (richest source), platelets and basophils</b></li> <li>Causes <b>vasodilation</b> (but vasoconstriction of large arteries), <b>increased permeability</b> (immediate transient response) &amp; <b>bronchoconstriction</b></li> </ul>
<b>Serotonin (5-HT)</b>	<ul style="list-style-type: none"> <li>Richest source is <b>platelets</b>; also present in enterochromaffin cells.</li> <li>Its primary function is as a <b>neurotransmitter</b> in the <b>gastrointestinal tract</b><sup>Q</sup></li> <li>It has actions similar to histamine</li> </ul>
<b>Lysosomal Enzymes</b>	<ul style="list-style-type: none"> <li>Present in lysosomes of neutrophils and monocytes.</li> <li>Function: Role in intracellular killing microbes and dead cells</li> </ul>
<b>Gelatinase &amp; Acid hydrolases</b>	<ul style="list-style-type: none"> <li>Source: Also have tertiary granules or C particles of neutrophils<sup>Q</sup></li> </ul>


## Newly Synthesized Cellular Mediators

### Nitric Oxide (NO)

<b>Formed from</b>	L-arginine with the help of enzyme nitric oxide synthase (NOS).
<b>Functions</b>	<ul style="list-style-type: none"> <li>Intracellular killing of microbes forming peroxynitrite anion</li> <li>Potent vasodilator, also known as <b>Endothelium-derived relaxing factor (EDRF)</b><sup>Q</sup></li> <li><b>Reduction of platelet aggregation</b><sup>Q</sup></li> </ul>

### Arachidonic Acid Metabolites

- Derived from dietary sources or by conversion from the **essential fatty acid: linoleic acid**.
- Released/mobilized from membrane phospholipids (PL) through the action of cellular phospholipases, mainly **phospholipase A2**
- Hallmark of acute inflammation is increased vascular permeability, thereby most of these mediators increase vascular permeability.**<sup>Q</sup>





### Platelet-Activating Factor (PAF)

- **Produced by:** Platelets, basophils, mast cells, neutrophils, macrophages and endothelial cells
- **Functions:** Platelet aggregation, vasoconstriction and bronchoconstriction.<sup>Q</sup>
- At low concentrations, it induces vasodilation and increased venular permeability

### Cytokines


- **Produced by:**
  - Activated lymphocytes, macrophages, and dendritic cells, but also **endothelial**, epithelial, and connective tissue cells
- **Mediate and regulate:**
  - Immune and inflammatory reactions
- **Action:**
  - Autocrine (same cell), paracrine (close proximity) and endocrine (long distance)
- **Features:**
  - **Pleiotropic:** One cytokine can have different effects on different cells

- **Redundant:** Different cytokines can have the same effect
- **Cascade effect:** Cytokines can stimulate the production of other cytokine



### High Yield Facts

- Most important cytokine responsible for systemic effects of inflammation are interleukin-1 (IL-1) and tumor necrosis factor – alpha (TNF- $\alpha$ )<sup>Q</sup>
- **Intrinsically pyrogenic cytokines-** IL-1 $\alpha$ ,<sup>Q</sup> IL-1 $\beta$ , TNF- $\alpha$ ,<sup>Q</sup> TNF- $\beta$ , IFN- $\alpha$ , and IL-6.<sup>Q</sup>
- **PLEASE NOTE: IL-18<sup>Q</sup> of IL-1 family** is not pyrogenic
- Chemokines mediate their actions through chemokine receptors (**CXCR or CCR**).<sup>Q</sup>
- CXCR4; CCR5- act as co-receptors for binding and entry of **HIV into CD4 cells**.<sup>Q</sup>
- **IL-6**, made by macrophages and other cells, which is involved in **local and systemic reactions**
- **IL-17**, produced mainly by T lymphocytes, which promotes **neutrophil recruitment**.



### Chemokines


- Family of cytokines that act primarily as **chemoattractants** for specific types of leukocytes.
- Classified **according to the arrangement of conserved cysteine residue<sup>Q</sup>** in mature proteins.

Family	$\alpha$ -chemokines (CXC)	$\beta$ -chemokines (C-C)	$\gamma$ -chemokines (C chemokines)	CX3C
Description	2 conserved cysteines separated by 1 amino acid	2 conserved cysteines separated by no amino acid	ONE conserved cysteine	2 conserved cysteines separated by 3 amino acid
Action on	Neutrophils	All WBCs <b>except neutrophils</b> <sup>Q</sup>	<b>Lymphocytes</b>	Monocytes & T cells
Examples	e.g. <b>IL-8</b> , IL-1 and TNF	<ul style="list-style-type: none"> <li>• MCP-1 (Monocyte chemoattractant protein)</li> <li>• <b>RANTES</b><sup>Q</sup>(regulated and normal T-cell expressed and secreted)</li> <li>• <b>Eotaxin (selectively recruits Eosinophils)</b><sup>Q</sup> MIP1<math>\alpha</math>.</li> </ul>	Lymphotoxin	<b>Fractalkine</b>

### Plasma Mediators

#### Complement System

- This system functions in **both innate and adaptive immunity**<sup>Q</sup>
- The **critical step** in complement activation is the proteolysis of **component C3**



\* MAC-membrane attack complex

Cleavage of C3 can occur by one of three pathways	
<b>Classical pathway</b>	Triggered by fixation of C1 to antibody (IgM or IgG) that has combined with antigen <sup>Q</sup>
<b>Alternative pathway</b>	Triggered by microbial surface molecules (e.g., endotoxin, or LPS), complex polysaccharides, cobra venom, other substances, in the absence of antibody
<b>Lectin pathway</b>	Plasma mannose-binding lectin binds to carbohydrates on microbes & directly activates C1. <sup>Q</sup>

All three pathways of complement activation lead to the formation of an active enzyme called the C3 convertase, which splits C3 into two functionally distinct fragments, C3a and C3b and finally leads to formation of membrane attack complex (MAC)<sup>Q</sup>

The complement system has three main functions

<b>Inflammation</b>	<ul style="list-style-type: none"> <li>Anaphylatoxins<sup>Q</sup>-C3a, C5a, C4a. Stimulate histamine release from mast cells<sup>Q</sup></li> <li>Chemotactic<sup>Q</sup>-C5a</li> </ul>
<b>Opsonization &amp; phagocytosis</b>	C <sub>3</sub> b, C <sub>4</sub> b and C <sub>5</sub> b
<b>Cell lysis</b>	<ul style="list-style-type: none"> <li>Deposition of MAC on cells makes them permeable to water &amp; ions → death (lysis) of cells</li> </ul>

### Regulation of Complement System

- C1 inhibitor(C1 INH)<sup>Q</sup> blocks binding of C1 to immune complex

- Decay accelerating factor (DAF) & CD59 are linked to plasma membranes by a glycoposphatidyl (GPI) anchor.
- DAF prevents formation of C3 convertases
- CD59 inhibits formation of the membrane attack complex.<sup>Q</sup>
- Factor H, factor I and CD46 prevent excessive alternate pathway activation.<sup>Q</sup>

### Disease Associated with Complement System


Refer immunity chapter

### Products of Coagulation

- Thrombin activates protease-activated receptors (PARs), which are expressed on platelets and leukocytes play a role in inflammation
- Major role of the PARs is in platelet activation during clotting<sup>Q</sup>

### Kinins

- Kinins are vasoactive peptides derived from plasma proteins, called kininogens, by the action of kallikreins.





## Functions of Bradykinin


- Increases vascular permeability, vasodilation
- Smooth muscle contraction **and pain** when injected into the skin.

## Mnemonic

### Bradykinin functions

- P** - Pain
- D** - Vasodilation
- P** - ↑ Permeability

## ACUTE INFLAMMATION




## Outcomes of Acute Inflammation

- Complete resolution**
- Healing by connective tissue replacement** (scarring, or fibrosis): A **process also called organization**.<sup>Q</sup>
- Progression** of the response to **chronic inflammation**<sup>Q</sup>

## CHRONIC INFLAMMATION

- Seen in:** Persistent infections, hypersensitivity diseases and prolonged exposure to exogenous or endogenous toxic agents- e.g silicosis, atherosclerosis

- Characterized by**
  - Infiltration with mononuclear cells-macrophages, lymphocytes, and plasma cells
  - **Tissue destruction- hallmark of chronic inflammation**<sup>Q</sup>
  - **Attempts at healing-via angiogenesis and fibrosis**
- Main Cells in Chronic Inflammation : Macrophages**
  - Macrophages are tissue cells **derived from hematopoietic stem cells in the bone marrow and from progenitors in the embryonic yolk sac and fetal liver during early development**<sup>Q</sup>
  - **Half-life of blood monocytes is about 1 day, whereas the life span of tissue macrophages is several months or years.**<sup>Q</sup>



Two types of macrophages:

- Activated macrophages
- Tissue macrophages





## Activated Macrophages: 2 Major Types

Type	Classically Activated Macrophages (M1)	Alternatively Activated Macrophages (M2)
<b>Induced by</b>	IFN- $\gamma$ , Microbial products-endotoxin.	IL-4 and IL-13: produced by T lymphocytes (Cytokines other than IFN $\gamma$ ) <sup>Q</sup>
<b>Releases</b>	Lysosomal enzymes, nitric oxide, IL-1 and IL-12, reactive oxygen species and NO	IL-10 <sup>Q</sup> , TGF-B <sup>Q</sup>
<b>Involved in</b>	Host defence against microbes and in many inflammatory reactions <sup>Q</sup>	Involved in anti-inflammatory actions, Angiogenesis, Tissue repair, fibrosis <sup>Q</sup> and collagen synthesis (Not microbiocidal) <sup>Q</sup>

## Tissue Macrophages

Tissue macrophages derived from progenitors in embryonic yolk sac and fetal liver during each development.


Organs	Liver	Spleen	CNS	Synovium	Bone	Lung	Lymph nodes	Placenta	Kidney
<b>Name of Macrophages</b>	Kupffer cell <sup>Q</sup>	Littoral cells <sup>Q</sup>	Microglia <sup>Q</sup>	Type A lining cells <sup>Q</sup>	Osteoclast	Alveolar macrophage or 'Dust cells' <sup>Q</sup>	Sinus histiocytes	Hoffbauer cells <sup>Q</sup>	Mesangial cells <sup>Q</sup>

### Role of Lymphocytes

- Dominant population in the chronic inflammation seen in autoimmune and other hypersensitivity diseases<sup>Q</sup>
- CD4+ T lymphocytes promote inflammation<sup>Q</sup> via different cytokines
- Activated B lymphocytes & antibody-producing plasma cells are also present

### Morphological pattern


- Characterized by formation of **granuloma**.
- Granuloma is an aggregation of **activated macrophages surrounded by mononuclear cells**<sup>Q</sup> principally lymphocytes. **Macrophages** may get activated to form **epithelioid cells** (epithelium-like cells). Some of the cells may fuse together to form a bigger cell called a **giant cell**.




## Unique Points of Certain Granulomas

- **Caseating granuloma:** TB, Histoplasma, Coccidioidomycosis, syphilis
- **Noncaseating granuloma:** Sarcoidosis, Hodgkin's, TB
- **Naked granuloma:** Granuloma without peripheral rim of lymphocytes e.g. **sarcoidosis**

- **Palisaded granuloma**—RA, Wegener's granulomatosis
- **Stellate granuloma:** Cat scratch diseases
- **Fibrin ring or "doughnut" granulomas:** Q fever
- **Malarial granulomas (Durck's granuloma):** P. falciparum




Caseating granuloma usually seen in TB  
(Caseous necrosis shown by arrow)




Non-caseating granuloma  
(Epithelioid cells shown by arrow)


## TYPES OF GIANT CELLS







## INCLUSIONS IN GRANULOMAS




The details about lymphocytes, formation of a granuloma is discussed later in the chapter of 'immunity'.


## SPECIAL TYPES OF GRANULOMA

### Granuloma Morphology

- **Caseating** epithelioid cell granuloma (Caseous necrosis (green arrow) and Langhans cell (yellow arrow))




- **Foreign body granuloma:** (Refractile FB seen with polarized light shown by arrows)




### Palisaded granuloma

- Rheumatoid nodule
- Wegener's granulomatosis




### Stellate granuloma

- Stellate granuloma-follicular hyperplasia with central stellate necrosis with neutrophils, surrounded by palisading histiocytes.
- Most commonly found in children
- Benign infectious disease caused by the bacterium *Bartonella henselae*.
- Seen in cat scratch disease






- A **fibrin ring** granuloma, also known as **doughnut** granuloma, is a histopathological finding that is characteristic of **Q fever**



#### Malarial granulomas (Durck's granuloma)

- Focus of reactive astrocytosis admixed with microglial cells and lymphocytes surrounding a focus of ischemic necrosis or hemorrhage.
- Seen in *P. Falciparum*



#### High Yield Facts

- Granulomas are not seen in ulcerative colitis<sup>Q</sup>
- Non Immune granulomas: Foreign body type granuloma e.g Silica granulomas<sup>Q</sup>
- Necrotic granulomas: TB, Syphilis, histoplasma, cat scratch disease, Wegners, granulomatosis, necrotizing sarcoidosis, bronchocentric granulomas
- Tuberculosis can show both caseating and non caseating granulomas<sup>Q</sup>
- Catscratch disease show stellate granuloma
- Sarcoidosis shows: (Non caseating granuloma, Schaumann and asteroid bodies<sup>Q</sup>)
- Caseating granulomas shows langhans cells: TB

#### TISSUE REPAIR

- **Repair/healing** refers to the **incomplete restoration** of tissue architecture and function **after an injury**<sup>Q</sup>
- **Repair** is often used for **parenchymal and connective tissues**,<sup>Q</sup> **healing** is used for **surface epithelia**<sup>Q</sup>

Repair of damaged tissues occurs by two types of reactions:

#### Regeneration

- **Proliferation of cells** and tissues to **replace lost structures**.
- It results in complete restitution of lost or damaged tissue.

On the basis of intrinsic proliferative capacity, tissues are divided into:

Tissue Subtype	Labile (Continuously Dividing) Tissues	Stable Tissues	Permanent Tissues
<b>Features</b>	Continuously being lost and <b>replaced by maturation from tissue stem cells</b> and by proliferation of mature cells <sup>Q</sup>	Cells are in <b>G<sub>0</sub> stage of the cell cycle</b> Minimal proliferative activity However, <b>capable of dividing in response to injury or loss of tissue mass</b> . <sup>Q</sup>	Cells are <b>terminally differentiated and nonproliferative</b> <sup>Q</sup> in postnatal life.
<b>Examples</b>	<ul style="list-style-type: none"> <li>• Hematopoietic cells<sup>Q</sup></li> <li>• Stratified squamous epithelia<sup>Q</sup> of</li> <li>• Cuboidal epithelia of ducts draining exocrine organs</li> <li>• Columnar epithelium of the GIT, FGS</li> <li>• Transitional epithelium</li> </ul>	<ul style="list-style-type: none"> <li>• Parenchyma of liver, kidney, and pancreas.</li> <li>• Endothelial cells, fibroblasts, and smooth muscle cells<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Neurons and cardiac<sup>Q</sup> muscle cells<sup>Q</sup></li> <li>• Skeletal muscle<sup>Q</sup></li> </ul>
<b>Regeneration</b>	Completely regenerate <sup>Q</sup>	Limited capacity to regenerate after injury except liver <sup>Q</sup>	Dominated by scar formation. <sup>Q</sup>

#### Connective Tissue Deposition (Scar Formation)

Occurs by the laying down of **connective (fibrous) tissue**:

- **Fibrosis**: extensive deposition of collagen<sup>Q</sup>
- **Organization**: fibrosis develops in a tissue space occupied by an inflammatory exudate


#### High Yield Facts

- Proliferation of **endothelial cells, fibroblasts, and smooth muscle cells** is important in **wound healing**.
- **Skeletal muscle** is usually classified as a **permanent tissue**
- **Satellite cells** attached to the **endomysial sheath** provide some **regenerative capacity for muscle**.
- **Granulation tissue**<sup>Q</sup> is the hallmark of the fibrogenic repair.
- Major cytokine involved in fibrosis is **Transforming growth factor-  $\beta$  (TGF-  $\beta$ )**<sup>Q</sup>
- TGF-  $\beta$  is the **most important cytokine**<sup>Q</sup> for the synthesis and deposition of connective tissue proteins
- Central role in repair of TGF-  $\beta$  is activation of **alternatively activated (M2) macrophages**<sup>Q</sup>





## Steps in Scar Formation



R<sup>9th</sup>

**Latest Update**

MMPs	Functions
• Interstitial collagenases	Cleave <b>fibrillar collagen</b> (MMP-1, -2 and -3);
• <b>Gelatinases<sup>Q</sup></b> (MMP-2 and 9)	<b>Degrade amorphous collagen and fibronectin</b>
• <b>Stromelysins<sup>Q</sup></b> (MMP-3, -10, and -11)	<b>Degrade a variety of ECM constituents</b> , including proteoglycans, laminin, fibronectin, and amorphous collagen

## WOUND HEALING

### Cutaneous Wound Healing

#### Three phases:

- Inflammation (early & late)
- Granulation tissue formation & re-epithelialization
- Wound contraction, ECM deposition & remodeling

#### Healing occurs in either of two ways:

By First Intention	By Second intention
<ul style="list-style-type: none"> <li>• <b>Clean and less damage to cell &amp; tissues, uninfected surgical incision<sup>Q</sup></b></li> <li>• <b>Less intense inflammatory reaction</b></li> <li>• <b>Limited granulation tissue is formed</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Extensive loss of cells and tissues &amp; infected<sup>Q</sup></b></li> <li>• <b>More intense inflammatory reaction<sup>Q</sup></b></li> <li>• <b>Much larger amounts of granulation tissue are formed<sup>Q</sup></b></li> <li>• <b>Wound contraction is present<sup>Q</sup></b></li> </ul>




## Healing of a Clean Uninfected Wound



Day (hrs)	Finding	
0	Blood clot SCAB	
1(24 hrs)	Neutrophils	
2(24–48 hrs)	Continuous thin epithelial layer <sup>q</sup>	
3	<ul style="list-style-type: none"><li>• Macrophages</li><li>• Appearance of <b>Granulation tissue</b></li></ul>	<ul style="list-style-type: none"><li>• Deposition of type III collagen but they <b>do not bridge the incision</b></li></ul>
5	<ul style="list-style-type: none"><li>• <b>Abundant granulation tissue</b></li><li>• <b>Collagen fibrils bridge the incision</b></li></ul>	<ul style="list-style-type: none"><li>• <b>Neovascularization</b> is maximum</li><li>• <b>Full epithelial thickness</b> with surface keratinization</li></ul>
14 (end of 2 weeks)	<ul style="list-style-type: none"><li>• <b>Accumulation of collagen</b>; fibroblast proliferation</li><li>• Disappearance of inflammation</li><li>• Blanching begins</li></ul>	
1 months	<ul style="list-style-type: none"><li>• <b>Replacement of collagen type III with collagen type I</b> (has greater tensile strength)</li><li>• <b>Dermal appendages are permanently lost</b></li></ul>	

## Wound Strength

- **Sutured wounds** have approximately **70% of the strength<sup>q</sup>** of normal skin due to sutures
- When sutures are removed, **at 1 week**, wound strength is **10%<sup>q</sup>** of that of unwounded skin<sup>q</sup>
- Wound strength reaches **approximately 70% to 80% of normal by 3 months<sup>q</sup>**
- Strength **never reaches 100%<sup>q</sup>**

## Abnormalities in Tissue Repair

Abnormalities in Tissue Repair	Conditions Associated	
Inadequate formation of granulation tissue	Wound dehiscence and ulceration <sup>q</sup>	 <p>Proud flesh</p>
Increased abdominal pressure <sup>q</sup>	Dehiscence or rupture of a wound	
Excessive formation of granulation tissue	Proud flesh <sup>q</sup> , Blocks re-epithelialization of wound <sup>q</sup>	
Excessive formation of collagen	Keloid, hypertrophic scar <sup>q</sup>	
Exuberant proliferation of fibroblasts	Desmoids, or aggressive fibromatoses	
Contractures: Exaggerated contraction	<ul style="list-style-type: none"> <li>• Seen in palms, soles, and the anterior aspect of the thorax</li> <li>• Commonly seen after serious burns</li> <li>• Can compromise the movement of joints</li> </ul>	
Delayed wound healing	Due to <b>foreign body</b> , ischemia, <b>diabetes<sup>q</sup></b> , malnutrition, hormones ( <b>glucocorticoids<sup>q</sup></b> ), <b>infection<sup>q</sup></b> , <b>zinc deficiency<sup>q</sup></b> or <b>scurvy<sup>q</sup></b>	

Disorders of Excessive Collagen	
Hypertrophic Scar	Keloid
<ul style="list-style-type: none"> <li>• Scar <b>localized</b> to the wound</li> <li>• Usually regresses</li> </ul>	<ul style="list-style-type: none"> <li>• Scar <b>beyond the boundaries</b> of the original wound</li> <li>• <b>Does not regress<sup>q</sup></b></li> <li>• <b>Genetically predisposed</b> condition, usually in African Americans<sup>q</sup></li> <li>• <b>Favoured sites-</b> sternum (<b>most common</b>), ear lobes<sup>q</sup></li> </ul>
 <p>Hypertrophic scar (scar to localized to wound)</p>	 <p>Keloid (scar beyond boundaries)</p>



## Fibrosis in Parenchymal Organs

- Fibrosis—**excessive deposition of collagen and other ECM components in a tissue**<sup>Q</sup>
- Major cytokine involved in fibrosis is **TGF- $\beta$** .<sup>Q</sup>
- Fibrotic disorders into liver cirrhosis

Liver Cirrhosis, systemic Sclerosis (Scleroderma), (idiopathic pulmonary fibrosis, pneumoconioses,), end-Stage Kidney disease and Constrictive Pericarditis.



### High Yield Facts

- **Epidermal appendages do not regenerate**<sup>Q</sup>
- In **superficial wounds**, the epithelium is reconstituted and there may be **little scar formation**<sup>Q</sup>.
- Predominant collagen in **adult skin** is **type I**<sup>Q</sup> whereas in **early granulation tissue**, it is **type III and I**<sup>Q</sup>
- Healing with secondary intension involves formation of a **network of actin-containing fibroblast**<sup>Q</sup> at the edges of the wound.<sup>Q</sup>
- **Permanent wound contraction** requires the **action of myofibroblasts**<sup>Q</sup> seen in healing by **secondary intention**<sup>Q</sup>
- **Myofibroblasts** are **altered fibroblasts** that have ultra-structural characteristics of smooth muscle cells.
- **Contraction of myofibroblasts**<sup>Q</sup> at the wound site **decreases the gap** between the dermal edges of the wound<sup>Q</sup>
- **Fetal cutaneous wounds heal without scar formation**<sup>Q</sup>
- These wounds show **little inflammation** and **practically no fibrosis**<sup>Q</sup>.
- **Vitamin C** is required for the **conversion of tropocollagen to collagen**<sup>Q</sup>
- **Vitamin C** helps in **hydroxylation of lysine and proline**<sup>Q</sup> residues providing **stability to collagen molecules**

R10<sup>th</sup>

## And Other Latest Update

### ACUTE PHASE REACTANTS

Inflammation is associated with cytokine-induced systemic reactions that are collectively called the acute-phase response.

Positive APPs	Negative APPs
<ul style="list-style-type: none"> <li>• <b>C-reactive protein (CRP)</b></li> <li>• <b>Serum Amyloid A (SAA)</b></li> <li>• <b>Haptoglobin (Hp)</b></li> <li>• <b>Ceruloplasmin</b></li> <li>• <b><math>\alpha</math>2-Macroglobulin</b></li> <li>• <b><math>\alpha</math>1-Acid glycoprotein (AGP)</b></li> <li>• <b>Fibrinogen</b></li> <li>• <b>Complement (C3, C4)</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Albumin</b></li> <li>• <b>Transferrin</b></li> <li>• <b>Transthyretin</b></li> <li>• <b>Retinol-binding protein</b></li> </ul>


### EMPERIPOLESIS

- The engulfed cell is viable without physiological or morphological changes
- Unlike in phagocytosis where the engulfed cell is killed by lysosomal enzymes of the macrophage, the cell exists as viable cell within another in emperipolesis and can exit at any time without any structural or functional abnormalities for either of them.

- Seen in:

Rosai—Dorfman disease

- Hematolymphoid disorders<sup>Q</sup> (Hodgkin's disease, leukemia, acute and chronic myeloid leukemia, Non-Hodgkin's lymphoma, myeloproliferative disorders)
- Non-hematological malignancies (neuroblastoma, Rhabdomyosarcoma)



MIC Showing Emperipolesis in megakaryocyte



## NEXT Pattern Questions



1. A 4-year-old child has a history of recurrent infections with pyogenic bacteria. The infections are accompanied by a neutrophilic leukocytosis. Microscopic examination of a biopsy specimen obtained from an area of soft tissue necrosis shows microbial organisms, but very few neutrophils. An analysis of neutrophil function shows a defect in adhesion. This child's increased susceptibility to infection is most likely caused by a defect involving which of the following molecules?

a. Complement C3b      b. Integrins      c. Leukotriene B4      d. Selectins


**Ans. (b) Integrins**

(Ref: Robins Basic Pathology 10th/pg 60)

- Leukocyte rolling is the first step in transmigration of neutrophils from the vasculature to the tissues. Rolling depends on interaction between selectins (P-selectin and E-selectin on endothelial cells, and L-selectin on neutrophils) and their sialylated ligands (e.g., sialylated Lewis X). Integrins (B1 and B2) are involved in the next step of adhesion during which there is firm adhesion between neutrophils and endothelial cells. Complement C3b acts as an opsonin to facilitate phagocytosis. Leukotriene B4 is a chemotactic agent.



2. A 50-year-old woman has had a high fever and cough productive of yellowish sputum for the past 2 days. Her vital signs include temperature of 37.8°C, pulse 103/min, respirations 25/min, and blood pressure 100/60 mm Hg. On auscultation of the chest, crackles are audible in both lung bases. The microscopic appearance of her lung is shown in the figure. Which of the following inflammatory cell types is most likely to be seen in greatly increased numbers in her sputum specimen?




a. Langhans giant cells      b. Macrophages      c. Mast cells      d. Neutrophils

**Ans. (d) Neutrophils**

(Ref: Robins Basic Pathology 10th/pg 63)


- These signs and symptoms suggest acute bacterial pneumonia. Such infections induce an acute inflammation dominated by neutrophils that fill alveoli, as shown in the figure, and are coughed up, which gives the sputum its yellowish, purulent appearance. Langhans giant cells are seen with granulomatous inflammatory responses. Macrophages become more numerous after initiation of acute events, cleaning up tissue and bacterial debris through phagocytosis. Mast cells are better known as participants in allergic and anaphylactic responses. Lymphocytes are a feature of chronic inflammation





## Image-Based Questions

1. A 5-year-old child presented with high grade fever and dyspnea. He also developed pleural effusion. Pleural tapping was done within 24 hours of presentation. Which is the predominant cell population seen?




- a. Neutrophils  
b. Lymphocytes  
c. Monocytes  
d. Eosinophils

3. Autopsy finding in a 45-year-old male adrenals. Diagnosis is:




- a. Caseous necrosis  
b. Fibrinoid necrosis  
c. Fat necrosis  
d. Liquefactive necrosis

2. Autopsy finding of 25-year-old male who died of chest pain. Diagnosis is:




- a. Fibrinous pericarditis  
b. Serous pericarditis  
c. Purulent pericarditis  
d. Caseous pericarditis

4. A 5-year-old male had fever and cervical lymphadenopathy. Histopathologic examination of cervical lymph nodes shows:



- a. Caseating granuloma  
b. Non caseating granuloma  
c. Stellate granuloma  
d. Fat necrosis



5. A 25-year-old male underwent a blunt trauma. On repair, following changes were noted:



- |                |                      |
|----------------|----------------------|
| a. Keloid      | b. Hypertrophic scar |
| c. Proud flesh | d. Desmoid           |



## Answers of Image-Based Questions

1. Ans. (a) **Neutrophils**
  - Neutrophils are predominant infiltrate within 24 hours of injury
2. Ans. (a) **Fibrinous pericarditis**
  - Adherent pericardium has been opened to reveal the surface of the heart. There are thin strands of fibrinous exudate that extend from the epicardial surface to the pericardial sac. This is typical for a fibrinous pericarditis.
3. Ans. (a) **Caseous necrosis**
  - We can see cheesy white necrosis, the patient probably died of disseminated tuberculosis
4. Ans. (a) **Caseating granuloma**
  - Caseating granuloma typical granuloma resulting from infection with Mycobacterium tuberculosis showing central area of caseous necrosis, activated epithelioid macrophages, giant cells, and a peripheral accumulation of lymphocytes.. We can also see langhans giant cell at 3, 0 clock position in image
5. Ans. (c) **Proud flesh**
  - Excessive formation of granulation tissue can be seen.




## Multiple Choice Questions


### ACUTE INFLAMMATION AND MECHANISMS

- In which of the following condition(s) erythrocyte sedimentation rate is increased:** (PGI May 2019)
  - Increased serum immunoglobulin level
  - Spherocytosis
  - Increased plasma viscosity
  - Sickle cell anemia
  - Increased level of C-reactive protein
- Match the following about the endothelial contraction:** (AIIMS May 2019)
 

Column-A	Column-B
1. Sunburn less than 24 hrs	a. Immediate sustained
2. Thorn prick and mast release	b. Delayed prolonged
3. Bacterial infection with necrosis	c. Immediate transient
a. 1-a, 2-b, 3-c	b. 1-b, 2-a, 3-c
c. 1-b, 2-c, 3-a	d. 1-a, 2-c, 3-b
- Which of the following causes vasodilation?** (AIIMS Nov 2019)
  - TXA<sub>2</sub>
  - Leukotriene C<sub>4</sub>
  - Histamine
  - Serotonin
- Identify the arrow marked cell in the given condition below:** (AIIMS May 16)



- Macrophage
  - Lymphocyte
  - Plasma cell
  - Eosinophil
- Which of the following cells will increase in case of parasite infection?** (AIIMS May 16)



- A
- B
- C
- D

- Vasoconstriction in acute inflammation shown by** (WB PGME 2016, Recent Question 2016)
  - Venules
  - Arterioles
  - Capillaries
  - Vein
- SIRS diagnostic criteria. Wrong statement is** (Recent Question 2016)
  - Band >10%
  - Leucocyte >12000 cells/mm<sup>3</sup>
  - Respi. rate >20
  - Band <5%
- Eosinophilia is found in:** (Recent Question 2016)
  - Cryptococcus
  - HPV
  - Strongyloides
  - Typhoid
- Severe infection will increase if absolute neutrophil count will become?** (Recent Question 2015)
  - <500
  - Less than 800
  - Less than 1000
  - Less than 2000
- Vasodilation in acute inflammation is first shown by:** (Recent Question 2015)
  - Venules
  - Arterioles
  - Capillaries
  - Vein
- Cellular infiltrate seen in late pseudomonas infection is formed mainly by** (Recent Question 2014-15)
  - Neutrophils
  - Lymphocytes
  - Monocytes
  - Plasma cells
- Which of the following is the mechanism of "late appearing sunburn"?** (Recent Question 2015)
  - Leucocyte mediated injury
  - Delayed prolonged leakage
  - Early prolonged leakage
  - Delayed transient leakage
- Increased permeability in acute inflammation is due to-** (Recent Question 2014)
  - Histamine
  - IL-2
  - TGFβ
  - FGF
- Sequence of events in acute inflammation-**
  - Vasodilation → Stasis → Transient vasoconstriction → Increased permeability (Recent Question 2014)
  - Transient vasoconstriction → Increased permeability → Stasis → Vasodilation →
  - Transient vasoconstriction → Vasodilation → Stasis → Increased permeability
  - Transient vasoconstriction → Vasodilation → Increased permeability → Stasis
- The following is not an adhesion molecule?** (JIPMER 2014)
  - Spectrin
  - Integrins
  - Selectins
  - Cadherin





## PATHOPHYSIOLOGY OF ACUTE INFLAMMATION

16. **The RBCs with schizonts of *P. Falciparum* are not visible on peripheral blood smear due to which of the following reason?** (AIIMS Nov 16)
  - a. Capillary adherence or sequestration of parasitized RBCs
  - b. ADCC mediated RBC destruction
  - c. Selective hemolysis of affected RBCs in spleen
  - d. Cellular lysis due to hemozoin produced by the parasites Erythrocyte Changes in Malaria
17. **Amino acid which is useful in Neutrophil extracellular traps (NETs) as to causes lysis of chromatin is?** (JIPMER 2016)
  - a. Arginine
  - b. Alanine
  - c. Phenyl alanine
  - d. Valine
18. **The following statements are true regarding neutrophil extracellular trap** (Recent Question 2015)
  - a. Produced by neutrophils in response to infectious pathogens and inflammatory mediators
  - b. Provide a high concentration of antimicrobial substances at sites of infection
  - c. Prevent the spread of the microbes by trapping them in the fibrils
  - d. All the above
19. **All of the following vascular changes are observed in acute inflammation, except:** (Recent Question 2015)
  - a. Vasodilation
  - b. Stasis of blood
  - c. Increased vascular permeability
  - d. Decreased hydrostatic pressure
20. **Correct sequence in extravasation of leukocytes is:** (Recent Question 2015)
  - a. Margination - rolling- adhesion - transmigration
  - b. Transmigration- margination - rolling-adhesion
  - c. Rolling- adhesion- transmigration- margination
  - d. Adhesion- transmigration- margination-rolling
21. **Transcytosis is:** (Recent Question 2015)
  - a. Transport of fluids and proteins through endothelial cells
  - b. Migration of neutrophils through endothelial cell
  - c. Migration of platelets through endothelial cell
  - d. Locomotion oriented along chemical gradient
22. **Transmigration of WBC across the endothelium is called:** (Recent Question 2015)
  - a. Margination
  - b. Rolling
  - c. Pavementing
  - d. Diapedesis
23. **All the statements are correct in autophagy, except:** (AP 2014)
  - a. It is a process in which a cell eats its own contents
  - b. It is a survival mechanism in times of nutrient deprivation
  - c. The starved cell lives by cannibalizing itself and recycling the digested contents
  - d. It is not regulated by any defined set of "genes"
24. **Most characteristic feature of acute inflammation:** (AI 11, AIIMS May 10)
  - a. Vasoconstriction
  - b. Vascular stasis
  - c. Vasodilatation and increased vascular permeability
  - d. Margination of leucocytes

25. **In acute inflammation due to the retraction of endothelial cell cytoskeleton, which of the following results -** (AIIMS Nov 11)
  - a. Delayed transient increase in permeability
  - b. Early transient increase in permeability
  - c. Delayed prolonged increase in permeability
  - d. Early permanent increase in permeability

## MOLECULAR & CELLULAR PATHOLOGY OF ACUTE INFLAMMATION

26. **Which of the following is the correct sequence of cellular events of acute inflammation?** (AIIMS May 18)
  - a. Rolling---Stable adhesion---Activation of integrins---migration via endothelium.
  - b. Rolling ----Activation of Integrins---- Stable Adhesion---migration via endothelium.
  - c. Stable adhesion---Rolling---Activation of integrins---migration via endothelium.
  - d. Activation of integrins--- migration via endothelium--stable adhesion--- Rolling.
27. **Which of the following involved in leukocyte attachment and emigration during inflammation?** (PGI May 18)
  - a. Complement C5a
  - b. Integrin
  - c. L-selectin
  - d. Leukotriene B4
  - e. Interleukin-8
28. **Esterase inhibitor deficiency causes:** (Recent Question 2015)
  - a. SLE
  - b. MPGN
  - c. Hereditary angioneurotic edema
  - d. Omen syndrome
29. **Which of the following is an alpha chemokine?** (Recent Question 2015)
  - a. IL-8
  - b. MCP-1
  - c. Eotaxin
  - d. Lymphotactin
30. **Role of hydroxyl-eicosatetraenoic acid (HETE) in inflammation** (Recent Question 2015)
  - a. Vasodilation
  - b. Vasoconstriction
  - c. Increased vascular permeability
  - d. Chemotaxis
31. **Leukocyte migration through endothelium is induced by** (Recent Question 2015)
  - a. Selectin
  - b. N CAM
  - c. CAM
  - d. PECAM
32. **Locomotion across chemical gradient is called**
  - a. Diapedesis
  - b. Chemotaxis
  - c. Pavementing
  - d. Margination
33. **Which selectin is stored in weibel palade bodies?** (Recent Question 2015)
  - a. Cadherins
  - b. P-Selectin
  - c. E-Selectins
  - d. L-Selectin
34. **Which of the following enzymes are responsible for generating 'oxygen burst' within neutrophils for killing intracellular bacteria?** (Recent Question 2014)
  - a. Superoxide dismutase
  - b. Glutathione peroxidase
  - c. Oxidase
  - d. Catalase
35. **Role of P-selectin in inflammation-** (Recent Question 2013)
  - a. Rolling
  - b. Adhesion
  - c. Homing
  - d. Transmigration





**36. Chemotaxis in response to activation of cells results in-**  
(PGI 2002, AIIMS May 10)

- a. Random multidirectional movement
- b. Unidirectional motion
- c. Adhesion to endothelium
- d. Augmented oxygen dependent bactericidal effect
- e. Phagocytosis

**37. Which among the following is not an adhesion molecule?**

- a. Integrin
- b. Selectin (DNB Dec 10)
- c. Interferon
- d. Transferrin

### MEDIATORS OF INFLAMMATION

**38. Which of the following is/are is endogenous pyrogen?**  
(Recent Question 2019)

- a. IL1
- b. TNF
- c. Lipopolysaccharide
- d. Both a&b

**39. Which of the following enzyme is in 1 granule of neutrophil?**  
(Recent Question 2019)

- a. Alkaline phosphatase
- b. Lactoferrin
- c. Elastase
- d. Lysozyme

**40. Which of the following is an Opsonin?**

- a. C3a
- b. C3b (Recent exam 2018)
- c. C5a
- d. LTC4

**41. Which of the cell derived chemical mediators of inflammation is/are preformed in secretory granules?**  
(PGI Nov 2017)

- a. Histamine
- b. Lysosomal enzymes
- b. Leukotrienes
- d. Prostaglandins
- e. Serotonin

**42. Which of the following is a negative acute phase reactant?**  
(AIIMS May 2017)

- a. Albumin
- b. Haptoglobin
- c. Ferritin
- d. C reactive protein

**43. What decreases during acute inflammation?**  
(PGI May 2017)

- a. Ceruloplasmin
- b. Transferrin
- c. CRP
- d. ESR
- e. Albumin

**44. Interleukin secreted by Th17 cells** (JIPMER 2016)

- a. IFN Gamma
- b. IL22
- c. IL6
- d. IL16

**45. 10-year-old male child complains of pain and swelling in the left foot. He gives history of injury while playing. Migration of leukocytes to the site of injury is mediated by:**  
(JIPMER 2016)

- a. Cytokines
- b. Histamine
- c. Chemokines
- d. Prostaglandins

**46. Which of the following is not an activator of alternate complement system?** (JIPMER 2016)

- a. Factor H
- b. IgA
- c. Bacteria
- d. Immune complex

**47. Integrins are involved in:** (JIPMER 2016)

- a. Adhesion
- b. Rolling
- c. Transmigration
- d. Opsonization

**48. Complement complex that attacks cell membrane is:**

- a. C12345
- b. C23456 (AIIMS May 16)
- c. C34567
- d. C56789

**49. Serotonin, a mediator of inflammation in our body, is secreted/released by:** (PGI May 16)

- a. Leukocytes
- b. Endothelial cell
- c. Mast cell
- d. Platelet
- e. Macrophage

**50. Th1 cells produce?** (Recent Question 2016-17)

- a. IL-1
- b. IL-2
- c. IL-4
- d. IL-5

**51. Primary granule of neutrophil has?**

(Recent Question 2016-17)

- a. Proteinase 3
- b. Alkaline phosphatase
- c. Acid protease
- d. Lactoferrin

**52. Interleukin responsible in surgical trauma.**

(Recent Question 2016)

- a. IL-1
- b. IL-2
- c. IL-3
- d. IL-4

**53. Late complement factor deficiency leads to:**

- a. Hereditary angioneurotic edema (Recent Question 2016)
- b. SLE
- c. Recurrent infections like Neisseria & Pneumococci
- d. HUS

**54. Major basic protein is found in? (Recent Question 2016)**

- a. Macrophage
- b. Eosinophils
- c. Basophil
- d. Neutrophil

**55. Most important pyrogenic Interleukin is?**

(Recent Question 2016)

- a. IL-2
- b. IL-1
- c. IL-6
- d. IL-8

**56. Membrane attack complex** (Recent Question 2016)

- a. C5-9
- b. C3
- c. C2
- d. C1

**57. Bronchospasm is initiated by? (Recent Question 2014)**

- a. C5a
- b. C3a
- c. Leukotrienes
- d. Acetylcholines

**58. Which of the following is the chief mediator associated with resetting the hypothalamic temperature set point at a higher level, resulting in fever? (Recent Question 2015)**

- a. PGF2 alpha
- b. PGE1
- c. PGE2
- d. PGI2

**59. Which of the following is a major pyrogenic cytokine**

(Recent Question 2015)

- a. IL-12
- b. TNF
- c. IFN-Y
- d. Bradykin

**60. The following is not a preformed chemical mediator of inflammation** (Recent Question 2015)

- a. Prostaglandins
- b. Histamine
- c. Serotonin
- d. Lysosomal enzymes

**61. Pain during inflammation is mediated by:**

(Recent Question 2015)

- a. Nitric oxide
- b. Leukotriene B4
- c. Bradykinin
- d. Chemokines



- 62. The following is not a chemoattractant in acute inflammation** (Recent Question 2015)  
a. IL-8                                      b. CSa  
c. LTB4                                      d. Kinins
- 63. The following is not a pyrogenic cytokine** (Recent Question 2015)  
a. IL-1                                      b. TNF  
c. Substance P                              d. Prostaglandins
- 64. Anti-inflammatory cytokine** (Recent Question 2015)  
a. IL-1                                      b. IL-4  
c. TNF                                      d. IFN- $\gamma$
- 65. All are major cytokines in chronic inflammation except** (Recent Question 2015)  
a. IL-6                                      b. IL-12  
c. IFN- $\gamma$                                       d. IL-17
- 66. The following enzyme is otherwise called TACE (TNF converting enzyme)** (Recent Question 2015)  
a. Matrix metalloproteinase  
b. Serine proteinase  
c. ADAM-17  
d. Interstitial collagenase
- 67. Most important cytokine for the synthesis and deposition of connective tissue proteins** (Recent Question 2015)  
a. TGF- $\alpha$                                       b. TGF- $\beta$   
c. FGF-1                                      d. FGF-2
- 68. Angiogenesis is stimulated by all except** (Recent Question 2015)  
a. PDGE                                      b. VEGF  
c. FGF                                      d. HGF
- 69. TNF and IL1 are produced by** (Recent Question 2014-15)  
a. Neutrophils                                      b. Monocytes  
c. Lymphocytes                                      d. Activated Macrophages
- 70. Which is not the action of TGF- $\beta$**  (Recent Question 2014-15)  
a. Anti-inflammatory  
b. Proliferation of fibrous tissue  
c. Inhibition of metalloproteinases  
d. Anaphylaxis
- 71. Which of the following doesn't belong to Interleukin-2 (IL-2) Subfamily** (Recent Question 2015)  
a. IL-2                                      b. IL-3  
c. IL-4                                      d. IL-1
- 72. Which of the following is not endogenous pyrogens?** (Recent Question 2015)  
a. IL 6                                      b. IL 1 beta  
c. IL 12                                      d. TNF alpha
- 73. Which of the following is NOT an anaphylatoxin?** (Recent Question 2014)  
a. C3a                                      b. C4a  
c. C3b                                      d. C5a
- 74. What is not caused by platelet activating factor?** (Recent Question 2014)  
a. Vasoconstriction  
b. Bronchodilation  
c. Causes platelet aggregation  
d. Transmits signals between cells
- 75. Which of the following chemical mediator is not a cell derived mediator?** (JIPMER 2014)  
a. Leukotrienes                                      b. Kinins  
c. Cytokines                                      d. Prostaglandins
- 76. Cytokines responsible for synthesis of acute phase reactant are all except ?** (Recent Question 2013)  
a. IL 1                                      b. IL 11  
c. IL 6                                      d. TNF
- 77. Which of the following is anti inflammatory?** (Recent Question 2013)  
a. IL-2                                      b. IL4  
c. IL-6                                      d. IL-10
- 78. Which interleukin is required for differentiation of eosinophils?** (DNB Aug. 12)  
a. IL-1                                      b. IL-3  
c. IL-4                                      d. IL-5
- 79. Important inflammatory mediators is/are:** (PGI Nov 2011)  
a. IL-2                                      b. IL-6  
c. TNF-alpha                                      d. Platelet activating factor  
e. Interferons
- 80. Acute phase reactant(s) in acute inflammation is/are:** (PGI Nov 2011, 08)  
a. Haptoglobin  
b. C-reactive protein  
c. Alpha-1 acid glycoprotein  
d. Prostaglandins  
e. Fibrinogen
- 81. Actions of bradykinin include all of the following, except-** (AI 10)  
a. Vasodilatation  
b. Bronchodilatation  
c. Increased vascular permeability  
d. Pain
- 82. All of the following are mediators of inflammation except** (AIIMS May 2005, Nov 10)  
a. Tumour necrosis factor- $\alpha$  (TNF- $\alpha$ )  
b. Interleukin-1  
c. Myeloperoxidase  
d. Prostaglandins
- 83. Resolution of inflammation caused by -** (DNB Dec 10)  
a. TNF Alfa, IL-1 and CRP  
b. TN F beta, IL-6 and CRP  
c. TNF Alfa, IL 10 and IL1 receptor antagonist  
d. TNF gamma
- 84. Histamine causes all except:** (DNB June 10)  
a. Arteriolar dilatation  
b. Increased permeability of venules  
c. Constriction of large arteries  
d. Platelet aggregation
- 85. C in CRP stands for -** (DNB June 10)  
a. Concanavalin A  
b. Chondroitin sulfate in series with ARP, BRP  
c. Capsular polysaccharide of pneumococcus  
d. Cellular
- 86. Endogenous chemoattractants are all except -** (MH 10)  
a. C5a  
b. Integrins  
c. LTB4  
d. IL8




87. Pro inflammatory Cytokines include all of the following except -
- |                  |                  |
|------------------|------------------|
| a. Interleukin 1 | b. Interleukin 2 |
| c. Interleukin 6 | d. TNF- Alpha    |

### MORPHOLOGIC PATTERNS OF ACUTE INFLAMMATION

88. All are true about serous inflammation except  
(Recent Question 2015)
- A pattern of acute inflammation
  - Exudation of cell-poor fluid
  - Fluid in infected by destructive organism
  - Fluid does not contain large numbers of leukocytes
89. Exudation of cell-poor fluid that is not infected by destructive organisms is characteristic of:  
(Recent Question 2015)
- Fibrinous inflammation
  - Purulent inflammation
  - Serous inflammation
  - Chronic inflammation
90. Commonest variety of acute inflammation is  
(Recent Question 2014)
- |                           |                          |
|---------------------------|--------------------------|
| a. Purulent inflammation  | b. Serous inflammation   |
| c. Catarrhal inflammation | d. Necrotic inflammation |
| e. Fibrinous inflammation |                          |

### CHRONIC INFLAMMATION

91. Histological picture of a lesion excised from the right cervical region is shown below. What is your diagnosis?  
(Recent Pattern Question 2020)



- Necrotizing granulomatous inflammation
  - Neurofibroma
  - Schwannoma
  - Hodgkin lymphoma
92. Stellate Granuloma are seen in  
(JIPMER 2016)
- |                        |                   |
|------------------------|-------------------|
| a. Cat scratch disease | b. Sarcoidosis    |
| c. LGV                 | d. Histoplasmosis |
93. 9-year-old boy is admitted with acute abdominal pain localized in the right iliac fossa. He is pyrexial with localized peritonism in RIF. The causative cell involved here  
(JIPMER 2015)
- |                |               |
|----------------|---------------|
| a. Lymphocytes | b. Neutrophil |
| c. Macrophages | d. Monocytes  |

94. True statement about alternative macrophage activation  
(WB PGEE 2016, Recent Question 2015)
- Induced by the cytokine IFN- $\gamma$
  - Kill ingested organisms
  - Secrete cytokines that stimulate inflammation
  - Main function is in tissue repair
95. Tissue macrophages are called  
(Recent Question 2014-15)
- |                 |                      |
|-----------------|----------------------|
| a. Monocytes    | b. Histiocytes       |
| c. Plasma cells | d. Epithelioid cells |
96. Biopsy of gastronemius muscle in an asymptomatic patient reveals non-caseating granuloma. Diagnosis  
(Recent Question 2015)
- |                 |             |
|-----------------|-------------|
| a. Tuberculosis | b. Syphilis |
| c. Sarcoidosis  | d. Leprosy  |
97. The principle cell in granuloma  
(Recent Question 2015)
- |               |                     |
|---------------|---------------------|
| a. Fibroblast | b. Epithelioid cell |
| c. Giant cell | d. Plasma cell      |
98. The following factor stimulate angiogenesis  
(Recent Question 2015)
- |                   |               |
|-------------------|---------------|
| a. Integrins      | b. Endostatin |
| c. Thrombospondin | d. Tumstatin  |
99. Durck granuloma is seen in  
(Recent Question 2015)
- |                        |                        |
|------------------------|------------------------|
| a. Congenital syphilis | b. Cat scratch disease |
| c. Histoplasmosis      | d. Cerebral malaria    |
100. Hallmark of chronic inflammation  
(Recent Question 2015)
- Increased vascular permeability
  - Vasodilation
  - Granulomas
  - Tissue destruction
101. Granulomatous inflammation seen in A/E  
(Recent Question 2015)
- |                |                 |
|----------------|-----------------|
| a. Syphilis    | b. Tuberculosis |
| c. Sarcoidosis | d. AIDS         |
102. Stellate granuloma is seen in  
(Recent Question 2015)
- |                |                        |
|----------------|------------------------|
| a. Leprosy     | b. Brucellosis         |
| c. Sarcoidosis | d. Cat-scratch disease |
103. Atherosclerosis is a type of?  
(Recent Question 2015)
- Chronic inflammation
  - Acute inflammation
  - Tissue repair
  - Hypertrophy in response to stimuli
104. Macrophages are converted into epithelioid cells by
- |                 |                 |
|-----------------|-----------------|
| a. IFN $\gamma$ | b. IL2          |
| c. TNF $\alpha$ | d. TGF- $\beta$ |

### TISSUE REPAIR

105. True about neutrophils role in wound healing is/are:  
(PGI May 2019)
- They prevent phagocytosis of bacteria
  - They are involved in early part of wound healing
  - Prominent role in chronic inflammation
  - They produce proteolytic enzymes
  - Appear within 24 hours on wound margin



- 106. A 12-year-old boy had a cut in his forearm 4 days ago. Now the bleeding has been stopped due to granulation tissue formation. While taking a skin biopsy a part of the granulation tissue was also included in the specimen. The histology of granulation tissue is shown below. Which type of collagen is found in this granulation tissue?** (AIIMS May 18)
- Type 1
  - Type 2
  - Type 3
  - Type 4
- 107. Cytokine involved in epithelial tissue migration and granulation tissue in healing?** (Recent Question 2016-17)
- TGF- $\beta$
  - IL-10
  - IL-4
  - Inf-gamma
- 108. Cell involved in the immediate phase of wound healing** (Recent Question 2016, JIPMER\_May 2015)
- Platelets
  - Fibroblasts
  - Basophils
  - Macrophages
- 109. Scar contraction is caused by?** (Recent Question 2016-17)
- Fibroblasts
  - Myofibroblasts
  - Epithelial cells
  - None
- 110. Which of the following promotes growth of fibroblasts and recruitment of macrophages in case of angiogenesis process of wound healing?** (Recent Question 2016)
- EGF
  - PDGF
  - VEGF
  - FGF
- 111. After an incised wound, new collagen fibrils are seen along with growing epithelium. The age of the wound is ?** (AIIMS May 2015)
- 4-5 days
  - About 1 week
  - 12-24 hours
  - 24-72 hours
- 112. Wound contraction occurs by the action of?** (Recent Question 2015)
- Macrophages
  - Myofibroblasts
  - Endothelial cells
  - Nyctes
- 113. Cell involved in the immediate phase of wound healing** (JIPMER 2015)
- Platelets
  - Fibroblasts
  - Basophils
  - Macrophages
- 114. The following factors impair wound healing except:** (Recent Question 2015)
- Diabetes mellitus
  - Glucocorticoids
  - Vitamin C deficiency
  - Clot formation
- 115. Not seen in the inflammatory stage of wound healing** (Recent Question 2015)
- Angiogenesis
  - Chemotaxis
  - Increased vessel permeability
  - Released of cytokines and chemokines
- 116. Find the false statement regarding wound healing** (Recent Question 2015)
- Macrophages are the key cellular constituents
  - TGF- $\beta$  is the most important fibrogenic agent
  - In remodeling type III collagen replaces type I collagen
  - Vitamin C is required for the hydroxylation of procollagen
- 117. Not true regarding primary union** (Recent Question 2015)
- Abundant granulation tissue to fill the wound gap
  - Clear margins
  - Uninfected
  - Neat linear scar
- 118. Find the false statement about wound healing** (Recent Question 2015)
- 3 phases in sequence are: Inflammation, proliferation and maturation
  - In healing by second intention, initially first matrix containing fibrin, plasma fibrin, plasma fibronectin, and type III collagen is formed
  - 70-80% tensile strength of unwounded skin is obtained in 3 months
  - After 2 months of wound healing, the increase in tensile strength is due to excess collagen synthesis
- 119. Which of the following growth factor is not involved in tissue repair** (Recent Question 2015)
- VEGF
  - EGF
  - TNF
  - TGF
- 120. In a sutured incised wound, re-epithelialization is complete by** (Recent Question 2015)
- 24 hours
  - 48 hours
  - 5 days
  - 2 weeks
- 121. Granulation tissue appear at the site of injury by** (Recent Question 2015)
- <24 hours
  - 24-72 hours
  - 48-96 hours
  - 5-7 days
- 122. Collagen fibres bridge the wound area by** (Recent Question 2015)
- <24 hours
  - 24-72 hours
  - 48-96 hours
  - 5-7 days
- 123. When suture are removed from an incisional surgical wound at the end of one week, the wound strength of the wounded skin when compared to unwounded skin is approximately** (Recent Question 2015)
- 1%
  - 10%
  - 50%
  - 80%
- 124. Myofibroblasts are seen in** (Recent Question 2015)
- Healing wounds
  - Cancerous site
  - Adipose tissue
  - Muscle septae
- 125. Wound contraction is due to** (Recent Question 2015)
- Myocyte
  - Fibroblast
  - Myofibroblast
  - Skeletal muscle fibre
- 126. Correct sequence of phases of wound injury** (Recent Question 2015)
- Inflammation-Maturation-Proliferation-Remodeling
  - Inflammation-Proliferation-Maturation-Remodeling
  - Inflammation-Proliferation-Remodeling-Maturation
  - Inflammation-Maturation-Remodeling-Proliferation
- 127. Primary intentional healing which is true** (Recent Question 2014-15)
- Neovascularization is maximum by day 5
  - Neovascularization is maximum by day 3
  - Neutrophils appear at wound margins on day 3
  - The epidermis recovers its maximum thickness by day 7
- 128. Formation of granulation tissue is due to** (MH PG 2014)
- Thrombosed vessels
  - Infiltration of cells
  - Budding of new capillaries
  - Mucosal proliferation





- 129. In wound injury sequence of appearance of cells is-** (Recent Question 2014)
- Macrophage → Platelet → Neutrophils → Fibroblast
  - Neutrophils → Macrophages → Platelet → Fibroblast
  - Platelet → Neutrophils → Macrophages → Fibroblast
  - Platelet → Macrophages → Neutrophils → Fibroblast
- 130. Which is not seen in inflammatory stage of wound healing?** (Recent Question 2013)
- Angiogenesis
  - Chemotaxis
  - Increased capillary permeability
  - Cytokine and chemotactic factor release
- 131. During angiogenesis recruitment of pericytes and periendothelial cells is due to-** (Recent Question 2013)
- VEGF & PDGF
  - Angiopoietins, TGF & PDGF
  - TGF, VEGF & PDGF
  - VEGF, IL-2, IL-6
- 132. Wound healing is the summation of following processes except -** (DNB 2012)
- Coagulation
  - Matrix synthesis
  - Angiogenesis
  - Fibrolysis

- 133. Complete restoration of tensile strength of the wound comparable to normal tissue takes as long as:** (DNB Nov 2012)
- Two weeks
  - Six weeks
  - Six months
  - Two years
- 134. Cells not involved in healing of clean wound:** (PGI Nov 2011)
- Macrophages
  - Platelet
  - Fibroblasts
  - Polymorphonuclear leukocytes
  - MyoFibroblast

#### TYPES OF CELLS

- 135. Permanent tissue is?** (Recent Question 2016-17)
- Heart
  - Liver
  - Kidney
  - Skeletal muscle
- 136. The following are labile cells except** (Recent Question 2015)
- Hepatocytes
  - Bone marrow
  - Intestinal mucosa
  - Epithelium of skin



## Answers with Explanations

- 1. Ans. (a, e); a. Increased serum immunoglobulin level; e. Increased level of C-reactive protein**

(Ref: Robbins 9th/pg 99)

- 2. Ans. (c) 1-b, 2-c, 3-a**

- 3. Ans. (c) Histamine** (Ref: to answer number 13)

- 4. Ans. (c) Plasma cell**

(Ref: Wintrob's 13ed/pg 303; Wintrob's Atlas)

Plasma cells are spherical or ellipsoid and range from 5 to 30  $\mu\text{m}$  in size. The cytoplasm is abundant, always is basophilic, and usually is deep blue; it may have a granular character. Plasma cells have a well-defined perinuclear clear zone that contains the Golgi apparatus.

- 5. Ans. (c) C** (Ref: Wintrob's 13th ed. Pg. 303; Wintrob's Atlas)

**Key to the figure:**

A: Lymphocyte, B: Neutrophil, C: Eosinophil, D: Basophil

- 6. Ans. (b) Arterioles**

(Ref: Essentials of Robbins Pathology 5th ed pg 19)

Transient vasoconstriction of arterioles at the site of injury is the earliest event in acute inflammation. It is usually

mediated by neurogenic and chemical mediators and usually resolves within seconds.

- 7. Ans. (d) Band <5%**

(Ref: <http://www.clevelandclinicmeded.com/medicalpubs/diseasemanagement/infectious-disease/sepsis>)

The term systemic inflammatory response syndrome (SIRS) describes the host response to a critical illness of infectious or non-infectious cause

Evidence of a systemic inflammatory response is indicated by at least two of the following:

- Fever or hypothermia: core body temperature  $38^{\circ}\text{C}$  or higher or  $36^{\circ}\text{C}$  or lower
- Tachypnea: 20 breaths/min or more, or need for mechanical ventilation for an acute process
- Tachycardia: heart rate 90 beats/min or more, unless the patient has a preexisting tachycardia
- White blood cell count:  $12,000$  cells/ $\text{mm}^3$  or higher,  $4,000$  cells/ $\text{mm}^3$  or less, or more than 10% bands on differential

- 8. Ans. (c) Stronglylodes**

(Ref: dacie and lewis practical ematology, 11th ed, pg 102)

Moderate eosinophilia occurs in allergic conditions; more severe eosinophilia ( $20-50 \times 10^9/\text{L}$ ) may be seen in parasitic infections



9. **Ans. (a) <500** (Ref: Robbins 9th/pg 582)  
Neutropenia is denoted as ANC (absolute neutrophil count)

$$\text{ANC} = \frac{(\% \text{neutrophils} + \% \text{bands}) \times (\text{WBC})}{100}$$

- Mild neutropenia ( $1000 \geq \text{ANC} < 1500$ ): minimal risk of infection
- Moderate neutropenia ( $500 \geq \text{ANC} < 1000$ ): moderate risk of infection
- Severe neutropenia ( $\text{ANC} < 500$ ): severe risk of infection.
- Agranulocytosis refers to a virtual absence of neutrophils in peripheral blood. It is usually applied to cases in which the ANC is lower than  $100/\mu\text{L}$ .

10. **Ans. (b) Arterioles** (Ref: Robbins 9th/pg 74; 8th/pg 47)

Vasodilation first involves the arterioles and then leads to opening of new capillary beds in the area.

The result is *increased blood flow*, which is the cause of heat and redness (*erythema*) at the site of inflammation.

11. **Ans. (a) Neutrophils** (Ref: 9th/pg 71; 8th/pg 44)

12. **Ans. (b) Delayed prolonged leakage** (Ref: R 9th/pg 74)

**Mechanism of increased vascular permeability:**

Mechanism	Caused by	Blood vessels affected	Type of response
Mild endothelial damage	<ul style="list-style-type: none"> <li>• Thermal and radiation injury<sup>Q</sup></li> <li>• Late-appearing Sunburn<sup>Q</sup></li> </ul>	Venules and capillaries	Reversible, Delayed & prolonged

13. **Ans. (a) Histamine** (Ref: Robbins 9th/pg 74; 8th/pg 47)

**Increased vascular permeability**

- Formation of endothelial gaps (**Immediate transient response**) is the **most common mechanism<sup>Q</sup>** for **increased permeability**.
- Most important immediate mediator responsible for **Immediate transient response<sup>Q</sup>** is **histamine<sup>Q</sup>**
- **Other immediate** mediators: bradykinin, leukotriene, substance P
- **Somewhat** delayed mediators: TNE, IL-1, IFN- $\gamma$

*Option b:* IL2: IL-2 is known to activate T-cells, especially CD4+ T-helper cells. Other minor function of IL2 are: IL2 can activate lipoxygenase pathway with release of lipoxygenase metabolites which play a role in pulmonary vascular permeability. (*Cytokines and Inflammation By Edward S. Kimball, pg 215*)

*Option c and d:* are mediators of chronic inflammation  
Here the answer is definitely histamine >> IL2

14. **Ans. (d) Transient vasoconstriction → Vasodilatation → Increased permeability → Stasis** (Ref: R 9th/pg 73-74)

15. **Ans. (a) Spectrin** (Ref: Robbins 9th/pg 75-76; 8th/pg 48-49)

Families of CAMs

- Ig (immunoglobulin) superfamily
- Integrins
- Cadherins
- Selectins.

16. **Ans. (a) Capillary adherence or sequestration of parasitized RBCs**

- In *P. falciparum* infections, **membrane protuberances** appear on RBC surface 12–15 h after the cell's invasion. These **"knobs"** extrude a membrane adhesive protein (**PfEMP1**) that mediates **attachment to receptors on venular and capillary endothelium—an event termed cytoadherence**.
- They result in the **sequestration of RBCs containing mature forms of the parasite in vital organs (particularly the brain), where they interfere with microcirculatory flow and metabolism**.

17. **Ans. (a) Arginine** (Ref: R 9th/pg 81)

NETs contain a framework of nuclear chromatin with embedded granule proteins, such as antimicrobial peptides and enzymes. The nuclear chromatin in the NETs, includes histones and associated DNA. Histone modification by peptidylarginine deiminase 4 (PAD4) is necessary for NET release.

18. **Ans. (d) All the above** (Ref: Robbins 9th/pg 81)

Neutrophil extracellular traps (NETs) are extracellular fibrillar networks that provide a high concentration of antimicrobial substances at sites of infection and prevent the spread of the microbes by trapping them in the fibrils. They consist of a viscous meshwork of nuclear chromatin that binds and concentrates granule proteins such as antimicrobial peptides and enzymes

19. **Ans. (d) Decreased hydrostatic pressure**

(Ref: R 9th/pg 73-75)

20. **Ans. (a) Margination - rolling- adhesion - transmigration** (Ref: Robbins 9th/pg 73-75)

Cellular events of inflammation is characterized by: **Margination - rolling- adhesion - transmigration**

21. **Ans. (a) Transport of fluids and proteins through endothelial cells** (Ref: Robbins 9th/pg 74)

Increased transport of fluids and proteins, called transcytosis, through the endothelial cell. This process may involve intracellular channels that may be stimulated by vascular endothelial growth factor (VEGF)

22. **Ans. (d) Diapedesis** (Ref: Robbins 9th/pg 76-77)

Migration of the leukocytes through the endothelium, is called transmigration or diapedesis. It is mediated by PECAM-1



23. Ans. (d) **It is not regulated by any defined set of "genes"**  
(Ref: Dev Cell. 2008 Sep; 15(3): 344-357)

24. Ans. (c) **Vasodilatation and increased vascular permeability** (Ref: Robbins 9th/pg 74; R8"47)

25. Ans. (b) **Early transient increase in permeability**

(Ref: Robbins 9th/pg 74; 8th/pg 47)

Contraction of endothelial cell cytoskeleton leads to Formation of endothelial gaps. This response is called Immediate transient response.

26. Ans. (b) **Rolling---Activation of Integrins---Stable Adhesion--- migration via endothelium**

27. Ans. (b, c) **b. Integrin; c. L-selectin**

28. Ans. (c) **Hereditary angioneurotic edema**

(Ref: Harrison 18:2666-67, 17th ed 2030, 2031)

Deficiency of complement component	Disease/Syndrome
1. C1 esterase Inhibitor	Hereditary angioneurotic edema

29. Ans. (a) **IL-8** (Ref: Robbins 8th/pg 62)

$\alpha$  Chemokines-IL8, IL, TNF

30. Ans. (d) **Chemotaxis** (Ref: Robbins 9th/pg 84)

Chemotaxis, leukocyte adhesion is mediated by Leukotrienes B<sub>4</sub>, HETE

31. Ans. (d) **PECAM** (Ref: Robbins 9th/pg 76; 8th/pg 49-50)

**Leukocyte Migration through Endothelium: Transmigration or diapedesis<sup>Q</sup>**

- Occurs mainly in **postcapillary venules**.<sup>Q</sup>
- Most important molecule: **PECAM-1** (platelet endothelial cell adhesion molecule) or CD31.<sup>Q</sup>

32. Ans. (b) **Chemotaxis** (Ref: Robbins 8th/pg 62)

**Migration in the tissues toward a chemotactic stimulus: Chemotaxis:**

- **Unidirectional movement<sup>Q</sup>** of the leukocytes **towards site of injury** along a **chemical gradient**.

33. Ans. (b) **P-Selectin** (Ref: Robbins 9th/pg 74-75; 8th/pg 48-49)

**P-selectin (CD62P):** Is stored in Platelets  $\alpha$  granules<sup>Q</sup> and endothelium weibel palade bodies<sup>Q</sup>

34. Ans. (c) **Oxidase** (Ref: Robbins 9th/pg 79; 8th/pg 53)

- **REACTIVE OXYGEN SPECIES (ROS)** - Produced by the rapid assembly and activation of a multicomponent oxidase, **NADPH oxidase**, which oxidizes **NADPH<sup>Q</sup>** and, in the process, **reduces oxygen to superoxide**.

- **NADPH oxidase** is also called **phagocyte oxidase** or **phagocyte NADPH oxidase** (**PHOX**) and is responsible for oxidative burst
- **All others options are antioxidants**

35. Ans. (a) **Rolling** (Ref: Robbins 9th/pg 74-75; 8th/pg 48-49)

36. Ans. (b) **Unidirectional motion**

(Ref: 9th/pg 77; 8th/pg 50)

**Unidirectional movement<sup>Q</sup>** of the leukocytes **towards site of injury** along a **chemical gradient**.

Exogenous <sup>Q</sup>	Bacterial products : N-formylmethionine terminal amino acid <sup>Q</sup> and some lipids
Endogenous products	<ul style="list-style-type: none"> <li>• Complement system-C5a<sup>Q</sup></li> <li>• Arachidonic acid (AA) metabolites-LTB<sub>4</sub><sup>Q</sup></li> <li>• Cytokines- IL-8<sup>Q</sup></li> </ul>

37. Ans. (d) **Transferrin**

(Ref: Robbins 9th/pg 74-75; 8th/pg 48-49)

Transferrin is iron carrying protein

38. Ans. (d) **Both a and b**

39. Ans. (c) **Elastase**

40. Ans. (b) **C3b** (Ref: Robbins 9th ed p 78)

The major opsonins are IgG antibodies, the C3b breakdown product of complement, and certain plasma lectins, notably mannosebinding lectin, all of which are recognized by specific receptors on leukocytes.

41. Ans. (a, b, e); a. **Histamine**; b. **Lysosomal enzymes**; e. **Serotonin**

42. Ans. (a) **Albumin** (Ref: R9th/ 99)

43. Ans. (b, e); b. **Transferrin**; e. **Albumin**

44. Ans. (b) **IL22** (Ref: Immunity 2008 Apr. 28(4): 454-467)

IL-22 is one of the IL-10 family cytokines, which also include IL-10, IL-19, IL-20, IL-24, and IL-26, as well as more distally related IL-28 and IL-29.

It has a role in autoimmune diseases, tissue-repair and wound-healing apart from being proinflammatory

45. Ans. (c) **Chemokines** (Ref: Robbins 9th/pg 87)

46. Ans. (d) **Immune complex**

#### Alternative pathway

- Triggered by **microbial surface molecules** (e.g., endotoxin, or LPS), complex polysaccharides, cobra venom, and other substances, **in the absence of antibody**
- IgA can directly activate this pathway



47. Ans. (a) **Adhesion** (Ref: Robbins 9th/pg 85)

48. Ans. (d) **C56789**

(Ref: Robbins and Cotran: Pathological basis of disease 8/e p64)

- Deposition of MAC (**C5-C9**) on cells makes them permeable to water & ions → death (lysis) of cells

49. Ans. (d) **Platelet** (Ref: Robbins 9th/pg 88)

50. Ans. (b) **IL-2**

(Ref: Inflamm Bowel Dis. 1999 Nov;5(4):285-94)

Type 1 T helper (Th1) cells produce interferon-gamma, interleukin (IL)-2, and tumour necrosis factor (TNF)-beta, which activate macrophages and are responsible for cell-mediated immunity and phagocyte-dependent protective responses. By contrast, type 2 Th (Th2) cells produce IL-4, IL-5, IL-10, and IL-13, which are responsible for strong antibody production, eosinophil activation, and inhibition of several macrophage functions, thus providing phagocyte-independent protective responses.

51. Ans. (c) **Acid protease** (Ref: R 9/88 Harrison)

52. Ans. (a) **IL-1** (Ref: Immunol Today, 1994;15:74-80)

The response starts at the trauma site where macrophages and monocytes stimulate cytokine release, especially IL-1 and TNF, which are considered primary interleukins. By themselves, they cause further cytokine release, especially IL-6, which has been considered a major acute phase liver protein stimulator and organic defense response mediator.

53. Ans. (c) **Recurrent infections like Neisseria and Pneumococci**

(Ref: Harrison 18:2666-67, 17th ed 2030, 2031)

Refer immunity chapter

- Deficiency of the terminal components of complement predisposes to **Neisseria infections**<sup>o</sup>

54. Ans. (b) **Eosinophils**

(Ref: Robbins 9th/pg 204, Ref: Anderson 10th ed: 397)

Eosinophils produce:

1. Major basic protein	4. PAF	6. Eosinophilic cationic protein
2. Peroxidase	5. Leukotrienes	7. Reactive oxygen species
3. Neurotoxin		

55. Ans. (b) **IL-1**

(Ref: J Infect Dis. (1999) 179 (Supplement 2): S294-S304. doi: 10.1086/513856, Harrison 18th ed:2666)

Following cytokines are intrinsically pyrogenic in that they produce a rapid-onset fever by acting directly on the hypothalamus without the requirement for the formation of another cytokine:

**IL-1 $\alpha$ , TNF- $\alpha$ , TNF- $\beta$ , IFN- $\alpha$ , and IL-6.**

PLEASE NOTE : **IL-18 of IL-1** family is not pyrogenic

56. Ans. (a) **C5-9** (Ref: Harrison 18:2666-67, 17th ed 2030, 2031)

57. Ans. (c) **Leukotrienes** (Ref: Robbins 9th/pg 89)

- LTE4 cause intense vasoconstriction, bronchospasm (important in asthma), and increased permeability of venules.
- C3a, C5a, and, to a lesser extent, C4a stimulate histamine release from mast cells and thereby increase vascular permeability and cause vasodilation. They are called anaphylatoxins

58. Ans. (c) **PGE2** (Ref: Robbins 9th/pg 99)

The increase in body temperature is caused by prostaglandins that are produced in the vascular and perivascular cells of the hypothalamus. In the hypothalamus, the prostaglandins, especially PGE2, stimulate the production of neurotransmitters that reset the temperature set point at a higher level.

59. Ans. (b) **TNF**

(Ref: J Infect Dis. (1999) 179 (Supplement 2): S294-S304. doi: 10.1086/513856, Harrison 18th ed:2666)

60. Ans. (a) **Prostaglandins** (Ref: Robbins 9th/pg 84)

61. Ans. (c) **Bradykinin** (Ref: Robbins 9th/pg 65, 89)

**Functions of bradykinin:**

- Increases vascular permeability, vasodilation
- Smooth muscle contraction **and pain** when injected into the skin.

62. Ans. (d) **Kinins** (Ref: Robbins 9th/pg 89, 65)

As we know: **LTB4**, -C5a and IL8 are strong chemotactic agents

Kinins are vasoactive peptides derived from plasma proteins, called **kininogens**, by the action of **kallikreins** – **not chemotactic**

63. Ans. (c) **Substance P**

(Ref: J Infect Dis. (1999) 179 (Supplement 2): S294-S304. doi: 10.1086/513856, Harrison 18th ed:2666)

IL1 AND TNF are well known pyrogenic cytokines

In the hypothalamus, the prostaglandins, especially PGE2, stimulate the production of neurotransmitters that reset the temperature set point at a higher level, hence pyrogenic

64. Ans. (b) **IL-4** (Ref: Robbins 9th/pg 94,95)

Anti-inflammatory cytokines are **IL-10, TGF-B, Lipoxins, IL-4**





**65. Ans. (a) IL-6**

(Ref: *Fundamental immunology by William E Paul*:1030, *Koj* -chapter 3)

IL6 IS INVOLVED IN ACUTE RESPONSES  
Others are involved in chronic inflammation

**66. Ans. (c) ADAM-17**

(Ref: *Immunology: Mucosal and Body Surface Defences*, 4.8)  
ADAM metalloproteinase domain 17 (ADAM17), also called TACE (tumor necrosis factor- $\alpha$ -converting enzyme), is a 70-kDa enzyme that belongs to the ADAM protein family of disintegrins and metalloproteases.

**67. Ans. (b) TGF- $\beta$**  (Ref: *Robbins 9th/pg 105*)

Transforming growth factor- $\beta$  (TGF- $\beta$ ) is the most important cytokine for the synthesis and deposition of connective tissue proteins.


**68. Ans. (d) HGF**

(Ref: *Angiogenesis: An Integrative Approach from Science to Medicine*, pg 82)

Pro- angiogenesis molecules

Stimulator	Mechanism
FGF	Promotes proliferation & differentiation of endothelial cells, smooth muscle cells, and fibroblasts
VEGF	Affects permeability
Ang1 and Ang2	Stabilize vessels
PDGF (BB-homodimer) and PDGFR	Recruit smooth muscle cells
TGF- $\beta$ , endoglin and TGF- $\beta$ receptors	$\uparrow$ extracellular matrix production
Integrins $\alpha V\beta 3$ , $\alpha V\beta 5$ and $\alpha 5\beta 1$	Bind matrix macromolecules and proteinases
VE-cadherin and CD31	Endothelial junctional molecules
Ephrin	Determine formation of arteries or veins
Plasminogen activators	Remodels extracellular matrix, releases and activates growth factors
Plasminogen activator inhibitor-1	Stabilizes nearby vessels

**69. Ans. (d) Activated Macrophages** (Ref: *R 9th/pg 86-87*)



**70. Ans. (d) Anaphylaxis**

(Ref: *Robbins 9th/pg 86-87, 105, Immunology: Mucosal and Body Surface Defences*, 4.8)

**Transforming growth factor- $\beta$  (TGF- $\beta$ ) is the most important cytokine for the synthesis and deposition of connective tissue proteins.** Other functions: Angiogenesis. Anti-inflammatory cytokine

**Up-regulation of tissue inhibitor of metalloproteinases-3 gene expression is done by TGF-beta in articular chondrocytes**

**71. Ans. (d) IL-1**

(Ref: *Curr. Opin. Immunol.* 23 (5): 598-604, *Harrison 18th ed*:2666)

**Interleukin-2 (IL-2) Subfamily<sup>Q</sup>:**

- **Interleukins:** IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-11, IL-12, IL-13, IL-15, IL-21, IL-23
- **Not called interleukins:** Colony-stimulating factor-1 (CSF1), granulocyte-macrophage colony-stimulating factor (CSF2) erythropoietin (EPO), thrombopoietin (THPO), leukocyte inhibitory factor (LIF)
- **Not interleukins:** Growth hormone (GH1), prolactin (PRL), leptin (LEP), **cytokine receptor-like factor 1<sup>Q</sup>** (CLC or CLF)
- **Interferon (IFN) subfamily:** IFN- $\beta$ , IFN- $\alpha$
- **IL-10 subfamily<sup>Q</sup>:** IL-10, IL-19, IL-20, IL-22, IL-24 and IL-26 IL-1 does not belong to IL2 family

**72. Ans. (c) IL 12**

(Ref: *J Infect Dis.* (1999) 179 (Supplement 2): S294-S304. doi: 10.1086/513856, *Harrison 18th ed*:2666)

**73. Ans. (c) C3b** (Ref: *Robbins 9th/pg 83; 8th/pg 57*)

**Anaphylatoxins-C3a, C5a, C4a.**

**74. Ans. (b) Bronchodilation** (Ref: *R 9th/pg 89; 8th/pg 59*)

PAF

- **Produced by:** platelets, basophils, mast cells, neutrophils, macrophages, and endothelial cells
- **Functions: platelet aggregation, vasoconstriction and bronchoconstriction.<sup>Q</sup>**
- At low concentrations it induces vasodilation and increased venular permeability
- Transmits signals between cells

**75. Ans. (b) Kinins** (Ref: *Robbins 9th/pg 82; 8th/pg 57*)

Plasma Derived	Cell Derived
Fibrin split products	Histamine
Kinins (bradykinin)	Serotonin
C3a, C5a	Prostaglandins, Leukotrienes
	Nitric oxide
	Platelet-activating factor

**76. Ans. (b) IL 11** (Ref: *Robbins 9th/pg 86-87*)

IL-6, tumour necrosis factor-alpha (TNF-alpha) and IL-1 are thought to be the key mediators of the acute phase response (Ref: *Clin Exp Immunol.* 1995 Oct; 102(1): 217-223.)

Coming to option b:



IL-11	gp 130	Bone marrow stromal cells	Megakaryocytes,	Induces megakaryocyte colony formation <sup>a</sup>
-------	--------	---------------------------	-----------------	---

#### 77. Ans. (d) IL-10

(Ref: Jolanta Jura and Aleksander Koj –chapter 3)

Anti inflammatory cytokines

IL-10, IL-13, IL-4 and TGF- $\beta$

Please note, both IL-4 and IL-6 are pro as well as anti inflammatory cytokines

#### 78. Ans. (d) IL-5

#### 79. Ans. (a, b, c, e); a. IL-2; b. IL-6; c. TNF-alpha; e. Interferons

(Ref: Fundamental immunology by William E Paul :1030, chapter 3)

#### 80. Ans. (a, b, e); a. Haptoglobin; b. C-reactive protein; e. Fibrinogen

(Ref: <http://www.uptodate.com/contents/acute-phase-reactants>)

Acute phase reactants are

CRP	Ferritin
Fibrinogen	Hepcidin
Serum Amyloid A	Alpha-1 antitrypsin
Haptoglobulin	Ceruloplasmin
Procalcitonin	

#### 81. Ans. (b) Bronchodilatation (Ref: Robbins 9th/pg 65, 89)

**Functions of bradykinin:**

- Increases vascular permeability, vasodilation
- Smooth muscle contraction **and pain** when injected into the skin.

#### 82. Ans. (c) Myeloperoxidase

(Ref: Robbins 9th/pg 83; 8th/pg 57)

MPO- lysosomal protein stored in azurophilic granules of the neutrophil.

#### 83. Ans. (c) TNF Alfa, IL 10 and IL1 receptor antagonist

(Ref: Fundamental immunology by William E Paul:1030, chapter 3)

**Ans none** but if to mark one, go for c. 2 options in c are correct and 1 wrong

Resolution of infection is by:

- IL10- anti inflammatory cytokine-CORRECT
- IL1- pro inflammatory cytokine and IL1 receptor antagonist will cause resolution of inflammation-CORRECT
- TNF- pro inflammatory cytokine-WRONG

#### 84. Ans. (d) Platelet aggregation (Ref: R 9th/pg 82; 8th/pg 57)

Mediator	Characteristics
Histamine	<ul style="list-style-type: none"> <li>Formed from the amino acid 'histidine'</li> <li>Sources: Mast cells (richest source), platelets and basophils</li> <li>Causes <b>vasodilation</b> (but vasoconstriction of large arteries), <b>increased permeability</b> (immediate transient response) &amp; <b>bronchoconstriction</b></li> </ul>

#### 85. Ans. (c) Capsular polysaccharide of pneumococcus

(Ref: Essential of pathology, 8th ed: 781)

#### 86. Ans. (b) Integrins (Ref: Robbins 9th/pg 77; 8th/pg 50)

#### 87. Ans. (b) Interleukin 2

(Ref: Robbins 9th/pg 85-86; 8th/pg 58)

All are Pro inflammatory Cytokines but if to mark one go for b. Since all other 3 are major Pro inflammatory Cytokines.

#### 88. Ans. (c) Fluid in infected by destructive organism

(Ref: Robbins 9th/pg 90)

### Serous Inflammation

A pattern of acute inflammation

Marked by the exudation of cell poor fluid into spaces created by cell injury or into body cavities lined by the peritoneum, pleura, or pericardium

Typically, the fluid in serous inflammation is not infected by destructive organisms

Does not contain large numbers of leukocytes

#### 89. Ans. (c) Serous inflammation (Ref: Robbins 9th/pg 90)

#### 90. Ans. (c) Catarrhal inflammation

(Ref: Chandrasoma taylor: 3ed: 45)

**Catarrhal inflammation**

- Commonest type of inflammation<sup>o</sup>**
- Increased mucus secretion**
- Seen in **common cold**

#### 91. Ans. (a) Necrotizing granulomatous inflammation

(Ref: Robbins 9th ed/pg 98)

#### 92. Ans. (a) Cat scratch disease

(Ref: Anderson 10th/pg 582)

#### 93. Ans. (b) Neutrophil

(Ref: Robbins 9th/pg 71; 8th/pg 44)

This is a case of acute appendicitis

The principal inflammatory cell in this case of acute appendicitis is the neutrophil.



94. Ans. (d) **Main function is in tissue repair**

(Ref: Robbins 9th/pg 94)

95. Ans. (b) **Histiocytes** (Ref: Anderson 10th ed pg 583)

- Macrophages are tissue cells **derived from hematopoietic stem cells in the bone marrow and from progenitors in the embryonic yolk sac and fetal liver during early development**<sup>Q</sup>
- Circulating cells of this lineage are known as monocytes. Tissue macrophages are called **histiocytes**. Activated macrophages are called epithelioid cells

96. Ans. (c) **Sarcoidosis** (Ref: Robbins 9th/pg 94; 8th/pg 71)

Leprosy and TB- Caseating granuloma  
Syphilis- plasma cell rich granuloma (gumma)  
Sarcoidosis- non caseating granuloma

97. Ans. (b) **Epithelioid cell** (Ref: Anderson 10th ed pg 583)

- Macrophages (also known as **histiocytes**) are the cells that define a granuloma
- The macrophages in granulomas are often referred to as "epithelioid"
- Epithelioid macrophages differ from ordinary macrophages in that they have **elongated nuclei** that often resemble the sole of a slipper or shoe.
- These changes are thought to be a consequence of "activation" of the macrophage by the offending antigen.

98. Ans. (a) **Integrins**

(Ref: Angiogenesis: An Integrative Approach from Science to Medicine, pg 123; **Angiogenesis molecules have been described at Ans no 82. Also students must know anti angiogenesis molecules**)

Antiangiogenic molecules are

- |                   |                 |
|-------------------|-----------------|
| 1. Endostatin     | 5. Endorepellin |
| 2. Thrombospondin | 6. Fibulin 5    |
| 3. Tumstatin      | 7. Canstatin    |
| 4. Restin         |                 |

99. Ans. (d) **Cerebral malaria** (Ref: Malaria - Page 76)

In malignant cerebral malaria caused by Plasmodium falciparum, brain vessels are plugged with parasitized red cells, causing ring hemorrhage which is accompanied by necrosis of surrounding parenchyma. The damage leads to formation of Durck's granuloma - collection of microglial cells surrounding area of demyelination

100. Ans. (d) **Tissue destruction** (Ref: Robbins 9th/pg 494)

101. Ans. (d) **AIDS** (Ref: Robbins 9th/pg 494; 8th/pg 500)

102. Ans. (d) **Cat-scratch disease** (Ref: Robbins 9th/pg 494)


103. Ans. (a) **Chronic inflammation**

(Ref: Robbins 9th/pg 494)

Atherosclerosis as a **chronic inflammatory and healing response** of the arterial wall to **endothelial injury**.

104. Ans. (a) **IFN $\gamma$**  (Ref: Anderson 10th ed pg 583)

- Epithelioid macrophages differ from ordinary macrophages in that they have **elongated nuclei** that often resemble the sole of a slipper or shoe.
- **IFN $\gamma$**  is important in activating macrophages and transforming them into epithelioid cells and multinucleate giant cells



105. Ans. (b, d, e); **b. They are involved in early part of wound healing; d. They produce proteolytic enzymes; e. Appear within 24 hours on wound margin**

(Ref: Robbin's 9th ed/pg. 107)

106. Ans. (c) **Type 3**

The early granulation tissue mainly has collagen type 3 >1

107. Ans. (a) **TGF-B** (Ref: Robbin's 9th ed/pg. 107)

108. Ans. (a) **Platelets** (Ref: Robbin's 9th ed/pg. 104)

1st phase is clot form action which involves platelets

109. Ans. (b) **Myofibroblasts**

- **Permanent wound contraction** requires the **action of myofibroblasts**<sup>Q</sup> seen in healing by **secondary intention**<sup>Q</sup>
- **Myofibroblasts** are **altered fibroblasts** that have ultra-structural characteristics of smooth muscle cells.

110. Ans. (d) **FGF** (Ref: Robbin's 9th ed/pg. 104)

- Fibroblast growth factors (FGFs), mainly FGF-2, stimulates the proliferation of endothelial cells. It also promotes the migration of macrophages and fibroblasts to the damaged area, and stimulates epithelial cell migration to cover epidermal wounds.



- PDGF-recruits smooth muscle cells
- TGF- $\beta$  suppresses endothelial proliferation and migration, and enhances the production of ECM proteins.
- Vascular endothelial growth factors (VEGFs), mainly VEGF-A, stimulates both migration and proliferation of endothelial cells

**111. Ans. (d) 24-72 hours** (Ref: Robbin's 9th ed/Pg 107)

**112. Ans. (b) Myofibroblasts** (Ref: R 9th/pg 108; 8th/pg 106)

**113. Ans. (a) Platelets** (Ref: Robbin's 9th ed/pg 107)

The earliest feature in wound repair is Presence of **blood clot** in the incision. This clot is formed by the initial effort of platelets and then the coagulation factors.

**114. Ans. (d) Clot formation** (Ref: Robbin's 9th ed/pg 106)

### Factors Impairing Healing

Infection  
Diabetes  
protein deficiency, vitamin C deficiency  
Glucocorticoids (steroids)  
Mechanical factors such as increased local pressure or torsion may cause wounds to dehiscence  
Poor perfusion, due either to arteriosclerosis and diabetes or to obstructed venous drainage (e.g., in varicose veins)  
Foreign bodies  
Blood clot formation is the earliest event in wound repair

**115. Ans. (a) Angiogenesis**

(Ref: Robbins 9th/pg 103; 8th/pg 101)

#### Three phases of wound healing:

- Inflammation (early & late)- option b, c, d
- Granulation tissue formation & re-epithelialization- option a
- Wound contraction, ECM deposition & remodeling

**116. Ans. (c) In remodeling type III collagen replaces type I collagen** (Ref: Robbins 9th/pg 107; 8th/pg 102-5)

In remodeling type I collagen replaces type III collagen

**117. Ans. (a) Abundant granulation tissue to fill the wound gap** (Ref: Robbins 9th/pg 108; 8th/pg 106)

**118. Ans. (d) After 2 months of wound healing, the increase in tensile strength is due to excess collagen synthesis**

(Ref: Robbins 9th/pg 108; 8th/pg 106)

**Option a-true-Three phases of wound healing-**  
Inflammation (early & late), Granulation tissue formation & re-epithelialization and Wound contraction, ECM deposition & remodeling

- Strength **never reaches 100%** Option d- The recovery of tensile strength results from the excess of collagen
- Synthesis over collagen degradation during the first 2 months of healing, and, at later times, from **structural modifications** of collagen fibers (cross-linking, increased fiber size) option D.

**119. Ans. NONE > TNF** (Ref: Robbins 9th/pg 107)

In Granulation tissue formation phase: Migration of fibroblasts to the site of injury is driven by chemokines, TNF, PDGF, TGF- $\beta$ , and FGF. Their subsequent proliferation is triggered by multiple growth factors, including PDGF, EGF, TGF- $\beta$ , and FGF, and the cytokines IL-1 and TNF.

**120. Ans. (c) 5 days** (Ref: Robbins 9th/pg 107; 8th/pg 102-5)

**121. Ans. (b) 24-72 hours** (Ref: R 9th/pg 107; 8th/pg 102-5)

**122. Ans. (d) 5-7 days** (Ref: Robbins 9th/pg 107)

**123. Ans. (b) 10%** (Ref: Robbins 9th/pg 108; 8th/pg 106)

**124. Ans. (a) Healing wounds** (Ref: 9th/pg 108)

**125. Ans. (c) Myofibroblast** (Ref: R 9th/pg 108; 8th/pg 106)

**126. Ans. (b) Inflammation-Proliferation-Maturation-Remodeling** (Ref: Robbins 9th/pg 108; 8th/pg 106)

**127. Ans. (a) Neovascularization is maximum by day 5**

(Ref: Robbins 9th/pg 107; 8th/pg 102-5)

**128. Ans. (c) Budding of new capillaries** (Ref: R 9th/pg 106)

**129. Ans. (c) Platelet  $\rightarrow$  Neutrophils  $\rightarrow$  Macrophages  $\rightarrow$  Fibroblast** (Ref: Robbins 9th/pg 107; 8th/pg 102-5)

Blood clot  $\rightarrow$  neutrophils  $\rightarrow$  macrophages  $\rightarrow$  collagen  $\rightarrow$  fibroblast

**130. Ans. (a) Angiogenesis**

(Ref: Robbins 9th/pg 103; 8th/pg 101)

#### Three phases of wound healing:

- Inflammation (early & late)- option b, c, d
- Granulation tissue formation & re-epithelialization- option a
- Wound contraction, ECM deposition & remodeling

**131. Ans. (b) Angiopoietins, TGF & PDGF**

(Ref: R 9th/pg 104)

Factors involved in angiogenesis:

Growth factors. • Mainly <b>VEGF-A</b> • Fibroblast growth factors mainly FGF-2	Structural maturation of new vessels <b>Angiopoietins 1 and 2 (Ang 1 and Ang 2)</b>	Stabilisation of vessels <b>PDGF and TGF-<math>\beta</math></b>	• <b>Notch signaling</b> • Regulates the sprouting and branching of new vessels • Ensures proper spacing of vessels.
---	--	--	--

Newly formed vessels need to be stabilized by the recruitment of pericytes and smooth muscle cells and by





the deposition of connective tissue-**Angiopoietins, PDGF and TGF- $\beta$** .

**132. Ans. (d) Fibrolysis** (Ref: R 9th/pg 107; 8th/pg 102-105)

**133. Ans. (c) Six months** (Ref: Robbins 9th/pg 108; 8th/pg 106)

Ans should be none. But if to mark one, it should be close to 3 months.

**134. Ans. (e) MyoFibroblast** (Ref: R 9th/pg 108; 8th/pg 106)

**Permanent wound contraction** requires the **action of myofibroblasts**<sup>Q</sup> seen in healing by **secondary intention**<sup>Q</sup>

- **Myofibroblasts** are **altered fibroblasts** that have ultra-structural characteristics of smooth muscle cells.

**135. Ans. (d) Skeletal muscle** (Ref: Robbins 9th/pg 101)

**Permanent Tissues**

- **Neurons and cardiac**<sup>Q</sup>
- **Muscle cells**<sup>Q</sup>
- **Skeletal muscle**<sup>Q</sup>

**136. Ans. (a) Hepatocytes**

(Ref: Robbins 9th/pg 101; 8th/pg 81)

**Labile Tissues**

- **Parenchyma** of liver, kidney, and pancreas.
- **Endothelial cells, fibroblasts, and smooth muscle cells**;<sup>Q</sup>

**Labile (Continuously Dividing) Tissues**

- **Hematopoietic cells**<sup>Q</sup>
- **Stratified squamous epithelia**<sup>Q</sup>
- **Cuboidal epithelia** of ducts draining exocrine organs (salivary glands, pancreas, biliary tract);
- **Columnar epithelium** of the **GIT, uterus**, and fallopian tubes;
- **Transitional epithelium** of the urinary tract.<sup>Q</sup>

# Hemodynamics

## Key Points

- » **Hyperemia** is an **active** process
- » **Nutmeg liver** is seen in chronic passive hepatic congestion
- » Hemosiderin-laden macrophages (**heart failure cells**) are seen in chronic pulmonary congestion
- » **Line of Zahn**-distinguish antemortem clots from the bland non-laminated postmortem<sup>o</sup> clots
- » Fat embolism syndrome is characterized by pulmonary insufficiency, neurologic symptoms, **anemia** & thrombocytopenia (petechial rash)
- » Arterial occlusions cause **white infarcts** whereas venous occlusion cause **red infarcts**<sup>o</sup>
- » Primary initiating factor in septic shock: **cytokine release**
- » Most common cytokine involved in Septic shock is **TNF- $\alpha$**


## Key Recent Updates

- » **Procalcitonin** is marker of sepsis
- » Most common cause of septic shock is **gram positive bacteria** (PAMP)
- » **Endotoxin** [lipopolysaccharide] and lipoteichoic acids are PAMPs (pathogen associated molecular patterns).



## HEMODYNAMICS

- **Vascular hydrostatic pressure** is **balanced** by **plasma colloid osmotic pressure**<sup>Q</sup>
- Accumulation of fluid in tissues is called **edema** or body cavities is called **effusions**.<sup>Q</sup>
- It occurs due to either **elevated**<sup>Q</sup> hydrostatic pressure or **diminished**<sup>Q</sup> colloid osmotic pressure



The excessive interstitial fluid can be either a **transudate** or an **exudate**

Characteristic	Transudate	Exudate
Appearance	Clear, Colorless	Yellow, turbid, purulent, bloody
Specific gravity	< 1.015 <sup>Q</sup>	> 1.015 <sup>Q</sup>
Protein	< 3 g/dL	> 3 g/dL
LDH	< 200 IU	> 200 IU
Cell count	< 1000/uL <sup>Q</sup>	> 1000/uL <sup>Q</sup>
Permeability	Normal <sup>Q</sup>	Altered <sup>Q</sup>
Conditions	Congestive Heart Failure	Infections, Malignancies

### Acute pulmonary congestion:

- Engorged alveolar capillaries<sup>Q</sup>
- Alveolar septal edema<sup>Q</sup>
- Focal intraalveolar hemorrhage<sup>Q</sup>

### Acute hepatic congestion:

- Centrilobular hepatocytes: ischemic necrosis<sup>Q</sup>
- Periportal hepatocytes - fatty change<sup>Q</sup>

### Chronic pulmonary congestion:

- Septa are thickened and fibrotic<sup>Q</sup>
- Hemosiderin-laden macrophages (heart failure cells)<sup>Q</sup> (Fig. 1)

### Chronic passive hepatic congestion:

- **Nutmeg liver**<sup>Q</sup>: centrilobular regions are red-brown against surrounding zones of uncongested tan liver (Fig. 2)
- Initially **centrilobular necrosis & hemosiderin laden macrophages**.<sup>Q</sup>
- Later: hepatic fibrosis called **cardiac cirrhosis**.<sup>Q</sup>

## HYPEREMIA AND CONGESTION


- **Increased volume of blood**<sup>Q</sup> within dilated vessels of tissue or organ.


Features	Hyperemia	Congestion
Process	Active process <sup>Q</sup>	Passive process <sup>Q</sup>
Cause	Arteriolar dilation	Impaired venous outflow
Edema	Absent	Present <sup>Q</sup>
Colour of the tissues	Red color <sup>Q</sup>	Blue red color <sup>Q</sup> (deoxyhemoglobin)
Seen in	Inflammation <sup>Q</sup>	Right heart failure




### High Yield Facts

- CVC liver—**Nutmeg liver**
- CVC spleen shown **Gamna Gandy Bodies**. (organized hemorrhage with dystrophic calcification and hemosiderin pigment in spleen)





**Figs 1A and B:** A. Normal lung; B. Microscopy showing heart failure cells s/o chronic pulmonary congestion



**Figs 2A and B:** Chronic passive hepatic congestion. A. Gross appearance: Liver shows dilated congested centrilobular regions s/o nutmeg liver. B. Microscopic preparation shows centrilobular hepatic necrosis.

## HEMOSTASIS

Explained in detail in Chapter Bleeding and its disorders (Chapter 11)


## THROMBOSIS

- **Pathologic formation of intravascular thrombus**
- **Virchow's triad** is required for **thrombus formation**.<sup>Q</sup>

Explained in detail in Chapter 11

### Types

There are three types of thrombosis









- Thrombi on heart valves are called **vegetations**: seen in **Rheumatic heart disease**, **infective endocarditis**; **nonbacterial thrombotic endocarditis** and **verrucous (Libman-Sacks) endocarditis**<sup>Q</sup> (**Fig. 3**).
- Thrombus has a **firm attachment**<sup>Q</sup> to underlying vessel or heart wall, which is **not seen in clot**<sup>Q</sup>
- Thrombi (**both arterial & venous**)<sup>Q</sup> have laminations, called **line of Zahn**<sup>Q</sup>
- Line of Zahn are due to **alternate pale layers of platelets with fibrin & darker layers with more RBCs** (**Fig. 4**). These lines **distinguish antemortem clots** from the bland non-laminated clots postmortem<sup>Q</sup> clots
- **Postmortem clot**: **Gelatinous & not attached**<sup>Q</sup> to the underlying wall (in contrast to thrombus<sup>Q</sup>), **Dark red dependent portion** where red cells have settled by gravity and yellow "**chicken fat**"<sup>Q</sup> upper portion (**Fig. 5**).



**Fig. 3:** Vegetations on heart valves



**Fig. 4:** Line of Zahn



**Fig. 5:** Postmortem clot

## Fate of the Thrombus

**Thrombus** undergoes four events in the ensuing days to weeks: **possible fates DOPE**:<sup>Q</sup>

- **Dissolution**-is the result of **fibrinolysis**
- **Organization & repair**-in older thrombi; by ingrowth of fibroblasts, endothelial & smooth muscle cells
- **Propagation**-by accumulation of additional platelets and fibrin
- **Embolization**-Thrombi dislodge and travel to other sites in the vasculature

## Pulmonary Emboli (Fig. 6)


- **Most common form of thromboembolic disease**.<sup>Q</sup>
- Most arise in the **deep leg veins above the level of knees**.<sup>Q</sup>
- 80% are **clinically silent**<sup>Q</sup> because of **dual circulation and small size**
- **Multiple emboli over time cause pulmonary hypertension and right ventricular failure**<sup>Q</sup>
- >60% obstruction in pulmonary circulation → sudden death or cor pulmonale
- **Paradoxical embolus**<sup>Q</sup> can pass through interatrial/inter-ventricular defect, thus entering the systemic circulation.<sup>Q</sup>

## DISSEMINATED INTRAVASCULAR COAGULATION (DIC)/CONSUMPTION COAGULOPATHY


Discussed in Chapter 11.

## EMBOLISM

- 'Embolus' is a **detached intravascular solid, liquid, or gaseous mass** that is carried by the blood from its point of origin to a distant site<sup>Q</sup>,
- The vast majority of emboli are dislodged thrombi, hence the term **thromboembolism**.<sup>Q</sup>



**Fig. 6:** Pulmonary emboli : Gross appearance



**Figs 7A and B:** A. Mic showing fat and hematopoietic elements; B. Fat emboli highlighted by oil red O

### Systemic Thromboembolism or Arterial Emboli

- Most systemic emboli (80%) arise from **intracardiac mural thrombi**<sup>Q</sup>
- 2/3rd associated with **left ventricular wall infarcts** and 1/4th with **left atrial dilation and fibrillation**
- Major sites:** **lower extremities** (75%) > brain (10%) > intestines > kidneys > spleen

### Fat and Marrow Embolism (Figs 7A and B)

- Occurs in **90% of individuals** with severe skeletal injuries
- MC cause is Fractures of **long bones (which contain fatty marrow)**
- Fat embolism syndrome is characterized by pulmonary insufficiency, neurologic symptoms, anemia & thrombocytopenia (petechial rash) typically seen **1 to 3 days after injury**
- Mechanical obstruction & free fatty acids causing **local toxic injury to endothelium**.<sup>Q</sup>
- Lab findings-** thrombocytopenia<sup>Q</sup>, anemia, fat **microglobulinemia**<sup>Q</sup> (not **macroglobulinemia**<sup>Q</sup>) & fat globules in urine<sup>Q</sup>

### Mnemonic

#### Fat embolism: findings

#### "Fat, Bat, Fract"

- Fat in urine, sputum
- Bat-wing lung x-ray
- Fracture history


Also, fracture of FEMur causes Fat EMboli.

### Air Embolism

- >**100 cc**, is necessary to produce a clinical effect in the **pulmonary circulation**
- Decompression sickness** occurs when **there is sudden decreases in atmospheric pressure**.
- Characterized by **pends and chokes**:
  - Bends:** **painful condition**<sup>Q</sup> due to rapid gas bubble formation within skeletal muscles & supporting tissues around joints
  - Chokes:** Respiratory distress due to gas bubbles in vessels causing edema, hemorrhage & focal atelectasis or emphysema
- Caisson disease (chronic form of decompression sickness)**- persistence of gas emboli in the skeletal system causes multiple foci of ischemic necrosis<sup>Q</sup>; the **more common sites** are the **femoral heads, tibia, and humeri**.<sup>Q</sup>

### Amniotic Fluid Embolism (Fig. 8)

- Fifth most common cause of maternal mortality** worldwide<sup>Q</sup>
- Ominous complication of **labor and immediate postpartum period**.
- Due to infusion of amniotic fluid or fetal tissue into the maternal circulation via **a tear in the placental membranes or rupture of uterine veins**
- Lead to severe dyspnea and pulmonary edema



**Fig. 8:** Mic shows amniotic membranes s/o amniotic fluid embolism in lung alveoli


### INFARCT

- An infarct is an area of **ischemic necrosis** due to occlusion of either arterial supply or venous drainage
- Nearly 99% of all infarcts result from **thrombotic or embolic events**.<sup>Q</sup>
- More in organs with a **single venous outflow** like testis and ovary
- The infarcts may be either red (hemorrhagic) or white (anemic) and may be either septic or bland.







Feature	Red Infarcts <sup>a</sup>	White infarcts <sup>a</sup>
Cause	<ul style="list-style-type: none"> <li>• <b>Dual circulations</b> (e.g., lung &amp; small intestine)</li> <li>• <b>Venous occlusion<sup>a</sup></b> (ovarian torsion)</li> <li>• Loose, spongy tissues (e.g., lung)</li> <li>• <b>Tissues previously congested by sluggish venous flow<sup>a</sup></b></li> <li>• When flow is <b>reestablished<sup>a</sup></b> to a site of previous arterial occlusion and necrosis (e.g., following angioplasty)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>End Arterial Circulation</b></li> <li>• <b>Arterial occlusions</b></li> <li>• <b>Solid organs</b> e.g., heart, spleen, and kidney</li> <li>• <b>Solid tissue<sup>a</sup></b> -tissue density limits the <b>seepage<sup>a</sup></b> of blood from adjoining capillary beds into the necrotic area.</li> </ul>
Organs	Loose organs <sup>a</sup> like Lung and small intestine ( <b>Figs 9 to 11</b> )	Solid organs <sup>a</sup> (heart, spleen, kidney) ( <b>Figs 12 to 14</b> )
Color	Hemorrhagic	Pale <sup>a</sup> & progressively paler with time
Margins	Ill-defined hemorrhagic margins, <sup>a</sup> brown in color	Well defined margins
Edema	Usually <b>present<sup>a</sup></b>	Usually <b>absent<sup>a</sup></b>




**Fig. 9:** White and red infarcts sites in different body organs




**Fig. 11:** Red infarct in intestine




**Fig. 12:** Gross showing white infarct in spleen



**Fig. 13:** White infarct in kidney



**Fig. 14:** White infarct in myocardium



**Fig. 10:** Red infarct in lung : Gross appearance



## High Yield Facts

- Liver can show both white as well as red infarcts
- Infarcts are **wedge-shaped<sup>Q</sup>**, with **occluded vessel at the apex and the periphery of the organ forming the base<sup>Q</sup>**
- All infarcts microscopically has features of **ischemic coagulative necrosis<sup>Q</sup>** except brain (**liquefactive necrosis<sup>Q</sup>**)

### Coralline thrombus

In veins thrombi form coral-like system with framework of platelets, fibrin and trapped white blood cells

### Septic infarctions

- Occur when infected **cardiac valve vegetations embolize or when microbes seed necrotic tissue**
- In these cases, the **infarct is converted into an abscess**, with a correspondingly greater inflammatory response

## SHOCK

A state in which **diminished cardiac output** or **reduced effective circulating blood volume** impairs **tissue perfusion** and leads to **cellular hypoxia**.

### Types

These are three main types of shock

Types of shock	Cardiogenic shock	Hypovolemic shock	Shock associated with systemic inflammation
<b>Cause</b>	Cardiac pump failure	Inadequate blood or plasma volume	Activation of <b>cytokine cascade<sup>Q</sup></b> ; peripheral vasodilation and pooling of blood <sup>Q</sup> ; endothelial activation/injury; <sup>Q</sup> leukocyte-induced damage <sup>Q</sup> , disseminated intravascular coagulation <sup>Q</sup>
<b>Associated conditions</b>	MI, cardiac arrhythmia, cardiac tamponade and pulmonary embolism	Hemorrhage, severe burns and severe dehydration	Microbial infections, burns, trauma, and or pancreatitis

### Other Types

- Neurogenic shock: seen with **anesthetic accident or a spinal cord injury<sup>Q</sup>**
- Anaphylactic shock: Due to generalized vasodilation (**type I hypersensitivity**)<sup>Q</sup>

### Stages of Shock

Stages	Phase	Features
I	<b>Non-progressive</b>	<b>Reflex compensatory mechanisms activated<sup>Q</sup></b> & perfusion of vital organs <b>maintained</b>
II	<b>Progressive<sup>Q</sup></b>	<b>Tissue hypoperfusion, worsening circulatory &amp; metabolic imbalances, lactic acidosis<sup>Q</sup></b>
III	<b>Irreversible<sup>Q</sup></b>	Having <b>irreversible tissue injury<sup>Q</sup></b> and multiple organ failure <sup>Q</sup>




## High Yield Facts

- **Primary initiating factor in septic shock: cytokine release** → **endothelial damage** → systemic inflammatory response → vasodilation, increased permeability & DIC
- Most common cause of septic shock is Gram-positive bacterial infections, followed by Gram-negative bacteria & fungi.
- Most common cytokine involved in Septic shock is **TNF-α**





## Pathogenesis of Septic Shock



## Features of Shock

Changes can manifest in any tissue—**Particularly evident in brain<sup>Q</sup>, heart<sup>Q</sup>, lungs<sup>Q</sup>, kidneys<sup>Q</sup>, adrenals, and gastrointestinal tract.**

<b>Adrenal</b> ↓ Cortical cell lipid depletion <sup>Q</sup>	<b>Kidneys</b> ↓ Acute tubular necrosis <sup>Q</sup>	<b>Heart</b> ↓ Coagulative necrosis or contraction band necrosis <sup>Q</sup>	<b>Brain</b> ↓ Ischemic encephalopathy <sup>Q</sup>	<b>Liver</b> ↓ Fatty change with hemorrhagic central necrosis; 'shock liver' <sup>Q</sup>	<b>GIT</b> ↓ Hemorrhagic Enteropathy <sup>Q</sup>
---	--	---	---	---	---

## Histological Features of Shock

Organ	Changes
<b>Adrenal</b>	Cortical cell lipid depletion. <sup>Q</sup>
<b>Kidneys</b>	Acute tubular necrosis <sup>Q</sup>
<b>Lungs</b>	Diffuse alveolar damage (DAD) <sup>Q</sup>
<b>Heart</b>	Contraction band necrosis <sup>Q</sup>
<b>Brain</b>	Ischemic encephalopathy <sup>Q</sup>
<b>Liver</b>	Fatty change with hemorrhagic central necrosis
<b>GIT</b>	Hemorrhagic Enteropathy <sup>Q</sup>

### *R10<sup>th</sup>* Latest Update

- LPS and lipoteichoic acids are PAMPs
- Procalcitonin is acute phase reactant → MARKER OF SEPSIS

### Definitions of sepsis


Criteria for SIRS<sup>a</sup>

Two or more of the following are required:

- Body temperature >38°C or <36°C
- Heart rate >90 beats/min
- Respiratory rate >20 breaths/min (or arterial pCO<sub>2</sub> <32 mmHg, indicating hyperventilation)
- White blood cell count >12.0 × 10<sup>9</sup>/L or <4.0 × 10<sup>9</sup>/L (or >10% immature forms)


Sepsis = Infection + SIRS

Severe sepsis = Sepsis + evidence of organ dysfunction



## Image-Based Questions

1. A 50-year-old male presents with chronic right heart failure. The succumbs to his illness. On autopsy, liver shows following changes:




- a. Nutmeg liver
- b. Normal liver
- c. Hemosiderosis liver
- d. Liver failure

3. A 64-year-old male died due to RTA. On examination, his vessels showed following clots. Diagnosis is:




- a. Lines of Zahn
- b. Chicken fat thrombus
- c. Red cells
- d. White cells

2. A 20-year-old male was found dead in his home. Family claimed it as dead due to sudden MI. What is the best test to distinguish antemortem from post-mortem clot?




- a. Lines of Zahn
- b. Chicken fat
- c. Red cells
- d. White cells

4. A 34-year-old male patient presented with left-upper-quadrant abdominal pain. On CT with contrast, diagnosis of splenic infarct was made. Identify the type of infarct.



- a. White infarct
- b. Red infarct
- c. Both
- d. None



## Answers of Image-Based Questions

### 1. Ans. (a) **Nutmeg liver**

- Pathologically, the term nutmeg liver refers to the speckled appearance of the cut liver in chronic venous congestion, due to dilated and congested red central veins surrounded by paler, unaffected liver tissue (resembling a grated nutmeg kernel)
- Note the dark red congested regions that represent accumulation of RBC's in centrilobular regions
- Microscopically, the nutmeg pattern results from congestion around the central veins
- Nutmeg liver is **most frequently** seen in **right heart failure**<sup>Q</sup>
- If the passive congestion is pronounced, then there can be centrilobular necrosis
- If chronic hepatic passive congestion continues for a long time, a condition called "**cardiac cirrhosis**"<sup>Q</sup>

### 2. Ans. (a) **Lines of Zahn**

- Line of Zahn are due to **alternate pale layers of platelets with fibrin & darker layers (marked with arrow) with more RBCs**.
- These lines **distinguish antemortem clots** from the bland non-laminated clots postmortem<sup>Q</sup> clots

### 3. Ans. (b) **Chicken Fat Thrombus/ Postmortem Clot**

A. Chicken fat


B. Current jelly (red cells)

- Postmortem clots are not attached to vessel walls
- Show separation of red cells and plasma so that the clotted plasma resembles "chicken fat" layered on top of a rubber gelatinous dark red mass of erythrocytes resembling "current jelly"

### 4. Ans. (a) **White Infarct**

- **Wedge-shaped**<sup>Q</sup>, with **occluded vessel at the apex and the periphery of the organ forming the base**<sup>Q</sup>
- **It is usually seen in solid organs** e.g., heart, spleen, and kidney
- **Solid tissue**<sup>Q</sup> density **limits the seepage**<sup>Q</sup> of blood from adjoining capillary beds into the necrotic area.






## Multiple Choice Questions

### HYPEREMIA AND CONGESTION

- Edema is due to:** (JIPMER 2016, Recent Question 2015)
  - Lymphatic obstruction
  - Decreased hydrostatic pressure
  - Increased oncotic pressure
  - Increased plasma proteins
- Liver biopsy in chronic RHF shows:** (Recent Question 2015)
  - Shrinking
  - Greasy fatty liver
  - Hypertrophy
  - Congestion
- Heart failure cells are** (Recent Question 2015)
  - Foam cells
  - Lipid laden macrophages
  - Hemosiderin laden macrophages
  - Type 1 pneumocytes
- The given figure shows which of the following?** (AIIMS Nov 2015)



- Amyloidosis-grew; viable white necrotic
  - Nutmeg liver-red areas are viable pericentral areas; white areas are periportal necrotic areas
  - Red areas are necrotic areas near central vein, white areas are viable, fibrotic periportal area
  - Amyloidosis-necrotic white periportal viable gray pericentral areas
- Nutmeg liver seen in** (Recent Question 2014-15)
    - Alcoholic liver disease
    - Chronic venous congestion
    - Hepatoma
    - Secondary carcinoma deposits in liver
  - Heart failure cells are seen in** (Recent Question 2014-15)
    - Heart
    - Lungs
    - Kidney
    - Liver
  - Edema occurs when plasma protein level is below:** (Recent Question 2015)
    - 8 g/dL
    - 2 g/dL
    - 5 g/dL
    - 10 g/dL

### THROMBOSIS

- Which of the following has a major role in thrombus formation?** (Recent Pattern Question 2020)
  - Endothelial injury
  - Vasoconstriction
  - Platelet activation
  - Coagulation cascade
- All the following are features of arterial thrombus except:** (Recent Question 2015)
  - Associated with endothelial injury
  - Lines of Zahn present
  - Also called white thrombi
  - Grows in an antegrade manner from the point of attachment
- Lines of Zahn are seen in:** (PGI Nov 2015)
  - Antemortem clots
  - Postmortem clots
  - Coralline thrombus
  - Line of Zahn are due to alternate pale layers of platelets with fibrin & darker layers with more RBCs
  - Lines of Zahn are thrombi (both arterial & venous) with laminations
- True about thrombus formation:** (PGI May 2015)
  - Arterial thrombus grow in direction toward heart
  - Venous thrombus grow in direction toward heart
  - Venous thrombus form chicken fat
  - Line of Zahn is seen microscopically in red thrombi
- Virchow's triad for thrombosis include all EXCEPT:** (MH PG 2014)
  - Endothelial injury
  - Stasis
  - Platelet aggregation
  - Hypercoagulability
- All are predisposing factors of Deep Vein thrombosis, except:** (Recent Question 2015)
  - HIT
  - Polycythemia
  - Malignancy
  - Hyperlipidemia
- All are hypercoagulable states, except:** (AIIMS June 12)
  - Protein C resistance
  - Protein S deficiency
  - Antiphospholipid antibody
  - Polycythemia
- Which of the following DOES NOT present with recurrent episodes of upper limb thrombosis:** (AIIMS June 11)
  - Prostatic Ca.
  - Pancreatic Ca,
  - Osteosarcoma
  - Acute pro myelocytic leukemia
- Both arterial and venous thrombosis occur in:** (PGI Nov 2011)
  - Antiphospholipid antibodies
  - Antithrombin III deficiency
  - Hyperhomocysteinemia
  - Protein C deficiency
  - Mutation in factor V gene






17. **Hypercoagulability due to defective factor V gene is called**  
 a. Lisbon mutation (AIIMS May 10)  
 b. Leiden mutation  
 c. Antiphospholipid syndrome  
 d. Inducible thrombocytopenia syndrome

### EMBOLISM

18. **Patient after trauma has respiratory discomfort. Ventilation tried but not helpful histopath of lung given.** (AIIMS Nov 2016)



- a. Diffuse alveolar hmg with pul edema  
 b. Diffuse damage due to ventilation pressure  
 c. FAT embolism  
 d. None
19. **Least affected organ in arterial thromboembolism is?** (MH PG 2016, Recent Question 2015)  
 a. Liver b. Kidney  
 c. Heart d. Brain
20. **Minimum quantity of air in pulmonary circulation to cause clinical effects** (Recent Question 2015)  
 a. 10 mL b. 50 mL  
 c. 100 mL d. 500 mL
21. **The most common source of embolism:** (Recent Question 2014)  
 a. DVT b. Trauma  
 c. Infection d. Surgery

### INFARCT

22. **White infarcts are seen in all the following organs except:** (Recent Question 2015)  
 a. Lung b. Heart  
 c. Spleen d. Kidney
23. **Red infarct is seen in:** (Recent Question 2015)  
 a. Lung b. Heart  
 c. Kidney d. Spleen
24. **Red infarct occur in:** (PGI May 2015)  
 a. In tissues with dual circulations  
 b. Occur only when both arterial and venous obstruction occurs simultaneously  
 c. Organs which are previously congested  
 d. Organs with loose tissue


25. **White infarct is seen in:** (Recent Question 2015)  
 a. Lung b. Intestine  
 c. kidney d. Ovary
26. **Pale infarcts are seen at all of the following sites except:** (AIIMS Nov 10/ AI 97)  
 a. Heart b. Spleen  
 c. Kidney d. Lung

### SHOCK

27. **Endothelium activation refers to:** (Kerela PG 2016, Recent Question 2014-15)  
 a. Aberration of anatomy of vessel wall  
 b. Irreversible changes in functional state of vessel wall  
 c. Smooth muscle proliferation  
 d. Increased expression of adhesion molecules for leukocyte recruitment
28. **Septic Shock is due to:** (Recent Question 2015)  
 a. Vasodilatation b. Decreased Cardiac output  
 c. Endothelial damage d. All of the above
29. **MC endogenous pyrogen and shock manifestation are:** (Recent Question 2015)  
 a. IL-6 b. IL-8  
 c. IL 1beta d. TNF alpha
30. **Procalcitonin is used as a marker for?** (DNB Aug 12)  
 a. Sepsis  
 b. Medullary carcinoma of thyroid  
 c. Vitamin D resistant rickets  
 d. Parathyroid adenoma
31. **Endotoxin shock is initiated by:** (AIIMS Nov 10)  
 a. Endothelial injury b. Peripheral vasodilation  
 c. Increased vascular permeability  
 d. Cytokines action

### MISCELLANEOUS

32. **In cornea stem cell present at?** (Recent Question 2016-17)  
 a. Limbus b. Descments membrane  
 c. Basement membrane d. Rods and cones
33. **Weibel-palade bodies are present in:** (AP 2012)  
 a. Vascular endothelial cells  
 b. Warthin finkeldey cells  
 c. Leydig cells  
 d. Dendritic cells
34. **Which one of the following factors is labelled as cytokine in the pathogenesis of systemic inflammatory response syndrome:** (Recent Question 2014)  
 a. Nitric oxide b. Complements  
 c. Leukotrienes d. Tumor Necrosis factor



## Answers with Explanations

1. **Ans. (a) Lymphatic obstruction** (Ref: Robbins 9th/pg 114)
2. **Ans. (d) Congestion** (Ref: Robbins 9th/pg 115; 8th/pg 113)
3. **Ans. (c) Hemosiderin laden macrophages**  
(Ref: Robbin's 9th/pg 115)  
**Chronic pulmonary congestion:** shows **Hemosiderin-laden macrophages (heart failure cells)**<sup>Q</sup>
4. **Ans. (c) Red areas are necrotic areas near central vein, white areas are viable, fibrotic periportal area**  
(Ref: Robbin's 9th/129)  
The given picture shows **Chronic passive hepatic congestion**, the centrilobular regions are red-brown & slightly depressed (because of cell death) & are prominently visible against surrounding zones of uncongested tan liver (**nutmeg liver**).
5. **Ans. (b) Chronic venous congestion** (Ref: R 9th/ 129)
6. **Ans. (b) Lungs** (Ref: Robbin's 9th/pg 115; 8th/pg 113)
7. **Ans. (c) 5 g/dL** (Ref: Harshmohan 5th pg 97)  
When total plasma proteins <5 gm/dl (normal 6-8 gm/dl) or albumin <2.5 gm/dl (normal 3.5-5 gm/dl) edema takes place
8. **Ans. (a) Endothelial injury**  
(Refer to answer 12)
9. **Ans. (d) Grows in an anterograde manner from the point of attachment**  
(Ref: Robbins 9th/pg 122; 8th/pg 121)
10. **Ans. (a, c, d, e) a. Antemortem clots; c. coralline thrombus; d. Line of Zahn are due to alternate pale layers of platelets with fibrin and darker layers with more RBCs; e. Lines of Zahn are thrombi (both arterial and venous) with laminations**  
(Ref: Robbins 9th/pg 122; 8th/pg 121)
  - Thrombi (**both arterial & venous**)<sup>Q</sup> have laminations, called **line of Zahn**<sup>Q</sup>
  - Line of Zahn are due to **alternate pale layers of platelets with fibrin & darker layers with more RBCs**.
  - These lines **distinguish antemortem clots** from the bland non-laminated clots that occur post-mortem

- In veins thrombi form coral-like system with framework of platelets, fibrin and trapped white blood cells this is a coralline thrombus. Lines of Zahn are seen in this.

11. **Ans. (b) Venous thrombus grow in direction toward heart** (Ref: Robbins 9th/pg 122; 8th/pg 121)

12. **Ans. (c) Platelet aggregation** (Ref: Robbin's 9th/pg 122)

13. **Ans. (d) Hyperlipidemia** (Ref: R 9th/pg 122; 8th/pg 121)

14. **Ans. (d) Polycythemia**

(Ref: Robbin's 9th/pg 122; 8th/pg 121)

Polycythemia is cause of hyperviscosity and not hypercoagulability

15. **Ans. (c) Osteosarcoma** (Ref: Robbin's 8th/pg 673)

Cancers causing thrombosis

Pancreas	Acute promyelocytic leukemia	Breast	Brain
Lung	Stomach	Prostate	

Peripheral venous thrombosis with visceral carcinoma esp pancreatic carcinoma is called trousseau syndrome or migratory thrombophlebitis.

16. **Ans. (a, c); a. Antiphospholipid antibodies; c. Hyperhomocysteinemia** (Ref: Harrison 17th pg 367)

Risk factor for thrombosis

	ARTERIAL + VENOUS
<b>Inherited</b>	Homocystinuria, Dysfibrinogenemia
<b>Acquired</b>	<ul style="list-style-type: none"> <li>• Cancer</li> <li>• Disseminated intravascular coagulation</li> <li>• Heparin-induced thrombocytopenia</li> <li>• Antiphospholipid antibody syndrome</li> <li>• PNH, TTP</li> <li>• ET (essential thrombocythemia)</li> <li>• PV (polycythemia vera)</li> </ul>
<b>Inherited + acquired</b>	Hyperhomocysteinemia

17. **Ans. (b) Leiden mutation**

Ref: Robbins 9th/pg 122; 8th/pg 121

- **Factor V mutation (called as Leiden mutation**, after the city in the Netherlands where it was discovered) is **most important cause** of primary hypercoagulability.



- The mutation results in a **glutamine to arginine** substitution at **position 506** that renders factor V **resistant to cleavage by protein C**.

18. Ans. (c) **FAT embolism** (Ref: Robbins 9th/pg 130)

19. Ans. (a) **Liver**

(Ref: Robbins 9th/pg 127; 8th/pg 126, MD Guidelines-Arterial Embolism and Thrombosis by Presley Reed, MD)

Arterial emboli often occur in the legs and feet. Some may occur in the brain, or heart.

Less common sites include the Liver, intestines, and eyes

20. Ans. (c) **100 mL** (Ref: Robbins 9th/pg 127; 8th/pg 126)

- **>100 cc**, is necessary to produce a clinical effect in the **pulmonary circulation**

21. Ans. (a) **DVT** (Ref: Robbins 9th/pg 127; 8th/pg 126)

- MC cause of venous embolism is DVT
- Most arise in the **deep leg veins above the level of the knee**.<sup>9</sup>

22. Ans. (a) **Lung** (Ref: Robbins 9th/pg 129; 8th/pg 128)

A. Red infarct- Lung, Ovary, Intestine

B. White infarct – **kidney, heart, spleen and brain**

Liver infarctions are rare because of liver's dual blood supply. Its infarction is described as dichotomic, i.e. it can show overlap between red to white infarct.

23. Ans. (a) **Lung** (Ref: Robbins 9th/pg 129; 8th/pg 128)

24. Ans. (a, c, d) **a. In tissues with dual circulations; c. Organs which are previously congested; d. Organs with loose tissue** (Ref: Robbins 9th/pg 129; 8th/pg 128)

25. Ans. (c) **Kidney** (Ref: Robbins 9th/pg 129; 8th/pg 128)

26. Ans. (d) **Lung** (Ref: Robbins 9th/pg 129; 8th/pg 128)

27. Ans. (d) **Increased expression of adhesion molecules for leukocyte recruitment**

(Ref: Thromb Res 123 (Suppl 4): S30–4)

**Endothelial activation** is a proinflammatory and procoagulant state of the endothelial cells lining the lumen of blood vessels. It is most characterized by **Increased expression of adhesion molecules for leukocyte recruitment**

28. Ans. (d) **All of the above** (Ref: R 9th/pg 131-132; 8th/pg 131)

29. Ans. (d) **TNF alpha** (Ref: R 9th/pg 131-132; 8th/pg 131)

30. Ans. (a) **Sepsis** (Ref: Critical care forum.com)

- Measurement of procalcitonin can be used as a **marker of severe sepsis** caused by bacteria and generally grades well with the degree of sepsis.

31. Ans. (d) **Cytokines action** (Ref: Robbins 9th/pg 131-132)

32. Ans. (a) **Limbus** (Ref: Robbins 9th/pg 28)

33. Ans. (a) **Vascular endothelial cells**

(Ref: Weibel ER, (October 1964). J. Cell Biol. 23 (1): 101–12)

**Weibel-Palade bodies** are the storage granules of endothelial cells. They store and release von Willebrand factor and P-selectin.

34. Ans. (d) **Tumor Necrosis factor** (Ref: R 9th/pg 131-132)

# Genetic Disorders

## Key Points

- » An **observed trait** is referred to as a **phenotype**<sup>o</sup>
- » **Genetic information** defining the phenotype is called the **genotype**<sup>o</sup>
- » **Locus** is the **position of gene** on a chromosome
- » **Alternative forms** of a gene or a genetic marker are referred to as **alleles**<sup>o</sup>
- » The **normal** or common allele is usually referred to as **wild type**<sup>o</sup>
- » When alleles at a given locus are **identical**, the individual is **homozygous**<sup>o</sup>
- » If alleles are different on maternal and paternal copy of the gene; individual is **heterozygous**<sup>o</sup>
- » If **two different mutant alleles** are inherited **at a given locus**, the individual is said to be a **compound heterozygote**<sup>o</sup>
- » **Hemizygous**<sup>o</sup> is used to describe males with a mutation in an X gene or a female with a loss of one X locus.

## Key Recent Updates

- » **CRISPR**—Knock out technology
- » Gaucher's disease is strongly linked with **Parkinson's disease**
- » **Zebra Bodies** are seen in Niemann Pick disease, metachromatic leukodystrophy and mucopolysaccharidoses.





## POLYMORPHISMS

**Physiological sequence variations<sup>Q</sup>**, have a frequency of at least 1%.

**Types depending on protein coding sequence alteration**

- **Synonymous polymorphism<sup>Q</sup>**: Single base-pair substitutions<sup>Q</sup> that **do not** alter the protein coding sequence
- **Non-synonymous polymorphism**: Alter mRNA stability, translation, or the amino acid sequence<sup>Q</sup>

**Types of Repeat length**

Microsatellite repeats (< 1 kb)<sup>Q</sup> and Mini-satellite repeats (1-3 kb)<sup>Q</sup>

**Polymorphisms detected by**

Linkage analysis<sup>Q</sup> or GWAS<sup>Q</sup>

R<sup>9th</sup>

**Latest Update**

### Human Genome Project

- Initiated in the mid-1980s to characterize the **human genome DNA sequence**.<sup>Q</sup>
- **23 pairs** of human chromosomes encode 23,000–30,000 genes<sup>Q</sup> (3 billion bp)
- **Single nucleotide polymorphisms (SNPs)** that are in **close proximity** are **inherited together** (i.e., they are **linked**) and are called **haplotypes<sup>Q</sup>**, hence the name **HapMap<sup>Q</sup>**
- **Genome-wide association studies (GWAS)<sup>Q</sup>** explains the complex **interactions** among **multiple genes** and **lifestyle factors** in multifactorial disorders.

## MUTATIONS

**Definition**

**Permanent change in DNA<sup>Q</sup>**

**Characteristics**


**Transmitted to the progeny** if germ cells are affected;<sup>Q</sup> give rise to **inherited diseases**

**Mechanisms**

Results from a change in nucleotide base-


- **Transitions** - One **purine** is replaced by **another purine<sup>Q</sup>** base (A → G) or a **pyrimidine** is replaced by **another pyrimidine** (C → T)
- **Transversions**- Changes from a **purine to pyrimidine<sup>Q</sup>**, or vice versa

**Flowchart 1:** Types of mutation





**Flowchart 2:** Classification of genetic disorders




## MENDELIAN DISORDERS

### Autosomal Dominant Disorders

- Manifestation of an allele in even **heterozygous**<sup>o</sup> state
- Its called **vertical inheritance** because of transmission from parent to offspring.

#### Characteristics

- When an **affected person** marries an **unaffected one**, every child has **50% chance of having the disease**<sup>o</sup>
- Incomplete **penetrance** and **variable expressivity**<sup>o</sup>
- Age of onset may be **delayed until adulthood**<sup>o</sup>
- May be **loss-of-function** (more common) or **gain-of-function mutations**



## Mnemonic

### Important Autosomal Dominant Diseases

#### "HEAVY DOMINANT"

- H** : Hypercholesterolemia<sup>o</sup>, Hereditary spherocytosis<sup>o</sup>, HNPCC<sup>o</sup>
- E** : Ehler Danlos syndrome (except type VI)
- A** : Adenomatous polyposis coli
- V** : Von willebrand disease<sup>o</sup>
- Y** : Pseudohypoparathyroidism
- D** : Dystrophia myotonica<sup>o</sup>
- O** : Osteogenesis imperfecta<sup>o</sup>
- M** : Marfan's syndrome<sup>o</sup>
- I** : Intermittent porphyria
- N** : Neurofibromatosis 1 and 2<sup>o</sup>
- A** : Achondroplasia<sup>o</sup>, Adult polycystic kidney disease
- N** : Noonan's syndrome
- T** : Tuberous sclerosis

### High Yield Facts

- **Codominance**<sup>o</sup>: When **both alleles** of a gene pair contribute to the phenotype. E.g., Blood group 'AB' and histocompatibility complex
- **Pleiotropism**<sup>o</sup>: Many end effects due to a **single mutant gene**
- **Genetic heterogeneity**<sup>o</sup>: Same trait produced by mutations at several loci
- **Penetrance**: % of individuals carrying the gene **who express the trait**<sup>o</sup>
- **Variable expressivity**<sup>o</sup>: Variable expression (severe or mild) of a disease, among individuals who carry the **same mutant**
- **Dominant negative**<sup>o</sup>: When a **mutant allele impairs the function of a normal allele**, causing profound deficiency of the final product. E.g.: Osteogenesis imperfecta
- **Gain of function** mutation: Huntingtons disease

### Autosomal Recessive Disorders

They occur only when **both alleles** at a given gene locus are **mutated (homozygous)**<sup>o</sup>

- It is called **horizontal inheritance** : single generation affected.

#### Features:

- Siblings have **one in four chance** of having the trait (**25% for each birth**)<sup>o</sup>
- If **mutant gene** is **uncommon**, affected individual can be the product of a **consanguineous marriage**<sup>o</sup>
- **Complete penetrance** is **common** with **onset early in life**<sup>o</sup>
- Uniform expression.
- All inborn errors of metabolism are the example.



## Mnemonic

### Important Autosomal Recessive Diseases

#### "ABCDEFGHI"

- A** : Albinism<sup>Q</sup>, Alkaptonuria<sup>Q</sup>, Ataxia Telangiectasia<sup>Q</sup>
- B** : Beta (thalassemia<sup>Q</sup>, Sickle cell anemia<sup>Q</sup>)
- C** : Cystic fibrosis<sup>Q</sup>, Congenital adrenal hyperplasia
- D** : Deafness (Sensorineural)
- E** : Emphysema ( $\alpha$ 1-antitrypsin deficiency)
- F** : Friedrich's Ataxia<sup>Q</sup>
- G** : Gaucher disease<sup>Q</sup>, Galactosemia
- H** : Homocystinuria, Hemochromatosis<sup>Q</sup>
- I** : Inborn errors of metabolism<sup>Q</sup>

R<sup>9th</sup>


### Latest Updates

#### Mutations within noncoding sequences

- Mutations in **promoter and enhancer** sequences which interfere with binding of transcription factors and thus lead to a marked reduction in, or total lack of transcription. E.g. Thalassemias
- **Point mutations within introns** lead to **defective splicing of intervening sequences** → **failure** to form **mature mRNA**.

## X-linked Disorders

- All **sex-linked disorders** are **X-linked**, and almost all are **recessive**.<sup>Q</sup>
- **Males** with **mutations affecting the Y-linked genes** are usually **infertile**, and hence there is **no Y-linked inheritance**<sup>Q</sup>



### X-Linked Recessive Disorders

#### Features

- **Affected male transmits** the disorder to **all his daughters** (**carriers**)<sup>Q</sup>
- Heterozygous female usually **does not express** the phenotype because of one normal X chromosome<sup>Q</sup>

- **Heterozygous female** transmits the disorder to **50% sons** (**affected**) and **50% daughters** (**carrier**)<sup>Q</sup>

## Mnemonic

### Important X-linked Recessive Disorders

#### "Lady Hardinge College Girls Don't Care About Foolish Words"

- Lady** : Lesch Nyhan syndrome
- Hardinge** : Hemophilia<sup>Q</sup>, Hunter syndrome<sup>Q</sup>
- College** : Chronic granulomatous disease
- Girls** : G6PD deficiency<sup>Q</sup>
- Don't** : Duchene muscular dystrophy<sup>Q</sup>, Diabetes insipidus
- Care** : Color blindness<sup>Q</sup>
- About** : Agammaglobulinemia
- Foolish** : Fragile X syndrome<sup>Q</sup>
- Words** : Wiskott Aldrich syndrome<sup>Q</sup>

## X-Linked Dominant

#### Features

- **Affected heterozygous female** transmits to **50% sons** and **50% daughters**
- **Affected male** transmits the disease to **all his daughters** (**100%**) but **none of his sons**

## Mnemonic

### Important X-linked Dominant Disorders

#### "Red Rose For All Children"

- Red** : X-linked hypophosphatemic **Rickets**<sup>Q</sup>
- Rose** : **Rett's** syndrome<sup>Q</sup>
- For** : **Fragile X** syndrome<sup>Q</sup> ( $X_R > X_D$ )
- All** : **Alport** syndrome<sup>Q</sup>
- Children** : **Charcot-Marie-Tooth** disease

## SOME IMPORTANT SINGLE GENE DISORDERS

### Disorders Associated with Defect in Structural Proteins

#### Marfan's Syndrome

R<sup>9th</sup>

### Latest Updates

Protein	Fibrillin-1	Fibrillin-2
<b>Gene (chr)</b>	FBN1(Chr 15q21.1)	FBN2 (Chr 5q23.31)
<b>Disease</b>	Marfan's syndrome	Congenital contractual arachnodactyly



## Mnemonic

### Features of Marfan's syndrome

#### "M-A-R-F-A-N-S"

- **M**itral valve prolapse<sup>a</sup>
- **A**ortic aneurysm<sup>a</sup>
- **R**etinal detachment
- **F**ibrillin deficiency<sup>a</sup>
- **A**rachnodactyly (long, tapering fingers)<sup>a</sup>
- **N**egative Nitroprusside test
- **S**uperotemporal subluxation of lens (ectopia lentis)<sup>a</sup>
- **S**keletal changes (Tall with long extremities, pectus excavatum/ carinatum, kyphoscoliosis, dolichocephaly)<sup>a</sup>

## Ehlers-Danlos Syndrome (EDS)

Results from some **defect in the synthesis or structure of fibrillar collagen; joint hyperextensibility and laxity<sup>a</sup>**

### Classification of Ehlers-Danlos Syndromes

EDS Type	Clinical Findings	Inheritance	Gene Defects
Classic (I/II)	Skin and joint hypermobility, atrophic scars, easy bruising	Autosomal dominant	COL5A1, COL5A2
Hypermobility (III)	Joint hypermobility, pain, dislocations	Autosomal dominant	Unknown
Vascular (IV)	Thin skin, arterial or uterine rupture, bruising, small joint hyperextensibility	Autosomal dominant	COL3A1
Kyphoscoliosis (VI)	Hypotonia, joint laxity, congenital scoliosis, ocular fragility	Autosomal recessive	Lysyl hydroxylase
Arthrochalasia (VIIa,b)	Severe joint hypermobility, skin changes (mild), scoliosis, bruising	Autosomal dominant	COL1A1, COL1A2
Dermatosparaxis (VIIc)	Severe skin fragility, cutis laxa, bruising	Autosomal recessive	Procollagen N-peptidase


## High Yield Facts

- All types are of **Ehlers-Danlos Syndromes** are **autosomal dominant** except types VI and VIIc
- **Ehlers-Danlos Syndromes Type IV: most dangerous<sup>a</sup> and type III: Most common<sup>a</sup>**
- Fibrillin is found in **aorta, ligaments and ciliary zonules** that support the lens
- **Cardiovascular lesions: Aortic dissections are most common cause of mortality in Marfan's syndrome**
- **Cystic medial necrosis** leads to **ascending aorta aneurysm<sup>a</sup> and aortic dissection in Marfan's syndrome**
- Waardenburg syndrome is due to mutation of **PAX-3 gene**

## Disorder with Defect in Receptor Proteins

### Familial Hypercholesterolemia

Familial hypercholesterolemia is a "**receptor disease**" that is the consequence of a mutation in the gene encoding the receptor for LDL, which is involved in the transport and metabolism of cholesterol.



## LYSOSOMAL STORAGE DISEASES

Disease	Enzyme def.	Cherry-red spot	Hepatosplenomegaly	Skeletal lesions
GM1 Gangliosidosis <sup>a</sup>	Beta galactosidase <sup>a</sup>	+	+	+
Gaucher's disease <sup>a</sup>	Glucocerebrosidase <sup>a</sup>	-	+	+
Niemann Pick disease	Sphingomyelinase <sup>a</sup>	+	+	-
Tay Sachs disease	Hexosaminidase A <sup>a</sup>	+	-	-






## Gaucher's Disease

Most common lysosomal storage disease<sup>Q</sup>


<b>Basic defect</b>	Accumulation of <b>cerebroside<sup>Q</sup></b> inside <b>mononuclear phagocytic cells<sup>Q</sup></b>
<b>Inheritance</b>	<b>Autosomal recessive<sup>Q</sup></b>
<b>Clinical features</b>	<ul style="list-style-type: none"> <li>• 3 types, of which <b>type I is most common<sup>Q</sup></b></li> <li>• <b>Splenohepatomegaly,<sup>Q</sup> Bone pains/pathologic fractures,<sup>Q</sup> Bruising (thrombocytopenia) and Anemia<sup>Q</sup></b></li> <li>• <b>Neurological features ±/- (present in types II and III)<sup>Q</sup></b></li> </ul>
<b>Diagnosis</b>	<ul style="list-style-type: none"> <li>• <b>Def. glucocerebrosidase<sup>Q</sup></b> in leukocytes/fibroblasts</li> <li>• <b>Gaucher cells<sup>Q</sup> (wrinkled paper appearance of cytoplasm) in bone marrow</b></li> <li>• X Ray long bones: <b>'Erlenmeyer flask deformity'<sup>Q</sup></b></li> </ul>
<b>Treatment</b>	Enzyme Replacement therapy ± Stem Cell transplantation



Crumpled Tissue Paper Appearance—Gaucher Cell

## Niemann-Pick Disease Light Microscopy

**Niemann-Pick disease (NPD)** is a lipid storage disorder that results from the deficiency of a lysosomal enzyme, acid sphingomyelinase. Compare foamy looking cell in Niemann-Pick disease.



Foamy looking cell in Niemann-pick disease.

## R10<sup>th</sup> Latest Updates

### Zebra bodies

Membrane-bound granules containing lamellae found in Schwann cells and macrophages. They are associated with **metachromatic leucodystrophy** and storage diseases such as **Niemann-Pick and mucopolysaccharidoses**. The storage diseases result in an accumulation of sphingomyelin and other phospholipids in the reticuloendothelial system.



Image shows Zebra bodies

## Mucopolysaccharidoses

All Mucopolysaccharidoses are **Autosomal recessive<sup>Q</sup>** except **Hunter's disease** which is **inherited as X linked recessive<sup>Q</sup>**

Type	Name	Clinical features
I	Hurler-Scheie	<b>Coarse face, Corneal Clouding,<sup>Q</sup> Mental retardation, Hepatosplenomegaly, Dysostosis multiplex</b>
II	Hunter	Same as above but <b>No corneal clouding<sup>Q</sup></b>
III	San Fillipo	Only Mental Retardation present
IV	Morquio	<b>Bony abnormalities most severe<sup>Q</sup></b> and corneal clouding may be present
VI	Maroteaux-Lamy	Same as Morquio + Coarse facies + Visceromegaly
VII	Sly	Hepatosplenomegaly + Bony abnormalities

## COMPLEX MULTIGENIC DISORDERS

- Disorders caused by interactions between variant forms of **genes and environmental factors**.
- Occur when many polymorphisms, each with a modest effect and low penetrance, are co-inherited


### Representative Diseases Associated with Multifactorial Inheritance

Adults	Children
Hypertension	Pyloric stenosis
Atherosclerosis	Cleft lip and palate
Diabetes type 2	Congenital heart disease
Allergic diathesis	Meningomyelocele
Psoriasis	Anencephaly
Schizophrenia	Hypospadias
Ankylosing spondylitis	Congenital hip dislocation
Gout	Hirschsprung's diseases








## CHROMOSOMAL DISORDERS

### Numerical Abnormalities of Chromosomes



### Structural Abnormalities of Chromosomes

<b>Deletion</b>	Loss of a portion <sup>o</sup> of a chromosome. E.g.: <b>46, XY, del(15) (p11.2p13.1)</b> describes breakpoints in short arm of chromosome 15 at 15p11.2 and 15p13.1	
<b>Ring chromosome</b>	Due to <b>break at both ends</b> <sup>o</sup> of a chromosome with <b>fusion</b> of the damaged ends. E.g.: <b>46, XY, r(10)</b>	
<b>Inversion</b>	Rearrangement that involves <b>two breaks within a single</b> chromosome with re-insertion of the inverted, intervening segment <ul style="list-style-type: none"> <li><b>Paracentric</b>: inversion involving <b>only one arm</b> of the chromosome</li> <li><b>Pericentric</b>: if the breaks are <b>on opposite sides</b> of the centromere</li> </ul>	
<b>Iso-chromosome<sup>o</sup></b>	1 arm of a chromosome is lost and the remaining arm is duplicated or when the axis of division occurs <b>perpendicular to the normal axis</b> <sup>o</sup> of division	
<b>Translocation</b>	Segment of one chromosome is transferred to another. <ul style="list-style-type: none"> <li><b>Balanced reciprocal</b>: Single breaks in two chromosomes, with exchange of material (<b>without any loss</b> of genetic material)<sup>o</sup> E.g. <b>46, XX, t(9;22)(q34;q11)</b></li> <li><b>Robertsonian translocation (or centric fusion)</b>: a translocation between <b>two acrocentric chromosomes</b>, leading to <b>one very large chromosome</b> and one <b>extremely small one</b>, along with <b>small product being lost</b>.<sup>o</sup></li> </ul>	



## IMPORTANT CYTOGENETIC DISORDERS INVOLVING AUTOSOMES

### Down's Syndrome

<b>Epidemiology</b>	Incidence increases with <b>increasing maternal age<sup>a</sup></b>
<b>Genetics</b>	<b>Trisomy 21;<sup>a</sup> 95% Nondisjunction (most common),<sup>a</sup> 4% Robertsonian-Translocation,<sup>a</sup> 1–2% Mosaicism<sup>a</sup></b>
<b>Clinical features</b>	Mnemonic: <b>“PROBLEMATIC Situation”</b>
<b>Diagnosis</b>	<b>Karyotype</b> shows <b>Trisomy 21<sup>a</sup></b> , t(21, 22), t(14, 21) <div style="display: flex; justify-content: space-around; width: 100%;"> <span>95% cases</span> <span>4% cases</span> </div>
<b>Antenatal screening</b>	<ul style="list-style-type: none"> <li><b>1<sup>st</sup> trimester:</b> <ul style="list-style-type: none"> <li><b>Noninvasive:</b> Nuchal thickness<sup>a</sup> on USG, Dual marker</li> <li><b>Invasive:</b> Karyotype on Chorionic villus sampling (9–11 wks)</li> </ul> </li> <li><b>2<sup>nd</sup> trimester:</b> <ul style="list-style-type: none"> <li><b>Noninvasive-</b> Triple test<sup>a</sup>, quadruple test<sup>a</sup></li> <li><b>Invasive:</b> Karyotype from sample obtained by amniocentesis (14–16 wks)</li> </ul> </li> <li><b>Latest antenatal screening tool:</b> Next generation sequencing of chromosome 21 linked genes in total cell free fetal DNA in maternal blood</li> </ul>



Clinical features S/o Down's syndrome

### Mnemonic

Important clinical features of Down's syndrome:  
**“PROBLEMATIC Situation”**

- **P**rotruding tongue
- **R**ound face
- **O**cciput flat/Open, wide fontanelle<sup>a</sup>
- **B**rushfield spots in iris<sup>a</sup>/Brachycephaly/Brachydactyly/Behavioural difficulties
- **L**ow (depressed) nasal bridge/Language problem/less tone (hypotonia)
- **E**picanthic fold<sup>a</sup>/Ears low-set and dysplastic<sup>a</sup> ± hearing problem
- **M**ongoloid slant<sup>a</sup> (oblique palpebral fissure)/Mental retardation<sup>a</sup>/Myoclonus
- **A**cute Leukemia (AML>ALL)/Alzheimer's disease<sup>a</sup>/Atlantoaxial instability<sup>a</sup>/Atresia of duodenum
- **T**risomy 21/thyroid problem (hypothyroidism)
- **I**ncurved 5th finger (**clinodactyly**)<sup>a</sup>/Intellectual disability
- **C**ongenital heart disease<sup>a</sup>/Cataracts
- **S**andle gap<sup>a</sup> (Increased gap between 1st and 2nd toes)/Simian palmar crease<sup>a</sup>



### High Yield Facts

- Down's syndrome is the most common chromosomal abnormality<sup>a</sup>
- Most common genetic cause of mental retardation<sup>a</sup> is Down syndrome
- Down Syndrome is most commonly caused by maternal nondisjunction<sup>a</sup>
- Klinefelter syndrome is most commonly caused by paternal nondisjunction<sup>a</sup>
- The most common congenital heart disease in Down syndrome is **Endocardial cushion Defect<sup>a</sup>** > VSD, ASD, PDA and PAH
- The most common cause of intestinal obstruction in Down's syndrome is **Duodenal atresia<sup>a</sup>**
- MC hematological abnormality in Down's syndrome is **TAM (Transient Abnormal Myelopoiesis)**
- MC leukemia in Down's syndromes < 4 yrs is **AML M7**
- MC leukemia in Down's syndromes > 4 yrs is **ALL**
- In cases of **translocation or mosaic** Down's syndrome, **maternal age is of no importance.**
- If a child with Down's syndrome has translocation (21; 21) recurrence risk is next child is 100%





## Mnemonic

### Other Trisomies

- **Trisomy 13: Patau syndrome<sup>Q</sup>** (Thirteen – PaTau)
- **Trisomy 18: Edward syndrome<sup>Q</sup>** (Eighteen – Edward)

### Important Deletions

**A. 22q11.2 deletion gives rise to: CATCH 22<sup>Q</sup>** (Cardiac anomaly, Anomalous face, Thymus hypoplasia/aplasia, Cleft palate, and Hypocalcemia)

**Previously classified as:** (now considered part of same spectrum)

- **Di-George syndrome** (thymic hypoplasia with diminished T-cell immunity, parathyroid hypoplasia with hypocalcemia) and
- **Velocardiofacial<sup>Q</sup> syndrome** (congenital heart disease, dysmorphism, developmental delay).

**B. Cri du Chat syndrome<sup>Q</sup>: 5p deletion<sup>Q</sup>**; Characteristic cry, developmental delay and behavioral problems.

## CYTOGENETIC DISORDERS INVOLVING SEX CHROMOSOMES

Genetic diseases involving the sex chromosomes are more common than autosomal aberrations.

### Lyon Hypothesis (Lyonisation)<sup>Q</sup>

- **Only one** of the **X chromosomes** is genetically **active**
- The **other X** of **either maternal or paternal** origin undergoes **heteropyknosis** and is rendered **inactive<sup>Q</sup>**
- Inactivation of either the maternal or paternal X **occurs at random<sup>Q</sup>** among all the cells of the blastocyst on or about **day 5.5<sup>Q</sup>** of embryonic life, and
- Inactivation of the same X chromosome **persists in all the cells<sup>Q</sup>** derived from each precursor cell
- The molecular basis of X inactivation involves a unique gene called **XIST<sup>Q</sup>**
- **21% of genes on Xp, and a smaller number (3%) on Xq** escape X inactivation **SRY** (sex-determining region on Y chromosome): dictates testicular development<sup>Q</sup>

### Turner's Syndrome (45, XO)<sup>Q</sup>

Always seen in **FEMALES<sup>Q</sup>**

### Karyotype

- Missing X chromosome in **57% (45, XO) (most common)**
- **Mosaics** in 29% (e.g. 45, XO/ 46, XX)
- Partial monosomy of X chromosome in 14%

## Mnemonic

### Features of Turners Syndrome

#### "C-L-O-W-N-S"

- **Cardiac abnormalities** (**coarctation of aorta**)<sup>Q</sup>, **Cubitus valgus<sup>Q</sup>**
- **Lymphedema**
- **Ovaries underdeveloped: "streak ovaries"<sup>Q</sup>** (causing infertility, amenorrhea)
- **Webbed neck**
- **Normal intelligence<sup>Q</sup>/Nipples widely spaced**
- **Short stature<sup>Q</sup>/Short 4<sup>th</sup> metacarpal<sup>Q</sup>**



### High Yield Facts

- Most important cause of **primary amenorrhea<sup>Q</sup>**: Turner's Syndrome
- **Infertility<sup>Q</sup>** due to rudimentary uterus and **streak ovaries<sup>Q</sup>** (ovaries are reduced to **atrophic fibrous strands without ova and follicles**)
- Reduced ovarian feedback results in significantly **elevated FSH and LH levels<sup>Q</sup>**
- **Most common CVS abnormality** in Turner's syndrome: **Bicuspid aortic valve (50%)<sup>Q</sup>** > Coarctation of Aorta
- In Turner's syndrome, there is increased incidence of **cystic hygroma and gonadoblastoma<sup>Q</sup>**
- **Noonan's syndrome: AD**, seen in **males as well as females**; **Mental retardation present<sup>Q</sup>**
- **Most common CVS abnormality** in **Noonan's syndrome** is **hypertrophic pulmonary stenosis<sup>Q</sup>**

### Klinefelter's Syndrome: 47, XXY (Seen in Males)

*Male hypogonadism that occurs when there are  $\geq 2$  X chr and  $\geq 1$  Y Chr<sup>Q</sup>*

### Genetic Abnormality

- **47, XXY karyotype (90%)<sup>Q</sup>**: results from **nondisjunction during the meiotic division**
- Maternal age is increased in the cases associated with errors in oogenesis.
- **Mosaic patterns (46, XY/47, XXY) (15%)<sup>Q</sup>** and **47, XXY/48, XXXY<sup>Q</sup>**

## Mnemonic

### "K-L-I-N-E-F-E-L-T-S" syndrome

- **Karyotype**, most common is 47, XXY<sup>Q</sup>
- **Long stature with long legs<sup>Q</sup>**
- **Infertility<sup>Q</sup>, Incidence: 1 in 1000 men**
- **Non-disjunction of paternal<sup>Q</sup> sex chromosomes**
- **Eunuchoid<sup>Q</sup> body proportions**
- **FSH elevated<sup>Q</sup>, scanty Facial and axillary hair**
- **Estradiol/testosterone ratio elevation** → Gynecomastia<sup>Q</sup>
- **LH elevated, Leukemias** (increased risk for AML)<sup>Q</sup> and breast tumors, **Learning disability**
- **Testosterone reduced, small Testes and penis<sup>Q</sup>**
- **Secondary sexual characters absent**






## Hermaphroditism and Pseudohermaphroditism

1. Hermaphrodite
2. Pseudohermaphrodite

### RECENT EXAM<sup>Q</sup>

1. The term true *hermaphrodite* implies the presence of both ovarian and testicular tissue. In contrast, a *pseudohermaphrodite* represents a disagreement between the phenotypic and gonadal sex.
2. Female pseudohermaphrodite has ovaries but male external genitalia.
3. Male pseudohermaphrodite has testicular tissue but female-type genitalia.



## DISEASES CAUSED BY TRINUCLEOTIDE-REPEAT MUTATIONS

- Disorder with expansion of Trinucleotides
- Can occur in coding or noncoding region.

## Trinucleotide Repeat Disorders

Region	Disease	Repeat	Inheritance	Gene Product
Coding region	X-chromosomal spinobulbar muscular atrophy	CAG <sup>Q</sup>	XR	Androgen receptor
	Huntington's disease (HD)	CAG <sup>Q</sup>	AD	Huntingtin
	Spinocerebellar ataxia type 1 (SCA1)	CAG	AD	Ataxin 1
Noncoding region	Fragile X-syndrome (FRAAX)	CGG <sup>Q</sup>	XR	FMR-1 protein <sup>Q</sup>
	Dystrophia Myotonica (DM)	CTG <sup>Q</sup>	AD	Myotonin protein kinase
	Friedreich's ataxia (FRDA1)	GAA <sup>Q</sup>	AR	Frataxin <sup>Q</sup>


## Fragile X Syndrome

- **Caused by:** A trinucleotide mutation in familial mental retardation-1 (FMR1) gene<sup>Q</sup>
- **Epidemiology:** Second most common genetic cause of mental retardation<sup>Q</sup> after Down's syndrome.
- **Clinical features:**
  - Long face with a large mandible, large ears, large testicles<sup>Q</sup> (macro-orchidism), hyperextensible joints, a high arched palate, and mitral valve prolapse
  - Most distinctive feature is **post-pubertal macro-orchidism<sup>Q</sup>**
- **Anticipation<sup>Q</sup>**
  - Clinical features **worsen with each successive generation<sup>Q</sup>** as the number of repeats increases with successive generations:
    - **Normal population:** 29–55 CGG repeats in the FMR1 gene.
    - **Carrier** males and females: Carry **pre-mutations<sup>Q</sup>** 55–200 CGG repeats that can expand during **oogenesis**.
    - **Full mutation** (fragile X syndrome): 4000 repeats
- **Diagnosis of choice:** PCR-detection of repeats

### High Yield Facts

#### Fragile X Tremor/Ataxia

- Toxic gain of function of CGG containing mRNA
- 20% –premature ovarian failure
- 50%–Progressive neurodegenerative illness




## Mutation in Mitochondrial DNA (mtDNA): Maternal Inheritance<sup>Q</sup>

### Mechanism

- **mtDNA** complement of the zygote is derived **entirely from the ovum<sup>Q</sup>**
- Thus, mutation in **mother's** mtDNA will be **transmitted to all her off springs<sup>Q</sup>**
- mtDNA encodes enzymes involved in **oxidative phosphorylation**



- Mutations of mtDNA primarily affect **organs** most dependent on **oxidative phosphorylation** such as **CNS, skeletal and cardiac muscles, liver, and kidneys.**<sup>Q</sup>



Pedigree shows all progeny of an affected male (shaded squares) are normal, but all children, male and female, of the affected female (shaded circles) manifest diseases

### Important Features of Mitochondrial Diseases

- **Heteroplasmy**<sup>Q</sup>: An individual may harbor **both wild-type and mutant**<sup>Q</sup> mt-DNA
- **Threshold effect**: **Minimum percentage** of mutant mtDNA that must be present in a cell **for the disease to occur**

### Important Mitochondrial Diseases: “K-L-M-N-O-P”

Disease	Phenotype
<b>Kearn-Sayre syndrome (KSS)</b> <sup>Q</sup>	External ophthalmoplegia, heart block, retinal pigmentation, ataxia
<b>Leber Hereditary Optic Neuropathy (LHON)</b> <sup>Q</sup>	Bilateral subacute or acute painless optic atrophy
<b>MELAS</b> <sup>Q</sup>	<b>Mitochondrial encephalomyopathy, lactic acidosis, and stroke like episodes</b> ; may manifest only as diabetes
<b>MERRF</b>	<b>Myoclonic epilepsy, ragged red fibers</b> in muscle, ataxia, increased CSF protein, sensorineural deafness, dementia
<b>NARP</b> <sup>Q</sup> , <b>Leigh disease</b>	Neurogenic weakness, ataxia, and retinitis pigmentosa (NARP)
<b>Chronic progressive external ophthalmoplegia (CPEO)</b> <sup>Q</sup>	Late-onset bilateral ptosis and ophthalmoplegia, proximal muscle weakness, and exercise intolerance
<b>Pearson syndrome</b>	Pancreatic insufficiency, pancytopenia, lactic acidosis

### Genomic Imprinting

- **Definition**
  - Inheritable **preferential expression**<sup>Q</sup> of **one**<sup>Q</sup> of the parental alleles.
- **Types**
  - **Maternal imprinting** refers to transcriptional **silencing of the maternal allele**<sup>Q</sup>

- **Paternal imprinting** implies that the **paternal allele is inactivated**<sup>Q</sup>
- **Site and Time**
  - Imprinting occurs **in the ovum or sperm, before fertilization**, and then is stably **transmitted** to all somatic cells through **mitosis**
- **Mechanism**
  - DNA **methylation** at CG nucleotides
  - Histone H4 **de-acetylation**
  - **Methylation**
- **Examples**
  - Prader-Willi syndrome, Angelman syndrome

### Mechanism of Imprinting: Epigenetic Alterations

<b>Definition</b>	Study of <b>heritable chemical modification of DNA or chromatin</b> that <b>do not alter the DNA sequence</b> <sup>Q</sup> itself.
<b>Examples</b>	<b>Methylation</b> <sup>Q</sup> of DNA, and methylation and acetylation of <b>histones</b> <sup>Q</sup>



### High Yield Facts

- Affected gene in Angelman syndrome = **UBE3A** (imprinted on paternal chromosome. Expressed on maternal chromosome especially in brain).
  - Affected genes in Prader Willi syndrome = **SNORP** family of gene (encode small nuclear RNA → modifies ribosomal RNA)
- Imprinted on maternal chromosome expressed on paternal chromosome
- Epigenetics play role in X chromosome inactivation and imprinting

### Prader-Willi Syndrome

- Characterized by **mental retardation**, short stature, **hypotonia**, profound hyperphagia, **obesity**, small hands and feet, and **hypogonadism**<sup>Q</sup>
- del(15)(q11.2q13); deletion affects the paternally derived chromosome 15

### Angelman syndrome: Mnemonic: “M-A-S-T”

Mentally retarded, Ataxic gait, Seizures, and inappropriate laughter “happy puppets.”


### Mnemonic

Mechanisms in Prader Willi and Angelman syndrome: <sup>Q</sup>		
PRADER WILLI	MECHANISM	ANGELMAN
PATERNAL	DELETION (70%)	MATERNAL
MATERNAL	DISOMY (20-25%)	PATERNAL



## Gonadal Mosaicism

<b>Mechanism</b>	<ul style="list-style-type: none"> <li>• Mutation that <b>occurs post-zygotically</b><sup>q</sup> during early (embryonic) development.</li> <li>• Mutation <b>affects only cells destined to form the gonads</b>, the <b>gametes carry the mutation</b><sup>q</sup>, therefore the somatic cells of the individual are completely normal.</li> </ul>
<b>Effect</b>	In some <b>autosomal dominant</b> disorders, like by <b>osteogenesis imperfecta</b> <sup>q</sup> , <b>phenotypically normal parents</b> have <b>more than one affected child</b> .




## DIAGNOSTIC TOOLS IN GENETICS

### Karyotype

<b>Obtained by</b>	Arranging each pair of autosomes <b>according to length</b> <sup>q</sup> , followed by sex chromosomes.				
<b>Normal</b>	<b>46, XX (females)</b> <sup>q</sup> and <b>46, XY (males)</b> <sup>q</sup>				
<b>Method</b>	Chromosomes are examined after <b>arresting dividing cells</b> in <b>metaphase</b> <sup>q</sup> with mitotic spindle inhibitors (e.g., N-diacetyl-N-methyl <b>colchicine</b> ), followed by staining				
<b>Samples used</b> <sup>q</sup>	<b>Antenatal</b>		<b>Postnatal</b>		
	<ul style="list-style-type: none"><li>• <b>Chorionic villi sampling (11-13 weeks)</b><sup>q</sup></li><li>• <b>Amniocentesis (14-16 weeks)</b><sup>q</sup></li><li>• Fetal umbilical blood</li><li>• <b>Circulating fetal DNA in maternal blood</b></li></ul>		<ul style="list-style-type: none"><li>• <b>Peripheral blood lymphocytes</b><sup>q</sup> (<b>Best</b>) (<b>not monocytes</b>)<sup>q</sup></li><li>• Bone marrow</li><li>• <b>Skin fibroblasts</b><sup>q</sup></li><li>• Lymph node tissue, Solid tumor sample</li></ul>		
<b>Resolution</b>	<b>2 to 5 million base (mb) pairs</b> <sup>q</sup> (in light microscope), Max resolution in <b>prophase</b> <sup>q</sup>				
<b>Staining methods</b>		<b>G-Banding</b> <sup>q</sup>	<b>R-Banding</b>	<b>Q- Banding</b>	<b>C-Banding</b>
	<b>Pre-treatment</b>	Digestion with Trypsin	Heat Denaturation	None	Chemically extract DNA
	<b>Stain used</b>	<b>Giemsa</b> <sup>q</sup>	Giemsa	<b>Quinacrine mustard</b> <sup>q</sup>	Giemsa
	<b>Microscopy</b>	Light	Light	Fluorescent	Light

### Description of a Karyotype

- **Total number** of chromosomes with **sex chromosome**
- Arrange chromosomes in **ascending**<sup>q</sup> **numerical order**
- **Short arm** of a chromosome is designated **p (for petit)**<sup>q</sup> and **long arm** is referred to as **q**<sup>q</sup>
- The **regions** are numbered (e.g., 1, 2, 3) from the centromere outward
- Each region is further subdivided into **bands and sub-bands**, and these are ordered numerically
- E.g. 6p12.3= short arm of chromosome 6, region 1, band 2, and sub-band 3



### Classification of Chromosomes


Group	Size and centromere position	Ideogram number
A or I	Large : Metacentric/submetacentric	1–3
B or II	Large : Submetacentric	4, 5
C or III	Medium : Submetacentric	6 – 12 and X

Contd...






D or IV	Medium : Acrocentric	13-15
E or V	Small : Metacentric/ Submetacentric	16-18
F or VI	Small : Metacentric	19-20
G or VII	Smallest : Acrocentric	21, 22 and Y







### 10<sup>th</sup> Latest Updates

- **CRISPR**—**Clustered regularly interspaced short palindromic repeats (CRISPR)**: Form the basis of a genome editing technology known as CRISPR-Cas9 that allows permanent modification of genes within organisms. (explained in detail in chapter 1)
- **CRISPR** is a molecule that finds a string of DNA code, locks on and makes a precision cut. And because scientists can tune it to target any genetic sequence, they can use it to turn genes off or replace them with new versions.
  - When viruses attack a bacterial cell, for instance, they inject a payload of their own DNA into the cell.
  - The cell responds by deploying CRISPR, which consists of a strand of ribonucleic acid, or RNA, hooked up to an enzyme called CRISPR associated protein, or Cas.
  - The RNA is primed to recognise and dock to the virus DNA, neatly encasing it in a pocket of the Cas enzyme. Cas, in turn, makes a cut in the DNA, which disables the virus' attack.
- **Gaucher** is strongly linked with Parkinson (enzyme glucocerebrosidase has reciprocal relation with alpha synuclein)
- **Marfan** syndrome is gain of function mutation in TGF- $\beta$  TYPE II receptor- can be treated with losartan (antihypertensive)



## NEXT Pattern Questions



Q's

1. A 3-year-old boy has a history of progressive developmental delay, ataxia, seizures, and inappropriate laughter since infancy. The child has a normal karyotype of 46,XY, but DNA analysis shows that he has inherited both of his number 15 chromosomes from his father. These findings are most likely to be indicative of which of the following genetic mechanisms?
  - a. Genomic imprinting
  - b. Maternal inheritance pattern
  - c. Mutation of mitochondrial DNA
  - d. Trinucleotide repeat expansion

Ans. (a) **Genomic imprinting**

(Ref: Robins Basic Pathology 10th ed/pg 271)

- This child has features of Angelman syndrome, and the DNA analysis shows uniparental disomy. The Angelman gene encoded on chromosome 15 is subject to genomic imprinting. It is silenced on the paternal chromosome 15, but is active on the maternal chromosome 15. If the child lacks maternal chromosome 15, there is no active Angelman gene in the somatic cells. This gives rise to the abnormalities typical of this disorder. The same effect occurs when there is a deletion of the Angelman gene from the maternal chromosome 15. The other listed options do not occur in uniparental disomy.



Q's

2. A 15-year-old girl has developed multiple nodules on her skin over the past 10 years. On physical examination, there are 20 scattered, 0.3–1 cm, firm nodules on the patient's trunk and extremities. There are 12 light brown macules averaging 2–5 cm in diameter on the skin of the trunk. Slit-lamp examination shows pigmented nodules in the iris. A sibling and a parent are similarly affected. Genetic analysis shows a loss-of-function mutation. Which of the following inheritance patterns is most likely to be present in this family?
- Autosomal dominant
  - Autosomal recessive
  - Mitochondrial
  - X-linked recessive

Ans. (a) **Autosomal dominant**

(Ref: Robins Basic Pathology 10th ed/pg 245)

- Neurofibromatosis type 1 (NF-1) is characterized by the development of multiple neurofibromas and pigmented skin lesions. Neurofibromas are most numerous in the dermis but also may occur in visceral organs. Patients with NF-1 also may develop a type of sarcomatous neoplasm known as a *malignant peripheral nerve sheath tumor (MPNST)*. NF-1 is a tumor suppressor that appears with an autosomal dominant pattern of inheritance, though some cases result from spontaneous new mutations (no prior family members with the mutation). NF-1 exhibits variable expressivity, because the manifestations (location and types of neoplasms) are not the same in all patients. The other forms of inheritance listed are not associated with tumor suppressor genes.




Q's

3. An 8-year-old girl experiences sudden severe dyspnea. On examination, she has upper airway obstruction from soft tissue swelling in her neck. A radiograph shows a hematoma compressing the trachea. She was then diagnosed to have hemophilia A. Both parents and two female siblings are unaffected by this problem, but a male sibling has experienced a similar episode. Which of the following genetic abnormalities is most likely to account for the findings in this girl?
- Autosomal dominant mutation
  - Genomic imprinting
  - Germline mosaicism
  - Random X inactivation

Ans. (d) **Random X inactivation**


(Ref: Robins Basic Pathology 10th ed/pg 267)

- This girl has features of hemophilia A. This X-linked recessive condition is expected to occur in males who inherit the one maternal X chromosome with the genetic mutation, and they do not have another X chromosome with a normal functional allele, as is the case in her brother. Hemophilia in a female can be explained by the Lyon hypothesis, which states that only one X chromosome in a female is active (the "turned off" X chromosome is the Barr body) for most genes, but this inactivation is a random event. Some unlucky females are out on the tail end of the Poisson distribution of random events and have few active X chromosomes with the normal allele, leading to markedly diminished factor VIII activity. The other choices do not explain this phenomenon.




## Image-Based Questions

1. The given pedigree shows which type of inheritance?




- a. Autosomal dominant
- b. Autosomal Recessive
- c. X-linked Dominant
- d. X-linked Recessive

2. The given pedigree shows which type of inheritance?




- a. Autosomal dominant
- b. Autosomal Recessive
- c. X-linked Dominant
- d. X-linked Recessive

3. Resolution of light microscope when viewing this is?




- a. 5 Kb
- b. 5 Mb
- c. 500 Kb
- d. 500 Mb

4. What is the locus of the gene marked as arrow in the given chromosome?




- a. 1p3.2
- b. 1q3.2
- c. 1q2.3
- d. 1q23.0

5. Identify the abnormality in the given karyotype?




- a. Normal
- b. Aneuploidy
- c. Triploidy
- d. Trisomy


6. Identify the chromosome in the given sequence?



- a. Telocentric, Metacentric, Acrocentric
- b. Telocentric, Acrocentric, Metacentric
- c. Metacentric, Sub-metacentric, Telocentric
- d. Telocentric, Metacentric, Acrocentric




7. The given figure shows a technique important for diagnosing chromosomal aberrations. What is the technique and abnormality seen?



- a. Fluorescence in-situ hybridization, Deletion
- b. Fluorescence in-situ hybridization, Translocation
- c. Comparative genomic hybridization, Deletion
- d. Chromosomal painting, Translocation

8. Identify the technique used?



- a. CGH
- b. FISH
- c. Microarray
- d. Next gen sequencing



## Answers of Image-Based Questions

1. Ans. (a) **Autosomal dominant**
  - One affected parent causing 50% affected progeny suggests autosomal dominant pattern
2. Ans. (b) **Autosomal Recessive**
  - Both carrier parents have 25% affected child, 50% carrier suggests autosomal recessive pattern
3. Ans. (b) **5Mb**
  - Karyotype is study of chromosomes in which each pair of autosomes are arranged according to length, followed by sex chromosomes.
4. Ans. (c) **1q2.3**
  - Locus of a gene is number, arm, region and band
5. Ans. (c) **Triploidy**
  - Triploidy is condition in which each set of chromosome has 3 chromosomes, so total is  $23 \times 3 = 69$
6. Ans. (c) **Metacentric, Sub-metacentric, Telocentric**
  - The category of chromosome is according to centromere position, middle is metacentric, middle to middle is submetacentric while near to one end is telocentric
7. Ans. (b) **Fluorescence in-situ hybridization, Translocation**
  - The given image shows study of chromosomes by tagging them with fluorescent probes called FISH. In this condition you can notice two colors coming together (left upper) which suggests a translocation among two different chromosomes with different colored probes.
8. Ans. (c) **Microarray**
  - This technique involves study/detection of genes, proteins, DNA or tissue by micro probes attached to a chip and causing reverse hybridization to it called microarray.





## Multiple Choice Questions

**1. Which of the following tumor(s) is/are related to DICER1 Gene mutation: (PGI May 2019)**

- Retinoblastoma
- Pleuropulmonary blastoma
- Cystic nephroma
- Thyroid carcinoma
- Sertoli-Leydig cell tumor

**2. Which of the following is/are chromosomal disorder(s): (PGI May 2019)**

- Noonan syndrome
- Turner syndrome
- Klinefelter's syndrome
- Fragile-X syndrome
- Prader-Willi syndrome

**3. In Down syndrome which is/correct? (PGI May 2018)**

- Most common cause of death is congenital heart disease
- 95% have an extra 21st chromosome
- Increased nuchal fold thickness on USG
- Karyotyping is required in all cases for prenatal diagnosis
- Advanced paternal age (>35 years) is a risk factor

**4. Which chromosome is responsible for the production of MIF? (Recent Question 2018)**

- Chromosome 16
- Chromosome 22
- X Chromosome
- Y Chromosome

**5. Which of the following is/are true about Turners Syndrome? (AIIMS May 2017)**

- Most common viable aneuploidy
- Long stature
- High hair line
- Narrow chest
- Mental retardation present


**6. Which of the following is/are not feature of Turner syndrome? (PGI Nov 2017)**

- Tall stature
- Associated with celiac disease
- 45XO karyotype
- 45XO/46XY karyotype
- Hypertension

**7. Chromosome 21 is which type of chromosome? (JIPMER 2016)**

- Metacentric
- Submetacentric
- Short acrocentric
- Medium acrocentric

**8. Karyotype of a male patient shows the following, what is the clinical abnormality that is expected? (Recent Question 2016)**



- Turner syndrome
- Kallman syndrome
- Androgen insensitivity syndrome
- Down syndrome


**9. Velocardiofacial defect is due to mutation in: (Recent Question 2016-17)**

- Chr 11
- Chr 13
- Chr 22
- Chr 3

**10. Presentation of Pierre-Robin syndrome includes: (Recent Question 2016-17)**

- Retrognathia
- Low set ear
- Prominent forehead
- Isolated cleft palate
- Glossoptosis


**11. Karyotype of a patient shows the following, what is the clinical abnormality that is expected? (AIIMS May 2015)**



- Gynecomastia with long thin limbs
- Short stature with polydactyly
- Webbed neck with widely spaced nipples
- Rocker bottom feet



- 12. In Down's syndrome there is?** (Recent Question 2016)  
 a. Translocation                      b. Mutations  
 c. Paternal nondisjunction  
 d. Maternal nondisjunction
- 13. A female presents with karyotype 45,XO and absent gonads. What is your diagnosis?** (Recent Question 2015)  
 a. Klinefelter's syndrome  
 b. Androgen insensitivity syndrome  
 c. Turner syndrome  
 d. Kallman's syndrome
- 14. Streak gonads are seen in -** (Recent Question 2014)  
 a. Turner syndrome                      b. Klinefelter's syndrome  
 c. Patau's syndrome                      d. Down's syndrome
- 15. True statement regarding chromosomes (JIPMER 2015)**  
 a. In females both X chromosomes are activated  
 b. Klinefelter syndrome results due to an extra Y chromosome in males  
 c. Germinal cells contain 23 chromosomes  
 d. Turner syndrome results due to an extra X chromosome in females
- 16. Cri du chat syndrome is:** (Recent Question 2015)  
 a. 22q-                                      b. 5q-  
 c. 22p-                                      d. 5p-
- 17. Most common cause of primary amenorrhoea with secondary sexual character development is?** (WBPG 16)  
 a. Turner syndrome  
 b. Kallmann syndrome  
 c. Androgen insensitivity syndrome  
 d. Down syndrome
- 18. The following karyotype in female is seen in which of the following syndromes?** (AIIMS Nov 14)



- a. Bloom syndrome  
 b. Fragile X syndrome  
 c. Angelman syndrome  
 d. Cri du chat syndrome
- 19. The structural abnormality of chromosome in which one arm is lost and remaining arm is duplicated is called:** (MA 2016)  
 a. Ring chromosome                      b. Isochromosome  
 c. Translocation                              d. Mutation


- 20. A 16-year-old female presents with primary amenorrhea and raised FSH. On examination, her height was 58 inches. What would be the histopathological finding in the ovary?** (AIIMS Nov 2013)  
 a. Absence of oocytes in the ovaries (streak ovaries)  
 b. Mucinous cystadenoma  
 c. Psammoma bodies  
 d. Hemorrhagic Corpus Luteum
- 21. If a chromosome divides in an axis perpendicular to usual axis of division it is going to form:** (AIIMS May 2013)  
 a. Ring chromosome  
 b. Isochromosome  
 c. Acrocentric chromosome  
 d. Subtelocentric chromosome
- 22. Condition not associated with increased risk of Cancer are all except?** (JIPMER 2013)  
 a. NF-1                                      b. Turner's syndrome  
 c. Down syndrome                      d. Chr 13 deletion
- 23. Most common microdeletion syndrome is?** (DNB Aug 2012)  
 a. WAGR syndrome  
 b. Velo-cardio-facial syndrome  
 c. PraderWilli syndrome  
 d. Angelman syndrome
- 24. Down's syndrome increases risk of:** (PGI May 2011)  
 a. Hirschprung's disease  
 b. Leukemia  
 c. Sensorineural hearing loss  
 d. Hyperthyroidism  
 e. Atlanto-occipital dislocation
- 25. Which of the following is not associated with Down's syndrome?** (AIIMS Nov 11)  
 a. Trisomy 21                              b. Mosaic 21  
 c. Translocation t(15;21), t(21;21)  
 d. Deletion of 21
- 26. The chromosomal karyotype in Patau syndrome is -** (DNB Dec 2008/ Karn 2011)  
 a. 47XX, +21                              b. 46XX/47XX,+18  
 c. 45XX, der(14;21)                      d. 47XX,+13
- 27. All the following are characteristic of Turner Syndrome EXCEPT-** (DNB Dec 10)  
 a. Webbing of Neck                      b. Cubitus valgus  
 c. Umbilical Hernia                      d. Coarctation of Aorta

#### MENDELIAN / SINGLE GENE DISORDERS

- 28. All of the following statement(s) is/are true about Marfan syndrome except:** (PGI May 2019)  
 a. It is autosomal dominant inheritance  
 b. Defect in type-1 collagen  
 c. Defect in type-2 collagen  
 d. Occur due to mutation in fibrillin 1 gene  
 e. There is defect in lysyl hydroxylase enzyme
- 29. CGG repeat sequence is found in:** (JIPMER 2019)  
 a. Fragile X syndrome  
 b. Huntington's chorea  
 c. Dentatorubropallidoluysian atrophy  
 d. Machado-Joseph disease



30. Identify the inheritance pattern? (AIIMS Nov 2018)



- a. Y linked recessive      b. Y linked dominant  
c. X linked recessive      d. X linked dominant

31. Which of the following dyads are correct? (PGI May 2018)

- a. Marfan syndrome: Autosomal recessive  
b. Phenylketonuria: Autosomal dominant  
c. Vit. D resistant rickets: Autosomal dominant  
d. Alkaptonuria: Autosomal recessive  
e. Duchene muscle dystrophy: X-linked recessive

32. Which of the following is Autosomal dominant disorder? (JIPMER 18)

- a. Bests disease  
b. Laurence-Moon Biedl syndrome  
c. Bassen-Kornzweig syndrome  
d. Usher syndrome

33. Which of the following disease is X- linked? (Recent Question 2018)

- a. Thalassemia      b. Color blindness  
c. Sickle cell anemia      d. Galactosemia

34. All of the following are autosomal dominant, except? (Recent Question 2018)

- a. Familial hypercholesterolemia  
b. Congenital adrenal hyperplasia  
c. Achondroplasia  
d. Acute intermittent porphyria

35. Which of the following is true regarding ataxia telangiectasia? (Recent Question 2018)

- a. It is X linked recessive disease  
b. It is autosomal recessive disorder  
c. It is associated with increased levels of IgM  
d. None of the above

36. Which of the following is the gene for Duchenne muscular dystrophy? (Recent Exam 2018)

- a. DMPK gene      b. STK11 gene  
c. PTCH gene      d. Dystrophin gene

37. Xeroderma pigmentosa all true except:

- a. Autosomal dominant  
b. Onset of squamous cell ca by 2nd-3rd decade  
c. Develops non melanotic cancer by the age of 9yrs  
d. Nucleotide excision repair defect

38. An obese women with T2 diabetes and hypertension had endometrial ca. Which of the following gene is involved? (JIPMER 2017)

- a. PTEN      b. P53  
c. B catenin      d. SMAD

39. Locations of gene which codes for alpha 1 anti-trypsin deficiency? (JIPMER 2017)

- a. 14q      b. 13p  
c. 17p      d. 14p

40. Find Correct match among the following? (PGI May 2017)

- a. Hypophosphatemic rickets- X linked dominant  
b. Duchenne muscular dystrophy- X linked recessive  
c. Sickle cell: AR  
d. Osteogenesis imperfecta-1 :AR  
e. Achondroplasia : AR

41. Autosomal recessive disease(s) is/are: (PGI May 2016)

- a. Sickle cell anaemia      b. Phenylketonuria  
c. Tuberous sclerosis      d. Familial polyposis coli  
e. Marfan syndrome

42. Chromosomal abnormalities in Down syndrome is/are due to: (PGI May 2016)

- a. Nondisjunctional of maternal chromosome  
b. Nondisjunctional of paternal chromosome  
c. Translocations between chromosome 21 & 14  
d. Disjunction of paternal chromosome  
e. Mosaicism

43. True about Lyonization of X chromosome: (Recent Question 2016-17)

- a. Inactivation of X chromosome only in somatic cell  
b. Inactivation of X chromosome only in germ cell  
c. Inactivation of X chromosome in somatic & germ cell both  
d. Maximum number of Barr body is equal to X chromosome

44. Which of the following pairs is not correctly matched? (Recent Question 2016-17)

- a. Turner's syndrome ..... 45, XO  
b. Down syndrome ..... 47, XY, +21  
c. Klinefelter's syndrome ..... 47, XXY  
d. Marfan's syndrome ..... 47, XYY

45. The clinical features of Turner Syndrome in girls include the following except: (Recent Question 2016-17)

- a. Severe mental retardation  
b. Webbing of the neck  
c. Delayed puberty  
d. Short stature

46. The law "Relative frequencies of each gene allele tends to remain constant from generation to generation", was given by? (Recent Question 2015)

- a. Henry Sigerist      b. Hardy Weinberg  
c. Doug Engelberg      d. Johanna Frank

47. ABO blood group is an example of? (AIIMS May 2015)

- a. Co-dominance  
b. AD  
c. AR  
d. Mitochondrial inheritance

48. A 48-year-old lady presented with hepatosplenomegaly with pancytopenia. On bone marrow examination, a tissue paper crumpled appearance is seen. Which is the most likely product to have accumulated? (AIIMS May 2015)

- a. Glucocerebroside      b. Sphingomyelin  
c. Sulfatide      d. Ganglioside



- 49. Frameshift mutation occurs due to?**  
(Recent Question 2016)  
a. Transition                      b. Transversion  
c. Insertion                      d. Point mutation
- 50. Hypophosphatemic Vit D Resistant Rickets is?**  
(Recent Question 2016)  
a. AD                                  b. AR  
c. XD                                  d. XR
- 51. Which of the following is X-linked recessive?**  
(Recent Question 2016)  
a. Duchenne muscular dystrophy  
b. Hypophosphatemic rickets  
c. Marfan's syndrome  
d. Down's syndrome
- 52. All the following are Autosomal Dominant EXCEPT**  
(Recent Question 2015)  
a. Cronkhite Canada syndrome  
b. Bannayan Ruvulcaba Riley syndrome  
c. Peutz Jegher's syndrome  
d. Gardner's syndrome
- 53. Waardenburg syndrome is due to mutation of**  
(Recent Question 2015)  
a. PAX2 gene                      b. PAX3 gene  
c. PAX6 gene                      d. PAX9 gene
- 54. Which among the following subtypes of Osteogenesis imperfecta is not associated with blue sclera?**  
(Recent Question 2015)  
a. Type I                              b. Type II  
c. Type III                            d. Type IV
- 55. Globoid cells is a diagnostic feature of**  
(Recent Question 2015)  
a. Krabbe's disease  
b. Progressive multifocal leukoencephalopathy  
c. Tay Sachs's disease  
d. Metachromatic leukodystrophy
- 56. Von Recklinghausen's disease of the bone is caused by**  
(Recent Question 2015)  
a. Paget's disease  
b. Severe hyperparathyroidism  
c. Renal osteodystrophy  
d. Severe osteomalacia
- 57. Which of the following is not due to defect in type II collagen?**  
(Recent Question 2015)  
a. Achondrogenesis II  
b. Hypochondrogenesis  
c. Stickler's syndrome  
d. Multiple epiphyseal dysplasia
- 58. Which of the following is inherited autosomal recessive?**  
(Recent Question 2015)  
a. Achondroplasia  
b. Tuberous sclerosis  
c. Hemochromatosis  
d. Osteogenesis imperfecta
- 59. RET proto-oncogene is located on which chromosome?**  
(Recent Question 2015)  
a. 9                                      b. 10  
c. 11                                    d. 12
- 60. Which of the following is autosomal recessive inherited cancer syndrome?**  
(Recent Question 2015)  
a. Ataxia telangiectasia              b. Cowden syndrome  
c. Retinoblastoma                      d. HNPCC
- 61. Inheritance of Beckers muscular dystrophy is?**  
(Recent Question 2016)  
a. X-linked recessive                  b. X-linked dominant  
c. Autosomal recessive                d. Autosomal dominant
- 62. Which of the following is an autosomal dominant metabolic disorder?**  
(Recent Question 2016)  
a. Hereditary hypercholesterolemia  
b. Tay Sachs disease  
c. Gaucher's disease                  d. Tyrosinemia
- 63. Gene responsible for embryogenesis of eye**  
(Recent Question 2016)  
a. PAX 6                                b. PAX 2  
c. PAX 5                                d. RAX
- 64. PRSS1 gene is on ?**  
(Recent Question 2016)  
a. Chr20q                              b. Chr 17p  
c. Chr 7q                                d. Chr1p
- 65. BRCA1 is the most common gene mutated in familial breast cancer. This gene is located on which chromosome?**  
(Recent Question 2015)  
a. 13                                      b. 17  
c. 20                                      d. 21
- 66. A 20-year-old male presents with mental retardation, large mandible, large everted ears and large testes. What is the most likely diagnosis?**  
(Recent Question 2015)  
a. Down syndrome                      b. Patau syndrome  
c. Fragile X syndrome                  d. Klinefelter syndrome
- 67. Y chromosome is ?**  
(Recent Question 2015)  
a. Metacentric                          b. Submetacentric  
c. Acrocentric                          d. Telocentric
- 68. Chromosome involved in friedreich ataxia**  
(Recent Question 2015)  
a. 9q                                      b. 19q  
c. 17p                                    d. X chromosome
- 69. For karyotyping, the dividing cells are arrested by the addition of colchicines in the following mitotic phase**  
(Recent Question 2015)  
a. Prophase                              b. Metaphase  
c. Anaphase                              d. Telophase
- 70. The best suited nucleated cell for chromosomal study**  
(Recent Question 2015)  
a. Polymorphs                          b. Lymphocytes  
c. Epithelial cells                      d. Langerhan's cell
- 71. F body is**  
(Recent Question 2015)  
a. Y chromatin                          b. X chromatin  
c. Chromosome 1                      d. Chromosome 21
- 72. An example for gain of function mutation is?**  
(Recent Question 2015)  
a. Osteogenesis imperfecta  
b. Ehler danlos syndrome  
c. Marfan syndrome  
d. Huntington's disease
- 73. All are true about Angelman syndrome except**  
(Recent Question 2015)  
a. Microdeletion on maternal chromosome 15  
b. Obesity  
c. Mental retardation  
d. Happy puppets






74. **Banding technique most commonly employed for cytogenetic analysis** (Recent Question 2015)
  - a. G banding
  - b. Q banding
  - c. R banding
  - d. C banding
75. **Germline mosaicism is seen in** (Recent Question 2015)
  - a. Fragile X syndrome
  - b. Angelman syndrome
  - c. Mitochondrial myopathy
  - d. Osteogenesis imperfecta
76. **Approximate number of genes in human genome** (Recent Question 2015)
  - a. 10000–15000
  - b. 20000–25000
  - c. 40000–50000
  - d. 100000
77. **Most lethal karyotype is?** (Recent Question 2016)
  - a. 45, XO
  - b. 45, YO
  - c. 47, XXY
  - d. 48, XYYY
78. **All the following are true regarding autosomal dominant inheritance except** (Recent Question 2015)
  - a. Usually defects of structural proteins or receptors
  - b. Many new mutations seem to occur in germ cells of relatively older fathers
  - c. Complete penetrance
  - d. Variable expressivity
79. **Arterial or uterine rupture occurs in which type of Ehlers Danlos syndrome** (Recent Question 2015)
  - a. Type I
  - b. Type II
  - c. Type III
  - d. Type IV
80. **The following feature is not common in 22q11.2 deletion syndrome** (Recent Question 2015)
  - a. Mental retardation
  - b. Schizophrenia
  - c. ADHD
  - d. Congenital heart defects
81. **Protein affected in spinocerebellar ataxia type 6** (Recent Question 2015)
  - a. Ataxin
  - b.  $\alpha$ 1A-Voltage-dependent calcium channel subunit
  - c. Atrophin
  - d. Androgen receptor
82. **Not a true statement about fragile X ataxia tremor syndrome** (Recent Question 2015)
  - a. Females carrying the premutations may have premature ovarian failure
  - b. Premutation carrying males cannot transmit the disease
  - c. Males exhibit a neurodegenerative syndrome characterized by intention tremors and cerebellar ataxia
  - d. May progress to parkinsonism
83. **The following technique is used to analyze DNA obtained from cancer biopsies, in which tumor cells are often contaminated with large numbers of admixed stromal cells** (Recent Question 2015)
  - a. Sanger sequencing
  - b. Single base primer extension
  - c. Pyrosequencing
  - d. Amplicon length analysis
84. **Submicroscopic deletions of any size can be detected by?** (Recent Question 2015)
  - a. Multiplex ligation-Dependent probe amplification (MLPA)
  - b. Southern blotting
  - c. Cytogenomic array technology
  - d. Chromosome painting
85. **True about autosomal dominant type of inheritance:** (PGI May 2015)
  - a. 25% affected & 50% carrier if one parent affected
  - b. 50% affected & 75% carrier if both parent affected
  - c. 75% affected if both parent affected
  - d. 50% affected if one parent affected
  - e. All carrier irrespective of either one parent affected or both parent affected
86. **Accumulation of cerebral gangliosides occurs due to deficiency of:** (Recent Question 2015)
  - a.  $\beta$  glucocerebrosidase
  - b.  $\beta$  galactosidase
  - c. Hexosaminidase A
  - d. Sphingomyelinase
87. **If parents are carrier for an autosomal recessive disorder. What are the chances of offspring to get affected-** (Recent Question 2014)
  - a. 1 : 1
  - b. 1 : 2
  - c. 1 : 3
  - d. 1 : 4
88. **Type of inheritance in Tuberous sclerosis -** (Recent Question 2014) (WB PG 2012)
  - a. Autosomal dominant
  - b. Autosomal recessive
  - c. X-linked dominant
  - d. X-linked recessive
89. **Enzyme deficiency in Hunter disease is:** (WB PG 2014)
  - a. L-iduronidase
  - b. Iduronate sulfatase
  - c. Heparan sulfamidase
  - d. Hyaluronidase
90. **Which of the following disease is caused by point mutation?** (PGI May 2013, May 2012)
  - a. Colon cancer
  - b. Diabetes mellitus type II
  - c. Cystic fibrosis
  - d. Sickle cell disease
  - e. Gauchers disease
91. **In Xeroderma Pigmentosum, defect is in?** (Recent Question 2013)
  - a. Base pair defect
  - b. Nucleotide excision Repair
  - c. Mismatch repair defect
  - d. Protein folding
92. **Hemophilia is associated with-** (Recent Question 2014)
  - a. X chromosome
  - b. Y chromosome
  - c. Chromosome 3
  - d. Chromosome 16
93. **If both parents have sickle cell anemia, then the likelihood of children having the disease is-** (Recent Question 2015)
  - a. 10%
  - b. 25%
  - c. 50%
  - d. 100%
94. **In an Autosomal Dominant disorder, mother is affected but is heterozygous. Father is normal. What are the chance of disease in children?** (Recent Question 2014)
  - a. 50% affected
  - b. 25% affected
  - c. 75% affected
  - d. All affected
95. **True statement about inheritance of an X linked recessive trait is -** (Recent Question 2015)
  - a. 50% of boys of carrier mother are affected
  - b. 50% of girls of diseased fathers are carriers
  - c. Father transmits disease to the son
  - d. Mother transmits the disease to the daughter
96. **In Marfan syndrome, the defect is in -** (Recent Question 2014, 2012), (DNB June 11)
  - a. Fibrillin I
  - b. Fibrillin II
  - c. Collagen
  - d. Elastin



97. Which of the following is not X linked condition?  
 a. Duchenne muscular dystrophy (Recent Question 13)  
 b. Emery-Dreifuss muscular dystrophy  
 c. Fascio-scapulo-humeral muscular dystrophy  
 d. Becker muscular dystrophy
98. In a X-linked recessive condition, what is the chance of an offspring being affected with an affected mother and normal father?  
 (Recent Question 2012)  
 a. 50% of daughters are carriers  
 b. 50% of sons are asymptomatic carriers  
 c. 50% of the off-springs are carriers  
 d. Males will never be affected
99. A normal couple has one daughter affected with Cystic fibrosis. They are now planning to have another child. What is the chance of siblings being affected by the disease -  
 (AI PGMEET 2010)  
 a. 0  
 b. 50%  
 c. 25%  
 d. 75%
100. Disease having autosomal recessive inheritance -  
 (PGI May 10)  
 a. Cystic fibrosis  
 b. Hydrocephalus  
 c. Duchene muscular dystrophy  
 d. Albinism  
 e. Vitamin D resistant rickets
101. Which is the most common tumor associated with type I neurofibromatosis?  
 a. Optic nerve glioma  
 b. Meningioma  
 c. Acoustic schwannoma  
 d. Low grade astrocytoma
102. Identify the inheritance pattern?  
 (JIPMER 2017)




- a. AD  
 b. AR  
 c. XD  
 d. XR
103. A 1-year-old boy presented with hepatosplenomegaly and delayed milestones. The liver biopsy and bone marrow biopsy revealed presence of histiocytes with PAS-positive Diastase resistant material in the cytoplasm. Electron microscopic examination of these histiocytes is most likely to reveal the presence of -  
 a. Birbeck granules in the cytoplasm  
 b. Myelin figures in the cytoplasm  
 c. Parallel rays of tubular structures in lysosomes  
 d. Electron dense deposit in the mitochondria

104. Which of the following is inherited as autosomal recessive form?  
 (DNB June 2010)  
 a. Sickle cell anemia  
 b. Hemophilia  
 c. Hereditary spherocytosis  
 d. Glucose 6-PO4 dehydrogenase deficiency

#### DISORDERS WITH NONCLASSIC INHERITANCE


105. Mutation in DNA Helicase causes:  
 (Recent Pattern Question 2020)  
 a. Werner syndrome  
 b. Werner syndrome  
 c. Sipple syndrome  
 d. Autoimmune lymphoproliferative syndrome
106. Which of the following mode of inheritance is shown below?  
 (Recent Pattern Question 2020)



- a. AD  
 b. AR  
 c. X linked dominant  
 d. X linked recessive
107. An 8-year-old child presented with history of recurrent infections. The child had rashes. Investigations revealed low platelets. What could be the probable cause?  
 (Recent Pattern Question 2020)  
 a. Job syndrome  
 b. Wiskott-Aldrich syndrome  
 c. Henoch-Schönlein purpura  
 d. Hyper IgM syndrome
108. True statement(s) about Mitochondrial Inheritance:  
 (PGI May 2019)  
 a. Only girls are affected  
 b. Mutation cause MELAS  
 c. Disease transfers from mother only  
 d. Commonly affects brain, heart, muscle, liver  
 e. Doesn't follow typical Mendelian inheritance pattern
109. An affected male does not have affected children but an affected female always has affected children. Type of inheritance?  
 (AIIMS May 2019)  
 a. X linked recessive  
 b. Autosomal recessive  
 c. X linked dominant  
 d. Mitochondrial




**110. Examine this pedigree chart carefully. What type of transmission does it depict?**



- a. AD inheritance      b. AR inheritance  
c. X-linked recessive      d. Holandric inheritance

**111. Examine this pedigree chart carefully. What type of transmission does this depict?**



- a. Autosomal dominant      b. Autosomal recessive  
c. X-linked recessive      d. X-linked dominant

**112. % of Gene's in mitochondria ?** (PGI Nov 2018)

- a. 1.5      b. 3.5  
c. 6      d. 10

**113. Child with growth retardation came with Tripod skull and clinodactyly?** (JIPMER 2017)

- a. Silver russel syndrome  
b. Beckwith wideman syndrome  
c. Angelman syndrome  
d. Prader willi syndrome

**114. Which of the following can change the gene expression by methylation and acetylation without changing content of the gene?** (AIIMS May 2017)

- a. Translocation      b. Inversion  
c. Mutation      d. Epigenetics

**115. True about mammalian mitochondrial DNA:** (PGI May 2016)

- a. Contains around 16500 nucleotide sequence  
b. Makes up around 3% total cellular DNA  
c. Makes up around 10% total cellular DNA  
d. Makes up around 0.3% total cellular DNA  
e. Makes up around 1% total cellular DNA

**116. % of individual who inherited the gene and will express the trait is known as?** (Recent Question 2016)

- a. Penetrance      b. Inheritance  
c. Co-dominance      d. Pleiotropism

**117. Trinucleotide repeat disorder is?**

- a. Mitochondrial myopathy (Recent Question 2016)  
b. Myotonia dystrophica  
c. Inflammatory myopathy  
d. Duchene's dystrophy

**118. Anticipation is a feature of** (Recent Question 2016)

- a. Tri nucleotide repeats  
b. Genomic imprinting  
c. Trisomy  
d. Mosaicism

**119. True about mitochondrial DNA:** (PGI May 2015)

- a. Linear      b. Circular  
c. Transmitted by mother only  
d. Transmitted by both parents  
e. Contains less gene than nuclear DNA

**120. The phenomenon where subsequent generations are at risk of earlier and more severe disease is known as?**

(AIIMS May 2015)

- a. Anticipation      b. Dominance  
c. Pleiotropy      d. Dominance

**121. Type of inheritance in MELAS-** (Recent Question 2014)

- a. AD      b. AR  
c. Mitochondrial      d. X-linked

**122. Paternal 15 chromosome deletion is seen in-**

(Recent Question 2014)

- a. Angelman syndrome  
b. PraderWilli syndrome  
c. Down syndrome  
d. Turner syndrome

**123. Uniparental disomy is seen in all except?**

(AIIMS May 2014, Nov 2013)

- a. Angelman syndrome  
b. Praderwilli syndrome  
c. Bloom's syndrome  
d. Silver Russell syndrome

**124. In Huntington Chorea, the causative mutation in the protein huntingtin is a** (AP PGME 2014)

- a. Point mutation  
b. Gene deletion  
c. Frameshift mutation  
d. Trinucleotide repeat expansion

**125. Fragile X syndrome is:** (WB PG 2014)

- a. AD      b. AR  
c. X-linked      d. Multifactorial

**126. All are mitochondrial disorders EXCEPT:** (MH 2016)

- a. Pearson Syndrome  
b. MERRF (Myoclonic Epilepsy Ragged Red Fibre)  
c. MELAS (Mitochondrial Encephalopathy Lactic Acidosis and Stroke-like episode)  
d. Fragile X Syndrome

**127. True about Mitochondrial inheritance is/are:**

(PGI May 2013)

- a. Mothers transmit their mtDNA to both their sons and daughters  
b. Both parents can transmit their mtDNA to both their children  
c. Mitochondrial DNA codes for 37 genes  
d. Mitochondrial disease commonly affect neuromuscular system  
e. Mutation cause Leber hereditary optic neuropathy



**128. True about mitochondrial DNA:** (PGI May 2011)

- a. Maternal inheritance
- b. Same evolutionary origin
- c. Not highly conserved and has high mutation rate
- d. Mitochondrial disease is associated mostly with mutation and some have deletion in mt-DNA
- e. Mostly encodes membrane protein

**129. Mitochondrial DNA (mt- DNA) is known for all except-** (DPG 11)

- a. Maternal inheritance
- b. Heteroplasmy
- c. Leber hereditary optic neuropathy is the prototype
- d. Nemesin myopathy results due to mutations in mt-DNA


**130. True about genomic imprinting -** (PGI Nov 10)

- a. Different expression of gene depending on parent of origin
- b. Prader-Willi syndrome is due to maternal deletion of chromosome 15
- c. Angelman syndrome is due to paternal deletion of chromosome 15
- d. Uniparental disomy is other name of genomic imprinting

**131. Two siblings with osteogenesis imperfecta, but their parents are normal. Mechanism of inheritance is -** (AIIMS May 10)

- a. Anticipation
- b. Genomic imprinting
- c. Germ line mosaicism
- d. New mutation

**132. Analyze the following pedigree and give the mode of inheritance:** (PGI May 10)



- a. Autosomal recessive
- b. Autosomal dominant
- c. Mitochondrial inheritance
- d. X linked dominant

**133. Which of the following disorders is due to maternal disomy of chromosome 15?** (AIIMS Nov 10)

- a. Prader-Willi syndrome
- b. Angelman syndrome
- c. Hydatidiform mole
- d. Klinefelter's syndrome

**134. True about mitochondrial diseases -** (PGI Nov 10)

- a. Paternal inheritance
- b. LHON & CPEO are typical examples
- c. Most common abnormality shown is neurological
- d. Li-Fraumeni syndrome is a mitochondrial disease
- e. Common mode of inheritance

**135. Mitochondrial abnormalities are seen in -** (PGI May 10)

- a. Oncocytoma
- b. Kearns-Sayre syndrome
- c. Faber disease
- d. Mitochondrial myopathy
- e. Leigh's disease

## DIAGNOSTIC TOOLS IN GENETICS

**136. Karyotyping is/are not necessary in which of the following conditions?** (PGI May 18)

- a. Recurrent abortions
- b. Child with multiple malformation
- c. Pregnancy in advance maternal age with previous pregnancy with an abnormal chromosomal complement
- d. Child with seizure
- e. Down syndrome

**137. Gene silencing means a gene which would be expressed under normal circumstances is switched off by cell machinery. Which one of the following cellular components is not involved in gene silencing?** (UPSC 2016)

- a. Micro-RNA
- b. Double stranded RNA
- c. Ribosomal RNA
- d. Si RNA

**138. With reference to "DNA Microarray" technology, which of the following statements is/are correct?** (UPSC 2016)

- 1. It helps in identifying the genes involved in a disease by comparing the gene expressions between the tissues from a healthy and an infected person.
- 2. It allows visualizing the activity of hundreds of genes simultaneously.

Select the correct answer using the code given below:

- a. 1 only
- b. 2 only
- c. Both 1 and 2
- d. Neither 1 nor 2

**139. True about array CGH is/are all except:** (PGI Nov 2014)

- a. Used for known Genetic loci
- b. Both duplication and deletion can be studied
- c. Uses 2 colour label
- d. Can detect multiple abnormalities at the same time
- e. Uses array hybridization technology

**140. Known gene loci is can be diagnosed by:** (AIIMS Nov 2013)

- a. FISH
- b. Comparative gene hybridization
- c. PCR
- d. Chromosomal painting

**141. Resolution of light microscope of viewing chromosome**

- a. 5 Kb
- b. 500 kb
- c. 5 mb
- d. 50 mb

**142. Real time polymerase chain reaction is done for:** (AIIMS May 2013)

- a. DNA detection only
- b. RNA detection only
- c. Both RNA and DNA detection
- d. Monitoring amplification of target nucleic acid

**143. Karyotyping is done in which phase of cell cycle:** (Recent Question 2014)

- a. Anaphase
- b. Metaphase
- c. Telophase
- d. S phase

**144. Microarray is -** (Recent Question 2014)

- a. Study of multiple genes
- b. Study of diseases
- c. Study of organisms
- d. Study of blood group





**145. Which of the following procedures is routine technique for karyotyping using light microscopy -**

(DNB Dec 2010, AI PGMEET 2003)

- a. C-banding
- b. G-banding
- c. Q-banding
- d. Brd V-staining

**146. Karyotyping is done with all, except:**

(AIIMS May 2011, June 98)

- a. Blood lymphocyte
- b. Blood monocyte
- c. Amnion
- d. Fibroblast

### MISCELLANEOUS

**147. BRCA 1 and BRCA 2 is responsible for all these Cancers except:**

(Recent Question 2015)

- a. Breast
- b. Ovary
- c. Intestinal lining cancer
- d. Prostate

**148. Multifactorial inheritance is most likely to play a role in-**

(Recent Question 2014)

- a. Cleft lip
- b. Marfan's syndrome
- c. Down's syndrome
- d. Erythroblastosis fetalis

**149. HOX gene mutation can cause all except-**

(Recent Question 2014)

- a. Syndactyly
- b. Polydactyly
- c. Fused carpal bones
- d. VSD

**150. Technique(s) used detecting for Gene Mutation is/are:**

(PGI May 2012)

- a. Real-time PCR
- b. Denaturing gradient gel electrophoresis
- c. DNA sequencing
- d. Restriction fragment polymorphism (RFLP)
- e. Single-strand conformational polymorphism

**151. Which of the following is a DNA repair defect?**

(DNB Dec 2010)

- a. Bloom syndrome
- b. Incontinentia pigmenti
- c. Aplastic anemia
- d. Tuberous sclerosis

**152. Karyopyknotic index is a method for?**

(DNB Aug 2012)

- a. Ovarian carcinoma
- b. Hormonal evaluation
- c. Dysplasia measurement
- d. Measurement of cells in active replication

**153. Regarding 'Davidson body', all are correct EXCEPT:**

(MH 2016)

- a. Present in males
- b. Present in neutrophils
- c. Present in 4 - 6 % of cells
- d. Drumstick appearance



## Answers with Explanations

**1. Ans. (b, c, d, e) b. Pleuropulmonary blastoma; c. Cystic nephroma; d. Thyroid carcinoma; e. Sertoli-Leydig cell tumor**

**2. Ans. (b, c) b. Turner syndrome; c. Klinefelter's syndrome** (Ref: Robbins 9th/pg 165)

**3. Ans. (a, b, c) a. Most common cause of death is congenital heart disease; b. 95% have an extra 21st chromosome; c. Increased nuchal fold thickness on USG**

**4. Ans. (d) Y Chromosome**

**5. Ans. (e) Mental retardation present**

**6. Ans. (a) Tall stature** (Ref: Robbins 9/e pg 166)

Left-sided cardiovascular abnormalities, particularly preductal coarctation of the aorta and bicuspid aortic valve, are seen. Hypertension may be a feature of coarctation of aorta.

**7. Ans. (c) Short acrocentric** (Ref: Emery Genetics)

**8. Ans. (a) Turner syndrome** (Ref: R 9th/pg 166; 8th/pg 165)

In the given karyotype, we can observe that there is only 1 X chromosomes and no Y chromosome. This suggests the karyotype of patient as 45 XO (Turners Syndrome)

**9. Ans. (c) Chr 22** (Ref: Robbins 9th/pg 163; 8th/pg 162)

**10. Ans. (a, d, e) Retrognathia d. Isolated cleft palate e. Glossoptosis**

(Ref: Oxford Handbook of Genetics pg 171)

**Pierre Robin sequence** is due to defective DNA near the SOX9 gene is a set of abnormalities affecting the head and face, The three main features are **cleft palate, retrognathia and glossoptosis**.

**11. Ans. (a) Gynecomastia with long thin limbs**

(Ref: Robbins 9th/pg 166; 8th/pg 165)

In the given karyotype, we can observe that there are two X chromosomes and single Y chromosome. This suggests the karyotype of patient as 47 XXY (Klinefelter Syndrome) Male hypogonadism that occurs when there are  $\geq 2$  X chrand  $\geq 1$  Y Chr.

**12. Ans. (d) Maternal nondisjunction** (Ref: R 9th/pg 161)

**Down's syndrome** is most commonly caused by **Maternal nondisjunction** > **Translocation** > **Mosaicism**

**13. Ans. (c) Turner syndrome** (Ref: Robbins 9th/pg 166-167)

**14. Ans. (a) Turner syndrome** (Ref: Robbins 9th/pg 166-167)

**15. Ans. (b) Klinefelter syndrome results due to an extra Y chromosome in males**

(Ref: Robbins 9th/pg 166; 8th/pg 165)

- False: Only one of the X chromosomes is genetically active, the other X of either maternal or paternal origin undergoes heteropyknosis and is rendered inactive called Barr body, so females have one activated chromosome
- True:** Klinefelter syndrome is best defined as male hypogonadism that occurs when there are **two or more X chromosomes and one or more Y chromosomes**
- False: Germ cells have haploid set of 23 chromosomes
- False: Turner's is due to missing X chr.

**16. Ans. (d) 5p-** (Ref: Robbins 9th/pg 163; 8th/pg 162)**17. Ans. (c) Androgen insensitivity syndrome****18. Ans. (d) Cri du chat syndrome** (Ref: Robbins 9th/pg 163)

This is karyotype showing an arrow on **deleted part of short arm of chromosome 5**, which is suggestive of **Cri du Chat syndrome: Chromosome 5p deletion**<sup>Q</sup>  
Features: Characteristic cry, developmental delay, behavioral problems.

**19. Ans. (b) Isochromosome** (Ref: R 9th/pg 160; 8th/pg 160)**20. Ans. (a) Absence of oocytes in the ovaries (streak ovaries)**

(Ref: Robbins 9th/pg 166-167; 8th/pg 165-166)

In this question, the patient is presenting with **primary amenorrhea** and **raised FSH**, along with **short stature** (Given Height = 58 inches, which is less than 5<sup>th</sup> percentile of expected at 16 years age). All these features are suggestive of Turner Syndrome.

**Infertility**<sup>Q</sup> due to **rudimentary uterus** and **streak ovaries** is an important feature<sup>Q</sup>, as ovaries are reduced to **atrophic fibrous strands without ova and follicles** in Turner syndrome

**21. Ans. (b) Isochromosome** (Ref: R 9th/pg 160; 8th/pg 160)**22. Ans. (b) Turner's syndrome**

(Ref: Robbins 9th/pg 166-167)

- Down Syndrome has increased risk of Leukemia
- Turner syndrome has very low risk of Wilms tumor and colorectal Ca**

So, it seems that the answer should be none, but to choose 1 of the answers, I would choose the least common one which is Turner syndrome.

**23. Ans. (b) Velo-cardio-facial syndrome**

(Ref: R 9th/pg 163)

**24. Ans. (a, b, c, e); a. Hirschprung's disease; b. Leukemia; c. Sensorineural hearing loss; e. Atlanto-occipital dislocation**

(Ref: Robbins 9th/pg 161; 8th/pg 161)

**25. Ans. (d) Deletion of 21**

(Ref: Robbins 9th/pg 161; 8th/pg 161)

**26. Ans. (d) 47XX,+13** (Ref: Robbins 9th/pg 161; 8th/pg 161)**27. Ans. (c) Umbilical Hernia** (Ref: Robbins 9th/pg 166-167)

Umbilical Hernia is usually not seen in Turner syndrome, but is commonly seen in Down syndrome

**28. Ans. (b, c, e); b. Defect in type-1 collagen; c. Defect in type-2 collagen; e. There is defect in lysyl hydroxylase enzyme** (Ref: Robbins 9th/pg 144)**29. Ans. (a) Fragile X syndrome** (Refer to answer 64)**30. Ans. (c) X linked recessive**

In the given pedigree, male are prominently affected so it goes in favour of X linked disease. Now since there is generation skip, so it will be recessive. So its, X-linked recessive.

**31. Ans. (d, e) d. Alkaptonuria: Autosomal recessive; e. Duchene muscle dystrophy: X-linked recessive****32. Ans. (a) Best's disease****33. Ans. (b) Color blindness**

(Ref: Robbins 9th ed and NCERT)

The most common cause of color blindness is an inherited fault in the development of one or more of the three sets of color sensing cones in the eye. Males are more likely to be color blind than females, as the gene responsible for the most common forms of color blindness are on the X chromosome.

**34. Ans. (b) Congenital adrenal hyperplasia**

(Ref: Robbins 9th ed 1128)

Congenital adrenal hyperplasia stems from several autosomal recessive, inherited metabolic errors, each characterized by a deficiency or total lack of a particular enzyme involved in the biosynthesis of cortical steroids, particularly cortisol.

**35. Ans. (b) It is autosomal recessive disorder**

(Ref: Robbins 9th ed p 315)

Several rare autosomal recessive cancer syndromes have been described that are characterized by hypersensitivity to certain kinds of DNA-damaging agents, such as ionizing radiation (Bloom syndrome and ataxia-telangiectasia), or DNA cross-linking agents, such as many chemotherapeutic drugs (Fanconi anemia). The phenotype of these diseases is complex and includes, in addition to predisposition to cancer, features such as neural symptoms (ataxia-telangiectasia), bone marrow aplasia (Fanconi anemia), and developmental defects (Bloom syndrome).



36. Ans. (d) **Dystrophin gene** (Ref: Robbins 9th ed p 1242)

Duchenne and Becker muscular dystrophy are caused by loss-of-function mutations in the dystrophin gene on the X chromosome. Dystrophin is one of the largest human genes, spanning 2.3 million base pairs and composed of 79 exons. The encoded protein, dystrophin, is a key component of the dystrophin glycoprotein complex (DGC)

37. (a) **Autosomal dominant**

Xeroderma pigmentosa is autosomal recessive disease.

38. Ans. (a) **PTEN**

39. Ans. (a) **14q**

SERPINA1 gene provides instructions for making a protein called alpha-1 antitrypsin, which is a type of serine protease inhibitor (serpin). Cytogenetic Location: 14q32.13, which is the long (q) arm of chromosome 14 at position 32.13

40. Ans. (a, b, c) **a. Hypophosphatemic rickets- X linked dominant b. Duchenne muscular dystrophy- X linked recessive d. Sickle cell: AR**

41. Ans. (a, b) **a. Sickle cell anaemia b. Phenylketonuria**

(Ref: Robbins 9th/141)

42. Ans. (a, b, c, e) **a. Nondisjunctional of maternal chromosome b. Nondisjunctional of paternal chromosome c. Translocations between chromosome 21 & 14 e. Mosaicism**

(Ref: Robbins (SEA) 9th/ 161-63)

43. Ans. (a) **Inactivation of X chromosome only in somatic cell**

(Ref: Robbins 9th/ 164-65; Genetics in Medicine by Thompson 8th/91)

44. Ans. (d) **Marfan's syndrome.. 47, XYY**

45. Ans. (a) **Severe mental retardation**

(Ref: R 9th/pg 166-167)

Turners syndrome has normal intelligence and not mental retardation.

46. Ans. (b) **Hardy weinberg**

(Ref: Modern Biology, pg 1-108)

47. Ans. (a) **Co-dominance** (Ref: R 9th/pg 141; 8th/pg 141)

When **both alleles** of a gene pair contribute to the phenotype.E.g.: Blood group 'AB'

48. Ans. (a) **Glucocerebroside**

(Ref: R 9th/pg 151; 8th/pg 151)

The clinical feature and bone marrow feature is typical of Gauchers disease. This results due to accumulation of

**cerebroside<sup>o</sup>** inside **mononuclear phagocytic cells** due to deficiency of **β-glucocerebrosidase**.

49. Ans. (c) **Insertion** (Ref: Robbins 9th/pg 160; 8th/pg 160)

NF1 is associated with hamartomas of the iris termed Lisch nodules.

50. Ans. (c) **XD** (Ref: Robbins 9th/pg 142; 8th/pg 142)

**Frame shift mutations occur due to nucleotide deletions or insertions cause a shift of codon reading frame.**

51. Ans. (a) **Duchenne muscular dystrophy**

(Ref: R 9th/pg 142)

a. Duchenne muscular dystrophy	X-linked recessive
b. Hypophostemic rickets	X-linked dominant
c. Marfans syndrome	Autosomal dominant
d. Downs syndrome	Chromosomal trisomy

52. Ans. (a) **Cronkhite Canada syndrome** (Ref: R 9th/pg 140)

**Cronkhite Canada syndrome** is non-inherited condition occurring in old age with features of loss of taste, intestinal polyps, hair loss and nail growth problem.

53. Ans. (b) **PAX3 gene** (Ref: Robbins 9th/pg 140; 8th/pg 140)

**Waardenburg syndrome (WS):**

- Autosomal dominant inherited condition
- Hypopigmentation of hair, skin, and/or iris of both eyes (partial albinism); and/or congenital deafness.
- 4 types: **Types I and II are the most common** forms of Waardenburg syndrome, while types III and IV are rare.
- Mutations in the **EDN3, EDNRB, MITE, PAX3, SNAI2, and SOX10** genes can cause Waardenburg syndrome.

54. Ans. (d) **Type IV** (Ref: Robbins 9th/pg 140; 8th/pg 140)

**Classification of Osteogenesis Imperfecta (OI)**

Type	Bone Fragility	Blue Sclera	Abnormal Dentition	Hearing Loss	Inheritance
I	Mild	Present	Present in some	Present in most	AD
II	Extreme	Present	Present in some	Unknown	S, rarely AR
III	Severe	Bluish at birth	Present in some	High incidence	AD, rarely AR
IV	Variable	Absent	Absent in IVA, present in IVB	High incidence	AD
V	Moderate to severe	Absent	Absent		AD
VI	Moderate to severe	Absent	Absent		Unknown
VII	Moderate	Absent	Absent	Absent	AR



**55. Ans. (a) Krabbe's disease**

(Ref: Harrison Chap 363)

Globoid cells are a unique and diagnostic feature of Krabbe disease which is actually an aggregation of engorged macrophages in the brain parenchyma and around blood vessels.

**56. Ans. (b) Severe hyperparathyroidism**

(Ref: R 9th/pg 141)

**57. Ans. (d) Multiple epiphyseal dysplasia**

(Ref: R 9th/pg 140)

**Diseases Caused by Mutations in Collagen Genes**

Gene or Enzyme	Disease
COL1A1/1A2	Osteogenesis imperfecta, type 1 Osteoporosis Ehlers-Danlos syndrome type VII
COL2A1	Severe chondrodysplasias <ul style="list-style-type: none"> <li>Achondrogenesis II</li> <li>Hypochondrogenesis</li> <li>Stickler's syndrome</li> <li>Osteoarthritis</li> </ul>
COL3A1	Ehlers-Danlos syndrome type IV
COL4A3–COL4A6	Alport syndrome
COL7A1	Epidermolysis bullosa, dystrophic
COL10A1	Schmid metaphysial chondrodysplasia
Lysyl hydroxylase	Ehlers-Danlos syndrome type VI
Procollagen N-proteinase	Ehlers-Danlos syndrome type VII
Lysyl hydroxylase	Menkes disease

**58. Ans. (c) Hemochromatosis**

(Ref: R 9th/pg 141; 8th/pg 141)

**59. Ans. (b) 10 (Ref: Robbins 9th/pg 166; 8th/pg 166)**

**60. Ans. (a) Ataxia telangiectasia**

(Ref: R 9th/pg 141; 8th/pg 141)

**61. Ans. (a) x linked recessive (Ref: R 9th/pg 142; 8th/pg 142)**

**62. Ans. (a) Hereditary hypercholesterolemia**

(Ref: Robbins 9th/pg 140; 8th/pg 140)

**Inheritance pattern of Progressive Muscular Dystrophies**

Type	Inheritance
Duchenne's	XR
Becker's	XR
Limb-girdle	AD/AR

Contd...

Emery-Dreifuss	XR/AD
Congenital	AR
Myotonia dystrophica	AD
Facioscapulohumeral	AD
Oculopharyngeal	AD

**63. Ans. (a) PAX 6 (Ref: Emery genetics)**

The PAX-6 gene locus is a transcription factor for the various genes and growth factors involved in eye formation. Eye formation in the human embryo begins at approximately 3 weeks into embryonic development and continues through the tenth week.

**64. Ans. (c) Chr 7q (Ref: Robbins 9th/pg 166; 8th/pg 166)**

- Hereditary pancreatitis is characterized by recurrent attacks of severe acute pancreatitis often beginning in childhood and ultimately leading to chronic pancreatitis.
- The disorder is genetically diverse, but **the shared feature of most forms is a defect that increases or sustains the activity of trypsin**. Three genes implicated in hereditary pancreatitis are *PRSS1*, *SPINK1*, and *CFTR*.
- Most hereditary cases are due to gain-of-function mutations in the *trypsinogen* gene (also known as *PRSS1*) on chr 7q34

**65. Ans. (b) 17 (Ref: Robbins 9th/pg 141; 8th/pg 141)**

**66. Ans. (c) Fragile X syndrome**

(Ref: R 9th/pg 170; 8th/pg 170)

**67. Ans. (c) Acrocentric (Ref: Emery Genetics) Refer to ans 1)**

**68. Ans. (a) 9q (Ref: Robbins 9th/pg 168; 8th/pg 168)**

**69. Ans. (b) Metaphase (Ref: Robbins 9th/pg 158; 8th/pg 159)**

**70. Ans. (b) Lymphocytes**

(Ref: Robbins 9th/pg 161; 8th/pg 161)

**71. Ans. (a) Y chromatin (Ref: Emery Genetics)**

Quinacrine, a fluorescence dye, binds strongly to the Y chromosome forming a bright fluorescent spot (F body). This is clearly visible in stained interphase cells from various tissues from the human male and in mature spermatozoa

**72. Ans. (d) Huntington's disease**

- In Huntington disease, the trinucleotide repeat mutation affecting the Huntington gene gives rise to an abnormal protein, called *huntingtin*, that is toxic to neurons causing neurologic deficit

**73. Ans. (b) Obesity**

(Ref: Clinical Embryology: A Color Atlas and Text; p 34)





74. Ans. (a) **G banding** (Ref: Robbins 9th/pg 160; 8th/pg 160)

75. Ans. (d) **Osteogenesis imperfecta**  
(Ref: Robbins 9th/pg 160)

In some **autosomal dominant** disorders, like **osteogenesis imperfecta**<sup>o</sup>, phenotypically normal parents have more than one affected child. This is due to germline mosaicism.

76. Ans. (b) **20000–25000**

(Ref: Robbins 9th/pg 160; 8th/pg 160)

77. Ans. (b) **45,YO** (Ref: Robbins 9th/pg 160; 8th/pg 160)

Karyotype 45, YO is never found in live born infants, presumably because the X chromosome carries genetic information essential for life.

78. Ans. (c) **Complete penetrance** (Ref: Robbins 9th/pg 140)

Complete penetrance is shown by autosomal recessive.

79. Ans. (d) **Type IV** (Ref: Robbins 9th/pg 140; 8th/pg 140)

80. Ans. (a) **Mental retardation** (Ref: Robbins 9th/pg 163)

Individuals with the 22q11.2 deletion syndrome have high risk for psychotic illnesses, such as **schizophrenia** (25% cases) and **bipolar disorders**.

In addition, **attention deficit hyperactivity disorder** is seen in 30–35% of affected children.

81. Ans. (b)  **$\alpha$ 1A-Voltage-dependent calcium channel subunit** (Ref: 19th<sup>o</sup>/pg 451)

Spinocerebellar ataxia type 6 is due to CACNA2A gene on chromosome 19p13.3, which codes for  $\alpha$ 1A-Voltage-dependent calcium channel subunit protein.

82. Ans. (b) **Premutation carrying males cannot transmit the disease** (Ref: Robbins 9th/pg 140; 8th/pg 140)

83. Ans. (c) **Pyrosequencing** (Ref: R 9th/pg 140; 8th/pg 140)

**Pyrosequencing:**

- **Highly sensitive** than Sanger sequencing and can detect even 5% mutated DNA
- For this reason, it is be used to analyze DNA obtained from cancer biopsies, in which tumor cells are often “contaminated” with large numbers of admixed stromal cells.

84. Ans. (a) **Multiplex ligation-Dependent probe amplification (MLPA)**

(Ref: Robbins 9th/pg 177; 8th/pg 179)

**Multiplex Ligation-Dependent Probe Amplification (MLPA):** MLPA is a technique which mixes DNA hybridization, DNA ligation, and PCR amplification to detect deletions and duplications of any size, including anomalies that are too large to be detected by PCR and too small to be identified by FISH.

85. Ans. (d) **50% affected if one parent affected**


(Ref: Robbins 9th/pg 140; 8th/pg 140)

86. Ans. (b)  **$\beta$  galactosidase** (Ref: Robbins 9th/pg 151–154)

**GM1-gangliosidosis:**

- Caused by a genetic deficiency of lysosomal acid  $\beta$ -galactosidase ( $\beta$ -gal); autosomal recessive inheritance;
- There is accumulation of **cerebral gangliosides**
- Manifests either as an **infantile, juvenile or adult form**.

87. Ans. (d) **1:4** (Ref: Robbins 9th/pg 141; 8th/pg 141)



Therefore, if parents are carriers for an autosomal recessive disorder, the chances of offspring to get affected is 1:4 or 25%

88. Ans. (a) **Autosomal dominant** (Ref: Robbins 9th/pg 140)

89. Ans. (b) **Iduronate sulfatase**

(Ref: Robbins 9th/pg 154–155)

All **Mucopolysaccharidoses** are **autosomal recessive** except **Hunter's disease** which is **X linked recessive**<sup>o</sup>

90. Ans. (c, d, e); c. **Cystic fibrosis**; d. **Sickle cell disease**; e. **Gauchers disease** (Ref: Robbins 9th/pg 140; 8th/pg 140)

Discussing the options one by one:

- Colon cancer –Mostly sporadic, can be Autosomal dominant
- Diabetes mellitus type II –Multifactorial** inheritance
- Cystic fibrosis –Point mutation on CFTR gene**
- Sickle cell disease- Point mutation at 6<sup>th</sup> position of  $\beta$ -chain of haemoglobin**
- Gaucher disease- Autosomal recessive** inheritance due to mutations in the GBA gene. Mutations can be point mutations or deletion in GBA gene


91. Ans. (b) **Nucleotide Excision Repair**

(Ref: Harrison 18th/Chapter 61)

Patients with **xeroderma pigmentosum** have **defects in DNA damage recognition or in the nucleotide excision and repair pathway**. Exposed skin is **dry and pigmented** and is extraordinarily sensitive to the mutagenic effects of ultraviolet irradiation.

92. Ans. (a) **X chromosome** (Ref: Robbins 9th/pg 141)

93. Ans. (d) **100%** (Ref: Robbins 9th/pg 141; 8th/pg 141)




Therefore, if both parents have sickle cell anemia, then the likelihood of children having the disease is 100%

**94. Ans. (a) 50% affected**

(Ref: Robbins 9th/pg 140; 8th/pg 140)

In an Autosomal Dominant disorder, mother is affected but is heterozygous and father is normal.




Therefore, the chances of disease in children is 50%

**95. Ans. (a) 50% of boys of carrier mother are affected**

(Ref: Robbins 9th/pg 141; 8th/pg 141)


In a X linked recessive disorder, if father is normal & mother is a carrier:



In such a situation,

- 50% of the sons are affected
- 50% of daughters are carriers, but none of the daughters are affected
- Mother transmits the disease to the sons

In a X linked recessive disorder, if father is affected & mother is normal:



In such scenario,

- 100% of daughters are carriers
- Father transmits the disease to the daughters and not to sons, as all sons of an affected father are normal

**96. Ans. (a) Fibrillin I**

(Ref: R 9th/pg 144-145; 8th/pg 144-145)

**Marfan syndrome**

- Due to an inherited defect in an extracellular glycoprotein called **fibrillin-1**<sup>o</sup>
- Affects **connective tissues**, manifested principally by changes **skeleton, eyes & cardiovascular system**.

- **Inheritance: Autosomal dominant** (75%)<sup>o</sup>, new mutations (25%)


**97. Ans. (c) Fascio-scapulo-humeral muscular dystrophy**

(Ref: Robbins 9th/pg 141; 8th/pg 141)

**98. Ans. (c) 50% of the off-springs are carriers**

(Ref: Robbins 9th/pg 141; 8th/pg 141)


In a X-linked recessive condition, if the father is normal & mother is affected:



50% of the off-springs are carriers (all daughters) & 50% of them are affected (all sons)

**99. Ans. (c) 25%** (Ref: Robbins 9th/pg 141; 8th/pg 141)

- **Cystic fibrosis** is an **autosomal recessive** condition.
- If one daughter is affected, genotype will be cc
- Therefore, both parents must have been carriers of cystic fibrosis.



Thus each child (other sibling) has a 1 in 4 (25%) chance of being affected.

**100. Ans. (a, d); a. Cystic fibrosis; d. Albinism**

(Ref: Robbins 9th/pg 141; 8th/pg 141)

Disease	Inheritance
A. Cystic fibrosis <sup>o</sup>	Autosomal Recessive
B. Hydrocephalus	Sporadic
C. Duchene muscular dystrophy <sup>o</sup>	X-linked recessive <sup>o</sup>
D. Albinism <sup>o</sup>	Autosomal Recessive <sup>o</sup>
E. Vitamin D resistant rickets <sup>o</sup>	Autosomal Dominant

**101. Ans. (a) Optic nerve glioma** (Ref: Robbins 9th/pg 298)

- NF1 patients have **increased risk** of **Meningioma**<sup>o</sup>, **Pheochromocytoma**<sup>o</sup> & **Wilm's Tumor**<sup>o</sup>
- Most common **tumor** in NF1 is **optic nerve glioma**<sup>o</sup>
- **Most common Leukemia** in NF1 is **JMML**<sup>o</sup> (Juvenile Myelomonocytic Leukemia)

**102. Ans. (c) XD**

The pattern of inheritance is X linked as males are not transmitting to sons and its dominant because there is no generation skipping.

**103. Ans. (c) Parallel rays of tubular structures in lysosomes**

(Ref: Robbins 9th/pg 153-154; 8th/pg 153-154)



This clinical scenario of a **1-year-old** boy with **hepatosplenomegaly** and **delayed milestones** (neurological involvement) and liver biopsy and bone marrow biopsy showing histiocytes with **PAS-positive Diastase-resistant** material in the cytoplasm is s/o **Gaucher disease**. In Gaucher disease, electron microscopic examination of histiocytes show parallel rays of tubular structures in lysosomes.

104. Ans. (a) **Sickle cell anemia**

(Ref: R 9th/pg 141; 8th/pg 141)

105. Ans. (b) **Werner syndrome**

106. Ans. (a) **AD**

107. Ans. (b) **Wiskott-Aldrich syndrome**

(Ref: Robbins 9th ed/pg 242)

108. Ans. (b, c, d, e); **b. Mutation cause MELAS; c. Disease transfers from mother only; d. Commonly affects brain, heart, muscle, liver; e. Doesn't follow typical Mendelian inheritance pattern** (Ref: Robbins 9th ed/pg 171-172)

109. Ans. (d) **Mitochondrial** (Ref: Robbins 9th ed/pg 171-172)

110. Ans. (d) **Holandric inheritance**

111. Ans. (d) **X-linked dominant**

112. Ans. (a) **1.5a**

113. Ans. (a) **Silver russel syndrome**

Russell Silver Syndrome which included short stature, relative macrocephaly, triangular facies and bilateral clinodactyly.

114. Ans. (d) **Epigenetics** (Ref: Robbins 9/e pg 5)

115. Ans. (a, e) **a. Contains around 16500 nucleotide sequence e. Makes up around 1% total cellular DNA**

(Ref: Robbins 9th/ 171-72; Genetic Medicine by Thomson 8th/246; Molecular Biology of Cell by Alberts 6th/802)

116. Ans. (a) **Penetrance**

(Ref: Robbins 9th/pg 140; 8th/pg 140)

117. Ans. (b) **Myotonia dystrophica** (Ref: Robbins 9th/pg 140)

118. Ans. (a) **Tri nucleotide repeats**

(Ref: Robbins 9th/pg 168)

119. Ans. (b, c, e); **b. Circular; c. Transmitted by mother only; e. Contains less gene than nuclear DNA**

(Ref: Robbins 9th/pg 170; 8th/pg 170)

120. Ans. (a) **Anticipation** (Ref: Robbins 9th/pg 168; 8th/pg 168)

121. Ans. (c) **Mitochondrial** (Ref: Robbins 9th/pg 171-172)

122. Ans. (b) **Prader-Willi syndrome**

(Ref: Robbins 9th/pg 171-172)

123. Ans. (c) **Bloom's syndrome** (Ref: Robbins 9th/pg 172-173)

**Bloom syndrome** is an **AR disorder** due to **defective DNA repair**

124. Ans. (d) **Trinucleotide repeat expansion**

(Ref: Robbins 9th/pg 168-171; 8th/pg 168-171)

125. Ans. (c) **X-linked** (Ref: R 9th/pg 168-171; 8th/pg 168-171)

**Fragile X syndrome** is **X linked**, as it is due to **Trinucleotide repeat expansion** involving **FRAXA** gene on X chromosome

126. Ans. (d) **Fragile X Syndrome** (Ref: R 9th/pg 171-172)

127. Ans. (a, c, d, e) **a. Mothers transmit their mtDNA to both their sons and daughters; c. Mitochondrial DNA codes for 37 genes; d. Mitochondrial disease commonly affect neuromuscular system; e. Mutation cause Leber hereditary optic neuropathy**

(Ref: Robbins 9th/pg 171-172)

128. Ans. (a, c); **a. Maternal inheritance; c. Not highly conserved and has high mutation rate**

(Ref: Robbins 9th/pg 171-172; 8th/pg 171; Harrison 18th/ chapter 61)

#### Mitochondrial DNA

- 16.6 kbp **circular dsDNA**
- Codes for **37 genes**: 2 rRNA, 22 tRNA & 13 protein **enzymes** that are components of the **respiratory chain** involved in **oxidative phosphorylation** and **ATP generation** (*option E*) and **affects muscles; CNS, liver and kidney**
- **Maternal inheritance** is seen in **mitochondrial diseases** (*option A*)
- Only **3% of mt-DNA** is **non-coding!**
- **Nuclear and mitochondrial DNA** are thought to be of **separate evolutionary origin**, with the mtDNA being derived from the circular genomes of the bacteria that were engulfed by the early ancestors of today's eukaryotic cells. This theory is called the **endosymbiotic theory** (*option B*)
- **mtDNA** is **not highly conserved** and has a **rapid mutation rate**, (*option C*)
- **Mitochondrial diseases** are sometimes (about 15% of the time) caused by **mutations in the mitochondrial DNA**. Other causes of mitochondrial disease are mutations in genes of the nuclear DNA, whose gene products are imported into the Mitochondria as well as acquired mitochondrial conditions. (*option D*)





129. Ans. (d) **Nemaline myopathy results due to mutations in mt-DNA**

(Ref: Robbins 9th/pg 171-172; 8th/pg 171)

130. Ans. (a) **Different expression of gene depending on parent of origin**

(Ref: R 9th/pg 172-173; 8th/pg 172-173)

**Genomic imprinting** refers to **different expression of gene depending on parent of origin**

131. Ans. (c) **Germ line mosaicism** (Ref: Robbins 9th/pg 174)

### Germ Line/ Gonadal Mosaicism

- Mutation that **occurs post-zygotically** during early (embryonic) development.
- If the mutation affects only cells destined to form the gonads, the **gametes carry the mutation, but the somatic cells of the individual are completely normal.**
- **Effect:** In some **autosomal dominant** disorders, e.g. **osteogenesis imperfecta**, **phenotypically normal** parents have **more than one affected child.**

132. Ans. (c) **Mitochondrial inheritance**

(Ref: Robbins 9th/pg 171-172)

The given pedigree shows:

**All children of the affected mother have the disease:** a characteristic of **Mitochondrial inheritance**

133. Ans. (a) **Prader-Willi syndrome**

(Ref: Robbins 9th/pg 172-173; 8th/pg 172-173)

**Discussing the options one by one:**

- A. **Prader Willi syndrome:** Uniparental disomy of maternal chromosome 15
- B. **Angelman syndrome:** Uniparental disomy of paternal chromosome 15
- C. **Hydatidiform mole:** Contains a normal number of diploid chromosomes, but they are all of **paternal** origin. The opposite situation occurs in **ovarian teratoma**, with 46 chromosomes of **maternal** origin. Note that **Hydatidiform mole & ovarian Teratoma are also due to uniparental disomy**
- D. **Klinefelter's syndrome: 47, XXY**

134. Ans. (b, c); **b. LHON & CPEO are typical examples; c. Most common abnormality shown is neurological**

(Ref: Robbins 9th/pg 171-172; 8th/pg 171)

- **Mitochondrial diseases** have **maternal inheritance:** **LHON & CPEO** are typical examples
- Most commonly involves **neuromuscular system**
- **Li-Fraumeni syndrome: p53 mutation;** Most common mutation associated with human cancers.

135. Ans. (a, b, d, e); **a. Oncocytoma; b. Kearn-Sayre syndrome; d. Mitochondrial myopathy; e. Leigh's disease**

(Ref: Robbins 9th/pg 171-172 & 953; 8th/pg 171 & 964)

### Oncocytoma

- **Benign tumors** with cells showing intense eosinophilia due to **characteristic accumulation of mitochondria;**
- May involve **salivary glands, thyroid, parathyroid, kidney, pituitary and lung**
- **Mitochondrial abnormality is seen, though it does not have mitochondrial inheritance.**

136. Ans. (d) **Child with seizure**

137. Ans. (c) **Ribosomal RNA** (Ref: Robbins 9th/pg 172-173)

- Two of the leading methods of gene silencing are RNA interference (RNAi), Si RNA and antisense oligonucleotides (ASOs), ds RNA
- It does not affect ribosomal RNA

138. Ans. (c) **Both 1 and 2** (Ref: Robbins 9th/pg 177-178)

- In Microarray, there is study of multiple genes, based on reverse hybridization technology. So it is highly possible to compare the DNA of patient and Normal DNA.
- Detects genomic abnormalities without prior knowledge of the genes involved, using microarray

139. Ans. (a) **Used for known Genetic loci**

(Ref: Robbins 9th/pg 177-178; 8th/pg 179-180)

In Array CGH (**Array-Based Comparative Genomic Hybridization**):

- Test DNA and a reference DNA are labeled with **two different fluorescent dyes**
- It is **used for unknown Genetic loci**
- **Primers for genes to be studied are hybridized on the chip;**
- **Both duplication and deletion** can be studied at the same time.
- It can **detect multiple abnormalities at the same time, by detecting difference in signal with duplication, deletion and normal genes;**

140. Ans. (a) **FISH**

(Ref: Robbins 9th/pg 177; 8th/pg 179)

141. Ans. (c) **5 mb**

(Ref: Robbins 9th/pg 158; 8th/pg 159)

Resolution of light microscope of viewing chromosome (Karyotype): 5 mb

**Resolution of FISH: 200 Kilobases (more sensitive than karyotype)**

142. Ans. (d) **Monitoring amplification of target nucleic acid**

(Ref: Robbins 9th/pg 175-176; 8th/pg 174)

**Real time polymerase chain reaction** is done for:

- **Monitoring amplification of target nucleic acid**
- **Quantitate amount of DNA in sample**

143. Ans. (b) **Metaphase**

(Ref: Robbins 9th/pg 158; 8th/pg 159)





## In Karyotype

- Chromosomes are examined after **arresting dividing cells in metaphase** with mitotic spindle inhibitors (e.g. N-diacetyl-N-methylcolchicine), followed by staining
- Resolution of Karyotyping: 2 to 5 million base pairs<sup>o</sup>**

### 144. Ans. (a) Study of multiple genes

(Ref: R 9th/pg 177-178)

In Microarray, there is study of multiple genes, based on reverse hybridization technology.

**Detects genomic abnormalities without prior knowledge of the genes involved, using microarray.**

### 145. Ans. (b) G-banding

(Ref: Robbins 9th/pg 158; 8th/pg 159)

### 146. Ans. (b) Blood monocyte

(Ref: R 9th/pg 158; 8th/pg 159)

- To produce a karyotype, **cells capable of growth and division is used**, as karyotyping is done by arresting mitosis in dividing cells in metaphase.
- None of the leukocytes in blood normally divide, but **lymphocytes can readily be induced to proliferate, providing a very accessible source of metaphase cells; but monocytes cannot be used.**

### 147. Ans. (c) Intestinal lining cancer

(Ref: Robbins 9th/pg 291)

Genes	Protein	Function	Associated Cancers
<b>BRCA1 (17q21)</b>	Breast cancer-1	Tumor suppressor, Transcriptional regulation,	<b>Ovarian, male breast cancer (but lower than BRCA2), prostate, pancreas, fallopian tube</b>
<b>BRCA2 (13q12-13)</b>	Breast cancer-2	Repair of double-stranded DNA breaks	Ovarian, male breast cancer, prostate, pancreas, <b>stomach, melanoma, gallbladder, bile duct, pharynx</b>

### 148. Ans. (a) Cleft lip (Ref: Robbins 9th/pg 158; 8th/pg 157)

### 149. Ans. (d) VSD

**HOX gene mutation can cause A. Syndactyly, B. Polydactyly, C. Fused carpal bones**

#### HOX genes (also known as 'homeotic' genes)

<b>Properties</b>	Their protein product is a <b>transcription factor</b> . They contain a <b>DNA</b> sequence known as the <b>homeobox</b> .
<b>Physiologic Role</b>	In the <b>development of the central nervous system, axial skeleton, limbs, gut, urogenital tract and external genitalia,</b>
<b>Examples</b>	<b>HOXD13</b> is mutated in synpolydactyly and <b>HOXA13</b> in Hand-Foot-Genital syndrome.

### 150. Ans. (a, b, c, d, e) a. Realtime PCR; b. Denaturing gradient gel electrophoresis; c. DNA sequencing; d. Restriction fragment polymorphism (RFLP); e. Single-strand conformational polymorphism

(Ref: R 9th/pg 177)

Techniques used detecting for Gene Mutation is/are given in pretexts

### 151. Ans. (a) Bloom syndrome

(Ref: R 9th/pg 314; 8th/pg 312)

### 152. Ans. (b) Hormonal evaluation

'Karyopyknotic index' is a method for **Hormonal evaluation; Has nothing to do with karyotyping**. Karyopyknotic index is % of **intermediate and superficial cells of squamous epithelium of vagina** which have pyknotic nuclei.

### 153. Ans. (a) Present in males

(Ref: Concepts of Genetics 7th ed/pg 208)

**Davidson bodies** are small nuclear buds of chromatin found in neutrophil leucocytes in females. Often shaped like a drumstick, they are found in up to six percent of cells. These bodies are absent in males (compare with Barr bodies)

# Diseases of the Immune System

## Key Points

- » **Innate immune** response is **immediate**, limited, **nonspecific**, **short lived** while **adaptive response** takes time, **specific & long lived**
- » Toll-like-receptors are pattern recognition receptors act by transcription of **NF- $\kappa$ B** from **nucleus**
- » NOD-like-receptors (NLRs) and the Inflammasome act via **caspase-1**
- » B-lymphocytes arise and develop in bone marrow while T-lymphocytes arise from **bone marrow** and develop in **Thymus**
- » **CD4** molecules bind to **class II<sup>o</sup>** MHC ( $4 \times 2 = 8$ ) and **CD8** molecules bind to class I<sup>o</sup> MHC ( $8 \times 1 = 8$ ) "**Rule of 8**"
- » **Innate lymphoid cells are lymphocytes that lack TCRs** but produce **cytokines similar T cells**
- » Hypersensitivity is poorly controlled, **excessive**, or **misdirected effector mechanisms** of defense against infectious pathogens
- » Substances secreted by **Mast cells** are Histamine, Heparin and Neutrophil chemotactic factor
- » LE or hematoxylin bodies<sup>o</sup> strongly indicative of SLE
- » **Class I Lupus Nephritis is the least common<sup>o</sup>** and **class IV is the most common pattern<sup>o</sup>**
- » **In transplant immunology, 50% HLA alleles matching** is required for kidney while 0% for cornea, liver and heart.
- » **The most Common Primary immunodeficiency is Isolated IgA immunodeficiency**
- » **HIV in CNS** infects macrophages and microglia, while **Neurons are not infected**
- » **Rectal biopsy is most specific** and **best site** for Amyloidosis diagnosis

## Key Recent Updates

- »  $\beta_2$  microglobulin of MHC I is coded by chromosome 15
- » **Natural IgM** antibodies are responsible for hyperacute rejection.



## THE NORMAL IMMUNE RESPONSE

Differences between Innate and Adaptive Immunity

Features	Innate	Adaptive
<b>Lag phase</b>	Absent, response is <b>immediate</b> <sup>a</sup>	Present, response <b>takes more time</b> <sup>a</sup> (a few days)
<b>Specificity</b>	<b>Limited</b>	<b>More specific</b> <sup>a</sup>
<b>Diversity</b>	<b>Limited</b> , hence limited specificity	<b>Extensive</b> , <sup>a</sup> a wide range of antigen receptors present
<b>Memory</b>	<b>Absent</b> , subsequent exposure to agent generate the same response	<b>Present</b> , <sup>a</sup> subsequent exposures to the same agent induce <b>amplified response</b> <sup>a</sup>

## INNATE IMMUNITY

### Components of Innate Immunity

- **Epithelial<sup>a</sup> barriers:** Skin, mucosa of GIT and respiratory tract
- Mucosa-associated lymphoid tissue (MALT)
- **Phagocytic cells** (mainly **neutrophils<sup>a</sup>** and **macrophages<sup>a</sup>**)
- **Dendritic cells<sup>a</sup>**
- **Natural killer (NK) cells<sup>a</sup>** **Complement system<sup>a</sup>**
- **C-reactive protein**
- **Innate lymphoid cells** <sup>a</sup>
- **Mannose-binding lectin** <sup>a</sup>
- **Lung surfactant** <sup>a</sup>


### Cellular Receptors (Pattern Recognition Receptors)<sup>a</sup>

Innate immunity cells are capable of recognizing certain microbial components that are often essential for infectivity.

<b>Types of pattern</b>	<ul style="list-style-type: none"> <li>• <b>Pathogen-associated molecular patterns<sup>a</sup></b>- recognize <b>microbial</b> components</li> <li>• <b>Damage-associated molecular patterns</b> - recognize <b>injured and necrotic cells</b></li> </ul>
<b>Location of receptor</b>	<ul style="list-style-type: none"> <li>• <b>Plasma membrane</b> receptors detect <b>extracellular</b> microbes</li> <li>• <b>Endosomal</b> receptors detect <b>ingested<sup>a</sup></b> microbes</li> <li>• <b>Cytosolic</b> receptors detect <b>microbes in the cytoplasm</b></li> </ul>
<b>Classes (examples)</b>	<ul style="list-style-type: none"> <li>• <b>Toll-like receptors (TLR)<sup>a</sup></b></li> <li>• <b>NOD-Like Receptors (NLR)</b></li> <li>• <b>C-type lectin receptors (CLRs)</b> detect <b>fungal glycans<sup>a</sup></b></li> <li>• <b>RIG-like receptors (RLRs)</b> detect <b>nucleic acids of viruses</b> <sup>a</sup></li> <li>• <b>G protein-coupled</b> receptors- recognize bacterial peptides with <b>N-formyl-methionyl residues<sup>a</sup></b></li> <li>• <b>Mannose</b> receptors- recognize <b>microbial sugars<sup>a</sup></b></li> </ul>

### Toll-Like Receptors (TLR)


- Found in **plasma membrane** and **endosomal vesicles<sup>a</sup>**
- **Mechanism of action:** Activation of transcription factors like-
  - **NF-κB<sup>a</sup>** stimulates expression of **adhesion molecules** and causes **activation of leukocytes<sup>a</sup>**
  - **Interferon regulatory factors (IRFs):** Stimulates production of antiviral type I interferons



### Types of Toll-Like Receptors (TLRS)

Receptor	Cellular Source	Ligand
<b>TLR2</b>	PMNs, Dendritic cells, Monocytes	<b><i>Mycobacterium tb</i><sup>a</sup></b> , Gram +ve bacteria
<b>TLR3</b>	Dendritic cells & NK cells	dsRNA of <b>viruses<sup>a</sup></b>
<b>TLR4</b>	<b>Macrophages</b> , Dendritic cells, Epithelial cells	<b><i>E. coli</i> LPS<sup>a</sup></b> (Gram-negative), <b>Chlamydia</b>
<b>TLR5</b>	Monocytes, immature Dendritic cells, Epithelial cells, NK, T cells	<b>Flagellin<sup>a</sup></b> , <b>Toxoplasma</b>
<b>TLR7/8</b>	B cells, plasmacytoid precursors dendritic cells	ssRNA, Imidazoquinolines
<b>TLR9</b>	Plasmacytoid precursor dendritic cells, B cells, macrophages, PMNs, NK cells, & microglia cells	CpG DNA

<b>Latest Update</b>	
<b>NOD-Like Receptors (NLRs) and the Inflammasome</b>	
<b>Location</b>	<b>Cytosolic receptors</b>
<b>Recognize</b>	Necrotic cells, ion disturbances (e.g., loss of K <sup>+</sup> ), urate crystal & some microbial products.
<b>Mechanism of action</b>	Activates Caspase-1 → activates IL-1 → recruits leukocytes & induces fever.
<b>Role in diseases</b>	<ul style="list-style-type: none"> <li>• <b>Periodic fever syndromes<sup>a</sup></b> ('autoinflammatory' syndromes): Due to Gain-of-function mutations</li> <li>• <b>Obesity-associated type 2 diabetes &amp; Atherosclerosis:</b> due to activation by lipids and cholesterol crystals</li> <li>• <b>Inflammation in Gout:</b> Due to recognition of urate crystals by a class of NLRs</li> </ul>



## Macrophages: A Part of Mononuclear-Phagocyte System

### Functions

- Antigen-presenting cells<sup>Q</sup> in T-cell activation.
- Cell-mediated immunity<sup>Q</sup>
- Efficiently phagocytose<sup>Q</sup> and destroy microbes that are opsonized (coated) by IgG or C3b.


## Natural Killer Cells: 5–10% of Peripheral Lymphocytes<sup>Q</sup>

- Derived from:** Large granular lymphocytes<sup>Q</sup>
- Markers**
  - CD16<sup>Q</sup> - An Fc receptor for IgG, makes NK cells lyse IgG-coated target cells: known as 'antibody-dependent cell-mediated cytotoxicity' (ADCC)<sup>Q</sup>
  - CD56<sup>Q</sup> - NCAM (Natural Killer Cell Adhesion Molecule)
- Function:** Destroy irreversibly stressed, virus-infected<sup>Q</sup> cells & tumor cells<sup>Q</sup> without prior exposure
- Activating Receptors:** NKG2D receptors: recognize surface molecules induced by various kinds of stress, such as infection & DNA damage
- Inhibiting Receptors:** Recognize self class I MHC<sup>Q</sup> molecules, prevent NK cells from killing normal cells<sup>Q</sup> (MHC Nonrestricted)
- Cytokines secreted**
  - IL-2<sup>Q</sup> and IL-15 stimulate proliferation of NK cells



### High Yield Facts

- NK cell is a large granular lymphocyte<sup>Q</sup>
- The primary function of Toll-like Receptors is activation of immune system<sup>Q</sup>
- As NK cells neither have B nor T-cell markers, hence called 'Null Cells'<sup>Q</sup>
- Super antigens bind directly to both MHC II and T cell receptor causing T cell activation and bind T cells directly to lateral aspect of T cell receptor irrespective of antigen specificity of TCR<sup>Q</sup>
- Toll-like receptors types II and IV are involved in action against bacterial endotoxins
- Interleukin characteristically produced in a Th1 response is IL-2<sup>Q</sup>
- Interleukin secreted by macrophages, stimulating lymphocytes is IL-1<sup>Q</sup>
- Dendritic cell expresses MHC 2<sup>Q</sup>



## ADAPTIVE IMMUNITY

### Types of Adaptive Immunity

Cell-mediated Immunity	Humoral Immunity
<ul style="list-style-type: none"> <li>Protects against intracellular<sup>Q</sup> microbes</li> <li>Mediated by T lymphocytes<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Protects against extracellular<sup>Q</sup> microbes &amp; their toxins</li> <li>Mediated by B – lymphocytes and Immunoglobulins<sup>Q</sup></li> </ul>

### Cells of Adaptive Immune System

#### T Lymphocytes (T Cells): 60 – 70% of Lymphocytes

- Arise from bone marrow (Hematopoietic stem cells) and develop in Thymus<sup>Q</sup>
- 60% of mature T cells are CD4+ and 30% are CD8+ ; so, CD4:CD8 ratio is 2:1<sup>Q</sup>

Type of T cell	Characteristics/functions
Helper T cells	Stimulate B cells <sup>Q</sup> to make antibodies & activate phagocytes to destroy microbes.
Naive CD4+ T cells	Recognize peptides displayed by dendritic cells → secrete IL-2 (autocrine growth factor) → stimulates proliferation of T cells → differentiation of T cells to Th1 or Th17 cells
Cytotoxic T cells	Kill virus infected and tumor cells <sup>Q</sup>

### Latest Update

#### Other types of T-cells

- Regulatory T cells:** Limit immune responses & prevent reactions against self-antigens. <sup>Q91</sup> They are CD4+, CD25+, Fox P<sub>3</sub>+
- NK-T cells:** T cells that express markers are also found on NK cells <sup>Q91</sup>
- γ δ T cells:** Aggregate at epithelial surfaces of skin & mucosa of GIT & genitourinary tract <sup>Q91</sup>

### T-cell Receptor (TCR)


- Antigen-specific
- 95% of TCR have a α and β polypeptide chain<sup>Q</sup>
- MHC restriction:** αβ TCR recognizes peptide antigens that are presented by major histocompatibility complex<sup>Q</sup> (MHC) molecules on the surfaces of antigen-presenting cells
- CD3 complex and ζ chain dimer** – also involved in signal transduction





## Helper T cells

### Types of Helper T Cells



Characteristics	Th <sub>1</sub> cells	Th <sub>2</sub> cells	Th <sub>17</sub> cells
<b>Activated by</b>	IFN $\gamma$ , IL-12 <sup>a</sup>	IL-4	TGF $\beta$ , <sup>a</sup> IL-6, IL-1, IL-23
<b>Cytokines produced</b>	IFN $\gamma$ <sup>a</sup>	IL-4, IL-5, IL-13 <sup>a</sup>	IL-17, IL-22
<b>Functions</b>	Activates Macrophages, <sup>a</sup> stimulates IgG production	Stimulates IgE <sup>a</sup> production, activates mast cells & eosinophils <sup>a</sup>	Recruitment of neutrophils, monocytes <sup>a</sup>
<b>Defense against</b>	Intracellular microbes	Helminthic parasites	Extracellular bacteria, fungi
<b>Role in disease</b>	Autoimmune diseases e.g. IBD, psoriasis & Chronic inflammation (granuloma)	Allergies	Autoimmune & chronic inflammatory diseases (e.g. IBD, psoriasis, MS)

### High Yield Facts

- CD4 molecules bind to class II<sup>a</sup> MHC (4X2=8) & CD8 molecules bind to class I<sup>a</sup> MHC (8X1=8) "Rule of 8"
- $\gamma\delta$  TCR does not require MHC<sup>a</sup> for antigen recognition
- Epstein-Barr virus (EBV) enters B cells via CR2<sup>a</sup>
- RBCs lack HLA antigen<sup>a</sup>

### High Yield Facts


- By CD40L- and helper T-cell actions, antibodies of different classes (IgG, IgA, IgE) produced
- Isotype switching is induced by IFN- $\gamma$  and IL-4.<sup>a</sup>
- Helper T cells also stimulate affinity maturation, that improves quality of humoral immune response<sup>a</sup>
- As these helper T-cells migrate to and reside in the germinal centers: called follicular helper T cells (TFH)

## Humoral Immunity: Activation of B Lymphocytes

<b>Acts against</b>	Extracellular Microbes <sup>a</sup>
<b>Mechanism</b>	On Activation $\rightarrow$ B cells proliferate & differentiate into plasma cells $\rightarrow$ secrete Antibodies

### B Lymphocytes: 10% – 20% of Lymphocytes

<b>Arise from</b>	Bone marrow (Hematopoietic stem cells) <sup>a</sup>
<b>Mature in</b>	Bone marrow
<b>Found in</b>	Peripheral lymphoid tissues such as lymph nodes, spleen & mucosa-associated lymphoid tissues
<b>Functions</b>	Mediates Humoral Immunity, Stimulation by Antigen $\rightarrow$ B cells develop into plasma cells <sup>a</sup> $\rightarrow$ produce antibodies
<b>Components of B-cell antigen receptor</b>	<ul style="list-style-type: none"> <li>• Ig receptors: IgM and IgD Isotypes<sup>a</sup></li> <li>• Ig<math>\alpha</math> (CD79a) and Ig<math>\beta</math> (CD79b)</li> <li>• Type 2 complement receptor (CR2 or CD21): recognize complement products</li> <li>• CD40-receives signals from helper T cells.</li> </ul>



### High Yield Facts

- **Homocytotropic Ab:** Higher affinity for receptors of cells in species in which they are produced, e.g. IgE
- All 4 IgG subclasses cross placenta, IgG 1 & IgG 3 predominate
- IgG form 15–18% of total plasma protein

Now, having understood T cells and B cells, let's now see how is antigen presented to T cells?

### Antigen Presenting cells; cells that present Ag to T-cells

- Antigen-presenting cells (APCs) initiate T-cell responses against protein antigens<sup>a</sup>
- Mature dendritic cells are the most potent<sup>a</sup> stimulator of Naive T-cells



### Types of Antigen Presenting Cells


Professional APC	Non-professional APC
Express MHC class II	Express MHC class I
Macrophages, dendritic cells and B-cells	Fibroblasts, thymic epithelial cells, thyroid epithelial cells, glial cells, pancreatic beta cells & vascular endothelial cells.

### Dendritic Cells: Professional Antigen-Presenting Cells

*Why do dendritic cells have a key role in antigen presentation?*

- Immature dendritic cells (Langerhans cells) within the epidermis are at **perfect site** to capture antigens
- Express many receptors like TLRs and lectins<sup>Q</sup> for capturing and responding to microbes.
- In response to microbes, they are **recruited to the T-cell zones** of lymphoid organs to present antigens to T cells.


- Dendritic cells **express high levels of MHC<sup>Q</sup>** for presenting antigens to and activating T cells.



## CELL-MEDIATED IMMUNITY: ACTIVATION OF T LYMPHOCYTES

### Activation of T-cell Requires Three Signals

- Signal 1:** TCR on T-cell with MHC on APC<sup>Q</sup>
- Signal 2:** CD28<sup>Q</sup> on T-cell with B7-1 (CD80) /B7-2(CD86) on APC
- Signal 3:** Cytokine **IL-1** from APC stimulates T-cell which in turn secretes **IL-2<sup>Q</sup>** (which has **autocrine** effect)



*R<sup>9th</sup>*

### Latest Update

- M cells (microfold cells) are APC
- Innate Lymphoid Cells (ILCs)**
- Lymphocytes that lack TCRs but produce **cytokines similar to those that are made by T cells**
- Example: NK cells

- Cytokines produced:** IFN- $\gamma$ , IL-5, IL-17, and IL-22.
- Functions:**
  - Early defense against infections
  - Recognition and elimination of stressed cells (so-called stress surveillance)<sup>Q</sup>
  - Providing cytokines that influence the differentiation of T lymphocytes.

## TISSUES OF THE IMMUNE SYSTEM

### Generative (Central) Lymphoid Organs

- Thymus:** Site of development of T cells<sup>Q</sup>
- Bone marrow:** Site of maturation of B lymphocytes<sup>Q</sup>

### Peripheral Lymphoid Organs

In which adaptive immune responses to microbes are initiated.



<b>Lymph nodes</b>	<ul style="list-style-type: none"> <li>• <b>B cells</b> are concentrated in <b>follicles</b>, located in <b>cortex</b> of lymph node<sup>Q</sup></li> <li>• <b>T lymphocytes</b> are concentrated in the <b>paracortex</b>, adjacent to the follicles<sup>Q</sup></li> <li>• Follicles contain <b>follicular dendritic cells</b> that are involved in <b>activation of B cells</b><sup>Q</sup>, &amp; the <b>paracortex</b> contains the <b>dendritic cells</b> that <b>present antigens</b> to T lymphocytes.</li> </ul>	<p>Structure of a lymph node</p>
<b>Spleen</b>	<ul style="list-style-type: none"> <li>• <b>T lymphocytes</b> are concentrated in <b>periarteriolar lymphoid sheaths</b><sup>Q</sup></li> <li>• <b>B cells</b> reside in the <b>follicles</b><sup>Q</sup></li> </ul>	
<b>Mucosal &amp; Cutaneous lymphoid tissues</b>	Epithelia of skin, GIT & respiratory tracts, pharyngeal tonsils & <b>Peyer's patches</b> <sup>Q</sup> of intestine ( <b>Mucosa-associated lymphoid tissue or MALT</b> ), also found in <b>thyroid, breast, lung, salivary glands, eye &amp; skin</b>	

**Latest Update**

**Lymphocyte Recirculation**


- Naive T lymphocytes traverse the **peripheral lymphoid organs** like lymph nodes through specialized postcapillary venules called **high endothelial venules (HEVs)** where **immune responses are initiated**, and **effector lymphocytes** migrate to sites of infection and inflammation.

## MAJOR HISTOCOMPATIBILITY COMPLEX (MHC)


- Display peptide fragments of protein antigens for recognition by antigen-specific T cells
- **On short arm of Chromosome 6 (6p)**<sup>Q</sup>
- Also called **human leukocyte antigens (HLA)**<sup>Q</sup> - detected on leukocytes by the binding of antibodies.
- **Highly polymorphic**<sup>Q</sup> so constitutes a formidable barrier in organ transplantation.

### Classification on the Basis of their Structure, Cellular Distribution and Function

Properties	Class I MHC molecules	Class II MHC molecules	MHC III
<b>Location</b>	All nucleated cells & platelets <sup>Q</sup>	APCs: B cells, dendritic cells, endothelial cells & fibroblasts <sup>Q</sup>	<ul style="list-style-type: none"> <li>• No direct role in immune system<sup>Q</sup></li> <li>• Codes for: <ul style="list-style-type: none"> <li>■ <b>Complement components</b> C2, C4, properdin, factor B<sup>Q</sup>, TNF, HSP-70, Tyrosine hydroxylase</li> </ul> </li> </ul>
<b>Encoded by</b>	HLA-A, HLA-B, and HLA-C <sup>Q</sup>	HLA-DP, HLA-DQ, and HLA-DR	
<b>Peptide binding cleft</b>	Comprised of $\alpha_1$ and $\alpha_2$	Comprised of $\alpha_1$ and $\beta_1$ domains	
<b>Antigens displayed</b>	<b>Viral &amp; tumor antigens</b> located <b>intracellularly</b> <sup>Q</sup>	<b>Extracellular microbes</b> <sup>Q</sup> and soluble proteins.	
<b>Recognized by</b>	<b>CD8+</b> T lymphocyte (MHC I-restricted)	<b>CD4+</b> T cells (MHC II-restricted)	
<b>Major role in</b>	<b>Graft rejection</b> <sup>Q</sup>	<b>GVHD</b> (graft vs host disease) <sup>Q</sup>	







## HLA Typing & HLA Matching

Methods of HLA typing	Uses of HLA matching
<ul style="list-style-type: none"> <li>Mixed lymphocyte reaction<sup>Q</sup></li> <li>Serological detection of HLA<sup>Q</sup></li> <li>Molecular detection methods: <ul style="list-style-type: none"> <li>Sequence specific primer PCR (SSP-PCR)</li> <li>Sequence specific oligonucleotide probes (SSOP)</li> <li>Direct DNA sequencing</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>Organ &amp; stem cell transplantation<sup>Q</sup></li> <li>Disease association studies</li> <li>Disputed paternity<sup>Q</sup></li> </ul>

## HYPERSENSITIVITY (HSN)


### General Features

- Elicited by **exogenous antigens** (microbial or nonmicrobial) or **endogenous self-antigens**
- Results from an **imbalance** between the effector and the control mechanisms
- Often associated with the **inheritance** of particular susceptibility genes
- Poorly controlled, **excessive, or misdirected effector mechanisms** of defense against infectious pathogens


## Immediate (Type I) Hypersensitivity

<b>What is it</b>	Rapid immunologic reaction occurring in a <b>previously sensitized</b> <sup>Q</sup> individual	
<b>Triggered by</b>	Binding of an <b>antigen to IgE antibody</b> on the surface of <b>mast cells</b> . <sup>Q</sup>	
<b>Also called</b>	<b>Allergy</b> , and the <b>antigens</b> that elicit them are <b>allergens</b> . <sup>Q</sup>	
<b>2 Phases</b>	<b>Immediate reaction</b>	<b>Late-phase reaction</b>
<b>Time taken</b>	Within <b>minutes to hours</b> <sup>Q</sup>	Sets in <b>2 to 24 hours to days</b> <sup>Q</sup>
<b>Characterized by</b>	<b>Vasodilation, vascular leakage</b> , and depending on the location, smooth muscle spasm or glandular secretions	Infiltration of tissues with <b>eosinophils</b> <sup>Q</sup> , neutrophils, basophils, monocytes & <b>CD4+T cells</b> <sup>Q</sup> , causing <b>tissue destruction &amp; mucosal epithelial cell damage</b>
<b>Mediators</b>	<b>A. Preformed Mediators:</b> <ul style="list-style-type: none"> <li>Vasoactive amines, e.g. <b>Histamine</b></li> <li>Enzymes: e.g. <b>Chymase, tryptase</b></li> </ul> <b>B. Lipid Mediators:</b> <ul style="list-style-type: none"> <li>Arachidonic acid-derived products: <b>Leukotrienes B4, C4 &amp; D4, Prostaglandin D2</b></li> <li><b>Platelet-activating factor (PAF)</b><sup>Q</sup></li> </ul> <b>C. Cytokines:</b> <b>TNF, IL-1, IL-4</b> <sup>Q</sup>	<ul style="list-style-type: none"> <li><b>Eotaxin</b></li> <li><b>IL-5</b></li> <li><b>MPB (Major Basic Protein)</b><sup>Q</sup></li> <li><b>ECP (Eosinophil Cationic Protein)</b></li> </ul>
<b>Examples of type I HSN</b>	<ul style="list-style-type: none"> <li><b>Local:</b> <b>Urticaria, angioedema, allergic rhinitis (hay fever), bronchial asthma, eczema</b><sup>Q</sup></li> <li><b>Systemic:</b> <b>anaphylactic shock</b><sup>Q</sup> (Initiated by histamine)</li> <li>Others: <b>Theobald Smith phenomenon, Prausnitz Kusnter reaction</b>,<sup>Q</sup> <b>Casoni's test</b>,<sup>Q</sup> <b>Schultz-Dale phenomenon</b></li> </ul>	

### Mechanism of Immediate Reaction



### Mechanism of Late-phase Reaction







## High Yield Facts

- **Histamine** causes **smooth muscle contraction**, **increased vascular permeability** & **increased mucus secretion**<sup>a</sup> by nasal, bronchial & gastric glands.
- **Leukotrienes C4 & D4** are the **most potent**<sup>a</sup> vasoactive & spasmogenic agents
- **Leukotriene B4** is highly **chemotactic**<sup>a</sup> for neutrophils, eosinophils, and monocytes.
- **Prostaglandin D2** is the **most abundant**<sup>a</sup> mediator produced in mast cells
- **Platelet-activating factor (PAF)**<sup>a</sup> causes platelet aggregation, release of histamine, bronchospasm, increased vascular permeability & vasodilation.
- **TNF, IL-1, chemokines**: promote **leukocyte recruitment**<sup>a</sup>
- **IL-4**: **increases TH2 response**<sup>a</sup>
- **IL-2** is a **growth & survival factor** for **activated & regulatory T cells**<sup>a</sup>


## Antibody-Mediated (Type II) Hypersensitivity

Caused by	Tissue specific antibodies that react with antigens present on cell surfaces & extracellular matrix		
Mechanisms	<pre> graph TD     IgM[IgM] --&gt; A[Activate complement system C3b, C4b]     IgG[IgG] --&gt; B[Recognized by phagocyte Fc receptors Opsonization]     A --&gt; C[Phagocytosis of opsonized cells &amp; destruction]     B --&gt; C           </pre>	Examples	Target Antigen
Opsonization & Phagocytosis		<b>Autoimmune hemolytic anemia</b> <sup>a</sup>  <b>Autoimmune thrombocytopenic purpura</b> <sup>a</sup>	<b>RBC membrane proteins</b> e.g., Rh blood group Antigen  <b>Platelet membrane proteins</b> e.g. GpIIb/IIIa <sup>a</sup>
Inflammatory damage	<pre> graph TD     A[Antibody to fixed tissues, e.g.; basement membranes &amp; ECM] --&gt; B[Activate complements- C5a -&gt; chemotaxis of neutrophils/monocytes, C3a, C5a -&gt; increase vascular permeability]     B --&gt; C[Damage tissues]           </pre>	<b>Pemphigus vulgaris</b> <sup>a</sup> <b>Vasculitis</b> caused by ANCA <b>Goodpasture syndrome</b> <sup>a</sup> <b>Acute rheumatic fever</b> <sup>a</sup>	<b>Epidermal cadherin</b> <sup>a</sup> <b>Neutrophil granule proteins</b> from activated neutrophils <b>Non-collagenous pr. in BM of glomeruli &amp; alveoli</b> <sup>a</sup> <b>Streptococcal</b> <sup>a</sup> wall Ag (Ab cross-reacts with myocardial Ag)
Cellular dysfunction	<b>Antibodies may dysregulate receptor function without causing cell injury or inflammation, such as:</b> <ul style="list-style-type: none"> <li>• <b>Block ACh receptors</b> in motor end plates of skeletal muscles in <b>myasthenia gravis</b><sup>a</sup></li> <li>• <b>Stimulate TSH receptor</b> causing <b>hyperthyroidism</b><sup>a</sup></li> <li>• <b>Long-acting thyroid stimulator (LATS)</b>: IgG that stimulates thyroid function similar to but slower than TSH (i.e. long-acting)</li> </ul>	<b>Myasthenia gravis</b> <sup>a</sup> Graves disease Insulin-resistant diabetes <b>Pernicious anemia</b> <sup>a</sup>	<b>Acetylcholine receptor</b> <sup>a</sup> <b>TSH receptor</b> <sup>a</sup> Insulin receptor <b>Intrinsic factor of gastric parietal cells</b> <sup>a</sup>



## Immune Complex–Mediated (Type III) Hypersensitivity

<b>Initiated by</b>	Nonspecific antibody towards an antigen	
<b>Morphology</b>	<ul style="list-style-type: none"> <li>Principal morphologic manifestation of immune complex injury is <b>acute vasculitis</b><sup>a</sup></li> <li><b>Fibrinoid necrosis</b><sup>a</sup> of vessel wall &amp; intense neutrophilic infiltration</li> </ul>	
<b>Mechanism</b>	<pre> graph TD     A[Interaction of antigen and antibody] --&gt; B[Formation of immune complexes (phase I) (locally at the site of injury or systemically in the circulation)]     B --&gt; C[Deposition of immune complexes (phase II)]     C --&gt; D[Activates complement]     D --&gt; E[Acute inflammation and injury (phase III) starts 10 days after antigen entry]           </pre>	
<b>Examples</b>	<b>Infections</b> <ul style="list-style-type: none"> <li>Subacute infective endocarditis<sup>a</sup></li> <li>Post-streptococcal glomerulonephritis<sup>a</sup></li> <li>Syphilis</li> </ul> <b>Malignant diseases</b> <ul style="list-style-type: none"> <li>Leukemias</li> <li>Hodgkin's disease<sup>a</sup></li> </ul>	<b>Autoimmune diseases</b> <ul style="list-style-type: none"> <li>Systemic lupus erythematosus<sup>a</sup></li> <li>Rheumatoid arthritis<sup>a</sup></li> <li>Polyarteritis nodosa</li> <li>Hashimoto's disease (thyroiditis)</li> </ul> <b>Drug reactions<sup>a</sup></b> <ul style="list-style-type: none"> <li>Serum sickness (systemic)</li> <li>Penicillamine toxicity</li> <li><b>Arthus reaction (localized)</b></li> </ul>





## T Cell-Mediated (Type IV) Hypersensitivity

<b>Mediated by</b>	Cells, not antibody: <b>Cell-mediated immunity</b>	
<b>Cells involved</b>	Inflammation due to <b>cytokines produced by CD4+ T cells<sup>a</sup></b> & <b>cell killing by CD8+ T cells<sup>a</sup></b>	
<b>Time taken</b>	Occurs <b>24–72 hours<sup>a</sup></b> after exposure: <b>delayed-type hypersensitivity (DTH)<sup>a</sup></b>	
<b>Mechanism</b>	<b>CD4+ T Cell–Mediated Inflammatory Reactions</b> <p><b>Activated macrophages:</b></p> <ul style="list-style-type: none"> <li>• ↑ <b>ability to phagocytose<sup>a</sup></b> &amp; kill microorganisms</li> <li>• ↑ <b>class II MHC</b> molecules on the surface</li> <li>• <b>Secrete TNF, IL-1 &amp; chemokines</b>, which promote inflammation</li> <li>• Produce <b>IL-12<sup>a</sup></b>, amplifying the <b>Th1 response</b></li> </ul>	<b>CD8+ T Cell–Mediated Cytotoxicity</b> <ul style="list-style-type: none"> <li>• Major role in <b>virus-infected cell &amp; tumor cells<sup>a</sup></b></li> <li>• Involves <b>perforins<sup>a</sup></b> &amp; <b>granzymes<sup>a</sup></b> (preformed mediators in granules of cytotoxic T cells)</li> <li>• <b>Perforins</b> facilitates release of <b>granzymes<sup>a</sup></b>.</li> <li>• Granzymes are <b>proteases<sup>a</sup></b> that cleave &amp; <b>activate caspases<sup>a</sup></b>, which induce <b>apoptosis<sup>a</sup></b> of target cells</li> <li>• <b>Activated CTLs</b> also express <b>Fas ligand<sup>a</sup></b>, with homology to <b>TNF</b>, which can <b>bind to Fas</b> expressed on target cells &amp; <b>trigger apoptosis<sup>a</sup></b></li> </ul>
<b>Examples of Type IV Hypersensitivity</b>	<b>Type 1 diabetes mellitus<sup>a</sup>, multiple sclerosis<sup>a</sup>, rheumatoid arthritis<sup>a</sup>, contact dermatitis<sup>a</sup></b> and <b>tuberculin reaction &amp; Granuloma formation</b>	

## IMMUNOLOGIC TOLERANCE

<b>Definition</b>	<b>Unresponsiveness<sup>a</sup></b> to an Ag induced by exposure of <b>lymphocytes to that antigen</b>
<b>Self-tolerance<sup>a</sup></b>	<b>Lack of responsiveness</b> to an individual's own antigens
<b>Mechanisms of self-tolerance</b>	Central tolerance & Peripheral tolerance

### Central Tolerance

<b>Mechanism</b>	Immature <b>self-reactive T- and B-lymphocyte</b> are killed in the <b>central</b> (or generative) lymphoid organs (the <b>thymus</b> for <b>T cells</b> and the <b>bone marrow</b> for <b>B cells</b> ) <sup>a</sup>
<b>T-cells</b>	<b>Negative selection or deletion:</b> Removal of self-reactive lymphocytes from the T-cell pool in <b>thymus<sup>a</sup></b>
<b>B-cells</b>	<b>Receptor editing:</b> Self-reacting B-cells edit their antigen receptor by <b>gene rearrangement<sup>a</sup></b> so that they are no longer reactive against self.

### High Yield Facts



- **Transfusion reaction** and **erythroblastosis fetalis** are **Type II hypersensitivity<sup>a</sup>**
- Main mediator in **Anaphylactic shock** is **Histamine<sup>a</sup>**
- Substances secreted by **Mast cells** are **Histamine, Heparin and Neutrophil chemotactic factor<sup>a</sup>**
- **Most important** cells in **type I hypersensitivity** are **Mast cells<sup>a</sup>**
- **Raji cell assay<sup>a</sup>** are used to identify **immune complexes<sup>a</sup>**
- **LATS** is a **IgG Ab<sup>a</sup>**



### Latest Update

- **AIRE (Autoimmune regulator) gene<sup>a</sup>** is responsible for removal of self-reacting immature T cells
- **Mutations in the AIRE gene** are the cause of an **autoimmune-polyendocrinopathy<sup>a</sup>**



## Peripheral Tolerance

<b>Definition</b>	Silencing of potentially auto-reactive T and B cells in peripheral tissues (Mainly for T cells) <sup>Q</sup>
<b>Mechanisms (S-A-D mnemonic)</b>	<p><b>Suppression by regulatory T cells. Regulatory T-cells<sup>Q</sup></b></p> <p>functions to prevent immune reactions against self-antigens. Regulatory T cells may play a role in the acceptance of the fetus.</p> <p>Regulatory T cells are CD4+ cells that express:</p> <ul style="list-style-type: none"> <li>• <b>CD25</b>: <math>\alpha</math> chain of the IL-2 receptor<sup>Q</sup></li> <li>• <b>FOXP3</b>: transcription factor of the forkhead family<sup>Q</sup></li> <li>• <b>CD25 &amp; FOXP3</b> → <b>Development &amp; maintenance of functional CD4+ regulatory T cells</b></li> <li>• FOXP3 mutations causes systemic autoimmune disease - <b>IPEX</b> (an acronym for <b>I</b>mmune <b>d</b>ysregulation, <b>P</b>olyendo-crinopathy, <b>E</b>nteropathy, <b>X</b>-linked)<sup>Q</sup></li> </ul> <p><b>Anergy<sup>Q</sup></b></p> <ul style="list-style-type: none"> <li>• Inhibition of self-reacting T-lymphocytes by certain receptors</li> </ul> <p><b>Mechanism: CTLA-4 &amp; PD-1</b> are inhibitory receptors &amp; are structurally homologous to CD28 which are the coreceptors for antigen recognition</p> <ul style="list-style-type: none"> <li>• Helper T cell have <b>CTLA-4<sup>Q</sup></b> (cytotoxic T-lymphocyte-associated protein 4) receptors which transmits an <b>inhibitory signal to T</b> cells. (CTLA-4 has higher affinity for B7 molecules than does CD28).</li> <li>• <b>PD-1 (programmed Death-1)<sup>Q</sup></b> receptor are upregulated on activated CD4 T-cells which binds to PD-L1 expressed on monocytes to induce IL-10 production for <b>immunosuppression</b>.</li> </ul> <p><b>Deletion by apoptosis</b></p> <ul style="list-style-type: none"> <li>• Self reacting T-cells are destroyed by apoptosis by either: <ul style="list-style-type: none"> <li>▪ <b>Overexpression of proapoptotic factor BIM<sup>Q</sup></b></li> <li>▪ <b>FAS-FAS Ligand pathway<sup>Q</sup></b></li> </ul> </li> </ul> <p><i>Mutations in the FAS gene results in <b>autoimmune lymphoproliferative syndrome (ALPS)<sup>Q</sup></b></i></p>

## AUTOIMMUNITY

<b>Definition</b>	Activation of self-reactive lymphocytes <sup>Q</sup> by combination of inheritance of <b>susceptibility genes</b> and <b>environmental triggers</b> such as infections & tissue damage
<b>Mechanisms</b>	<ul style="list-style-type: none"> <li>• <b>Defective tolerance or regulation</b>: failure of mechanisms that maintain self-tolerance<sup>Q</sup></li> <li>• Abnormal display of self-antigens<sup>Q</sup></li> <li>• <b>Inflammation</b> or an initial innate immune response which may induce autoimmunity</li> </ul>



### High Yield Facts

#### Immune-privileged sites:<sup>Q</sup>

- **Testis, eye, and brain<sup>Q</sup>**
- Tissues in which these antigens are located **do not communicate** with the blood and lymph
- **Difficult to induce immune responses** to antigens introduced into these sites
- Prolonged tissue inflammation on injury & release of antigen from these sites: **post-traumatic orchitis & uveitis<sup>Q</sup>**

### Role of Susceptibility Genes: Association of HLA Alleles with Disease

#### Association of non-MHC genes with Diseases

Gene	Diseases	Function of encoded protein & role in Disease
<b>Genes involved in immune regulation</b>		
<b>PTPN 22<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• Rheumatoid Arthritis<sup>Q</sup></li> <li>• Type I Diabetes<sup>Q</sup></li> <li>• Inflammatory Bowel Ds<sup>Q</sup></li> </ul>	<b>Protein tyrosine phosphatase<sup>Q</sup></b> affects signaling of lymphocytes & alter negative selection or activation of self- reactive T cells
<b>IL23R</b>	<ul style="list-style-type: none"> <li>• Inflammatory Bowel Ds</li> <li>• Psoriasis &amp; Ankylosing spondylitis</li> </ul>	<b>Receptor for T<sub>H</sub> 17-induces IL-23;<sup>Q</sup></b> alters differentiation of CD4+T cell into pathogenic T <sub>H</sub> 17 effector cells
<b>CTLA4</b>	<ul style="list-style-type: none"> <li>• Type I Diabetes</li> <li>• Rheumatoid Arthritis</li> </ul>	<b>Inhibits T cells responses<sup>Q</sup></b> by terminating activation & promoting activity of regulatory T cells; Interferes with self-tolerance
<b>IL2RA</b>	<ul style="list-style-type: none"> <li>• Multiple sclerosis</li> <li>• Type I Diabetes</li> </ul>	$\alpha$ chain of receptors for IL-2 may affect development of <b>effector cells</b> and/or regulation of immune responses

Contd...





Gene	Diseases	Function of encoded protein & role in Disease
<b>Genes involved in immune response to microbes</b>		
<b>NOD2<sup>a</sup></b>	Inflammatory Bowel Disease	<b>Cytoplasmic sensor of bacteria<sup>a</sup></b> expressed in paneth & other intestinal epithelial cells; controls resistance to gut commensals
<b>ATG 16</b>	Inflammatory Bowel Disease	Involved in <b>autophagy<sup>a</sup></b> ; Role in defense against microbes & maintenance of epithelial barrier function
<b>IRF5, IRH1</b>	Systemic Lupus Erythematosus	Role in <b>type 1 IFN<sup>a</sup></b> production, which is involved in the pathogenesis of SLE

### Role of Infections

- Infections may **up-regulate** the expression of **costimulators on APCs**
- **Molecular mimicry<sup>a</sup>**: Some infections may express antigens that have the same amino acid sequences as self antigens. Immune responses against the microbial antigens may result in the activation of self-reactive lymphocytes.

## AUTOIMMUNE DISEASES

### Systemic Lupus Erythematosus (SLE)

- Caused by **deposition of immune complexes** and **binding of antibodies** to various cells and tissues.
- **Characteristic lesions due to immune complex deposition in**: blood vessels, kidneys, connective tissue, skin
- **Blood Vessels**: Acute necrotizing vasculitis with **fibrinoid deposits<sup>a</sup>**

### High Yield Facts



#### Renal Involvement in SLE

Classification of Lupus Nephritis (International Society of Nephrology)

- **Class I** : Minimal Mesangial<sup>a</sup>
- **Class II** : Mesangial Proliferative
- **Class III** : Focal Lupus Nephritis<sup>a</sup>
- **Class IV** : Diffuse Lupus Nephritis<sup>a</sup>
- **Class V** : Membranous Lupus Nephritis<sup>a</sup>
- **Class VI** : Advanced Sclerotic Lupus Nephritis



### High Yield Facts

#### Drug Induced Lupus

- Less female predilection than SLE
- Anti histone Ab: Common
- Rarely involves kidney & brain
- Anti-ds DNA ±

R10<sup>th</sup>

**Latest Update**

### Systemic lupus international collaborating clinic criteria for classification of systemic lupus erythematosus

Clinical manifestations	Immunologic manifestations
<ul style="list-style-type: none"> <li>• Skin, acute, subacute cutaneous LE and chronic cutaneous LE</li> <li>• Oral ulcers</li> <li>• Alopecia</li> <li>• Synovitis (nonerosive)</li> <li>• Renal: Prot/Cr <math>\geq 0.5</math>, RBC casts and biopsy*</li> <li>• Neurologic: Seizures, psychosis, mononeuritis, myelitis</li> <li>• Hemolytic anemia: Leukopenia (<math>&lt; 4000</math>) or, lymphopenia (<math>&lt; 1000</math>) and thrombocytopenia (<math>&lt; 100,000</math>)</li> </ul>	<ul style="list-style-type: none"> <li>• ANA &gt; reference negative value</li> <li>• Anti-ds DNA</li> <li>• Anti-Sm</li> <li>• Antiphospholipid</li> <li>• Low serum complement</li> <li>• Positive direct Coombs test</li> </ul>

\*Renal biopsy read as systemic lupus qualities for classification as SLE even if none of the other above features are present

Interpretation: Presence of any 4 criteria (must have at least 1 in each category) qualifies patient to be classified as having SLE with 93% specificity and 92% sensitivity.



## Autoantibodies in Systemic Lupus Erythematosus (SLE)<sup>Q</sup>

Antibody	Antigen Recognized	Clinical Utility
<b>Antinuclear antibodies<sup>Q</sup></b>	Multiple nuclear	<b>Best screening test/ Most sensitive<sup>Q</sup></b>
<b>Anti-dsDNA<sup>Q</sup></b>	DNA (double-stranded)	<b>Specific,<sup>Q</sup> correlate with disease activity, nephritis, vasculitis</b>
<b>Anti-Sm<sup>Q</sup></b>	U1 RNA	<b>Most Specific for SLE<sup>Q</sup></b>
<b>Anti-RNP</b>	U1 RNA	Not specific for SLE
<b>Anti-Ro/La (SS-A)<sup>Q</sup></b>	hY RNA,	Sicca syndrome, neonatal lupus with congenital heart block <sup>Q</sup>
<b>Anti-histone<sup>Q</sup></b>	Histones	<b>Drug-induced lupus<sup>Q</sup></b>



### High Yield Facts

- **50%** patients with SLE have clinically significant **renal involvement<sup>Q</sup>**
- **Class I Lupus Nephritis is least common<sup>Q</sup> and class IV is the most common pattern<sup>Q</sup>**
- **"Wire loop lesions"<sup>Q</sup> are characteristically seen in Class IV<sup>Q</sup> Lupus Nephritis** due to **sub-endothelial** immune complex deposits, on **light microscopy**-Also seen in **Class III/V Lupus nephritis<sup>Q</sup>**

## Sjogren Syndrome

- **Characterized by:** Dry eyes (**keratoconjunctivitis sicca<sup>Q</sup>**) and **dry mouth (xerostomia)<sup>Q</sup>** due to immunologically mediated destruction of the lacrimal and salivary glands.
- **Two forms**
  - **Primary form (sicca syndrome)<sup>Q</sup>**: as an isolated disorder
  - **Secondary form:** in association with other autoimmune disease (e.g. RA, SLE, scleroderma)
- **Autoantibodies**
  - **Anti SS-A (Ro) and SS-B (La)<sup>Q</sup>**: Most important, present in 90% patients
  - High titers of Anti SS-A → more likely to have **early disease onset, longer disease duration, and extraglandular manifestations** (e.g. cutaneous vasculitis and nephritis)
- **Morphology:**
  - **Main targets:** Lacrimal and salivary glands<sup>Q</sup>
  - **Earliest histologic finding:** Periductal and perivascular **lymphocytic infiltration**
  - **Lymphoid follicles** with **germinal centers** in larger salivary glands may be seen
  - **Late changes:** **atrophy** of acini, fibrosis, and hyalinization;
  - At high risk for development of **B-cell Marginal zone lymphoma<sup>Q</sup>**
- **Diagnosis: Biopsy of the lip** (to examine minor salivary glands) is required for diagnosis

## Scleroderma (Systemic Sclerosis) (SSc)

<b>Characterized by</b>	<ul style="list-style-type: none"> <li>• <b>Chronic inflammation,<sup>Q</sup></b> as a result of autoimmunity</li> <li>• <b>Widespread damage</b> to small blood vessels</li> <li>• <b>Progressive interstitial &amp; perivascular fibrosis<sup>Q</sup></b> in skin &amp; multiple organs.</li> </ul>	
<b>2 major categories</b>	<b>Limited Cutaneous SSc</b>	<b>Diffuse Cutaneous SSc</b>
<b>Skin involvement</b>	Limited to fingers, forearms, face <sup>Q</sup>	Diffuse: fingers, extremities, face, trunk; <sup>Q</sup>
<b>Onset &amp; course</b>	Indolent & slow	Rapid
<b>Visceral involvement</b>	Late	Early
<b>Auto-antibodies seen</b>	Anti-centromere <sup>Q</sup>	Anti-topoisomerase I (Scl-70), <sup>Q</sup> anti-RNA polymerase III <sup>Q</sup>
<b>Crest syndrome</b>	More common <sup>Q</sup>	Less common <sup>Q</sup>



### High Yield Facts

- **CREST syndrome:** Calcinosis, Raynaud's phenomenon, Esophageal dysmotility, Sclerodactyly & Telangiectasia<sup>Q</sup>
- **Mikulicz syndrome** refers to lacrimal & salivary gland enlargement from any cause, including sarcoidosis, lymphoma & other tumors<sup>Q</sup>
- **Most common organ involved in Scleroderma is Skin<sup>Q</sup>**
- **Antitopoisomerase I (Scl-70)<sup>Q</sup> is most specific<sup>Q</sup> for Scleroderma**
- **ANA positivity is seen in virtually all patients with SSc (Systemic sclerosis)**

## Mixed Connective Tissue Disease

- Clinical features are a mixture of the features of **SLE, systemic sclerosis, and polymyositis<sup>Q</sup>**
- Elevated **anti-U1 ribonucleoprotein Ab<sup>Q</sup>**
- Presentation: Synovitis of fingers, Raynaud phenomenon, mild myositis, renal involvement

## REJECTION OF TISSUE TRANSPLANTS



<b>Types of grafts<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• <b>Autografts<sup>Q</sup>:</b> Transplant of individual own organ</li> <li>• <b>Isograft:</b> Graft from identical twin</li> <li>• <b>Allografts<sup>Q</sup>:</b> Between individuals of the same species</li> <li>• <b>Xenografts<sup>Q</sup>:</b> Grafts from one species to another species</li> </ul>
<b>Types of Rejection Pathway</b>	<ul style="list-style-type: none"> <li>• <b>Direct pathway:</b> <b>CD4 &amp; CD8-T cells<sup>Q</sup></b> of the transplant recipient recognize allogenic (donor) MHC molecules on the surface of APCs in the graft.</li> <li>• <b>Indirect pathway:</b> Recipient <b>CD4-T lymphocytes<sup>Q</sup></b> recognize <b>MHC antigens</b> of the graft donor after they are presented by the recipient's own APCs.</li> </ul>



## High Yield Facts

- In kidney transplants, polymorphic HLA alleles are **at least 50% matched** (HLA-A, -B & DR)<sup>Q</sup>
- HLA matching is **NOT required for Cornea transplant**<sup>Q</sup>
- HLA matching is usually **NOT done** for transplants of liver, heart, and lungs, because other considerations, such as **anatomic compatibility, severity** of the underlying illness, and the **need to minimize the time** of organ storage, **override the potential benefits** of HLA matching
- “000 mismatch” means **no mismatch** in HLA-A, B, D
- Most important HLA is HLA DRB1 for transplant survival

## Types of Renal Graft Rejection: (On the Basis of the Morphology & Underlying Mechanism)

Features	Hyperacute rejection	Acute Rejection	Chronic Rejection
<b>Time to occur</b>	Minutes to hours <sup>Q</sup>	Days (months or even years) <sup>Q</sup>	Months or even years <sup>Q</sup>
<b>Mechanism (Type of hypersensitivity)</b>	<ul style="list-style-type: none"> <li>• <b>Type 2 HSN</b>:<sup>Q</sup> due to preformed antibodies</li> <li>• Type 3 HSN</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Type 4 HSN</b>-Cellular rejection<sup>Q</sup></li> <li>• Type 2-mediated by anti-donor antibodies</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Type 4-HSN</b> Cellular rejection<sup>Q</sup></li> </ul>
<b>Histology</b>	<ul style="list-style-type: none"> <li>• <b>Neutrophilic infiltration</b><sup>Q</sup> &amp; thrombotic occlusion of capillaries</li> <li>• <b>Fibrinoid necrosis</b> occurs in arterial walls.</li> </ul>  <p>Showing fibrin thrombi in glomerulus along with ischemic necrosis</p>	<p>In <b>cell-mediated rejection</b>:</p> <ul style="list-style-type: none"> <li>• Tubulo-interstitial pattern (<b>tubulitis</b>)</li> <li>• Vascular pattern (<b>endotheliitis</b>)</li> </ul> <p>In <b>antibody mediated rejection</b>:</p> <p>Inflammation of glomeruli &amp; peritubular capillaries with <b>deposition of C4d</b><sup>Q</sup></p>  <p>Showing C<sub>4</sub>d deposition in peritubular capillaries</p>	<p><b>Intimal thickening</b> with inflammation; Glomerulopathy; <b>peritubular capillaritis</b><sup>Q</sup> <b>Interstitial fibrosis</b><sup>Q</sup> &amp; tubular atrophy</p>

## IMMUNODEFICIENCY SYNDROMES

### Types

- **Primary (or congenital)** : Genetically determined<sup>Q</sup>

- **Secondary (or acquired)**: Complications of **cancers, infections, malnutrition**, or side effects of immunosuppression, **irradiation**, or **chemotherapy** for cancer and other diseases<sup>Q</sup>

### Primary Immunodeficiency Diseases


#### Defects in Innate Immunity

Disease	Defect
<b>Defects in Leukocyte function</b>	
<b>Leukocyte adhesion deficiency 1</b>	Defective WBC adhesion because of <b>mutations in <math>\beta</math> chain of CD11/CD18 integrins</b> <sup>Q</sup>
<b>Leukocyte adhesion deficiency 2</b>	Defective WBC adhesion due to <b>mutations in fucosyltransferase</b> <sup>Q</sup> required for synthesis of sialylated oligosaccharide (receptor for selectins)
<b>Chediak – Higashi syndrome</b>	Decreased <b>leukocyte functions</b> due to mutations of <b>protein involved in lysosomal membrane traffic</b> <sup>Q</sup>
<b>Chronic granulomatous disease</b>	Decreased <b>oxidative burst</b> <sup>Q</sup> Phagocyte oxidase (membrane component) Phagocyte oxidase (cytoplasmic components)
<ul style="list-style-type: none"> <li>• X-linked</li> <li>• Autosomal recessive</li> </ul>	
<b>Myeloperoxidase deficiency</b>	Decreased microbial killing because of <b>defective MPO-H<sub>2</sub>O<sub>2</sub> system</b> <sup>Q</sup>



## Complement Deficiencies & Associated Diseases

Component	Associated Diseases
<b>Classic Pathway</b>	
Clq,Clr,Cls,C4,C2	Immune-complex syndromes <sup>a</sup> , pyogenic infections
C1 esterase Inhibitor	Hereditary angioneurotic edema <sup>a</sup>
<b>C3 and Alternative Pathway C3</b>	
C3	Immune-complex syndromes, pyogenic infections
D	Pyogenic infections <sup>a</sup>
Properdin	<i>Neisseria</i> infections <sup>a</sup>
I	Pyogenic infections <sup>a</sup>
H	Hemolytic uremic syndrome <sup>a</sup>
<b>Membrane Attack Complex</b>	
C5, C6, C7, C8	Recurrent <i>Neisseria</i> infections, <sup>a</sup> immune-complex disease
C9	Rare <i>Neisseria</i> infections







## Defects in Adaptive Immune System: (Primary Immunodeficiency Disorders)

Disease	Primary Defect	Secondary Defect	Infections/Features	Morphology
B cell (Humoral) defects				
Bruton's X-Linked Agammaglobulinemia	Bruton tyrosine kinase(Btk); Xq21.22 gene mutations <sup>Q</sup>	<ul style="list-style-type: none"><li>Failure of pro B-cells to mature</li><li>Plasma cells absent<sup>Q</sup></li><li>T-cells normal</li></ul>	H. influenzae, S. pneumoniae, S. aureus	Underdeveloped: <ul style="list-style-type: none"><li>Germinal centers of L. nodes<sup>Q</sup></li><li>Peyer's patches of appendix<sup>Q</sup></li><li>Tonsils<sup>Q</sup></li></ul>
Hyper-IgM Syndrome	X-linked: Mutations in CD40L (Xq26) <sup>Q</sup> Autosomal recessive: Mutations in CD40 & AID (activation-induced cytidine deaminase)	<ul style="list-style-type: none"><li>Defect in Ig class switching<sup>Q</sup> and affinity maturation</li><li>Inability to produce IgG, IgA, and IgE Antibodies</li></ul>	<ul style="list-style-type: none"><li>Recurrent</li><li>Pyogenic infections<sup>Q</sup></li><li>Pneumocystis jiroveci<sup>Q</sup></li></ul>	Normal
Isolated IgA Deficiency	Impaired differentiation of naive B cells to IgA-producing plasma cells	<ul style="list-style-type: none"><li>Deficient IgA</li><li>Concomitant defect in IgG2 and IgG4<sup>Q</sup></li></ul>	<ul style="list-style-type: none"><li>Infections</li><li>Respiratory tract allergy<sup>Q</sup></li><li>Autoimmune: SLE and RA<sup>Q</sup></li></ul>	Normal
Common Variable Immunodeficiency	Defect in receptor for a cytokine called BAFF <sup>Q</sup>	Defect in survival and differentiation of B cells	<ul style="list-style-type: none"><li>Recurrent sino-pulmonary &amp; pyogenic infections</li><li>Herpes virus</li><li>Enteroviral meningoencephalitis</li><li>G. lamblia diarrhea<sup>Q</sup></li></ul>	B-cell areas of the lymphoid tissues are hyperplastic <sup>Q</sup>
	Defect in ICOS <sup>Q</sup> (inducible costimulator)	Defect in T-cell activation & interactions b/w T & B cells <sup>Q</sup>		
Predominant T cell defect				
DiGeorge Syndrome (Thymic Hypoplasia) <sup>Q</sup>	Familial cases: failure of development of 3 <sup>rd</sup> & 4 <sup>th</sup> pharyngeal pouches <sup>Q</sup> Nonfamilial cases: 22q11 deletion <sup>Q</sup> syndrome (50%)	<ul style="list-style-type: none"><li>T-cell defect</li><li>Tetany<sup>Q</sup></li><li>Congenital defects of heart and great vessels</li><li>Mucocutaneous Candidiasis</li></ul>	<ul style="list-style-type: none"><li>Fungal</li><li>Viral infections</li></ul>	Depleted: <sup>Q</sup> <ul style="list-style-type: none"><li>Paracortical areas of L. nodes<sup>Q</sup></li><li>Periarteriolar sheaths of spleen</li></ul>
Bare lymphocyte syndrome	Defect in class II MHC gene expression.	Abnormal development of CD4+ T cells.		Normal
Combined T and B cell defect				
Severe Combined Immunodeficiency	X-linked:γ-chain (γc) of cytokine receptor mutations	Defect in signaling of IL-2, 4, 7, 9, 11, 15 & 21	<ul style="list-style-type: none"><li>Candida<sup>Q</sup></li><li>Pneumocystis<sup>Q</sup></li><li>Pseudomonas</li><li>CMV</li><li>Varicella</li></ul>	Thymus is small <sup>Q</sup> and devoid of lymphoid cells
	Autosomal recessive: adenosine deaminase (ADA) deficiency <sup>Q</sup>	Accumulation of deoxyadenosine-toxic to rapidly dividing WBCs		
Ataxia Telangiectasia	Autosomal-recessive mutation in ATM gene <sup>Q</sup> (ataxia telangiectasia mutated) on chr 11	<ul style="list-style-type: none"><li>Combination of T &amp; B cell defects</li><li>Defective isotype switching</li><li>Low IgA &amp; IgG-2</li></ul>	<ul style="list-style-type: none"><li>Ataxia</li><li>Vascular telangiectases<sup>Q</sup></li><li>Neurologic deficits<sup>Q</sup></li><li>Tumors<sup>Q</sup></li><li>Immunodeficiency</li></ul>	Normal

Contd...



Disease	Primary Defect	Secondary Defect	Infections/Features	Morphology
<b>Hyper IgE syndrome</b>	Autosomal dominant: <sup>Q</sup> mutation in <b>STAT3</b> gene: Defective signaling pathways	<ul style="list-style-type: none"> <li>• Elevated serum IgE levels</li> <li>• Defective TH17 effector responses<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Recurrent skin abscesses<sup>Q</sup></li> <li>• Lung infections &amp; pneumatoceles<sup>Q</sup></li> <li>• Pyogenic bacteria &amp; fungi</li> <li>• Dysmorphism</li> <li>• Defective teeth</li> <li>• Hyperextensibility</li> <li>• Scoliosis, Osteoporosis.</li> </ul>	Normal
<b>X-linked Lymphoproliferative disease</b>	Mutations in <b>SLAM-associated protein (SAP)</b> <sup>Q</sup>	Activation of NK, T & B cells, including Signalling Lymphocyte Activation Molecule (SLAM)	Epstein-Barr virus (EBV)	
<b>Wiskott-Aldrich Syndrome</b>	X-linked: Mutations in Wiskott-Aldrich syndrome protein (WASP), Xp11.23 <sup>Q</sup>	<ul style="list-style-type: none"> <li>• CD8+T-cell deficiency</li> <li>• IgM-low<sup>Q</sup></li> <li>• IgA &amp; IgE-↑ed<sup>Q</sup></li> <li>• IgG-normal<sup>Q</sup></li> </ul>	Thrombocytopenia <sup>Q</sup> Eczema <sup>Q</sup> Recurrent infection <sup>Q</sup>	Normal

Immune Regulatory Defects	
Innate immunity	<ul style="list-style-type: none"> <li>• Autoinflammatory syndromes (outside the scope of this chapter)</li> <li>• Severe colitis</li> </ul>
Adaptive immunity	<ul style="list-style-type: none"> <li>• Hemophagocytic lymphohistiocytosis (HLH)</li> <li>• Autoimmune lymphoproliferation syndrome (ALPS)<sup>Q</sup></li> <li>• Autoimmunity and inflammatory diseases (IPEX, APECED)</li> </ul>



### High Yield Facts


#### SCID

- Most severe immunodeficiency in children
- Presents soon after Birth
- Graft rejection not seen
- Risk of severe GVHD with non-irradiated blood products

### Hemophagocytic Lymphohistiocytosis (HLH)

- **Characterized by:**
  - Activation of CD8+ T lymphocytes and macrophages leading to organ damage
  - Impaired T and NK lymphocyte cytotoxicity.
- **Triggered by: Viruses (Most commonly: EBV)Q**
- **Clinical features**
  - Fever
  - Edema
  - Hepatosplenomegaly
  - Neurologic diseases
  - **Blood cytopenia**
  - Hypofibrinogenemia
  - Increased liver enzymes
  - **High triglyceride levels**
- **Lab findings**
  - Elevated markers of T cell activation
  - Hemophagocytic features in the bone marrow or cerebrospinal fluid
  - **Functional assays** of post-activation cytotoxic granule exocytosis (CD107 fluorescence at cell membrane) can suggest **genetically** determined HLH.
- **Subsets of HLH**
  - **Familial HLH (A.R)**
    - Perforin deficiency<sup>Q</sup>
    - Munc13-4 deficiency

- Syntaxin 11 deficiency
- HLH with partial **albinism**
- **Chediak-Higashi syndrome**<sup>Q</sup>
- **Griscelli syndrome**<sup>Q</sup>
- Hermansky Pudlak syndrome type II.
- **X-linked** proliferative syndrome
- Following EBV infection
- **Main Defects include:**
  - SH2DIA gene
  - Low 2B4-mediated NK cell cytotoxicity
  - **Impaired differentiation of NK T cells**
  - Defective antigen-induced T cell death
  - Defective T cell helper activity for B cells



Hemophagocytosis in bone marrow aspirate



- Nitrobluetetrazolium test<sup>Q</sup> is used for Phagocytes
- The commonest Primary immunodeficiency is Isolated IgA immunodeficiency<sup>Q</sup>
- In agammaglobulinemia there is loss of germinal centre in lymph node<sup>Q</sup>
- Commonest fungal infection in neutropenia is Candida<sup>Q</sup>
- HIV was isolated in 1983
- HIV-2: 1<sup>st</sup> case in 1987

## High Yield Facts

## Secondary (Acquired) Immunodeficiencies

Cause	Mechanism
Human immunodeficiency virus (HIV) infection <sup>Q</sup>	Depletion of CD4+ helper T cells
Irradiation & chemotherapy <sup>Q</sup> treatments for cancer	Decreased bone marrow precursors <sup>Q</sup> for all leukocytes
Involvement of bone marrow by cancers <sup>Q</sup> (Metastases, Leukemias)	Reduced site of leukocyte development
Protein- calorie malnutrition	Metabolic derangements inhibit lymphocyte maturation & function
Removal of spleen	Decreased phagocytosis of microbes

## Acquired Immunodeficiency Syndrome (AIDS)

Causative Organism: HIV (Human Immunodeficiency Virus)

<b>Family</b>	Non-transforming human retrovirus of lentivirus family <sup>Q</sup>	<p>Structure of HIV virus</p>
<b>Classification</b>	Two types: HIV-1 & HIV-2	
<b>Groups of HIV1</b>	3 Groups: <ul style="list-style-type: none"> <li>• Major group (HIV-1 M)-Most common world wide<sup>Q</sup></li> <li>• Outlier group (HIV-1 O)</li> <li>• New/Neither M nor O group (HIV-1 N)</li> </ul>	
<b>Subtypes/clades of HIV-1 M group</b>	HIV-1 M group is further classified into subtypes or clades: A through H, J and K <ul style="list-style-type: none"> <li>• HIV-1 group M subtypes C predominant in India, fastest spreading<sup>Q</sup></li> <li>• HIV-1 group M subtypes B predominant in U.S</li> </ul>	
<b>Structure of HIV</b>	<p>Viral core is surrounded by a matrix protein (p17)<sup>Q</sup> followed by host cell derived lipid envelope<sup>Q</sup></p> <p>A. Core protein</p> <ul style="list-style-type: none"> <li>• Major Capsid protein (p24) (most abundant)<sup>Q</sup></li> <li>• Nucleocapsid protein (p7/p9)</li> <li>• Two copies of single stranded RNA</li> <li>• Viral enzymes:<sup>Q</sup> Reverse transcriptase (p51), Integrase (p31) &amp; Protease (p9)</li> </ul> <p>B. Matrix (p17)<sup>Q</sup></p> <p>C. Lipid Envelope with gp 120 and gp 41<sup>Q</sup></p> <ul style="list-style-type: none"> <li>• gp-120: Protrudes out on the surface of the virus<sup>Q</sup> &amp; binds to CD4 molecule &amp; co-receptors<sup>Q</sup></li> <li>• gp-41: Embedded in the lipid matrix &amp; helps in fusion with target cells<sup>Q</sup></li> </ul>	



- Difference between HIV-1 and 2:
- HIV-2 Infection:
- Not transmitted as efficiently as HIV-1<sup>Q</sup>
  - Progression to AIDS takes longer
  - Slower rate of CD4 cell decline and viral replication
  - Rarely causes vertical transmission

## High Yield Facts



## Genetics of HIV

### Structural Genes



### High Yield Facts

#### AIDS defining illness CDC stage 3 Includes:

- Esophageal candidiasis
- Extrapul cryptococcosis
- CMV Retinitis
- Diss. Histoplasmosis 1B (any site)
- MAC
- Pneumocystis jiroveci
- Toxoplasmosis of brain
- Invasive cervical Ca
- Kaposi sarcoma
- Burkitt's lymphoma
- Primary CNS lymphoma
- Immunoblastic lymphoma

Infection

Neoplasms

## Regulatory/Accessory Genes

### Pathogenesis of HIV Infection and AIDS (Steps)

- **gp120 binds to host CD4 receptors** on <sup>o</sup>T-cells, Macrophages, Dendritic cells & other Antigen presenting cells
- Binding to **co-receptor molecules CCR-5/CXCR4<sup>o</sup>**
  - **R5 strains** use CCR5: **M-tropic**, preferentially infects monocyte/macrophages (**MC in EARLY infection**)<sup>o</sup>
  - **X4 strains** use CXCR4: **T-tropic**, preferentially infects T-cells (**MC in LATE infection**)
  - R5X4 strains: Dual tropic
- Exposure of **gp-41** leading to **fusion of the virus with the host cell**
- **Uncoating** of viral genome
- **Reverse transcription** of ssRNA to dsDNA
- ds DNA along with enzymes (**Pre-integration complex**)<sup>o</sup> transported to nucleus
- **Viral DNA gets integrated** with host cell DNA (**provirus**)<sup>o</sup>
- **Direct viral protein synthesis** (transcription & translation)
- **Virus assembly and budding**



### High Yield Facts


#### Modes of Transmission of HIV

- **Sexual** contact – heterosexual/homosexual<sup>o</sup>
- Receipt of **infected blood** and its products<sup>o</sup>
- **Mother to child (vertical)** transmission<sup>o</sup>
- **Percutaneous/mucosal exposures** to blood and body fluids of HIV infected
- Accidental – **occupational** (health care workers)
- I/V drug abuse



### High Yield Facts

- **Most common fungal** infection in patients with AIDS: **Candida<sup>o</sup>**
- **Most common sites of Candida** infection in HIV: **oral cavity, vagina, and esophagus<sup>o</sup>**
- In asymptomatic HIV, **oral candidiasis** is a sign of **immunologic decompensation<sup>o</sup>**
- **Invasive candidiasis** is **INFREQUENT** in patients with AIDS<sup>o</sup>
- Majority (50%) of **CNS mass lesions** is caused by: **Toxoplasma gondii<sup>o</sup>**



### High Yield Facts

- **Most common malignancy** in patients with AIDS: **Kaposi Sarcoma<sup>o</sup>**
- **HHV-8 (Kaposi sarcoma virus)** causes **Kaposi sarcoma, primary effusion lymphoma and multicentric Castleman's disease** in HIV<sup>o</sup>
- **Most common hematologic abnormality** in HIV-infected patients: **Anemia<sup>o</sup>**
- **Most common lymphoma** seen in HIV: **Immunoblastic lymphoma (DLBCL)<sup>o</sup> > Burkitt's lymphoma > primary CNS lymphoma**
- **Most common Kidney biopsy finding** in HIV is : **Collapsing variant of FSGS<sup>o</sup>**
- CD4 T cells <200 cells/nm<sup>3</sup> predispose to **P. jirovecii**
- CD4 T cells <100 cells/nm<sup>3</sup>—**cryptococcus**
- CD4 T cells <50 cells/nm<sup>3</sup>—**CMV, MAC., IRIS**
- CD4 T cells >300 cells/nm<sup>3</sup>—**TB.**





### Natural History of HIV Infection

- **Acute retroviral syndrome**- 3-6 wks after infection, usually resolves spontaneously<sup>Q</sup>
- **Middle, chronic phase (clinical latency)**<sup>Q</sup>
- **Clinical AIDS**

### Immune Reconstitution Inflammatory Syndrome (IRIS)

- **Paradoxical worsening**<sup>Q</sup> of clinical condition is seen following the initiation of ART (antiretroviral therapy)
- Occurs weeks to months following the initiation of antiretroviral therapy<sup>Q</sup>

- Is **most common** in patients starting therapy with a **CD4+ T cell count < 50/uL**<sup>Q</sup>
- Is frequently seen in the setting of **tuberculosis**<sup>Q</sup>
- Can be fatal

### Morphology-Lymph node Biopsy in HIV infection


- **Early HIV:** Marked **hyperplasia of follicles** with attenuated mantle zones<sup>Q</sup>
- **Late HIV:** **Atrophic Lymph nodes**<sup>Q</sup> with depletion of lymphocytes and disruption of follicular dendritic cells

## AMYLOIDOSIS

Definition	Amyloidosis is a pathological proteinaceous substance deposited between cells <sup>Q</sup> in various tissues and organs of the body in a variety of clinical settings.																				
Sites of biopsy	<ul style="list-style-type: none"><li>• <b>Rectal biopsy</b>-Most specific and best site</li><li>• <b>Abdominal fat aspirate:</b> Most sensitive &amp; easy site</li><li>• Gingival biopsy</li><li>• Organ specific biopsy for localized amyloidosis</li></ul>																				
Diagnosis	<ul style="list-style-type: none"><li>• <b>Grossly:</b> Affected organ is <b>enlarged</b> and grey with a waxy, firm consistency</li><li>• <b>Histologically:</b> <b>Extracellular pink</b> homogenous deposit</li><li>• On special staining:</li></ul> <table><tr><th>Technique</th><th>Stain</th><th>Feature</th></tr><tr><td>Light microscope</td><td>Congo red</td><td>Salmon-red color amyloid deposits<sup>Q</sup></td></tr><tr><td>Polarized light</td><td>Congo red</td><td>Apple-green birefringence<sup>Q</sup></td></tr><tr><td>Fluorescent staining</td><td>Thioflavin T</td><td>Yellow color<sup>Q</sup></td></tr><tr><td>Electron microscope</td><td>—</td><td>Non-branching fibrils<sup>Q</sup>, 7.5 to 10 nm<sup>Q</sup> diameter</td></tr><tr><td>X-ray crystallography &amp; infrared spectroscopy</td><td>—</td><td>Characteristic <b>cross-β-pleated sheet</b><sup>Q</sup></td></tr></table>			Technique	Stain	Feature	Light microscope	Congo red	Salmon-red color amyloid deposits <sup>Q</sup>	Polarized light	Congo red	Apple-green birefringence <sup>Q</sup>	Fluorescent staining	Thioflavin T	Yellow color <sup>Q</sup>	Electron microscope	—	Non-branching fibrils <sup>Q</sup> , 7.5 to 10 nm <sup>Q</sup> diameter	X-ray crystallography & infrared spectroscopy	—	Characteristic <b>cross-β-pleated sheet</b> <sup>Q</sup>
Technique	Stain	Feature																			
Light microscope	Congo red	Salmon-red color amyloid deposits <sup>Q</sup>																			
Polarized light	Congo red	Apple-green birefringence <sup>Q</sup>																			
Fluorescent staining	Thioflavin T	Yellow color <sup>Q</sup>																			
Electron microscope	—	Non-branching fibrils <sup>Q</sup> , 7.5 to 10 nm <sup>Q</sup> diameter																			
X-ray crystallography & infrared spectroscopy	—	Characteristic <b>cross-β-pleated sheet</b> <sup>Q</sup>																			


### Classification of Amyloidosis

Clinicopathologic category	Associated Diseases	Fibril protein	Precursor protein
<b>Systemic (Generalized) Amyloidosis</b>			
• <b>Primary amyloidosis</b>	Multiple myeloma	<b>AL</b>	Immunoglobulin light chain (λ)
• <b>Secondary amyloidosis</b>	Chronic inflammatory conditions	AA	SAA
• <b>Hemodialysis- associated amyloidosis</b>	Chronic renal failure	A β <sub>2</sub> m	β <sub>2</sub> -microglobulin
<b>Localized Amyloidosis</b>			
• <b>Senile cerebral</b>	Alzheimer's disease	A β	APP
• <b>Medullary carcinoma of thyroid</b>		A Cal	Calcitonin
• <b>Islets of Langerhans</b>		AIAPP	Islet amyloid peptide
• <b>Isolated atrial amyloidosis</b>		AANF	Atrial natriuretic factor
<b>Hereditary Amyloidosis</b>			
• <b>Familial Mediterranean fever</b>		AA	SAA
• <b>Familial amyloidotic neuropathies</b>		ATTR	Transthyretin
• <b>Systemic senile amyloidosis</b>		ATTR	Transthyretin



## AA (Amyloid Associate Protein)

- It is unique **non-immunoglobulin protein** synthesized by reticuloendothelial cells of liver<sup>Q</sup>.
- Associated with **secondary amyloidosis & reactive systemic amyloidosis**<sup>Q</sup>.



- Seen in:

Connective tissue disorders	Nonimmune derived tumors
<ul style="list-style-type: none"> <li>Rheumatoid Arthritis (MC)<sup>Q</sup></li> <li>Ankylosing spondylitis<sup>Q</sup></li> <li>Primary biliary cirrhosis<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Renal cell carcinoma<sup>Q</sup></li> <li>Hodgkin's lymphoma<sup>Q</sup></li> </ul>

## Individual organ Amyloidosis



Kidney	1st deposits in mesangium
Spleen	Sago → deposit in follicles Lardaceous-deposit in red pulp
Heart	1st deposit in subendocardial areas of atrium
Liver	1st deposit in space of Disse
Blood	Deposit inactivate factor X

## Hemodialysis-Associated Amyloidosis

- Patients on long-term dialysis for renal failure develop amyloidosis due to deposition of  $\beta$ 2-microglobulin
- This protein is present in high concentrations in the serum of persons with renal disease and is retained in the circulation because it **cannot be filtered through dialysis membranes**.
- Patients often present with **carpal tunnel syndrome because of  $\beta$ 2-microglobulin deposition**.
- In some patients on long-term dialysis, amyloid deposits in the **synovium, joints, or tendon sheaths**.

## Involvement of Heart in Amyloidosis

- Heart is the second most commonly affected organ**, in 50% of patients, in primary amyloidosis
- Cardiac failure is the leading cause of mortality in Amyloidosis**.
- Early on, the electrocardiogram may show low voltage in the limb leads, with a **pseudo-infarct pattern**.
- Eventually, the echocardiogram will display **concentrically thickened ventricles and diastolic dysfunction**, leading to a **restrictive cardiomyopathy**; systolic function is preserved until late in the disease.



## Amyloidosis of Kidney

- It is the **most common (affected in 70%) and potentially the most serious** form of organ involvement.
- Amyloid is deposited primarily in glomeruli, it starts first in mesangium.
- Interstitial peritubular tortuous arteries and arterioles are affected.
- Usually manifests as **proteinuria, often in nephrotic range** and associated with significant hypoalbuminemia, secondary hypercholesterolemia, and edema or anasarca.
- Tubular rather than glomerular deposition of amyloid can produce **azotemia** without significant proteinuria



## High Yield Facts

Scintigraphy with radiolabeled serum amyloid P is rapid and specific test. It can tell extent of amyloidosis



## BIOLOGICAL PROPERTIES OF DIFFERENT ANTIBODY CLASSES

<b>IgG</b>	<ul style="list-style-type: none"> <li>• <b>Most abundant</b> of all the Ig classes</li> <li>• <b>Longest half-life</b> of all the Ig classes</li> <li>• Agglutinates particulate antigens &amp; Precipitates soluble antigens</li> <li>• The only antibody class that <b>crosses the placenta</b></li> <li>• IgG bound to cells or aggregated in antigen-antibody complexes can <b>activate complement</b>.</li> </ul>	<ul style="list-style-type: none"> <li>• IgG is good at <b>neutralizing toxins</b></li> <li>• IgG can interfere with bacterial virulence features such as motility and adherence to tissues</li> <li>• IgG can <b>block virus ability to adhere to target cells</b></li> <li>• <b>IgG is the main antibody found in secondary immune response</b></li> </ul>
<b>IgM</b>	<ul style="list-style-type: none"> <li>• IgM monomer is the <b>B-cell receptor</b> of the mature B-cell</li> <li>• <b>First antibody made in a primary immune response</b>, 7-10 days after initial exposure</li> <li>• IgM is a very good agglutinating antibody (<b>Natural isohemagglutinins are also of IgM class</b>)</li> </ul>	<ul style="list-style-type: none"> <li>• IgM is the <b>first antibody made in life</b> (about 5 months in utero)</li> <li>• IgM is the antibody made to T-independent antigens polysaccharides</li> <li>• IgM attached to antigen is a very <b>good activator of complement</b></li> <li>• IgM <b>does not cross</b> the placenta</li> </ul>
<b>IgA</b>	<ul style="list-style-type: none"> <li>• IgA is the <b>secretory antibody</b> found in all secretions including colostrum and milk.</li> <li>• IgA is important for protection against <b>respiratory and gastrointestinal</b> infectious agents.</li> <li>• IgA is important for <b>passively acquired immunity</b> of nursing baby.</li> </ul>	<ul style="list-style-type: none"> <li>• IgA makes lysozyme work better especially against gram-negative bacteria</li> <li>• IgA <b>neutralizes virus</b></li> <li>• IgA is transported across membranes via the poly-Ig receptor</li> </ul>
<b>IgE</b>	<ul style="list-style-type: none"> <li>• Homocytotropic antibody</li> <li>• IgE is made in response to <b>parasitic worms</b>.</li> </ul>	<ul style="list-style-type: none"> <li>• IgE is the antibody that <b>triggers allergy</b></li> <li>• IgE binds to FcεRI receptors on the surface of mast cells</li> </ul>

R10<sup>th</sup>

### Latest Update

#### IgG4-Related Disease (IgG4-RD)

- A **newly recognized** constellation of disorders characterized by **tissue infiltrates by IgG4 antibody-producing plasma cells** and lymphocytes, particularly T cells, **storiform fibrosis**, **obliterative phlebitis**, and usually **increased serum IgG4**.
- Spectrum includes:

- Mikulicz syndrome (enlargement and fibrosis of salivary and lacrimal glands)
- Riedel thyroiditis
- Idiopathic retroperitoneal fibrosis
- Autoimmune pancreatitis
- Inflammatory pseudotumors of the orbit, lungs and kidneys

**\*Note:** IgG4 levels may or may not be high.


R10<sup>th</sup>

### Latest Update

- B2 microglobulin protein of MHC 1 is coded by chr 15. beta-2-microglobulin (coded chromosome 15) which plays an important role in the structural support of the heavy chains
- Natural IgM antibodies are also responsible for hyperacute reaction

#### HLA Typing

Serology	<ul style="list-style-type: none"> <li>• Complement Dependent Cytotoxicity (CDC)</li> <li>• Microlymphocytotoxic test</li> </ul>
Molecular methods (Diagnosis of choice)	<ul style="list-style-type: none"> <li>• PC-I SSP (Sequence Specific Priming)</li> <li>• PCR SSOP (Sequence Specific Oligonucleotide Probes)</li> <li>• Sequence Based Typing (BT)</li> </ul> <p>Reference Strand Conformational Analysis (RSCA) Luminex technology-SSOP Based Method</p>



## NEXT Pattern Questions



Q's

### 1. Scenario Type Question

1. CD8+ cells
2. Dendritic cells
3. Natural killer cells
4. Neutrophils
5. Plasma cells
6. CD4+ lymphocyte
7. Mast cell

**Scenario A.** An epidemiologic study is conducted to determine risk factors for HIV infection. The study documents that individuals with coexisting sexually transmitted diseases such as chancroid are more likely to become HIV-positive. It is postulated that an inflamed mucosal surface is an ideal location for the transmission of HIV during sexual intercourse. Which of the following cells in these mucosal surfaces is most instrumental in transmitting HIV to CD4+ T lymphocytes?


**Scenario B.** A 48-year-old woman has fingers that are tapered and claw-like, with decreased motion at the small joints. There are no wrinkle lines on her facial skin. The microscopic appearance of the skin is shown in the figure. The patient also has diffuse interstitial fibrosis of the lungs, with pulmonary hypertension and cor pulmonale. Which of the following dermal inflammatory cells is the most likely initiator of the process that is the cause of her skin disease?

**Ans. A-2, B-6**



Q's

2. A 40-year-old man complains of cough with sputum and evening rise of temperature for the past 3 months. The lung lesions were biopsied and the following image was obtained. What is the type of hypersensitivity reaction occurring in this patient?



- a. Hypersensitivity Type I    b. Hypersensitivity Type II    c. Hypersensitivity Type III    d. Hypersensitivity Type IV

**Ans. (d) Hypersensitivity Type IV**


(Ref: Robins Basic Pathology 10th ed/pg142)

Picture given: Caseating granulomatous inflammation → Tuberculosis


Examples of:

- **Type I HS:** Anaphylaxis, allergy, bronchial asthma
- **Type II HS:** Autoimmune hemolytic anemia, Good pasture syndrome
- **Type III HS:** SLE, serum sickness, Arthus reaction
- **Type IV HS:** Contact dermatitis, multiple sclerosis, type I diabetes, tuberculosis





3. The type of WBC shown in the picture plays a major role in which of the following hypersensitivity reactions?



- a. Hypersensitivity Type I      b. Hypersensitivity Type II      c. Hypersensitivity Type III      d. Hypersensitivity Type IV

Ans. (a) **Hypersensitivity Type I**

(Ref: Robins Basic Pathology 10th ed/ pg 136)


Picture – orange red granules and a spectacle shaped nucleus → Eosinophil

- **Type I—Immediate hypersensitivity:** Production of IgE antibody – release of vasoactive amines from mast cells. TH2 response occurs in late phase → IL5 release → Eosinophil activation
- **Type II—Antibody mediated hypersensitivity:** Production of IgG and IgM – binds to target and leads to phagocytosis; recruitment of WBCs
- **Type III—Immune-complex mediated hypersensitivity:** Deposition of antigen- antibody complexes – complement activation
- **Type IV—Cell mediated hypersensitivity:** Activated T lymphocytes – release of cytokines, inflammation and macrophage activation



## Image-Based Question

1. The following are features of which disease?



- a. SLE      b. RA  
c. Autoimmune hepatitis      d. HIV




## Answer of Image-Based Question

1. Ans. (a) **SLE**

- The first figure shows butterfly malar rash and the second one shows a neutrophil engulfing a lymphocyte (L.E body) which are highly suggestive features of SLE.



## Multiple Choice Questions

1. **Antibody independent cell mediated toxicity is seen in:** (AIIMS Nov 18)
  - a. NK cell
  - b. NK cell only
  - c. Macrophage
  - d. NK cells, neutrophils & macrophages
2. **Gama Delta-T cells have role in?** (AIIMS May 18)
  - a. First line of defence against bacterial peptide.
  - b. CD 5
  - c. CD 4
  - d. CD 8
3. **Macrophage engulfs different cells as shown in the image, the process is called:** (AIIMS Nov 18)
 
  - a. Necrosis
  - b. Phagocytosis
  - c. Emperipolesis
  - d. Autophagy
4. **Immune privilege site is:** (AIIMS Nov 18)
  - a. Seminiferous tubule
  - b. Optic nerve
  - c. Area postrema
  - d. Spinal canal
5. **CD markers for natural killer cells is/are:**
  - a. CD 16
  - b. CD 117
  - c. CD 56
  - d. CD 1
  - e. CD 4
 (PGI Nov 2017)
6. **Which CD molecule is important for presentation of lipid antigen?** (JIPMER 2016)
  - a. CD4
  - b. CD8
  - c. CD1
  - d. CD16
7. **Interleukin secreted by Th1 cells include?**
  - a. IL-2
  - b. IL-4
  - c. IL-10
  - d. IL-13
 (JIPMER 2016)
8. **Which of the following is a secondary lymphoid organ?**
  - a. Liver
  - b. Spleen
  - c. Bone marrow
  - d. Thymus
 (JIPMER 2016)
9. **Which of the following are/is not an activator of alternate complement system?** (JIPMER 2016)
  - a. Factor H
  - b. IgA
  - c. Bacteria
  - d. Toxin
10. **Which antibody indicates intrauterine infection in baby?** (Recent Question 2016-17)
  - a. IgA
  - b. IgM
  - c. IgG
  - d. IgD
11. **Which antibody protects from intestinal infection?** (Recent Question 2016-17)
  - a. IgA
  - b. Ig M
  - c. Ig G
  - d. IgE
12. **Immunoglobulins differ with each other in all except** (Recent Question 2016-17)
  - a. Heavy chain
  - b. Light chain
  - c. Electrophoresis
  - d. Molecular weight
 (Recent Question 2016-17)
13. **Stellate Granuloma are seen in** (Recent Question 2016-17)
  - a. Cat scratch disease
  - b. Sarcoidosis
  - c. LGV
  - d. Histoplasmosis
14. **Castleman syndrome is associated with?** (Recent Question 2016-17)
  - a. EBV
  - b. CMV
  - c. HHV 8
  - d. Herpes virus
15. **Role of macrophages in antibody production is?** (Recent Question 2016-17)
  - a. Antigen presentation
  - b. B cell production
  - c. Class switching
  - d. B cell activation
16. **Which virus causes abnormal proliferation of B cells?** (Recent Question 2016-17)
  - a. CMV
  - b. HHV-8
  - c. HTLV
  - d. EBV
17. **Which of the following immunoglobulin is called homocytotropic antibody?** (Recent Question 2015)
  - a. IgA
  - b. IgG
  - c. IgM
  - d. IgE
18. **Which Antibody crosses placenta?** (Recent Question 2016)
  - a. IgG1
  - b. IgG4
  - c. IgA
  - d. IgD
19. **IgG is what % of total plasma proteins?** (Recent Question 2016)
  - a. 5-10
  - b. 10-15
  - c. 15-20
  - d. 25-30
20. **EBV enters the cells through** (Recent Question 2016)
  - a. CR1
  - b. CR2
  - c. CR3
  - d. CR4
21. **About Natural killer cells, all are true except?** (PGI NOV 2015)
  - a. Large granular cells
  - b. Have perforins
  - c. Need thymus for development
  - d. CD16 is FcR of IgG
  - e. Kills intracellular bacteria
22. **Conditions with high ESR and normal CRP is/are?** (PGI Nov 2015)
  - a. Pregnancy
  - b. Tuberculosis
  - c. Multiple myeloma
  - d. Polycythemia
  - e. SLE
23. **Which is a large granular lymphocyte?** (Recent Question 2015)
  - a. NK cell
  - b. B-lymphocyte
  - c. T-lymphocyte
  - d. Macrophage
24. **Which of these is CD marker of cytotoxic cells?** (Recent Question 2015/2014)
  - a. CD8
  - b. CD4
  - c. CD1a
  - d. CD1



- 25. Cells involved in humoral immunity:** (Recent Question 2014)  
 a. B-cells b. T-cells  
 c. Helper cells d. Dendritic cells
- 26. Macrophages are converted to epithelioid cells by which cytokine:** (Recent Question 2014)  
 a. IL-2 b. IFN- $\gamma$   
 c. TNF-alpha d. TGF-beta
- 27. Which cytokine activate macrophages:** (Recent Question 2014)  
 a. IL-8 b. IFN- $\gamma$   
 c. PAF d. Leukotriene B4
- 28. Phagocytosis is the function of:** (Recent Question 2014)  
 a. Astrocytes b. Oligodendrocytes  
 c. Microglia d. Schwann cells
- 29. MALT is most commonly present in** (AIIMS Nov 14)  
 a. Duodenum b. Jejunum  
 c. Ileum d. Stomach
- 30. Which of the following are examples of innate immunity?** (PGI MAY 2014)  
 a. NK cells b. Macrophages  
 c. B-cell d. T cell  
 e. Complements
- 31. Null cells constitute what percentage of peripheral lymphocytes?** (JIPMER 2014)  
 a. 0-1 b. 2-5  
 c. 5-10 d. 15-20
- 32. All of the following are true about type 1 HLA except?** (Recent Question 2016)  
 a. Present on APC  
 b. Activate cytotoxic T cell and kill virus infected cell  
 c. Present on nucleated cells  
 d. First line defense mechanism
- 33. EBV action in nasopharynx through?** (Recent Question 2015)  
 a. CD 3 b. CD 4  
 c. CD 8 d. CD 21
- 34. Macrophages- false statement is?** (JIPMER 2013)  
 a. Derived from monocytes  
 b. Harbor Mycobacteria  
 c. Involved in Type 3 HSN  
 d. Produces TNF and interleukins
- 35. Antibody that is produced rapidly and in high amounts during secondary response?** (JIPMER 2013)  
 a. IgM b. IgG  
 c. IgA d. IgE
- 36. Cells that are identified by presence of immunoglobulins on the surface are?** (JIPMER 2013)  
 a. Neutrophils b. B-cells  
 c. NK cells d. Monocytes
- 37. Not true regarding NK cells is?** (JIPMER 2013)  
 a. Active against malignant cells  
 b. Involve MHC antigen for killing micro-organisms  
 c. First line defense against viral infections  
 d. No prior sensitization required

- 38. Which of the following are activators of T lymphocyte:** (PGI May 12)  
 a. CD 79b b. CD 3  
 c. CD 28 d. CD 14  
 e. LCK
- 39. All of the following are functions of CD4 helper cells, except:** (AIIMS May 11, 10, AI 09)  
 a. Immunogenic memory b. Produce immunoglobulins  
 c. Activate macrophages d. Activate cytotoxic cells
- 40. B-cell markers are:** (PGI Nov 2011)  
 a. CD-15 b. CD-19  
 c. CD-20 d. CD-56  
 e. CD-79a
- 41. Virus infected cells die by the help of:** (WB PG 2011)  
 a. IL1 b. BIL-2  
 c. B cell with antibody d. Cytotoxic T cell

### HLA, MHC & ANTIGEN PRESENTING CELLS

- 42. Naive T cell gets activated by?** (AIIMS Nov 18)  
 a. NK cell b. Dendritic cell  
 c. Macrophage d. B- lymphocytes
- 43. MHC-II is present on?** (Recent Question 2019)  
 a. All nucleated cells b. Antigen presenting cells  
 c. RBCs d. CD4T cells
- 44. Which of the following cells have MHC class II?** (AIIMS MAY 2017)  
 a. RBC  
 b. B cells, Dendritic cells, NK cells  
 c. All nucleated cells d. Platelets
- 45. Antigen presenting cells are?** (AIIMS May 2017)  
 a. Langerhans cells of skin b. Kupffer  
 c. Macrophages d. Monocytes  
 e. Thymocytes
- 46. Complement complex that attacks cell membrane is:** (AIIMS Nov 2016)  
 a. C12345 b. C23456  
 c. C34567 d. C56789
- 47. HLA is absent in** (AIIMS Nov 2016)  
 a. RBC b. Monocyte  
 c. Neutrophils d. Thrombocytes
- 48. Which of the following is not an antigen presenting cell?** (AIIMS May 2016)  
 a. M cell b. Thymocyte  
 c. Macrophage d. Langerhans cells
- 49. Which hepatitis virus is associated with spontaneous remission with HLA DR4 ?** (JIPMER 2016)  
 a. HAV b. HBV  
 c. HCV d. HDV
- 50. Dendritic cell expresses?** (Recent Question 2015)  
 a. MHC 1 b. MHC 2  
 c. MHC 3 d. MHC 4
- 51. MHC III codes for** (Recent Question 2015)  
 a. TNF alpha b. IL 1  
 c. HLA A d. HLA B
- 52. The professional antigen presenting cells are:** (Recent Question 2014)  
 a. B cells b. Dendritic cells  
 c. T cells d. NK cells



- 53. HLA is located on** (Recent Question 2014/AIIMS Nov 09, 08)
- Short arm of chr-6
  - Long arm of chr-6
  - Short arm of chr-3
  - Long arm of chr-3
- 54. True about MHC:** (Recent Question 2014)
- Present on chromosome 5
  - Class II comprises A,B,C
  - Class III codes for complement
  - Class I is involved in mixed leucocyte reaction
- 55. The role played by Major Histocompatibility Complex-I and -2:** (AIIMS Nov 14/MAY 2012)
- Transduce the signal to T cells following antigen recognition
  - Mediate immunogenic class switching
  - Present antigens for recognition by T cell antigen receptors
  - Enhance the secretion of cytokines
- 56. Class II MHC is expressed in all except?** (JIPMER 2013)
- Activated T cells
  - Dendritic cells
  - Langerhans cells
  - Neutrophils
- 57. MHC II is/are presented by:** (PGI May 12)
- Macrophage
  - Dendritic cells
  - Lymphocyte
  - Epithelial cell
  - Platelets

#### HYPERSENSITIVITY REACTIONS

- 58. B cells are induced to produce IgE by which of the following?** (Recent Pattern Question 2020)
- IL-2
  - IL-4
  - IL-1
  - IL-6
- 59. Basophils are activated by:** (AIIMS Nov 2019)
- IL-5
  - IgE fixed cell
  - Complement factor
  - TNF
- 60. Variation-2: Basophils are activated by:** (AIIMS Nov 2019)
- IL-1
  - IL-2
  - IL-4
  - IL-10
- 61. Interleukin involved in early phase of hematopoiesis is:** (JIPMER 2019)
- IL-1
  - IL-2
  - IL-3
  - IL-4
- 62. Which of the following is secreted by TH-1 helper cells?** (JIPMER 2019)
- IL-1
  - IL-2
  - IL-4
  - IL-3
- 63. Which of the following is a major cell infiltrate in rheumatoid arthritis?** (Recent exam 2018)
- CD4+ helper cell
  - Macrophage
  - Lymphocyte
  - Dendritic cells
- 64. Rh incompatibility is an example of** (Recent Question 2016)
- Immune complex reaction
  - Ag Ab reaction
  - Delayed hypersensitivity
  - Immediate type
- 65. IgE receptor present on ?** (AIIMS May 2015)
- Mast cell
  - NK cell
  - B cell
  - T cell
- 66. A 45-year-old patient presents with fever, night sweats, weight loss. On X ray a mass in apical lobe of lung is seen. On histopathology found to have caseous necrosis. What is the underlying process?** (AIIMS May 2015)
- Enzymatic degeneration
  - Hypersensitivity reaction with modified macrophages, lymphocytes and giant cells
  - Acute decrease in blood supply
  - decreased growth factors
- 67. Delayed hypersensitivity is initiated by** (MH PG 2014)
- CD3+T cells
  - CD4 + T cells
  - CD8+Tcells
  - CD10+Tcells
- 68. Schwartzman reaction is an example of:** (APPGMEE 2015)
- None of the options
  - Type I hypersensitivity reaction
  - Type III hypersensitivity reaction
  - Type II hypersensitivity reaction
- 69. Which factor is secreted by macrophage and acts on neutrophil and endothelium in hypersensitivity reaction?** (Recent Question 2015)
- IL-1
  - Tumor necrosis factor
  - Insulin like Growth factor
  - Thromboxane A2
- 70. Type V hypersensitivity reaction is a subtype of:** (Recent Question 2015)
- Type I
  - Type II
  - Type III
  - Type IV
- 71. Major basic protein is found in?** (Recent Question 2015)
- Macrophage
  - Eosinophils
  - Basophil
  - Neutrophil
- 72. Most important mediator of late phase of immediate type of hypersensitivity reaction is:** (Recent Question 2015)
- Histamine
  - Major basic protein
  - Platelet activating factor
  - PGE2
- 73. Antibodies in ITP are:** (Recent Question 2014)
- IgG
  - IgM
  - IgE
  - IgD
- 74. Transfusion reaction and erythroblastosis fetalis are:** (Recent Question 2014, DNB June 11)
- Type I hypersensitivity
  - Type II hypersensitivity
  - Type III hypersensitivity
  - Type IV hypersensitivity
- 75. 48 hours after taking penicillin, a patient complains of body ache. He gives no history of any old allergy to penicillin. Blood tests show anti penicillin antibody and hemolysis. Which type of hypersensitivity reaction is this?** (AIIMS Nov 14)
- Type I HSN
  - Type II HSN
  - Type III HSN
  - Type IV HSN





- 76. Which is the example of type 2 hypersensitivity reaction?**  
a. Serum sickness      b. PAN (WB PG 2014)  
c. SLE      d. Rh incompatibility
- 77. An adult develops swelling of tongue and neck after ingestion of peanut. What is your diagnosis?**  
(JIPMER 2014)  
a. Foreign body larynx  
b. Angioneurotic edema  
c. Parapharyngeal abscess  
d. Inflammatory response
- 78. Type I lepra reaction is an example of which type of hypersensitivity according to coombs and gel classification?**  
(JIPMER 2014), (MAHA 2016)  
a. I      b. II  
c. III      d. IV
- 79. Which of the following immune hypersensitivity reaction is responsible for Myasthenia Gravis?** (AI 12)  
a. Type I Hypersensitivity  
b. Type II Hypersensitivity  
c. Type III Hypersensitivity  
d. Type IV Hypersensitivity
- 80. LATS is a?** (DNB Aug 12 Pattern)  
a. IgMAB      b. IgGAb  
c. Glycoprotein      d. IgA Ab
- 81. Substance that can cause Anaphylactic shock is?**  
(Recent Question 12)  
a. Histamine      b. Adrenaline  
c. Nor-adrenaline      d. Glucocorticoids
- 82. Raji cell assay are used to identify?** (DPG 11)  
a. Complement levels      b. Immune complexes  
c. T- cells      d. IFN levels

#### AUTOIMMUNE SYSTEM

- 83. Which of the following is/are involved in etiopathogenesis of SLE?** (PGI May 2019)  
a. Interferon-1 (IFN-1)  
b. Interferon-1 (IFN-β)  
c. Exposure to UV rays  
d. Deficiency of early complement factors  
e. Failure of self-tolerance in B cells
- 84. A 29-year-old female with history of polyarthralgia and back ache was investigated. She had nucleolar pattern in IF. Which of the following is the best fit in her case?**  
a. Glomerulonephritis and heart failure (AIIMS MAY 2017)  
b. Raynaud phenomenon and sclerodactyly  
c. Genital and oral painful ulcers  
d. Joint pains with nodules on skin.
- 85. During development, all the antigens of self are introduced to thymic cells in-order to be removed to prevent autoimmunity. Which of the following genes is involved in the process?** (AIIMS MAY 2017)  
a. NOTCH1      b. AIRE  
c. RB gene      d. CPK gene
- 86. Which of the following gene is responsible for Autoimmune polyendocrinopathy which gene?**  
a. FOX P3      b. AIRE (JIPMER 2016)  
c. CD 25      d. PD-1

- 87. Schaumann bodies are not seen in :**  
a. Sarcoidosis (Recent Question 2016-17)  
b. Histoplasmosis  
c. Cryptococcosis  
d. Hypersensitivity pneumonitis  
e. Tuberculosis
- 88. Which of the following is a Immune privileged region?** (AIIMS May 2015)  
a. Area postrema      b. Seminiferous tubules  
c. Kidney      d. Optic nerve
- 89. Which of the following is not an auto immune disorder?** (AIIMS May 2015)  
a. Ulcerative colitis      b. Grave's disease  
c. Rheumatoid arthritis      d. SLE
- 90. Most sensitive antibody for SLE** (Recent Question 2015)  
a. Anti-nuclear      b. Anti-ds DNA  
c. Anti-Smith      d. Anti-histone
- 91. Antibody in drug induced lupus**  
(Recent Question 2015)  
a. Anti-nuclear      b. Anti-ds DNA  
c. Anti-Smith      d. Anti-histone
- 92. All of the following are features of SLE except**  
a. Shrinking lung syndrome (Recent Question 2015)  
b. Erosive arthritis  
c. Libman sack's endocarditis  
d. Cognitive dysfunction
- 93. Site of biopsy in Sjogren syndrome**  
(Recent Question 2015)  
a. Kidney      b. Abdominal fat  
c. Lip      d. Rectum
- 94. Antibody in diffuse scleroderma**  
a. Anti-DNA topoisomerase (Recent Question 2015)  
b. Anti-Centromere  
c. Anti-RNA polymerase  
d. Anti-Smith
- 95. Most common antibody in sjogren syndrome**  
(Recent Question 2015)  
a. Anti-DNA topoisomerase  
b. Anti-Centromere  
c. Anti-RNA polymerase  
d. Anti-Ribonucleoprotein
- 96. Class IV lupus nephritis** (Recent Question 2015)  
a. Mesangial lupus nephritis  
b. Proliferative lupus nephritis  
c. Membranous lupus nephritis  
d. Diffuse sclerosing lupus nephritis
- 97. Shrinking lung syndrome is a feature of**  
(Recent Question 2015)  
a. Systemic lupus erythematosus  
b. Systemic sclerosis  
c. AIDS  
d. ARDS
- 98. Anti-smith antibody is specific for**  
(Recent Question 2015)  
a. Systemic sclerosis  
b. Sjogren syndrome  
c. Systemic lupus erythematosus  
d. Drug induced lupus



- 99. Commonest renal lesion in SLE is:**  
(Recent Question 2015)  
a. Focal proliferative      b. Diffuse Proliferative  
c. Membranous              d. Minimal change disease
- 100. Joint erosions are typically absent in**  
(Recent Question 2015)  
a. Rheumatoid arthritis  
b. Systemic lupus erythematosus  
c. Psoriatic arthritis  
d. Osteoarthritis  
e. Mesangial Proliferative
- 101. Antibody seen in neonatal lupus with congenital heart block**  
(Recent Question 2015)  
a. Anti-ribosomal              b. Anti-Ro (SS-A)  
c. Anti-neuronal              d. Anti-histone
- 102. Most severe form of lupus nephritis**  
(Recent Question 2015)  
a. Mesangial lupus lomerulonephritis  
b. Focal proliferative glomerulonephritis  
c. Diffuse proliferative glomerulonephritis  
d. Membranous glomerulonephritis
- 103. Anti-histone antibodies are specific for**  
(Recent Question 2015)  
a. CNS lupus  
b. Drug induced lupus  
c. Neonatal lupus  
d. Cutaneous lupus
- 104. False statement regarding musculoskeletal manifestations of SLE**  
(Recent Question 2015)  
a. Intermittent polyarthritis  
b. Commonly involves hands, wrist, knees  
c. Erosions seen on X-rays  
d. Ischemic necrosis of bone can occur, when the patient is on glucocorticoids
- 105. All are true about limited cutaneous systemic sclerosis except**  
(Recent Question 2015)  
a. Anti-Topoisomerase I antibody  
b. Digital ischemia  
c. Calcinosis  
d. Isolated pulmonary arterial hypertension
- 106. Most common cause of death in SLE after first few years of diagnosis**  
(Recent Question 2015)  
a. CNS disease  
b. Opportunistic infections  
c. Accelerated atherosclerosis  
d. Nephritis
- 107. All the following about drug induced SLE are true except**  
(Recent Question 2015)  
a. More common in females  
b. Renal and CNS involvement is uncommon  
c. Complement level are normal  
d. The disease remits after withdrawal of the offending drug
- 108. Most common autoimmune disease associated with sjogren syndrome**  
(Recent Question 2015)  
a. Scleroderma  
b. Rheumatoid arthritis  
c. Inflammatory bowel disease  
d. SLE

- 109. Which part of GIT is most severely affected in systemic sclerosis**  
(Recent Question 2015)  
a. Esophagus                      b. Jejunum  
c. Sigmoid colon              d. Rectum
- 110. Digital ischemia is common in** (Recent Question 2015)  
a. Diffuse scleroderma  
b. Diffuse cutaneous scleroderma  
c. Limited scleroderma  
d. Systemic lupus erythematosus
- 111. Match the above HLA associations and choose the best match:** (APPGMEE 2015)
- |   |         |   |                          |
|---|---------|---|--------------------------|
| p | HLA DR2 | w | Dermatitis herpetiformis |
| q | HLA DR3 | x | Celiac disease           |
| r | HLA DR4 | y | Dermatitis herpetiformis |
| s | HLA DQ2 | z | Pemphigus vulgaris       |
- a. pqrs = wxzy                      b. pqrs = ywzx  
c. pqrs = xwys                      d. pqrs = zxwy
- 112. Which is not included in Mixed connective tissue disorder?** (Recent Question 2015)  
a. Systemic sclerosis              b. SLE  
c. Rheumatoid arthritis              d. Polymyositis
- 113. Which antibody is associated with Mixed connective tissue disease (MCTD)?** (Recent Question 2015)  
a. Anti U1-RNP Ab              b. Anti DNA Ab  
c. Anti histoneAb              d. Anti ds-DNA Ab
- 114. Band test is done in:** (Recent Question 2014)  
a. RA                                      b. SLE  
c. Scleroderma                      d. PAN
- 115. Antinuclear antibody specific for SLE is:** (Recent Question 2014)  
a. Anti ds DNA  
b. Anti nuclear antibodies  
c. Anti centromere antibody  
d. Anti histoneAb
- 116. HLA associated with psoriasis:** (Recent Question 2014, 2013)  
a. HLA-B27                              b. HLA-DR4  
c. HLA-CW6                              d. HLA-B8
- 117. Drug induced lupus antibodies are:** (Recent Question 2014)  
a. Anti-Rho                              b. ds-DNA  
c. Anti-Sm                              d. Anti-histone antibody
- 118. HLA marker of Bechet's syndrome:** (Recent Question 2014)  
a. HLA-B27                              b. HLA-DR5  
c. HLA-B51                              d. HLA-CW6
- 119. HLA associated with rheumatoid arthritis:** (Recent Question 2014, 2013)  
a. HLA-B27                              b. HLA-DR4  
c. HLA-CW6                              d. HLA-B8
- 120. Thromboangitis obliterans is associated with:** (Recent Question 2014)  
a. HLA-B27                              b. HLA-DR4  
c. HLA-B5                              d. HLA-DR2
- 121. ANA (antinuclear antibody) is seen in all except:** (Recent Question 2014)  
a. SLE                                      b. RA  
c. Sjogren's syndrome              d. Systemic sclerosis



- 122. HLA B-27 has > 90% association with?** (AIIMS Nov 14)  
 a. Enteropathic                      b. Reactive arthritis  
 c. Rheumatoid arthritis              d. Ankylosing spondylitis
- 123. Miller-Fisher syndrome associated with:** (WB PG 2014)  
 a. HLA DQBI                      b. Antibodies to GQ1B  
 c. Antibodies to peroxidase              d. HLAB5
- 124. Thrombosis is seen in which stage of lupus nephritis?** (Recent Question 2013)  
 a. Class I                      b. Class II  
 c. Class III                      d. Class IV
- 125. A 30-year-old lady presents to the outpatient department with an erythematous butterfly rash on her cheeks. Which of the following antibodies should be assayed initially for her suspected condition:** (AI 12)  
 a. Anti-ds-DNA  
 b. Anti-Ro-Antibody  
 c. Anti-Centromere-Antibody  
 d. Anti-mitochondrial-Antibody
- 126. In SLE, active nephritis is/are characterized by:** (PGI Nov 2011)  
 a. Proliferation of endothelium  
 b. Glomerular leukocyte infiltration  
 c. Mesangial cell proliferation  
 d. Epithelial cell proliferation  
 e. Tubulitis
- 127. In SLE, pulmonary involvement is characterized by:** (PGI Nov 2011)  
 a. Interstitial Fibrosis  
 b. Shrinking lung syndrome  
 c. Alveolar hemorrhage  
 d. Pulmonary arterial hypertension  
 e. Cavitation
- 128. Nucleolar pattern on immunofluorescence is seen in:** (PGI May 2011)  
 a. Antibody to DNA                      b. Antibody to RNA  
 c. Antibody to histone                      d. Systemic sclerosis  
 e. SLE
- 129. Biopsy of the parotid gland in a patient with Sjogren's syndrome shows:** (Jipmer 11)  
 a. Neutrophils                      b. Lymphocytes  
 c. Eosinophils                      d. Basophils
- 130. HLA associated with rheumatoid arthritis:** (PGI May 10)  
 a. DRB1                      b. DR1  
 c. DR2                      d. DR3
- 131. Best marker of SLE?** (DNB June 10)  
 a. Anti Sm antibodies                      b. Anti-ds DNA antibodies  
 c. Anti-Histone antibodies                      d. Anti Ro (SS-A) antibodies

#### REJECTION OF TISSUE TRANSPLANTS

- 132. Graft between identical twins is:** (Recent Pattern Question 2020)  
 a. Allograft                      b. Xenograft  
 c. Isograft                      d. Autograft
- 133. Organ transplantation between mother to child is a type of?**  
 a. Autograft                      b. Isograft  
 c. Allograft                      d. Xenograft

- 134. Cells involved in GVHD are?** (JIPMER 2017)  
 a. Recipient B cells                      b. Recipient T cells  
 c. Donor B cells                      d. Donor T cells
- 135. Hyperacute rejection occurs within?** (Recent exam 2018)  
 a. 12 hours                      b. 2 weeks  
 c. 1 month                      d. 3 months
- 136. A patient of cirrhosis with liver failure comes to you for stem cell transplantation your method will be?** (PGI Nov 2016)  
 a. Transfer of stem cells from other persons liver  
 b. Taking patient skin stem cell and transferring into liver  
 c. Tranfer hepatocytes from the same person for regeneration  
 d. Transfer hepatic progenitor cells (HPCs) of same person for regeneration
- 137. Graft vs Host reaction can be reduced by?** (JIPMER 2016)  
 a. Irradiation  
 b. Leukoreduction/leuckofiltration  
 c. Immunosuppression  
 d. Buffy coat removal
- 138. After a solid organ transplantation, which of the following is responsible for acute graft rejection?** (AIIMS May 2016)  
 a. C3a  
 b. C3b  
 c. C5a  
 d. C4d
- 139. For transplantation which HLA requires minimum matching is required** (Recent Question 2016-17)  
 a. HLA A  
 b. HLA B  
 c. HLA DR  
 d. HLA DP
- 140. In a case of kidney allograft rejection, what is the best diagnostic feature ?** (Recent Question 2016-17)  
 a. Increased neutrophil  
 b. Increased basophils  
 c. Biopsy  
 d. Complements levels
- 141. A patient requires liver transplant. He plans to receive it from his brother. They are not twins. On HLA typing, HLA is matched at the A, B, and DRB1 loci. Siblings are:** (Recent Question 2016)  
 a. Matched, unrelated donors  
 b. Matched, related donors  
 c. Mismatched, related donors  
 d. Mismatched, unrelated donors
- 142. For a successful transplantation how many HLA should be matched?** (Recent Question 2016)  
 a. 1-4                      b. 5-10  
 c. 11-15                      d. 15-20
- 143. Which of the following is not a characteristic feature of GVHD?** (Recent Question 2015)  
 a. Skin involvement  
 b. Renal involvement  
 c. Liver involvement  
 d. Intestinal involvement



**144. "000" mismatch in graft transplanatation means?**  
(Recent Question 2016)

- a. Partial mismatched
- b. Completely mismatched
- c. Matched
- d. Partially matched

**145. True about hyperacute rejection in renal transplant:**

- a. Occur within few days of transplant (PGI May 2015)
- b. T cell involvement
- c. Blood vessel thrombosis
- d. Eosinophilic infiltration
- e. B cell infiltration

**146. Graft rejection is:** (Recent Question 2014)

- a. Cell mediated
- b. Humoral
- c. Both
- d. None

**147. HLA-I is present on:** (Recent Question 2014)

- a. All nucleated cells
- b. Only on cells of immune system
- c. Only on B-cells
- d. Only on T-cells

**148. In acute transplant after 6 months, the rejection is because of?**  
(Recent Question 2015)

- a. Neutrophil
- b. T-lymphocyte
- c. B-lymphocyte
- d. Macrophage

**149. Following conditions must be fulfilled in a person before taking him as a kidney donor EXCEPT** (APPGMEE 14)

- a. ABO compatibility with recipient
- b. Presence of two normally functioning kidneys
- c. No HIV infection
- d. Zero HLA mismatch with recipient

**150. Allograft rejection is an example of?** (JIPMER 2014)

- a. GVHD
- b. Delayed types hypersensitivity
- c. Immediate hypersensitivity
- d. Acute rejection

**151. Hyperacute rejection is due to:** (AIIMS May 2013)

- a. Preformed antibodies (AIIMS Nov 2012)
- b. Cytotoxic T-lymphocyte mediated injury
- c. Circulating macrophage mediated injury
- d. Endothelitis caused by donor antibodies

**152. Most commonly involved organs in graft versus host disease are all except:** (Recent Question 12)

- a. Gut
- b. Liver
- c. Skin
- d. Kidney

**153. Graft from identical twin is defined as:** (WB PG 2012)


- a. Allograft
- b. Isograft
- c. Xenograft
- d. Autograft

**154. In GVH disease all are involved except:**  
(WB PG 2011, AIIMS May 07)

- a. Liver
- b. Lungs
- c. Intestine
- d. Skin

## IMMUNODEFICIENCY SYNDROMES

**155. In the given below flow cytometry graphs; If A and B are normal scatter plots for B cells. Interpret images C and D and come to diagnosis.**  
(AIIMS Nov 2019)



- a. Bare lymphocyte syndrome
- b. SCID
- c. Hyper IgM syndrome
- d. Chronic granulomatous disease

**156. Which is not a cause/involved in SCID?** (AIIMS Nov 2019)

- a. ZAP70
- b. IL2R
- c. JAK3
- d. BTK

**157. Which is not a feature of Lofgren syndrome?**

(JIPMER 2019)

- a. Uveitis
- b. Erythema nodosum
- c. Polyarthralgia
- d. Bilateral hilar adenopathy

**158. Which of the following is/are results from defect in innate immunity:**  
(PGI May 2019)

- a. DiGeorge syndrome
- b. Severe combined immunodeficiency
- c. Common variable immunodeficiency
- d. Chediak-Higashi syndrome
- e. Bruton agammaglobulinemia

**159. True regarding 'Bare lymphocytes syndrome'?**

(PGI May 2018)

- a. Autosomal recessive
- b. Part of SCID
- c. MHC overexpression
- d. Leads to abnormal CD4T cell development

**160. Feature(s) of DiGeorge syndrome is/are all except?**

(PGI Nov 2017)

- a. Results from failure of development of the third and fourth pharyngeal pouches
- b. Absent thyroid
- c. Absent parathyroid glands
- d. B cell defect
- e. Enhanced susceptibility of bacterial infection






- 161. In SCID which DNA mechanism is defective?**  
 a. Non homologous disjunction repair (AIIMS Nov 2017)  
 b. Homologous disjunction repair  
 c. Base excision repair  
 d. NER
- 162. Combined B & T cell immunodeficiency is/are seen in?** (PGI Nov 2017)  
 a. Severe Combined Immunodeficiency  
 b. Adenosine deaminase deficiency  
 c. Wiskott-Aldrich syndrome  
 d. Ataxia telangiectasia  
 e. DiGeorge syndrome
- 163. Common variable immunodeficiency shows?** (Recent Question 2016-17)  
 a. B cell defect  
 b. T cell defect  
 c. Both  
 d. None
- 164. Lymphocyte phenotype test done for?** (Recent Question 2016-17)  
 a. Agammaglobulinemia      b. SCID  
 c. Sepsis      d. Acute leukemia
- 165. All of the following are not seen in SCID except?** (Recent Question 2016-17)  
 a. Autoimmune ds      b. Granuloma formation  
 c. GVHD      d. Graft rejection
- 166. Both B and T cell defect is present in:** (PGI May 12)  
 a. SCID  
 b. Common Variable immunodeficiency  
 c. Wiskott-Aldrich syndrome  
 d. X-linked Agammaglobulinemia  
 e. Chronic mucocutaneous candidiasis
- 167. Chediak Higashi syndrome is characterised by the following except:** (MH 11)  
 a. Neutrophilia  
 b. Defective degranulation  
 c. Delayed microbial killing  
 d. Giant granules

#### HIV/AIDS

- 168. Identify the parasite in the intestinal biopsy of a HIV positive patient.** (Recent Pattern Question 2020)



- a. Giardia      b. CMV  
 c. Amoebic colitis      d. Cryptosporidium

- 169. HIV-1 differs with HIV-2 in?** (PGI Nov 2016)  
 a. HIV-2 is more dangerous than HIV-1  
 b. Enfuvirtide is not active against HIV-2  
 c. HIV-2 is more common than HIV-1 in India  
 d. More commonly transmitted from mother to child  
 e. HIV-1 worsens faster than HIV-2
- 170. Which of the following malignancy is commonly found in AIDS patients?** (MH PG 2014)  
 a. Kaposi's sarcoma      b. Fibrosarcoma  
 c. Cavernous hemangioma      d. Melanoma
- 171. HIV was discovered in which year?** (AIIMS Nov 2014)  
 a. 1983      b. 1979  
 c. 1969      d. 1990
- 172. A person died of HIV infection. Lung Autopsy performed in this person showed intranuclear basophilic inclusions. His CD4 count was less than 100/ uL. Which is the most probable diagnosis?** (AIIMS May 2014)  
 a. CMV  
 b. Herpes infection  
 c. ARDS  
 d. Pneumocystis carinii
- 173. Which of the following are AIDS defining cancers?** (PGI May 2014)  
 a. Burkitt's lymphoma  
 b. Non Hodgkin's Lymphoma  
 c. Ca esophagus  
 d. Primary lymphoma of brain  
 e. Invasive cancer of uterine cervix
- 174. HIV affects?** (JIPMER 2014)  
 a. B-cells  
 b. Helper T cells  
 c. Suppressor T-cells  
 d. Cytotoxic T-cells
- 175. Immunity in HIV true is?** (Recent Question 2016)  
 a. Cellular immunity is lost  
 b. Antibody mediated is lost  
 c. Not lost  
 d. Both lost
- 176. Family of HIV virus** (Recent Question 2016)  
 a. Lentivirus  
 b. Alpha retrovirus  
 c. Beta retro virus  
 d. none
- 177. What is false about HIV?** (Recent Question 2016)  
 a. HIV 2 is more pathogenic than HIV 1'  
 b. HIV 2 was discovered in 1987  
 c. HIV 1 is the most common type  
 d. HIV-1 group M subtype C is predominant in India
- 178. WHO AIDS stage III criteria are all except?** (PGI Nov 2015)  
 a. Candidiasis of esophagus  
 b. Kaposi's sarcoma  
 c. Pneumocystis jirovecii  
 d. Cryptococcosis, extrapulmonary  
 e. Ankylostoma duodenale infection
- 179. Most common HIV subtype in India is?** (Recent Question 2013)  
 a. HIV-1 M      b. HIV-1 N  
 c. HIV-2      d. HIV-1 O



## AMYLOIDOSIS

- 180. Amyloidosis is/are associated with which of the following feature(s)?** (PGI May 2019)
- Hepatosplenomegaly
  - Congestive heart failure
  - Proteinuria
  - Lytic bone lesions
  - Large fiber neuropathy in initial presentation
- 181. True or false regarding amyloidosis:** (AIIMS May 2019)
- SAA is most common Mediterranean fever
  - On Congo red staining, amyloid shows apple green birefringence
  - AL has kappa light chain.
  - Senile amyloidosis is due to  $\beta 2$ -microglobulin
- 182. Secondary amyloidosis is seen in?** (PGI Nov 2018)
- Bronchiectasis
  - Pulmonary TB
  - Lung abscess
  - Malignancy
  - Myeloma
- 183. Congo Red stains due to ?** (Recent Question 2016-17)
- B pleated sheets
  - $\alpha$  helix
  - Isoelectric Ph
  - Amyloid fibrils
- 184. Characteristic feature of amyloid on staining** (Recent Question 2015)
- Apple green birefringence on infrared microscopy
  - Apple green birefringence on X-ray crystallography
  - Apple green birefringence on polarizing microscopy
  - Apple green birefringence on light microscopy
- 185. Most widely used stain for amyloidosis:** (Recent Question 2015)
- Oil red O
  - Congo red
  - PAS
  - Thioflavin T
- 186. Chemical nature of amyloid in hemodialysis associated amyloidosis** (Recent Question 2015)
- AA
  - AL
  - A $\beta 2$
  - TTR
- 187. True regarding familial amyloidotic polyneuropathy** (Recent Question 2015)
- Autosomal recessive
  - Mutation in pyrin gene
  - Mutant form transthyretin is deposited
  - Deposited in the heart
- 188. Site of biopsy in amyloidosis** (Recent Question 2015)
- Kidney
  - Abdominal fat
  - Lip
  - Rectum
- 189. Amyloid deposited in the heart of aged** (Recent Question 2015)
- Normal transthyretin
  - Mutant transthyretin
  - Beta 2 microglobulin
  - Amyloid light chain
- 190. A 70-year-old patient presents with features of cardiac failure. Abdominal fat aspirate is stained with congo red and on examination with polarizing microscope exhibit apple green birefringence, Diagnosis** (Recent Question 2015)
- Marfan syndrome
  - Familial hypercholesterolemia
  - Amyloidosis
  - SLE
- 191. Which is the most striking and specific test to diagnose amyloid in tissue?** (APPGMEE 2015)
- Congo red + polarized microscopy
  - Congo red stain+ light microscopy
  - Toluidine stain
  - Methyl violet stain
- 192. All of the following statements are true about properties of amyloid, except:** (AP 2012)
- By electron microscope, it is made up largely continuous non-branching fibrils with a diameter of approximately 7.5 to 10 nm.
  - This electron microscope structure is not identical in all types of amyloid.
  - X-ray crystallography and infrared spectroscopy demonstrates a characteristic cross-beta-pleated sheet conformation
  - Congo red staining shows apple green birefringence under polarizing microscope
- 193. Which thyroid carcinoma is associated with calcitonin amyloid deposition?** (Recent Question 2014)
- Papillary
  - Follicular
  - Anaplastic
  - Medullary
- 194. Which one of the following stains is specific for Amyloid?** (Recent Question 2014)
- Periodic Acid schif (PAS)
  - Alzerian red
  - Congo red
  - Von - Kossa
- 195. Gingival biopsy is used for diagnosis of:** (Recent Question 2014)
- Scurvy
  - Sarcoidosis
  - Amyloidosis
  - SLE
- 196. Major fibril protein in Primary Amyloidosis is?** (Recent Question 2014)
- AL
  - AA
  - Transthyretin
  - Procalcitonin
- 197. False statement about Amyloidosis is?** (JIPMER 2013)
- Extracellular eosinophilic hyaline material
  - Made of calcified proteins
  - Apple green birefringence
  - Complication of chronic infection
- 198. Which of the following amyloid forms is seen in secondary amyloidosis associated with chronic diseases:** (AI 12)
- Amyloid Associated Protein
  - Amyloid light chain
  - Beta 2 Amyloid
  - ATTR
- 199. Amyloid deposition in patients with long term hemodialysis usually takes place in?** (JIPMER 2012)
- Renal vessels
  - Peripheral nerve
  - Knee joint
  - Carpal tunnel
- 200. Pinch purpura is seen in?** (JIPMER 2012)
- Primary systemic amyloidosis
  - Vitamin C deficiency
  - Purpura fulminans'
  - Kawasaki disease



- 201. A 60-year-old female is suffering from renal failure and is on hemodialysis since last 8 years. She developed carpal tunnel syndrome. Which of the following will be associated?** (AIIMS Nov 11)
- AL
  - AA
  - ATTR
  - Beta 2 microglobulin
- 202. Protein deposited in familial amyloid neuropathy:** (PGI May 2011)
- Mutated transthyretin
  - Normal transthyretin
  - Mutated beta-1
  - Mutated beta-2
  - Mutated beta-3
- 203. Most common cause of death in primary amyloidosis is?** (DNB June 11)
- Respiratory failure
  - Cardiac failure
  - Renal failure
  - Septicemia
- 204. Deposition of protein A beta 2 microglobulin is seen in which clinico-pathologic category of amyloidosis:** (Karnataka 11)
- Familial Mediterranean fever
  - Hemodialysis associated
  - Senile cerebral
  - Systemic senile
- 205. Best investigation for diagnosing amyloidosis:** (AIIMS May 10, AI 07, DNB 10)
- Rectal biopsy
  - Colonoscopy
  - CT scan
  - Upper GI endoscopy
- 206. Which of the following is/are Heredofamilial amyloidosis** (PGI May 10)
- Alzheimer's disease
  - Multiple myeloma
  - Familial Mediterranean fever
  - RA
  - Systemic senile amyloidosis
- 207. Stains used in amyloidosis:** (PGI May 10)
- Congo red
  - Thioflavin
  - Reticulin
  - Grams Iodine
  - PAS



## Answers with Explanations

### 1. Ans. (d) NK cells, neutrophils and macrophages

Antibody-dependent cell-mediated cytotoxicity (ADCC) is the killing of an antibody-coated target cell by a cytotoxic effector cell through a non-phagocytic process, characterised by the release of the content of cytotoxic granules or by the expression of cell death-inducing molecules. ADCC is triggered through interaction of target-bound antibodies (belonging to IgG or IgA or IgE classes) with certain Fc receptors (FcRs), glycoproteins present on the effector cell surface that bind the Fc region of immunoglobulins (Ig). Effector cells that mediate ADCC include natural killer (NK) cells, monocytes, macrophages, neutrophils, eosinophils and dendritic cells

### 2. Ans. (a) First line of defence against bacterial peptide

### 3. Ans. (c) Emperipolesis

Emperipolesis is the active penetration of one cell by another which remains intact. It differs from phagocytosis in that an engulfed cell exists temporarily within another cell and with an intact normal structure while in phagocytosis, the engulfed cell is destroyed by the protective action of lysosomal enzymes.

### 4. Ans. (a) Seminiferous tubules

In testis, there occurs segregation of antigens in the seminiferous tubules from immune cells in the interstitial space by a layer of Sertoli cells connected by impermeable tight junctions which form a blood-testis barrier.



## High Yield Facts

### Immune-privileged sites:<sup>a</sup>

- **Testis, eye and brain<sup>a</sup>**
- Tissues in which these antigens are located **do not communicate** with the blood and lymph
- **Difficult to induce immune responses** to antigens introduced into these sites
- Prolonged tissue inflammation on injury & release of antigen from these sites: **post-traumatic orchitis & uveitis<sup>a</sup>**

### 5. Ans. (a, c); a. CD 16; c. CD 56

### 6. Ans. (c) CD1 (Ref: Wintrob's 13th/2499)

Expression of CD1 in foam cells of atherosclerotic plaques may present lipid antigens to CD1-restricted T cells and contribute to inflammation of these lesions

### 7. Ans. (a) IL-2 (Ref: Robbins 9th/198)

Cells of the TH1 subset secrete IL-2 for self activation & cytokine IFN- $\gamma$ , which is a potent macrophage activator.

### 8. Ans. (b) Spleen (Ref: Robbins 9th/193)

### 9. Ans. (a) Factor H (Ref: Robbins 9th/pg 162-164)

- The alternative pathway, can be triggered by microbial surface molecules (e.g., endotoxin, or LPS), complex polysaccharides, cobra venom, and other substances, in the absence of antibody
- **Factor I and H inhibits complement activation**



**10. Ans. (b) IgM** (Ref: Robbins 9th/pg 199)

Since IgM is a pentameric Ig, Maternal IgM will not pass through the placenta into the fetus. However in case of fetus being infected, it will produce its own IgM molecule against the infection.

**11. Ans. (a) IgA** (Ref: Kuby immunology pg 419)

IgA antibodies are found in circulation, they are the major isotype found in secretions, including mucus in the gut, milk from mammary glands, tears, and saliva. In these secretions, IgA can neutralize both toxins and pathogens, continually interacting with the resident (commensal) bacteria that colonize our mucosal surfaces and preventing them from entering the bloodstream

**12. Ans. (b) Light chain** (Ref: Kubys Immunology pg 84)

- All antibodies share a common structure of four polypeptide chains consisting of two identical **light (L) chains** and two identical **heavy (H) chains**.
- The two major light chain constant region sequences are referred to as **(kappa) or (lambda) chains** common in all the Antibodies.

**13. Ans. (a) Cat scratch disease** (Ref: Robbins 9th/98)

Cat-scratch disease causes rounded or stellate granuloma containing central granular debris and recognizable neutrophils; giant cells

**14. Ans. (c) HHV 8** (Ref: Harrison 19th/697)

Human herpesvirus 8 is associated with primary effusion lymphoma in HIV-infected persons and multicentric Castleman's disease, a diffuse lymphadenopathy associated with systemic symptoms of fever, malaise, and weight loss.

**15. Ans. (a.) Antigen presentation** (Ref: Robbins 9th/198)

Macrophages that have phagocytosed microbes and protein antigens process the antigens and present peptide fragments to T cells. **Upon activation, B lymphocytes proliferate and then differentiate into plasma cells that secrete different classes of antibodies with distinct functions**

**16. Ans. (d) EBV** (Ref: Robbins 9th/pg 192-194)

**17. Ans. (d) IgE** (Ref: Kuby Immunology pg 298)

**Homocytotropic antibodies** are antibodies which have a higher affinity to Fc-receptors of the cells of the animal species in which they are produced than to Fc-receptors of the cells of other animal species.

In humans, IgE antibodies belong to this group.

**18. Ans. (a) IgG1**

(Ref: Maternal, Fetal, & Neonatal Physiology: A Clinical Perspective; pg 484)

- All four **IgG subclasses cross**, although the IgG1 and IgG3 subclasses are predominate.
- IgG1 crosses earliest in pregnancy & is the primary immunoglobulin transferred before 28 weeks.
- IgG3 crosses later & does not reach maternal levels until after 32-33 weeks

**19. Ans. (c) 15-20**

(Ref: Chemical and Cellular Architecture edited by N.S. Abel Lajtha pg 423)

- Albumin constitutes 52-67% of serum protein
- IgG accounts for 15-18% of total plasma proteins but only 5-12% of total CSF protein

**20. Ans. (b) CR2** (Ref: Robbins 9th/pg 191; 8th/pg 186)

**Epstein-Barr virus (EBV) enters B cells via CR2 (CD21)**

**21. Ans. (b, c, e); b. Have perforins; c. Need thymus for development; e. Kills intracellular bacteria**

(Ref: Robbins 9th/pg 191; 8th/pg 186)

- True, because of its morphology
- False, CD8 T cells have perforins
- False, T lymphocytes need thymus for development
- True, CD16 is FcR of IgG
- False, NK cells kill viruses & tumor infected cells

**22. Ans. (a, c, d, e); a. Pregnancy c. Multiple myeloma d. Polycythemia e. SLE** (Ref: Harrison 19th)

Patients with a high ESR (erythrocyte sedimentation rate) and normal C-reactive protein are those conditions without systemic inflammation such as malignancy.

- Some low-grade bone and joint infections by coagulase negative staphylococci
- Systemic lupus erythematosus. (high levels of type 1 interferons which inhibit the production of C-reactive protein in hepatocytes)

**Other conditions:** Leukaemia, anaemia, polycythaemia, viral infection, ulcerative colitis, pregnancy, oestrogens or steroids.

**23. Ans. (a) NK cell**

(Ref: Wintrobe's clinical hematology - 12th ed, pg 300; Robbins 9th/pg 192; 8th/pg 188)

- Most lymphocytes in blood are small ( $\leq 10 \mu\text{m}$ ), while some are large, known as **large granular lymphocytes (LGL)**, as they contain **azurophilic granules** in their cytoplasm.
- These cells are LGL type of **natural killer (NK) cells**.

**24. Ans. (a) CD8** (Ref: Robbins 9th/pg 191; 8th/pg 186)

**CD8**

- Is a marker for **cytotoxic T cells**
- Acts as a **co-receptor** for TCR
- Binds to HLA-I to cause **cytotoxic effect** on **virus infected and tumor cells**


**25. Ans. (a) B-cells** (Ref: Robbins 9th/pg 191; 8th/pg 187)

**26. Ans. (b) IFN- $\gamma$**  (Ref: Robbins 9th/pg 210; 8th/pg 207)

- **TH1 secrete IFN- $\gamma$   $\rightarrow$  activates macrophages** ("classically activated"). Macrophages transform to **epithelioid cells which** are large epithelium-like cells with abundant cytoplasm.

**27. Ans. (b) IFN- $\gamma$**  (Ref: Robbins 9th/pg 210; 8th/pg 207)





28. Ans. (c) **Microglia** (Ref: Robbins 9th/pg 192; 8th/pg 188)

Microglial cells are **modified macrophages** of CNS, hence they take part in **phagocytosis**.

29. Ans. (c) **Ileum**

(Ref: Wheeler's Functional Histology: A Text and Colour Atlas 6th ed Pg 216)

• **MALT is most commonly** present in Peyer's patches in lamina propria of **ileum** & throughout small intestine

Components of MALT are sometimes subdivided into:

<b>GALT</b>	Gut-associated lymphoid tissue; Eg Peyer's patches found in the lining of the small intestines
<b>BALT</b>	Bronchus-associated lymphoid tissue
<b>NALT</b>	Nasal-associated lymphoid tissue
<b>CALT</b>	Conjunctival-associated lymphoid tissue
<b>O-MALT</b>	Organized mucosa-associated lymphatic tissue; Eg tonsils of Waldeyer's tonsillar ring
<b>D-MALT</b>	Diffuse mucosa-associated lymphatic tissue
<b>LALT</b>	Larynx-associated lymphoid tissue
<b>SALT</b>	Skin-associated lymphoid tissue

30. Ans. (a, b, e); **a. NK cells; b. Macrophages; e. Complements**

(Ref: Robbins 9th/pg 186-188; 8th/pg 184)

31. Ans. (c) **5-10** (Ref: Robbins 9th/pg 192; 8th/pg 188)

- **Natural Killer Cells or Null cells constitute 5% to 10%** of peripheral lymphocytes<sup>Q</sup>
- As NK cells neither have B nor T-cell markers, hence also called **Null Cells**
- NK cells destroy irreversibly stressed, virus-infected cells & tumor cells **without prior exposure**

32. Ans. (d) **First line defense mechanism** (Ref: R 9th/pg 191)

- True; It is present on all nucleated cells including antigen presenting cells as well as platelets
- True; It activates cytotoxic T cell and kill virus infected cell
- True; It is present on platelets also, which are non-nucleated cells
- False; It is a part of adaptive immunity (second line defense mechanism)

33. Ans. (d) **CD21** (Ref: Robbins 9th/pg 191; 8th/pg 186)

34. Ans. (c) **Involved in Type 3 HSN**

(Ref: Robbins 9th/pg 210)

A. True	Monocytes in tissues are known as Macrophages
B. True	Macrophages engulf Mycobacteria but are not able to kill it; So they harbor Mycobacteria
C. False	Macrophages are involved in Type 4 Hypersensitivity & not type 3, which is immune-complex mediated
D. True	<ul style="list-style-type: none"> <li>• Macrophages <b>secrete TNF, IL-1, and chemokines</b>, which promote inflammation</li> <li>• Macrophages also produce <b>IL-12</b>, thereby amplifying the <b>TH1 response</b></li> </ul>

35. Ans. (b) **IgG** (Ref: Kuby Immunology)

36. Ans. (b) **B-cells** (Ref: Robbins 9th/pg 191; 8th/pg 187)

B-cells have IgM and IgG on the surface of cells, which are used to identify them

37. Ans. (b) **Involve MHC antigen for killing micro-organisms** (Ref: Robbins 9th/pg 192; 8th/pg 188)

- **Natural Killer Cells Destroy** virus-infected cells and tumor cells **without prior exposure**;
- **NK cells do not require MHC for killing micro-organisms**
- **Surface molecules of NK cells are CD16 & CD56**

38. Ans. (b, c, e); **b. CD 3; c. CD 28; e. LCK**

(Ref: Robbins 9th/pg 191; 8th/pg 186)

<b>A. CD 79b</b>	B-cell development and B-cell function
<b>B. CD 3</b>	Activates T cells & helps in signal transduction
<b>C. CD 28</b>	Co-stimulatory receptor for T cells
<b>D. CD 14</b>	LPS-induced <b>activation of monocytes</b> ; innate immunity
<b>E. LCK</b>	Tyrosine phosphorylation causing T Cell Receptor stimulation & activation

The table suggests that the activators of lymphocyte are: **CD3, CD28 and LCK**

39. Ans. (b) **Produce immunoglobulins** (Ref: R 9th/pg 210)

40. Ans. (b, c, e) **b. CD-19; c. CD-20; e. CD-79a**

(Ref: Robbins 9th/pg 191; 8th/pg 187)

B cell markers are **CD19 (most specific)**, CD20, CD21, CD22, CD23, CD24, CD10, CD79a

41. Ans. (d) **Cytotoxic T cell** (Ref: Robbins 9th/pg 191)

Cytotoxic T lymphocytes (CTLs) kill **virus infected and tumor cells**

42. Ans. (b) **Dendritic cell**

Antigen Presenting cells; cells that present Ag to T-cells

- **Antigen-presenting cells (APCs) for initiating T-cell responses against protein antigens<sup>Q</sup>**
- **Mature dendritic cells are the most potent<sup>Q</sup> stimulator of Naive T-cells**

43. Ans. (b) **Antigen presenting cells**

44. Ans. (b) **B cells, Dendritic cells, NK cells**

(Ref: R 9/p 191)

45. Ans. (a, c); **a. Langerhan cells of skin; c. Macrophages**

46. Ans. (d) **C56789**

(Ref: Robbins and Cotran: Pathological basis of disease 8/e p64,89)

The deposition of the MAC (C5-C9) on cells makes these cells permeable to water and ions and results in death (lysis)



47. Ans. (a) **RBC**

As RBC lacks nucleus so lacks HLA

48. Ans. (b) **Thymocyte** (Ref: Robbins 9th / pg 195-196)

Actually thymic epithelial cells are Antigen presenting cells not thymocytes

About M cells (in gut mucosa) or microfold (M) cells

- Unique morphological features include:
  - Presence of a **reduced glycocalyx**
  - Irregular brush border and reduced microvilli.
  - Highly specialized for the **phagocytosis and transcytosis** of gut lumen macromolecules, particulate antigens and pathogenic or commensal microorganisms across epithelium

49. Ans. (b) **HBV**

(Ref: Harrison 19th/2010; J Med Virol. 2016 Mar; 88(3):371-9)

This Question is based on the concept that HBV-HLA-specific cytolytic T cell responses of the adaptive immune system are felt to be responsible for recovery from HBV infection

The observed findings from various studies are:

- HLA-DR\*03 and HLA-DR\*07 were associated with an increased risk of persistent HBV infection
- HLA-DR\*04 and HLA-DR\*13 were associated with clearance of HBV infection.

50. Ans. (b) **MHC 2** (Ref: Robbins 9th/pg 192; 8th/pg 187)

- **Dendritic cells** are the **most important antigen-presenting cells (APCs) for initiating T-cell responses against protein antigens**<sup>Q</sup>, they express MHC-II

51. Ans. (a) **TNF alpha** (Ref: Robbins 9th/pg 195; 8th/pg 191)

MHC III: No direct role in immune system<sup>Q</sup>

- Codes for:
- **complement components** C2, C4, properdin, factor B<sup>Q</sup>
- TNF, HSP-70, Tyrosine hydroxylase

52. Ans. (b) **Dendritic cells** (Ref: Robbins 9th/pg 192)

53. Ans. (a) **Short arm of chr-6** (Ref: Robbins 9th/pg 195)

54. Ans. (c) **Class III codes for complement**

(Ref: Robbins 9th/pg 195)

55. Ans. (c) **Present antigens for recognition by T cell antigen receptors** (Ref: Robbins 9th/pg 195; 8th/pg 191)

HLA I and II display **peptide fragments** of protein antigens for recognition by antigen-specific T cells.

56. Ans. (d) **Neutrophils**

(Ref: Robbins 9th/pg 195; 8th/pg 191)

57. Ans. (a) **Macrophage; b. Dendritic cells; d. Epithelial cell** (Ref: Robbins 9th/pg 195-196; 8th/pg 191-192)

58. Ans. (b) **IL-4**

(Ref: Robbins 9th ed/pg 201)

59. Ans. (b) **IgE fixed cell**

(Ref: Robbins 9th ed/pg 201)

60. Ans. (c) **IL-4**

(Ref: Robbins 9th ed/pg 201)

61. Ans. (c) **IL-3**

(Ref: Robbins 9th ed/pg 198)

62. Ans. (b) **IL-2**

(Ref: Robbins 9th ed/pg 198)

63. Ans. (a) **CD4+ helper cell** (Ref: Robbins 9th ed p 1209)

CD4+ T helper (TH) cells may initiate the autoimmune response in Rheumatoid arthritis by reacting with an arthritogenic agent, perhaps microbial or a self-antigen. The T cells produce cytokines that stimulate other inflammatory cells to effect tissue injury.

64. Ans. (b) **Ag Ab reaction** (Ref: Robbins 9th/205-206)

65. Ans. (a) **Mast cell** (Ref: Robbins 9th/pg 200; 8th/pg 198)

Mast cells have Fc receptor for IgE; Antigen binding to it results in the activation of mast cells leading to type I hypersensitivity.

66. Ans. (b) **Hypersensitivity reaction with modified macrophages, lymphocytes and giant cells**

(Ref: Robbins 9th/pg 208-211; 8th/pg 204-208)

The given clinical features and radiological findings are suggestive of Tuberculosis in which histopathology of the affected organ shows granuloma, that consists of modified macrophages, lymphocytes and giant cells.

67. Ans. (b) **CD4 + T cells**

(Ref: Robbins 9th/pg 208; 8th/pg 204)

68. Ans. (a) **None of the options** (Ref: Robbins 9th/pg 208)

Shwartzman reaction is a **non-immunologic** phenomenon in which **endotoxin (lipopolysaccharide)** induces local & systemic reactions; It can either be local or systemic type.

69. Ans. (b) **Tumor necrosis factor**


(Ref: Robbins 9th/pg 198)

- **TNF and IL-1** are two of the major cytokines that mediate inflammation & are produced mainly by **activated macrophages**.

70. Ans. (b) **Type II**

(Ref: Textbook of Microbiology & Immunology by Subhash Chandra Parija, 2nd edition, 2012, pg 155)

- **Type V hypersensitivity** reaction is a **subtype of type II** hypersensitivity
- In type V hypersensitivity, antibodies combine with antigens on cell surface which induce cells to proliferate & enhances activity of effector cells; **E.g., Graves disease**.



**71. Ans. (b) Eosinophils**

(Ref: Robbins 9th/pg 200; 8th/pg 198)

**72. Ans. (b) Major Basic Protein**

(Ref: Robbins 9th/pg 200-204)

Out of the given options, most **important mediator of late phase of immediate type of hypersensitivity reaction** is Major basic protein.

**73. Ans. (a) IgG** (Ref: Robbins 9th/pg 205-206)

- Antibodies in Immune thrombocytopenic purpura are of IgG type directed against gp Ib/IX and IIb/IIIa.
- These antibodies cause Type II hypersensitivity reaction.

**74. Ans. (b) Type II hypersensitivity**

(Ref: R 9th/pg 205-206)

Transfusion reaction and erythroblastosis fetalis are examples of Type II (antibody) mediated reactions.

**75. Ans. (b) Type II HSN** (Ref: Robbins 9th/pg 205-206)

In the question, there are some clues to the diagnosis:

- Reaction **started 48 hrs after** taking penicillin: Rules out Immediate type I HSN
- **No prior history of allergy**: Again rules out Immediate HSN
- **Anti-penicillin antibody**: Shows Antibody (Humoral) type of reaction
- There are two type of HSN based on Antibodies: Type II and Type III
- **Hemolysis occurs in Type II HSN**
- So, clearly the answer is Type II HSN

**76. Ans. (d) Rh incompatibility**

(Ref: Robbins 9th/pg 205-206)

Examples of Antibody-Mediated Diseases (Type II Hypersensitivity) Refer to preface of this chapter

**77. Ans. (b) Angioneurotic edema** (Ref: R 9th/pg 200-204)

Hereditary angioedema or Angioneurotic edema

- **Autosomal dominant** disorder due to an underlying **deficiency of C1 inhibitor**
- C1 inhibitor is a protease inhibitor whose target enzymes are C1r & C1s of complement cascade, factor XII of the coagulation pathway, and the kallikrein system.
- Patients have episodes of edema affecting skin and mucosal surfaces such as larynx & GIT, swelling of lips
- May result in life-threatening asphyxia or nausea, vomiting & diarrhea after minor trauma or emotional stress.

**78. Ans. (d) IV** (Ref: Harrison 18th/chapter 166)

Type I Lepa Reactions (Downgrading and Reversal Reactions)

- When type I lepra reactions **precede the initiation of antimicrobial therapy**, they are termed 'downgrading' reactions, and the case becomes histologically more lepromatous;

- When they occur **after the initiation of therapy**, they are termed '**reversal**' reactions, and the case becomes more tuberculoid.

- **Edema** is the most characteristic microscopic feature of type I lepra lesions

- **Reversal reactions** are typified by a TH1 cytokine profile, with an influx of **CD4+ T helper cells** and increased levels of **IFN-γ & IL-2**; So it is an example of **type IV hypersensitivity**

Type 2 Lepa Reactions: Erythema Nodosum Leprosum

- It is **immune complex mediated (type III hypersensitivity)**
- Occurs exclusively in patients near the **lepromatous** end of the leprosy spectrum (**BL-LL**)

**79. Ans. (b) Type II Hypersensitivity**

(Ref: R 9th/pg 205-206)

**80. Ans. (b) IgG Ab** (Ref: Robbins 9th/pg 200-204)

**Long-acting thyroid stimulator (LATS)**: IgG that stimulates thyroid function similar to but slower than TSH (i.e. long-acting)

**81. Ans. (a) Histamine** (Ref: Robbins 9th/pg 200-204)

Histamine can cause Anaphylactic shock

**82. Ans. (b) Immune complexes**

Raji cell assay:

- A sensitive test for **detection & quantitation of soluble complement fixing immune complexes** in sera of patients.
- **Raji cells lack membrane-bound immunoglobulin** but have **receptors for IgG Fc**, AHG is used as an in vitro model of human immune complexes.
- The uptake by Raji cells is quantitated by **I<sup>125</sup>-radiolabeled antihuman IgG**.

**83. Ans. (a, b, c, d, e) a. Interferon-1 (IFN-1); b. Interferon-1 (IFN-1); c. Exposure to UV rays; d. Deficiency of early complement factors; e. Failure of self-tolerance in B cells** (Ref: Robbins 9th ed/pg 218)

**84. Ans. (b) Raynaud phenomenon and sclerodactyly**

(Ref: R9/p 226)

Patterns in Anti-Nuclear Antibody (ANA) testing

Diffuse or homogenous	dsDNA, histone	SLE, DLE
Rim/peripheral	Ds DNA	SLE
Speckled	Sm	SLE
	SS-A, SS-B	Sjogren syndrome
Nucleolar	Nucleolar RNA	Scleroderma

**85. Ans. (b) AIRE** (Ref: R 9/p 213)

The process in the questions refers to Central tolerance, where the self reacting T cells are presented to the



thymus to prevent autoimmunity. A protein called AIRE (autoimmune regulator) stimulates expression of few “peripheral tissue-restricted” self antigens in the thymus. This causes deletion of immature T cells specific for these antigens and so is able to prevent autoimmunity.

**86. Ans. (b) AIRE** (Ref: Robbins 9th/216)

Mutations in the AIRE gene are the cause of an autoimmune-polyendocrinopathy

**87. Ans. (b, c) b. Histoplasmosis c. Cryptococcosis**

(Ref: Harshmohan 7th/155; Robbins (SEA) 9th/ 693)

Schaumann bodies are calcium and protein inclusions inside of Langhans giant cells as part of a granuloma.

Seen in:

- Sarcoidosis,
- Hypersensitivity pneumonitis, and
- Berylliosis.
- Crohn's disease and tuberculosis

**88. Ans. (b) Seminiferous tubules**

(Ref: Robbins 9th/pg 214)

**89. Ans. (a) Ulcerative colitis** (Ref: Robbins 9th/pg 214-215)

**90. Ans. (a) Anti-nuclear** (Ref: Robbins 9th/pg 218-221)

**91. Ans. (d) Anti-histone** (Ref: Robbins 9th/pg 218-221)

**92. Ans. (b) Erosive arthritis** (Ref: Robbins 9th/pg 218)

Non-erosive arthritis is a feature of SLE

**93. Ans. (c) Lip** (Ref: Robbins 9th/pg 218; 8th/pg 213)

**94. Ans. (a) Anti-DNA topoisomerase**

(Ref: Robbins 9th/pg 218)

**95. Ans. (d) Anti-Ribonucleoprotein**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

Autoantibodies in Sjogren's syndrome:

- **Anti SS-A (Ro) and SS-B (La)**<sup>Q</sup>: Most important, present in 90% patients;
- High titers of Anti SS-A → more likely to have **early disease onset, longer disease duration, and extra-glandular manifestations** (eg cutaneous vasculitis and nephritis)
- **Rheumatoid factor** (an antibody reactive with self IgG): in 75%<sup>Q</sup>, **ANA**: in 50% to 80%

**96. Ans. (d) Diffuse sclerosing lupus nephritis**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

**97. Ans. (a) Systemic lupus erythematosus**

(Ref: R 9th/pg 218)

**98. Ans. (c) Systemic lupus erythematosus**

(Ref: Robbins 9th/pg 218)

**99. Ans. (b) Diffuse Proliferative** (Ref: Robbins 9th/pg 218)

**100. Ans. (b) Systemic lupus erythematosus**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

**101. Ans. (b) Anti-Ro (SS-A)** (Ref: Robbins 9th/pg 218)

**102. Ans. (c) Diffuse proliferative glomerulonephritis**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

**103. Ans. (b) Drug induced lupus** (Ref: Robbins 9th/pg 218)

**104. Ans. (c) Erosions seen on X-rays**

(Ref: Robbins 9th/pg 218)

**105. Ans. (a) Anti-Topoisomerase I antibody**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

**106. Ans. (d) Nephritis**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

Nephritis is usually the most serious manifestation of SLE, particularly because nephritis and infection are the leading causes of mortality in the first decade of disease.

**107. Ans. (a) More common in females**

(Ref: Robbins 9th/pg 218; 8th/pg 213)

### Drug-Induced Lupus


- It is a syndrome of positive ANA associated with symptoms such as fever, malaise, arthritis or intense arthralgias/myalgias, serositis, and/or rash.
- It appears during therapy with certain medications like antiarrhythmics procainamide, disopyramide, and propafenone; antihypertensive hydralazine; antithyroid propylthiouracil; antipsychotics chlorpromazine and lithium; anticonvulsants carbamazepine and phenytoin; antibiotics isoniazid, minocycline & nitrofurantoin; antirheumatic sulfasalazine; diuretic hydrochlorothiazide;
- It is predominant in whites & has less female predilection than SLE
- It rarely involves kidneys or brain, is rarely associated with anti-dsDNA, is commonly associated with antibodies to histones
- It usually resolves over several weeks after discontinuation of the offending medication.

**108. Ans. (b) Rheumatoid arthritis** (Ref: Robbins 9th/pg 218)

**109. Ans. (a) Esophagus** (Ref: Robbins 9th/pg 218; 8th/pg 213)

**110. Ans. (c) Limited scleroderma** (Ref: Robbins 9th/pg 228)





111. Ans. (b) **pqrs = ywzx**

(Ref: Robbins 9th/pg 228; 8th/pg 223)

112. Ans. (c) **Rheumatoid arthritis** (Ref: Robbins 9th/pg 231)

Mixed connective tissue disease

- Clinical features are a mixture of the features of **SLE, systemic sclerosis, and polymyositis**.
- Elevated **anti-U1 ribonucleoprotein Ab<sup>q</sup>**
- Presentation: Synovitis of fingers, Raynaud phenomenon, mild myositis, renal involvement

113. Ans. (a) **Anti U1-RNP Ab** (Ref: Robbins 9th/pg 231)

114. Ans. (b) **SLE** (Ref: Dubois' Lupus Erythematosus pg 542)

Lupus Band test

- A diagnostic procedure used to detect deposits of **immunoglobulins and complement** components along the **dermo-epidermal junction** in patients with SLE & cutaneous lupus
- It is **positive in about 70%–80% of sun-exposed** & 55% of sun-protected **non-lesional skin specimens** obtained from patients with SLE

115. Ans. (a) **Anti ds DNA** (Ref: Robbins 9th/pg 218-221)

116. Ans. (c) **HLA-CW6** (Ref: Robbins 9th/pg 215)

Role of Susceptibility Genes: Association of HLA Alleles with Disease; Refer to pretext of this chapter

Disease	Marker
<ul style="list-style-type: none"> <li>• Ankylosing spondylitis<sup>q</sup></li> <li>• Reiter's syndrome<sup>q</sup></li> <li>• Acute anterior uveitis</li> <li>• Reactive arthritis<sup>q</sup></li> <li>• Psoriatic spondylitis</li> </ul>	<b>B27<sup>q</sup></b> <b>B27<sup>q</sup></b> B27 <b>B27<sup>q</sup></b> B27
<ul style="list-style-type: none"> <li>• Juvenile arthritis, pauciarticular</li> <li>• Rheumatoid arthritis<sup>q</sup></li> <li>• Sjögren's syndrome<sup>q</sup></li> </ul>	DR8, DR5 <b>DR4 (DRB1*04)<sup>q</sup></b> <b>DR3<sup>q</sup></b>
<ul style="list-style-type: none"> <li>• (celiac disease)<sup>q</sup></li> <li>• Chronic active hepatitis</li> <li>• Dermatitis herpetiformis</li> <li>• Psoriasis vulgaris</li> <li>• Pemphigus vulgaris</li> </ul>	<b>DQ2/DQ8<sup>q</sup></b> DR3 DR3 <b>Cw6<sup>q</sup></b> <b>DR4, DQ1<sup>q</sup></b> DQ7
<ul style="list-style-type: none"> <li>• Type 1 diabetes mellitus</li> <li>• Hyperthyroidism (Graves')</li> <li>• Adrenal insufficiency</li> </ul>	<b>DQ8, DR4, DR3,</b> <b>DR2<sup>q</sup></b> B8, DR3 DR3
<ul style="list-style-type: none"> <li>• Myasthenia gravis</li> <li>• Multiple sclerosis</li> </ul>	<b>B8, DR3<sup>q</sup></b> DR2
<ul style="list-style-type: none"> <li>• Behçet's disease</li> <li>• Congenital adrenal hyperplasia</li> <li>• Good-pasture's syndrome</li> <li>• Thromboangitis obliterans</li> </ul>	B51 B47 <b>DR2<sup>q</sup></b> B5

117. Ans. (d) **Anti-histone antibody** (Ref: R 9th/pg 218-221)

118. Ans. (c) **HLA-B51** (Ref: Robbins 9th/pg 215)

119. Ans. (b) **HLA-DR4** (Ref: Robbins 9th/pg 215)

HLA-DR4 (DR4) is an HLA-DR serotype that recognizes the DRB1\*04 gene products.

120. Ans. (c) **HLA-B5** (Ref: Robbins 9th/pg 215)

121. Ans. (b) **RA** (Ref: Harrison 18th/chapter 321, 323)

Antibodies seen in RA are **rheumatoid factors (RFs)** and **anti-cyclic citrullinated peptides (CCP)** antibodies  
ANA is seen in:

- Almost **all (>95%) cases of SLE**
- In virtually **all patients with SSc** (Systemic sclerosis)
- **50% to 80% cases of Sjogren syndrome**

122. Ans. (d) **Ankylosing spondylitis**

(Ref: Harrison 18th/chapter 325)

Prevalence of HLA B27 is **90% in patients with Ankylosing Spondylitis**, independent of disease severity.

123. Ans. (b) **Antibodies to GQ1B**

(Ref: Harrison 18th/chapter 385)

Miller Fisher syndrome (MFS)

- It is a **variant of GBS**, which presents as **rapidly evolving ataxia and areflexia** of limbs **without weakness**
- **Ophthalmoplegia**, often with **pupillary paralysis** is present.
- Accounts for **5% of all cases of GBS (Guillain Barre syndrome)**
- Is strongly associated with **antibodies to the ganglioside GQ1b**

124. Ans. (c) **Class III** (Ref: Robbins 9th/pg 222-223)

**Focal proliferative glomerulonephritis** (class III lupus nephritis)

- It is seen in **20-35% of patients**, and is defined by **fewer than 50% involvement of all glomeruli**.
- Lesions **may be segmental** (affecting only a portion of glomerulus) or **global** (involving entire glomerulus).
- Affected glomeruli may exhibit **crescent formation, fibrinoid necrosis, proliferation of endothelial & mesangial cells**, infiltrating **leukocytes**, and **eosinophilic deposits** or **intracapillary thrombi**, which often correlate with **hematuria and proteinuria**.

125. Ans. (a) **Anti-ds-DNA** (Ref: Robbins 9th/pg 218-221)

The given clinical features are suggestive of SLE;

- **Specific test** for diagnosis of SLE is **anti ds DNA**;
- This antibody also **correlates with disease activity, nephritis, vasculitis**.
- So Anti-dsDNA should be initially assayed in this case.



**126. Ans. (a, b, d); a. Proliferation of endothelium; b. Glomerular leukocyte infiltration; d. Epithelial cell proliferation (Ref: Harrison 18th/chapter 319)**

Active nephritis in SLE includes:

- **Focal proliferative** lupus nephritis
- **Diffuse segmental proliferative** lupus nephritis &
- **Diffuse global proliferative** lupus nephritis
- So, **proliferation of endothelium, glomerular leukocyte infiltration & epithelial cell proliferation** are features of active nephritis;
- Mesangial alterations may or may not be present;
- **Tubulitis is not a feature of lupus nephritis;**

**127. Ans. (a, b, c, d); a. Interstitial Fibrosis; b. Shrinking lung syndrome; c. Alveolar hemorrhage; d. Pulmonary arterial hypertension**

(Ref: Robbins 9th/pg 218; 8th/pg 214) Refer to pretexts.

**128. Ans. (b, d) b. Antibody to RNA; d. Systemic sclerosis**

(Ref: Robbins 9th/pg 219; 8th/pg 215)

**Antinuclear antibodies:**

- They are **directed against nuclear antigens**.
- The most widely used method for detecting ANAs is **indirect immunofluorescence**, which can identify antibodies that bind to a variety of nuclear antigens, including **DNA, RNA, and proteins**
- The **pattern of nuclear fluorescence suggests the type of antibody present** in the patient's serum.

**Four basic patterns** are recognized:

Staining Pattern	Antibodies & their Characteristics
Homogeneous or diffuse nuclear staining	Antibodies to <b>chromatin, histones &amp; occasionally, double-stranded DNA</b> .
Rim or peripheral staining	Antibodies to <b>double-stranded DNA</b>
Speckled pattern	Presence of uniform or variable-sized speckles. Antibodies to <b>non-DNA nuclear constituents</b> . Eg Antibody to <b>Sm antigen, ribonucleoprotein, SS-A &amp; SS-B antigens</b> .
Nucleolar pattern	Presence of a few discrete <b>spots of fluorescence within the nucleus</b> Represents <b>antibodies to RNA</b> Seen most often in patients with <b>systemic sclerosis</b> .

**129. Ans. (b) Lymphocytes (Ref: Robbins 9th/pg 226-227)**

**130. Ans. (a, b, e) a. DRB1; b. DR1; e. DR4**

(Ref: Harrison 18th/chapter 321)

In **Rheumatoid Arthritis**, some of the **HLA-DRB1 alleles** bestow a **high risk of disease (\*0401)**, whereas others confer a more **moderate risk (\*0101, \*0404, \*1001 & \*0901)**. **HLA-DR4 (DR4)** is a **HLA-DR serotype** that recognizes the **DRB1\*04 gene products**.

**131. Ans. (b) Anti-ds DNA antibodies**

(Ref: Harrison 18th/chapter 319)

- **Anti dsDNA is specific for SLE**, only in **high titers**;
- But still, **anti ds DNA is the best marker for SLE** because its **prevalence in SLE is 70%**;
- Whereas, **Anti Sm antibody**, which is **specific for SLE**, but is **seen in only 25% patients with SLE**

**132. Ans. (c) Isograft**

**133. Ans. (c) Allograft**

Transplantation between same species is allograft

**134. Ans. (d) Donor T cells**

Immunocompetent Donor T cells destroy immunosuppressed recipient cells in GVHD

**135. Ans. (a) 12 hours (Ref: Robbins 9th ed p 233)**

Hyperacute rejection occurs when preformed antidonor antibodies are present in the circulation of the recipient. This form of rejection occurs within minutes or hours after transplantation. A hyperacutely rejecting kidney rapidly becomes cyanotic, mottled, and flaccid, and may excrete a mere few drops of bloody urine. Immunoglobulin and complement are deposited in the vessel wall, causing endothelial injury and fibrin-platelet thrombi.

**136. Ans. (d) Transfer hepatic progenitor cells (HPCs) of same person for regeneration**

(Ref: Nature Cell Biology 17, 971-983 (2015))

Hepatocytes and cholangiocytes self-renew following liver injury. But after severe injury like **cirrhosis** and acute liver failure hepatocytes are increasingly senescent. In this case there are two options: first diseased-donor liver and living-donor transplantation, in which a just portion of a healthy donor's liver is used for transplantation or second hepatic progenitor cells (HPCs), which contribute significantly to restoration of liver parenchyma, regenerating hepatocytes and biliary epithelia, highlighting their *in vivo* lineage potency. The best option is D among the options.

**137. Ans. (a) Irradiation (Ref: Wintrob's 13th/704-705)**

- Irradiation with a dose of 2,500 cGy at the center of the irradiation field, with a minimum dose of 1,500 cGy at any point in the field **inhibits proliferation of donor lymphocytes** but has no significant adverse effect on red cell, platelet, or granulocyte function.
- It should be noted that though the other options are also effective in reducing the leukocytes but they do not completely deplete the lymphocytes and so can cause GVHD.

**138. Ans. (d) C4d (Ref: Robbins 9th/234)**

**Acute antibody-mediated rejection** is manifested mainly by damage to glomeruli and small blood vessels. Typically, the lesions consist of inflammation of glomeruli and peritubular capillaries, associated with deposition of the complement breakdown product C4d, which is produced during activation



of the complement system by the antibody dependent classical pathway

**139. Ans. (d) HLA DP** (Ref: Harrison 18th/chapter 282)

- In kidney transplants, polymorphic HLA alleles are at least 50 % matched (HLA-A, -B & DR)

**140. Ans. (c) Biopsy** (Ref: Harrison 18th/chapter 282)

**141. Ans. (b) Matched, related donors**

(Ref: AIIMS, Transplant immunology Protocol; Kidney transplant principle & practise pg 156)

**Important terminologies for Donors in transplant immunology:**

- **Matched Donor:** HLA allelic identity at HLA A, B, C and DRB1 loci which is referred as 8/8match. (These 4 HLA antigens have 2 alleles each that makes it 8)
- **Related Donor:** Siblings from same parents who is HLA matched for HLA A, B, DRB1 minimum at the antigen level.

So obviously, the answer here is matched related donor.

**142. Ans. (b) 5-10**

(Ref: AIIMS, Transplant immunology Protocol)

For different types of transplantation, different criteria are followed for HLA matching:

Organs	Minimum Matching	Matching done at
Bone Marrow	100% (8/8)	A,B,C DRB1
Kidney	50% (3/6)	A,B,DR
Liver/Heart	0% (0/6)	—
Cornea	0% (0/6)	—

**Note:** In liver, heart transplantation; practical considerations (ischemic times, availability of donors, clinical need of recipients) makes HLA matching less important.

- **HLA-DR mismatches** are the **most important** in the **first 6 months** after transplantation.
- **HLA-B** effect emerges in the **first 2 years**.
- **HLA-A** mismatches have a deleterious effect on **long-term graft survival**.

**143. Ans. (b) Renal involvement** (Ref: Robbins 9th/pg 231)

**144. Ans. (c) Matched**

(Ref: AIIMS, Transplant immunology Protocol)

Best tissue matches are HLA A, B, DR identical referred to as “000 mismatch” (100% matching which is 6/6 match). Any mismatch is referred to as 1 like “100, 010, 110”

**145. Ans. (c) Blood vessel thrombosis**

(Ref: Robbins 9th/pg 231; 8th/pg 221); Refer to pretexts

**146. Ans. (c) Both** (Ref: Robbins 9th/pg 231-234)

**147. Ans. (a) All nucleated cells** (Ref: Robbins 9th/pg 195)

Properties	Class I MHC Molecules	Class II MHC Molecules
Location	All nucleated cells and platelets <sup>a</sup>	APCs: B cells, dendritic cells, endothelial cells & fibroblasts <sup>a</sup>

**148. Ans. (b) T-lymphocyte** (Ref: Robbins 9th/pg 189)

**149. Ans. (d) Zero HLA mismatch with recipient**

(Ref: Harrison 18th/chapter 282)

**Donor for kidney transplantation:**

- Donors can be **deceased or volunteer living donors**.
- Living donors are usually family members who have **at least partial compatibility for HLA** antigens.
- **At least 50% HLA match (3/6) is a must requirement** for kidney transplant
- Living volunteer donors should be **normal on physical examination**
- Should have the **same major ABO blood group**, because crossing major blood group barriers prejudices survival of the allograft.
- Should have **two normally functioning kidneys**
- Donor **should be free of malignant neoplastic disease, hepatitis, and HIV**

**150. Ans. (b) Delayed type hypersensitivity**

(Ref: Robbins 9th/pg 231-234; 8th/pg 221-229; Refer Ans 200)

**Types of Rejection Pathway:**

- **Direct pathway:** **CD4 and CD8-T** cells of the transplant recipient recognize allogeneic (donor) MHC molecules on the surface of APCs in the graft.
- **Indirect pathway:** recipient **CD4-T lymphocytes** recognize MHC antigens of the graft donor after they are presented by the recipient's own APCs.

As both the pathways are **cell mediated**, hence they are **delayed hypersensitivity** reactions.

**151. Ans. (a) Preformed antibodies** (Ref: R 9th/pg 231-234)

**152. Ans. (d) Kidney** (Ref: Robbins 9th/pg 236; 8th/pg 230)

**In Acute GVH disease:**

Immune system and epithelia of the skin, liver & intestines are mainly involved.

**153. Ans. (b) Isograft** (Ref: Robbins 9th/pg 231-234)

**Types of grafts<sup>o</sup>**

- **Autografts<sup>o</sup>:** transplant of individual own organ
- **Isograft:** Graft from identical twin
- **Allografts<sup>o</sup>:** between individuals of the same species
- **Xenografts<sup>o</sup>:** grafts from one species to another species

**154. Ans. (b) Lungs** (Ref: Robbins 9th/pg 236)

**155. Ans. (c) Hyper IgM syndrome**

(Ref: Robbins 9th/pg 241; Wintrobe 13th/pg 342)

**156. Ans. (d) BTK**

(Ref: Robbins 9th/pg 239)



**157. Ans. (a) Uveitis**

**158. Ans. (d) Chediak-Higashi syndrome**

(Ref: Robbins 9th/pg 237)

**159. Ans. (a, b, d) a. Autosomal recessive; b. Part of SCID; d. Leads to abnormal CD4T cell development**

Bare lymphocyte syndrome is a condition caused by mutations in certain genes of the major histocompatibility complex or involved with the processing and presentation of MHC molecules, so there is abnormal T cell development. It is a form of severe combined immunodeficiency

**160. Ans. (d) B cell defect (Ref: R 9TH/P 24)**

DiGeorge syndrome is a T-cell deficiency that results from failure of development of the third and fourth pharyngeal pouches.

**161. Ans. (a) Non homologous disjunction repair**

(Ref: Wintrob's 14th/pg 333)

DNA-PK (DNA-dependent protein kinase) complex consists of a catalytic subunit (DNA-Pkcs) and a DNA-binding complex called Ku, which binds altered DNA structures such as double-strand breaks, nicks, or hairpin loops. Recombination of broken DNA strands occurs either between strands that have long stretches of homology (homologous recombination) or between DNA strand breaks without relying on the presence of considerable homology between them. This recombination is known as nonhomologous end joining (NHEJ). Mutations in this system result in the accumulation of V(D)J-specific double-strand breaks, indicating a defective repair mechanism, which causes SCID.

**162. Ans. (a, b, c, d) a. Severe combined immunodeficiency; b. Adenosine deaminase deficiency; c. Wiskott-Aldrich syndrome; d. Ataxia telangiectasia**

**163. Ans. (c) Both**

(Ref: Robbins 9th/pg 239-240; 8th/pg 234-235)

Defect in Common Variable Immunodeficiency is mainly Defect in receptor for a cytokine called BAFF causing defect in survival and differentiation of B cells defect in T-cell activation & interactions b/w T & B cells

**164. Ans. (b) SCID**

(Ref: HENRY'S Clinical Diagnosis and Management by Laboratory Methods; 22nd ed/ pg 944-945)

Lymphocyte phenotyping for assessing immune deficiency includes CD3 (pan T cell), CD4 (T helper subset), CD8 (T cytotoxic/suppressor subset), CD19 (pan B cell), CD 16/56 (pan NK cell) and other cell surface markers as required based on clinical data. This analysis is indicated in diagnosis and monitoring of immunodeficiencies and immunotherapy in children and adults

**165. Ans. (c) GVHD (Ref: Robbins 9th/pg 236; 8th/pg 230)**

- Patients with SCID have the most severe immunodeficiency.

- Affected infants also lack the ability to reject foreign tissue and are therefore at risk for severe or fatal graft-versus host disease (GVHD) from T lymphocytes in non-irradiated blood products or in allogeneic stem cell transplants.

**166. Ans. (a) SCID; c. Wiskott-Aldrich syndrome**

(Ref: Robbins 9th/pg 239-240; 8th/pg 234-235)

*Mucocutaneous candidiasis is seen in DiGeorge Syndrome (Thymic Hypoplasia), which is a predominant T cell defect;*

**167. Ans. (a) Neutrophilia**

(Ref: Nelson 19 pg 744; Robbins 9th/pg 238; 8th/pg 55)

**Neutropenia rather than neutrophilia is seen in Chediak-Higashi Syndrome**

**168. Ans. (d) Cryptosporidium**

**169. Ans. (b) Enfuvirtide is not active against HIV-2, (d) More commonly transmitted from mother to child, (e) HIV-1 worsens faster than HIV-2**

(Ref: Robbins 9th/pg 243)

In the United States, the first case of HIV-2 infection was diagnosed in 1987 in a West African woman who presented with central nervous system toxoplasmosis

**Difference between HIV-1 and 2: HIV-2 infection –**

- **Not transmitted as efficiently** as HIV-1<sup>o</sup>
- Progression to AIDS **takes longer**
- **Slower rate of CD4 cell decline** and viral replication
- Rarely causes vertical transmission

**170. Ans. (a) Kaposi's sarcoma (Ref: Robbins 9th/pg 253)**

**171. Ans. (a) 1983 (Ref: Robbins 9th/pg 243; 8th/pg 236)**

**History of HIV:**

Year	HIV linked developments
1981	AIDS was first recognized in the United States by U.S. Centers for Disease Control and Prevention (CDC).
1983	HIV isolated from a patient with lymphadenopathy (HIV 1 <sup>st</sup> discovered)
1984	HIV demonstrated as causative agent of AIDS
1985	Enzyme-linked immunosorbent assay (ELISA) was developed to diagnose HIV

**172. Ans. (a) CMV (Ref: Robbins 9th/pg 246; 8th/pg 239)**

In HIV infection with:

CD4+ T cell counts <200/uL → high risk of disease from *P. jiroveci*

CD4+ T cell counts <50/uL → high risk of disease from CMV, *M. avium* complex (MAC), and/or *T. gondii*

*Intramucular parophyllic inclusion is seen with CMV*

**173. Ans. (d, e); d. Primary lymphoma of brain; e. Invasive cancer of uterine cervix (Ref: Robbins 9th/pg 253)**





The neoplastic diseases considered to be AIDS-defining conditions are **Kaposi's sarcoma, non-Hodgkin's lymphoma, and invasive cervical carcinoma.**

In addition, there is also an increase in the incidence of a variety of **non-AIDS-defining malignancies:**

- Hodgkin's disease;
- Multiple myeloma;
- Leukemia;
- Melanoma; and
- Cervical, brain, testicular, oral, lung, gastric, liver, renal, and anal cancers

**174. Ans. (b) Helper T cells** (Ref: Robbins 9th/pg 246-249)

gp120 receptors on HIV binds to host cell receptors (CD4) on<sup>Q</sup>

- T-Lymphocytes (helper T cells)
- Macrophages
- Dendritic cells
- Other Antigen presenting cells

**175. Ans. (d) Both lost** (Ref: Robbins 9th/pg 246)

**176. Ans. (a) Lentivirus** (Ref: Robbins 9th/pg 246; 8th/pg 240)

**177. Ans. (a) HIV 2 is more pathogenic than HIV 1**

(Ref: Robbins 9th/pg 246; 8th/pg 240)

**178. Ans. (e) Ankylostoma duodenale infection**

(Ref: Harrison 19th/pg 1215)

**179. Ans. (a) HIV-1M** (Ref: Robbins 9th/pg 246-249)

Causative organism: HIV (*Human immunodeficiency virus*)

- HIV-1 group M subtypes C predominant in India, fastest spreading<sup>Q</sup>
- HIV-1 group M subtypes B predominant in U.S

**180. Ans. (a, b, c, d); a. Hepatosplenomegaly; b. Congestive heart failure; c. Proteinuria; d. Lytic bone lesions**

(Ref: Robbins 9th/pg 257-60)

**181. Ans. (a, b, c, d); a. True, b. True, c. True, d. False, it shows wild type of ATTR.**

(Ref: Robbins 9th/pg 257-60)

**182. Ans. (a, b, c, d.) a. Bronchiectasis; b. Pulmonary TB; c. Lung abscess; d. Malignancy**

**183. Ans. (d) Amyloid fibrils**

(Ref: Histopathology Guy Orchard, Brian Nation ; pg 70)

Congo Red dye forms non-polar hydrogen bonds with amyloid and red to green birefringence occurs when viewed by polarised light due to parallel alignment of the dye molecules on the linearly arranged amyloid fibrils.

**184. Ans. (c) Apple green birefringence on polarizing microscopy** (Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

**185. Ans. (b) Congo red** (Ref: Robbins 9th/pg 257-260)

**186. Ans. (c) A $\beta$ 2**

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

**187. Ans. (c) Mutant form transthyretin is deposited**

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

**188. Ans. (d) Rectum** (Ref: Robbins 9th/pg 257-260)

Sites of biopsy for diagnosis of Amyloidosis:

- Rectal biopsy-Most specific and best site
- Abdominal fat aspirate
- Gingival biopsy
- Organ specific biopsy for localized amyloidosis

**189. Ans. (a) Normal transthyretin** (Ref: R 9th/pg 257-260)

This is an example of senile amyloidosis in which unmutated (normal) transthyretin is deposited.

**190. Ans. (c) Amyloidosis** (Ref: Robbins 9th/pg 257-260)

**191. Ans. (a) Congo red + polarized microscopy**

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

**192. Ans. (b) This electron microscope structure is not identical in all types of amyloid.**

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

**193. Ans. (d) Medullary** (Ref: Robbins 9th/pg 260; 8th/pg 253)

Microscopically, medullary carcinomas:

- Composed of polygonal to spindle-shaped cells, which may form nests, trabeculae, and even follicles.
- Acellular **amyloid deposits**, derived from altered calcitonin polypeptides, are present in the adjacent stroma in many cases.
- Calcitonin is readily demonstrable within the cytoplasm of the tumor cells as well as in the stromal amyloid by immunohistochemical methods.

**194. Ans. (c) Congo red** (Ref: Robbins 9th/pg 257-260)

**195. Ans. (c) Amyloidosis** (Ref: Robbins 9th/pg 257-260)

**196. Ans. (a) AL** (Ref: Robbins 9th/pg 257-260)

- AL protein is made of complete **Ig light chains**, amino-terminal fragments of light chains, or both.
- Most of the AL proteins are composed of  $\lambda$  light chains or their fragments, but in some cases  $\kappa$  chains
- The amyloid fibril protein of the AL type is produced from free Ig light chains secreted by a monoclonal population of plasma cells, like multiple myeloma.

**197. Ans. (b) Made of calcified proteins**

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

- **Amyloid is a pathological proteinaceous substance deposited between cells** in various tissues and organs of the body in a variety of clinical settings.



In polarized light, amyloid shows Apple-green birefringence on Congo Red staining

- **Chronic infections like Tuberculosis & Osteomyelitis may show Amyloidosis**

---

**198. Ans. (a) Amyloid Associated Protein**

---

(Ref: Harrison 18th/chapter 112)

- In **secondary amyloidosis associated with chronic diseases**, there is deposition of SAA or AA
- **AA does not have structural homology to immunoglobulins.**
- **Derived by proteolysis** from a larger precursor in the serum called **SAA (serum amyloid-associated)** protein that is **synthesized in liver** and circulates in association with high density lipoproteins.
- Production of SAA protein is **increased in inflammatory states** as part of the “**acute phase response**,” and is often called **secondary amyloidosis**.

---

**199. Ans. (d) Carpal tunnel** (Ref: Harrison 18th/chapter 112)

---

---

**200. Ans. (a) Primary systemic amyloidosis**

---

(Ref: Harrison 18th/chapter 112)

- **Cutaneous lesions** associated with **primary systemic amyloidosis** are often **pink** in color and translucent.
- Common locations are **face, especially periorbital and perioral** regions, and **flexural areas**.
- On biopsy, **homogeneous deposits of amyloid are seen in the dermis** and in the **walls of blood vessels**;
- These deposits lead to an **increase in vessel wall fragility**.
- As a result, petechiae and purpura develop in clinically normal or abnormal skin following minor trauma, hence the term **pinch purpura**.

---

**201. Ans. (d) Beta 2 microglobulin**

---

(Ref: Harrison 18th/chapter 112)

---

**202. Ans. (a) Mutated transthyretin**

---

(Ref: Robbins 9th/pg 259; 8th/pg 252)

---

**203. Ans. (b) Cardiac failure** (Ref: Harrison 18th/chapter 112)

---

---

**204. Ans. (b) Hemodialysis associated** (Ref: R 9th/pg 259)

---

---

**205. Ans. (a) Rectal biopsy** (Ref: Robbins 9th/pg 257-260)

---

---

**206. Ans. (c) Familial Mediterranean fever; e. Systemic senile amyloidosis** (Ref: Harrison 18th/chapter 112)

---

**Familial forms of amyloidosis: Familial Mediterranean fever:**

- **Autosomal recessive** condition
- “**Autoinflammatory**” syndrome associated with abnormally high production of the cytokine IL-1
- **Attacks of fever** accompanied by inflammation of serosal surfaces (peritoneum, pleura & synovial membrane).
- The gene for familial Mediterranean fever encodes a **protein called pyrin** (for its relation to fever)
- The amyloid fibril proteins are made up of **AA proteins**, suggesting that this form of amyloidosis is related to the recurrent bouts of inflammation.

**Familial amyloidotic polyneuropathies:**

- **Autosomal dominant** familial disorder
- Characterized by deposition of amyloid predominantly in **peripheral and autonomic nerves**.
- In these genetic disorders, the fibrils are made up of **mutant TTRs**

---

**207. Ans. (a) Congo red; b. Thioflavin; e. PAS**

---

(Ref: Robbins 9th/pg 257-260; 8th/pg 250-253)

[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slightly textured appearance and is set against a dark background.

# Neoplasia

## Key Points

- » **Neoplasia** means “new growth”, and new growth is called a neoplasm
- » **Oncology** (Greek oncos = tumor) is the study of tumors or neoplasms
- » **Anlage**: Mass of primitive cells from which an organ develops
- » **Hamartoma**<sup>o</sup>: **Disorganized**, benign mass, composed of cells **indigenous** to involved site<sup>o</sup>
- » **Choristoma**<sup>o</sup>: **Heterotopic** rest of cells<sup>o</sup>, e.g., pancreatic tissue in submucosa of stomach, duodenum, or small intestine
- » **Metaplasia**<sup>o</sup>: **Replacement of one type of tissue with another** type of tissue<sup>o</sup>
- » **Benign tumor**: A tumor with **relatively innocent** gross and microscopic appearances, which remain localized and is amenable to local surgical removal
- » **Malignant tumor**: Tumor that tends to **adhere** to any part that they seize on, can **invade** and destroy adjacent structures and **spread to distant sites (metastasize)**.<sup>o</sup>
- » **Desmoplasia**<sup>o</sup>: Abundant **collagenous stroma** (fibrosis) in a tumor, stimulated by **parenchymal cells**<sup>o</sup>

## Key Recent Updates


- » CART cells is called live drug
- » Tumor evolution follows Darwin's finches.





## SOME IMPORTANT TERMINOLOGIES

### Nomenclature of Neoplasms of Differentiated Cells



### Neoplasms with More than One Type of Neoplastic Cells

Organ involved	Benign	Malignant
<b>Mixed tumors<sup>q</sup>:</b> Derived from <b>one germ cell layer</b>		
• Salivary glands	Pleomorphic adenoma <sup>q</sup>	Malignant mixed tumor
• Renal anlage <sup>q</sup>	–	Wilms tumor <sup>q</sup>
<b>Teratomas<sup>q</sup>:</b> Derived from <b>more than one germ cell layer</b>		
Totipotential cells in embryonic rests	Mature teratoma, <sup>q</sup> Dermoid cyst	Immature teratoma, Terato-carcinoma

#### Basic Component Soft Tumors

- **Neoplastic Cells:** Constitute the tumor parenchyma
- **Reactive Stroma:** Made up of connective tissue, blood vessels and cells of the immune system.

## CHARACTERISTICS OF BENIGN AND MALIGNANT NEOPLASMS

### Comparison between Benign and Malignant Tumors

Characteristics	Benign	Malignant
<b>Differentiation</b>	Well differentiated <sup>q</sup>	Poorly differentiated (anaplasia) <sup>q</sup>
<b>Rate of growth</b>	Slow	Erratic and rapid
<b>Mitotic figures</b>	Rare and normal <sup>q</sup>	Numerous and abnormal <sup>q</sup>
<b>Local invasion</b>	Do not invade <sup>q</sup> or infiltrate surrounding tissue	Locally invasive, infiltrating surrounding tissue <sup>q</sup>
<b>Metastasis</b>	Absent <sup>q</sup>	Frequently present <sup>q</sup>

### Differentiation

Dysplasia	Anaplasia
Disordered growth Partially reversible	Lack of differentiation Irreversible
Common features	
• Pleomorphism	• High N:C ratio
• Nuclear irregularity	• Loss of polarity
• Hyperchromatism	• Atypical, increased Mitosis

- When dysplasia is severe and involves full epithelial thickness without penetrating basement membrane, it is called carcinoma in situ
- Once the tumor cells breach the basement membrane, the tumor is said to be invasive
- **ANAPLASIA is the hallmark of malignancy**

First let's understand the cell cycle

#### Cell Cycle

- **Definition:** Cell cycle is the **sequence of events that results in cell division<sup>q</sup>**
- **Sequence:**

$G_0 \rightarrow G_1 \rightarrow S \rightarrow G_2 \rightarrow M$

Cells can enter G<sub>1</sub> either from G<sub>0</sub> quiescent cell pool, or after completing a round of mitosis

- **Phases**

<b>G<sub>1</sub></b>	Pre-synthetic growth <sup>q</sup>
<b>S</b>	DNA synthesis <sup>q</sup> ( <b>most radioresistant phase</b> ) <sup>q</sup>
<b>G<sub>2</sub></b>	Pre-mitotic growth
<b>M</b>	Mitotic phase ( <b>most radiosensitive phase</b> ) <sup>q</sup>
<b>G<sub>0</sub></b>	<b>Quiescent</b> cells that are <b>not actively</b> cycling are said to be in G <sub>0</sub> state



- **G<sub>1</sub>/S checkpoint:**
  - Checks for **DNA damage before replication in S phase** → **arrests cell cycle if damage is present** → **activates DNA repair mechanisms**
  - DNA can be **repaired** only as long as the **chromatids have not separated**.<sup>Q</sup>
- **G<sub>2</sub>/M checkpoint:**
  - Monitors the **completion of DNA replication**<sup>Q</sup>
  - Checks whether the cell can safely initiate mitosis and separate sister chromatids.
  - Defects in this checkpoint give rise to **chromosomal abnormalities**.<sup>Q</sup>
  - Cells exposed to **ionizing radiation** → Cell cycle **arrested in G<sub>2</sub>** and **repair mechanisms activated**<sup>Q</sup>


### Components of Cell Cycle Cyclins and Cyclin-dependent Kinases (CDK)<sup>Q</sup>

- **Cyclins<sup>Q</sup>:** Proteins (with cyclic production and degradation) that **drive cell cycle progression**
- **Cyclin-dependent kinases (CDKs):** Cyclin-associated **enzymes** that acquire the ability to **phosphorylate** protein substrates by **forming complexes with the relevant cyclins**<sup>Q</sup>
- **CDK inhibitors (CKIs):** Proteins that **block the cell cycle** by **binding to cyclin-CDK complexes**.

Check Points	Cyclin	CDK
G <sub>1</sub> -S	Cyclin D	CDK4/CDK6 <sup>Q</sup>
	Cyclin E	CDK2
G <sub>2</sub> -M	Cyclin A	CDK2/CDK1 <sup>Q</sup>
	Cyclin B	CDK1 <sup>Q</sup>

### Cell Cycle Inhibitors<sup>Q</sup>

Family	Checkpoints	Proteins	Functions
<b>CIP/KIP</b> <sup>R99<sup>Q</sup></sup> (CDKN1)	G1-S & G2-M	p21 p27 p57	<ul style="list-style-type: none"> <li>• <b>p21</b> is induced by <b>p53</b><sup>Q</sup></li> <li>• <b>p27</b> responds to <b>TGF-β</b><sup>Q</sup></li> </ul>
<b>INK4/ARF</b> <sup>R99<sup>Q</sup></sup> (CDKN2)	G1-S	p15 p16 p18 p19	<ul style="list-style-type: none"> <li>• <b>p16/INK4a</b> binds to CyclinD-CDK4 &amp; promotes inhibitory effects of RB<sup>Q</sup></li> <li>• <b>p14/ARF</b> increases <b>p53</b> levels by inhibiting <b>MDM2</b><sup>Q</sup></li> </ul>



### Hallmarks of Cancer ("S-T-E-A-L A-T-M")

- **Self-sufficiency<sup>Q</sup>** in growth signals
- **Tumor suppressor gene inactivation:** Insensitivity to growth inhibitory signals<sup>Q</sup>
- **Evasion of apoptosis**
- **Altered cellular metabolism** (aerobic glycolysis-Warburg effect)<sup>Q</sup>
- **Limitless replicative potential** (immortality)
- **Sustained Angiogenesis**<sup>Q</sup>
- **Evasion of immunity**
- **Metastasis**















### Self-sufficiency in Growth Signals (By Genetic Mutations and Oncogene Activation)



- **Proto-oncogenes:** Unmutated counterparts of oncogenes found **normally in a cell**
- **Functions of Proto-oncogenes:** Cell growth<sup>Q</sup>, inhibition of apoptosis<sup>Q</sup> and nuclear transcription<sup>Q</sup>
- **Oncogenes<sup>Q</sup>:** Mutated or over expressed version of **proto-oncogenes** that **function autonomously**
- **Oncogenes have no<sup>Q</sup> dependence** on normal growth promoting signals and **growth factors**



- Proto-oncogenes can be activated to oncogene by translocation (Most common in hematologic neoplasms), amplification and point mutation
- Proto-oncogenes can be activated by virus<sup>Q</sup> to oncogenes
- Oncoproteins: Proteins encoded by oncogenes that promote cell growth in absence of growth promoting signals

### Oncogenes and Associated Tumors

Category	Proto-oncogene	Associated Human Tumor
<b>Growth Factors</b>		
<i>PDGF-β chain</i>	<i>SIS</i> <sup>Q</sup> ( <i>PDGFB</i> )	Astrocytoma <sup>Q</sup> 
<i>Fibroblast growth factors (FGF)</i>	<i>HST1</i>	Osteosarcoma
	<i>INT2</i> ( <i>FGF3</i> )	Stomach, Bladder
<i>TGF-α</i>	<i>TGFA</i>	Astrocytomas
<i>HGF</i>	<i>HGF</i>	Hepatocellular Carcinoma,  Thyroid cancer
<b>Growth Factor Receptors</b>		
EGF-receptor family	<i>ERBB1</i> ( <i>EGFR</i> ) <i>ERBB2</i> ( <i>HER</i> )	Adenocarcinoma Lung  Breast Ca 
<i>FMS-like tyrosine kinase3</i>	<i>FLT3</i>	Leukemia (ALL)
<i>Receptor for neurotrophic factors</i>	<i>RET</i> <sup>Q</sup>	MEN 2A & 2B, medullary thyroid <sup>Q</sup>
<i>PDGF receptor</i>	<i>PDGFRB</i> <sup>Q</sup>	Gliomas, <sup>Q</sup> Leukemias <sup>Q</sup>
<i>Receptor for KIT</i>	<i>KIT</i> <sup>Q</sup>	Gastrointestinal stromal tumors,
<i>ALK receptor</i> 	<i>ALK</i> 	AdenoCa Lung, Anaplastic lymphoma, Neuroblastoma 
<b>Proteins Involved In Signal Transduction</b>		
<i>GTP-binding</i>	<i>KRAS</i> <sup>Q</sup>	Colon, Lung Ca
	<i>HRASQ</i>	Bladder and Kidney tumors
	<i>NRASQ</i>	Melanomas, Hematologic malignancies
	<i>GNAQ</i> 	Uveal melanoma 
	<i>GNAS</i> 	Pituitary adenoma 
<i>Non-receptor tyrosine kinase</i>	<i>ABL</i>	CML, <sup>Q</sup> ALL
<i>RAS signal transduction</i>	<i>BRAF</i> <sup>Q</sup>	Melanoma, Hairy Cell Leukemia, ColonCa
<i>Notch signal transduction</i>	<i>NOTCH1</i>	Leukemias,
<i>JAK/STAT signal transduction</i> 	<i>JAK2</i> 	Myeloproliferative disorders, ALL 
<i>WNT signal transduction</i>	β-catenin <sup>Q</sup>	Hepatoblastomas, hepatocellular carcinoma
<b>Nuclear-Regulatory Proteins</b>		
<i>Transcriptional activators</i>	<i>C-MYC</i> <sup>Q</sup>	Burkitt lymphoma <sup>Q</sup>
	<i>N-MYC</i> <sup>Q</sup>	Neuroblastoma, small-cell Ca lung <sup>Q</sup>
	<i>L-MYC</i>	Small-cell of lung <sup>Q</sup>




### High Yield Facts

#### Oncogene Addition

- Tumor cells are highly dependent on the activity of one or more oncogenes Seen in CML. Hence inhibition of its activity is a highly effective therapy.

Contd...



Category	Proto-oncogene	Associated Human Tumor
<b>Cell Cycle Regulators</b>		
<b>Cyclins</b>	<b>Cyclin D1<sup>Q</sup></b> (CCND1)	<b>Mantle cell lymphoma<sup>Q</sup></b> , 
<b>Cyclin-dependent kinase</b>	<b>CDK4</b>	Glioblastoma, Melanoma

## Tumor Suppressor Genes

Genes that prevent uncontrolled proliferation of cell We will discuss few important ones


### RB (Retinoblastoma) Gene: on Chromosome 13q14<sup>Q</sup>

- **Also called: Governor of proliferation<sup>Q</sup>**
- **Function:** Key **negative<sup>Q</sup>** regulator of **G1/S<sup>Q</sup>** cell cycle transition
- **Tumors associated:** **Retinoblastoma<sup>Q</sup>**, **Osteosarcoma<sup>Q</sup>**, Glioblastomas, Small-cell Ca of lung, Breast Ca & Bladder Ca

### Knudson's "2 hit"<sup>Q</sup> hypothesis of Oncogenesis for Retinoblastoma:

- In **sporadic** form, **mutations** in **both RB** genes in the retinal cell are **acquired<sup>Q</sup>**
- In **familial** form, all somatic cells **inherit one mutated copy** of RB gene (heterozygous), and **one additional RB mutation** is required for complete loss of RB function. (**loss of heterozygosity<sup>Q</sup>**)

### Regulation of Cell Cycle by RB Gene







## TP53: on chromosome 17p13.1<sup>Q</sup>

Also called	Guardian of the Genome <sup>Q</sup>
Functions	<ul style="list-style-type: none"> <li>p53 is <b>universally expressed</b> in all cells &amp; <b>encodes a 53 kDa protein</b><sup>Q</sup></li> <li><b>Regulates cell cycle progression, DNA repair, cellular senescence &amp; apoptosis</b></li> <li><b>p53 activates CDK inhibitor (p21) → inhibits cyclin-CDK complexes &amp; hence inhibits phosphorylation of RB → arrests cell-cycle at G1-S phase.</b></li> </ul>
TP53 mutation causes	Brain Tumors, Breast Ca, <sup>Q</sup> Leukemia, Adrenal Ca, Sarcomas,
Other members of P53 family	<ul style="list-style-type: none"> <li><b>p63</b> is essential for the differentiation of <b>stratified squamous epithelia</b><sup>RG</sup></li> <li><b>p73</b> has strong <b>pro-apoptotic effects</b> after DNA damage by <b>chemotherapeutic agents</b><sup>RG</sup></li> </ul>
Li-Fraumeni syndrome <sup>Q</sup>	<ul style="list-style-type: none"> <li>Individuals who have <b>inherited one mutated TP53 allele</b><sup>Q</sup></li> <li><b>Predisposed to malignancy</b><sup>Q</sup> as only <b>1 additional “hit”</b><sup>Q</sup> in the lone normal allele is <b>needed</b></li> <li><b>Have 25-fold greater chance of developing a malignancy by age 50</b></li> <li><b>Develop cancer at younger ages</b><sup>Q</sup> and may develop <b>multiple</b><sup>Q</sup> primary tumors</li> <li><b>Common Tumors: Sarcomas, Breast cancer,<sup>Q</sup> Leukemias,<sup>Q</sup> Brain tumors,<sup>Q</sup> Adrenal Ca.<sup>Q</sup></b></li> </ul>

## Adenomatous Polyposis Coli (APC) Gene: Chr 5q21<sup>Q</sup>

Also called	Gate-keeper of Colonic Neoplasia <sup>Q</sup>
Functions	<ul style="list-style-type: none"> <li>Component of the <b>WNT signaling pathway</b><sup>Q</sup></li> <li>Controls cell fate, adhesion &amp; cell polarity during embryonic development.</li> <li>It also <b>controls oncogenic effects of β-catenin</b> so prevents carcinomas</li> </ul>
Tumors due to APC mutation	<ul style="list-style-type: none"> <li>Germline <b>loss-of-function</b> mutation → <b>Familial adenomatous polyposis colon Cancer (AD)</b><sup>Q</sup>, <b>Hepatoblastomas, Hepatocellular Ca</b></li> </ul>



## High Yield Facts

- Phosphorylated form is the deactivated form of RB gene
- Knudson’s “2 hit” hypothesis of oncogenesis is for Retinoblastoma
- Wild form of TP53 gene is the non-mutated form,**<sup>Q</sup> while mutated form causes tumors
- TP53 is the most commonly mutated gene in human cancers**<sup>Q</sup>
- TGF-β is a potent inhibitor**<sup>Q</sup> of proliferation in most normal epithelial, endothelial, and hematopoietic cells.
- TGF-β type II receptor mutations** are seen in cancers of the colon, stomach, endometrium & pancreas.<sup>Q</sup>
- GAPs**<sup>Q</sup> (GTPase-activating proteins) function as “brakes” that prevent uncontrolled RAS activity.

## List of Important Tumor Suppressor Genes and Associated Cancers

<div> <div>RG<sup>th</sup></div> <div>Latest Update</div> </div>		
Gene	Protein	Cancers
<b>Inhibitors of Mitogenic Signaling Pathways</b>		
<b>APC</b>	APC protein	<b>Familial colonic polyps &amp; Carcinoma</b> <sup>Q</sup>
<b>NFI</b>	Neurofibromin 1	<b>NF type 1</b> <sup>Q</sup> (Neurofibromas & peripheral nerve sheath tumors)
<b>NF2</b>	Merlin	Neurofibromatosis type 2 (acoustic <b>schwannoma &amp; meningioma</b> ) <sup>Q</sup>
<b>PTCH</b>	Patched	<b>Gorlin syndrome</b> <sup>Q</sup> (Basal cell Ca, Medulloblastoma)
<b>PTEN</b>	Phosphatase & tensin homologue	<b>Cowden Syndrome</b> <sup>Q</sup> (Breast, endometrial, and prostate carcinoma)
<b>SMAD2, SMAD4</b>	SMAD2, SMAD4	<b>Juvenile polyposis</b> <sup>Q</sup> , <b>pancreatic Ca</b>
<b>Inhibitors of Cell Cycle Progression</b>		
<b>RB</b>	Retinoblastoma (RB) Protein	<b>Familial retinoblastoma syndromes</b> (Retinoblastoma <sup>Q</sup> & Osteosarcoma <sup>Q</sup> )
<b>CDKN2A</b>	P16/INK4 a and p14/ARF	<b>Familial melanoma</b> <sup>Q</sup>
<b>Inhibitors of “pro-growth” Programs of Metabolism and Angiogenesis</b>		
<b>VHL</b>	VHL protein	<b>Von Hippel-Lindau syndrome</b>

Contd...



<b>STK11</b>	Liver kinase B1 (LKB1) or STK11	<b>Peutz-Jeghers syndrome<sup>Q</sup></b> (GI polyps, GI cancers, pancreatic carcinoma) <sup>Q</sup>
<b>SDHB, SDHD</b>	Succinate dehydrogenase B & D	Familial paraganglioma, pheochromocytoma
<b>Inhibitors of Invasion and Metastasis</b>		
<b>CDH1</b>	E-cadherin <sup>Q</sup>	Familial gastric cancer <sup>Q</sup>
<b>Enablers of Genomic Stability</b>		
<b>TP53</b>	P53 protein	Li–Fraumeni syndrome <sup>Q</sup> (see above)
<b>DNA Repair Factors</b>		
<b>BRCA1, BRCA2</b>	BRCA1 and BRCA2	Familial breast and ovarian Ca; <sup>Q</sup>
<b>Unknown Mechanisms</b>		
<b>WT1</b>	Wilms tumor-1 (WT1)	Familial Wilms' tumor <sup>Q</sup>
<b>MEN1</b>	Menin	MEN1 <sup>Q</sup> syndrome

## Warburg Effect: Aerobic Glycolysis/Glucose Hunger<sup>Q</sup>

- **Otto Warburg** was awarded Nobel Prize in 1931 for this theory
- **Cancer cells tend to convert most glucose to lactate<sup>Q</sup> even in presence of ample oxygen (aerobic glycolysis)<sup>Q</sup>**
- Even though **ATP production is low** with formation of lactate compared to mitochondrial oxidative phosphorylation, it **provides metabolic intermediates** that are needed for the synthesis of **cellular components**.
  - Warburg Effect is the Basis of Positron Emission Tomography (PET)<sup>Q</sup>
- Patients are injected with **18F-fluorodeoxyglucose (FDG)**, a non-metabolizable derivative of glucose, that is **preferentially taken up into tumor cells** (due to glucose hunger), which can be detected by PET scanning

## Evasion of Programmed Cell Death (Apoptosis)

- **Intrinsic apoptotic pathway** (mitochondrial pathway) **most frequently disabled in cancer.<sup>Q</sup>**
- **BCL-2:** mutated in **85% of follicular B-cell lymphomas<sup>Q</sup>**
- **Autophagy:** Tumor cells in **severe nutrient deficiency**, arrest their growth and also **cannibalize** their own **organelles, proteins, and membranes** as carbon sources for **energy production.<sup>Q</sup>**

## Limitless Replicative Potential

- **Cancer stem cells:** Stem cell-like cells that are **immortal** and have limitless replicative potential, seen in all cancers.
- May arise through **transformation of a normal stem cell** or through **acquired genetic lesions**
- **Immortality of cancer cells is due to:**
  - **Evasion of senescence:** By disruption of **RB-dependent G1/S cell cycle checkpoint<sup>Q</sup>**
  - **Evasion of mitotic crisis:** By **telomere maintenance,<sup>Q</sup> up-regulation of telomerase<sup>Q</sup> or alternative lengthening of telomeres**

- **Capacity for self-renewal<sup>Q</sup>:** Self-renewal means that **each time a stem cell divides at least one of the two daughter cells remains a stem cell** (also called **Asymmetric cell division**)<sup>Q</sup>

## Angiogenesis

- Tumor cannot enlarge beyond **1–2 mm<sup>Q</sup>** in diameter unless it has the capacity to **induce angiogenesis**.
- **Angiogenesis** is an important requirement for tumors to **undergo metastasis**

Pro-angiogenic factors	Anti-angiogenic factors
• <b>Angiogenin</b>	• Endostatin <sup>Q</sup>
• <b>Vascular Endothelial Growth Factor (VEGF)<sup>Q</sup></b>	• Angiostatin <sup>Q</sup>
• <b>Fibroblast Growth Factor (FGF)<sup>Q</sup></b>	• Interferon-Alpha
• <b>Transforming Growth Factor-β (TGF-β)<sup>Q</sup></b>	• Thrombospondin-1 <sup>Q</sup>
• <b>Platelet activating factor (PAF)</b>	

## Metastasis

- **Definition:** Spread of a tumor to sites that are **physically discontinuous** with primary tumor.
- **Pathways of metastasis:**
  - **Direct** seeding of body cavities or surfaces, e.g., ovarian Ca
  - **Lymphatic** spread e.g., Carcinomas<sup>Q</sup>
  - **Hematogenous** spread e.g., sarcomas<sup>Q</sup>
- **Steps involved:**
  - Invasion of the extracellular matrix (ECM):
    - “**Loosening up**” of tumor cell–tumor cell interactions: **E-cadherin mutations**
    - **Degradation of ECM:** **Metalloproteinases** (MMPs type 2 and 9)<sup>Q</sup> also known as **Type IV collagenase<sup>Q</sup>**, cathepsin D, & urokinase plasminogen activator
    - **Attachment** to novel ECM components by **Fibronectin<sup>Q</sup>**
    - **Migration and invasion** of tumor cells

- Vascular Dissemination and Homing of Tumor Cells:
  - Expression of **CD44<sup>Q</sup>** in solid tumors which binds to **hyaluronate** on high endothelial venules which enhances their spread to lymph nodes and other metastatic sites.

R10<sup>th</sup>

## Latest Update

## Metastasis oncogenes:

- **SNAIL & TWIST**: encode transcription factors which promote **epithelial-to-mesenchymal transition (EMT)**
- **EMT: Down-regulation of epithelial markers** (e.g., E-cadherin) and **up-regulation of mesenchymal markers** (e.g., vimentin and smooth muscle actin)
- **Pro-migratory phenotype** is essential for metastasis.

## Evasion of Host-Immune Defense

Mechanisms by which immune system is evaded in immune-competent hosts:

- Selective outgrowth of **antigen-negative<sup>Q</sup>** variants
- Loss or **reduced** expression of **histocompatibility antigens<sup>Q</sup>**
- **Immunosuppression<sup>Q</sup>** mediated by expression of certain factors (e.g., **TGF- $\beta$ , PD-1 ligand, galectins**) by the tumor cells

RECENT EXAM<sup>Q</sup>

- The 2018 Nobel Prize in Physiology or Medicine has been awarded jointly to two cancer immunotherapy researchers, James P. Allison, PhD, of The University of Texas MD Anderson Cancer Center, and Dr. Tasuku Honjo of Kyoto University in Japan.
- In his laboratory at the University of California, Berkeley, Allison studied the T cell protein CTLA-4. When this protein attaches to another protein (called B7) found on the surface of some cancer cells, it signals the T cell that this cell is functioning properly. As long as a regular or cancer cell is sending the message that it is functioning well, the immune system will not destroy the cell. By blocking the CTLA-4 protein and thereby the message that the cell is working as it should, the immune system can

recognize cancer cells and attack them. This discovery eventually led to the drug Yervoy (ipilimumab), which is used to treat melanoma skin cancer and some other cancers.


- Honjo at Kyoto University in Japan discovered PD-1, which is another protein found on the surface of some T cells. When this protein attaches to a protein called PD-L1 on cancer cells, it can prevent the T cells from recognizing the cancer cells, so the immune system won't destroy them. Blocking the PD-L1 protein on cancer cells, or the corresponding PD-1 protein on immune cells, allows the immune system to recognize the cancer cells as foreign and attack them.
- Drugs that target PD-1 include Keytruda (pembrolizumab) and Opdivo (nivolumab), while drugs that target PD-L1 include atezolizumab (Tecentriq), avelumab (Bavencio), and durvalumab (Imfinzi). These drugs are now used to treat people with many different cancer types, including melanoma skin cancer, non-small cell lung cancer, kidney cancer, bladder cancer, head and neck cancers, and Hodgkin lymphoma.

## High Yield Facts


- **All malignant tumors can metastasize except Gliomas and Basal cell Ca<sup>Q</sup>**
- **Decrease of telomerase activity cause antitumor effects<sup>Q</sup>**
- **Most common site of metastasis is Lungs > Liver<sup>Q</sup>**
- **'Anoikis'** is **apoptosis stimulated by detachment of epithelial cells from basement membranes & from cell-cell interactions<sup>Q</sup>**
- **Immunosuppressed patients have increased risk for development of cancer**, particularly caused by **oncogenic DNA viruses**.
- **Fibromatoses** are apparently **autonomous proliferation of myofibroblasts**, occasionally forming **tumor like masses**


## GENETIC BASIS OF CANCERS

- Genetic aberrations that **increase mutation rates** fastens **driver mutations** that are required for transformation and subsequent tumor progression
- Selection of fittest cells can explain not only tumor evolution but also changes in tumor behaviour following therapy



Tumor evolution resembles Darwin's finches






## Dysregulation of Cancer-Associated Genes

Chromosomal Abnormalities like deletions, translocation, gene amplifications can cause cancer (Refer to annexures)

### Defect in DNA Repair Mechanism



### Ataxia Telangiectasia

- **Immunodeficiency:** Thymic hypoplasia<sup>Q</sup> (most consistent defect) with cellular and humoral (IgA and IgG2)<sup>Q</sup> immunodeficiency
- **Clinical features:** Recurrent pulmonary infections,<sup>Q</sup> premature aging, Type 1 diabetes mellitus
- **Tumors seen:** Lymphomas,<sup>Q</sup> Hodgkin's disease,<sup>Q</sup> T cell ALL & Breast cancer
- **Neuropathologic changes:**
  - Loss of Purkinje, granule, and basket cells in the cerebellar cortex<sup>Q</sup> (most striking change) & deep cerebellar nuclei.

## Epigenetic Changes

Epigenetic Changes are factors other than the sequence of DNA that regulate gene expression

Play major role in:

- Expression of cancer genes<sup>Q</sup>, Control of differentiation, Self renewal, Drug sensitivity and drug resistance

### Epigenetic Alterations in Cancers

- Exhibiting **abnormal DNA methylation:** hypomethylation<sup>Q</sup> or hypermethylation<sup>Q</sup>
- **Silencing of tumor suppressor genes<sup>Q</sup>** by local hypermethylation of DNA
- Changes in histones near genes that **influence cellular behavior<sup>Q</sup>**

### Examples of Epigenomic Regulatory Genes that are Mutated in Cancer

Gene(s)	Tumor
ARID1A	Ovarian clear cell Ca (60%), Endometrial Ca
SNF5	Malignant rhabdoid tumor (100%)
DNMT3A	AML (20%)
MLL1	ALL
MLL2	Follicular lymphoma (90%) <sup>Q</sup>
CREBBP/EP300	Diffuse large B Cell lymphoma (40%)

## MicroRNAs (miRs) and Cancer

- **What are miRs?**
  - MicroRNAs (miRs) are small non-coding, ssRNA, approx 22 nucleotides in length
- **Function:**
  - Mediate sequence-specific **inhibition of messenger RNA (mRNA) translation** through the action of the RNA-induced silencing complex (RISC)
- **Mechanism of Oncogenesis:**
  - **Decreased expression of** tumor suppressive miRs → increases oncogenic mRNA translation
  - **Overexpression of** oncogenic miRNAs → repress the expression of tumor suppressor genes
- **Examples:**
  - **OncomiRs** (oncogenic miRNAs)
    - miR-200- promote epithelial-mesenchymal transitions, invasiveness & metastasis
    - miR-155- upregulates genes that promote proliferation (MYC) → B cell lymphomas
  - Tumor suppressive miRs:
    - miR-15 and miR-16 upregulates BCL-2 in CLL
  - **Tumor suppressive properties of miR processing factors**
    - DICER: a gene that encodes an endonuclease that is required for the processing and production of functional miRs.
    - Ovarian and testicular tumors

R10<sup>th</sup>

Latest Update

Some other Non-coding RNAs		
Type	Full name	Function
piRNA	piwi-interacting RNA	Most common type of small noncoding RNA, have a role in post-transcriptional gene silencing
snoRNA	small nucleolar RNA	Maturation of rRNA and the assembly of ribosome
lncRNA	long intervening non-codingRNA	Regulate histones and thereby control gene expression

## Chromothripsis

Literally means, **chromosome shattering**.

- Single event in which several **chromosome breaks** occur within a single chromosome or multiple chromosomes.
- Later **DNA repair mechanisms<sup>Q</sup>** are activated in affected cells that **stitch the pieces** together in a **disorganized** way
- This may create **chromosome rearrangements<sup>Q</sup>** and also result in the **loss of some chromosome** segments.
- This may also **activate oncogenes** and **inactivate tumor suppressors**,
- Finally these changes **accelerate the process of carcinogenesis<sup>Q</sup>**.





## CARCINOGENIC AGENTS

### Chemical Carcinogens (Occupational Cancers)

#### Major Chemical Carcinogens

Chemical	Types of Cancer
<b>Polycyclic hydrocarbons</b>	
Soot (benzopyrene, dibenzanthracene)	Skin; scrotal cancer
Tobacco	Lung, bladder, oral cavity, larynx, esophagus
<b>Aromatic amines</b>	
Benzidine, 2-naphthylamine	Bladder <sup>Q</sup>
Aflatoxins	Liver <sup>Q</sup>
Nitrosamines	Esophagus, stomach
<b>Cancer chemotherapeutic agents</b>	
Cyclophosphamide, <sup>Q</sup> chlorambucil, busulfan	Leukemias
Asbestos <sup>Q</sup>	Lung cancer, mesothelioma <sup>Q</sup>
<b>Heavy metals</b>	
Nickel, chromium, cadmium	Lung
Arsenic	Skin
Vinyl chloride <sup>Q</sup>	Liver (angiosarcoma) <sup>Q</sup>

### Radiation Carcinogenesis

Ultraviolet B Rays	Ionizing Radiation
<ul style="list-style-type: none"> <li>Leads to formation of pyrimidine dimers in DNA<sup>Q</sup></li> <li>Produces skin cancers like Squamous cell Ca, Basal cell Ca, and melanoma of skin</li> </ul>	<ul style="list-style-type: none"> <li>Particulate radiation (<math>\alpha</math> &amp; <math>\beta</math> particles, protons, neutrons) are all carcinogenic</li> <li>Electromagnetic (x-rays, <math>\gamma</math> rays)</li> <li>X-ray causes DNA mutation by Pyrimidine dimer breakdown</li> </ul>

Most **radiosensitive** cells are those that are **undifferentiated**, **well nourished**, **rapidly dividing** and **highly active** metabolically. e.g., cells of buccal mucosa > skin



#### High Yield Facts

- Lethal dose of radiation for humans is **250–400 rads<sup>Q</sup>**
- Most radiosensitive tissue are **spermatogonia<sup>Q</sup>**, **erythroblasts**, **epidermal & GI stem cells**
- Least radiosensitive are **Cartilage<sup>Q</sup>** > Bone, nerve cells and muscle fibers.
- Radiation increases the risk of leukemias<sup>Q</sup> and solid tumors in several organs eg thyroid,<sup>Q</sup> breast, and lungs.<sup>Q</sup>
- In thyroid, papillary Ca of thyroid is common due to radiation<sup>Q</sup>

### Microbial Carcinogenesis

Organism	Gene	Mechanism	Tumor
<b>Oncogenic RNA Viruses</b>			
HTLV-1	Tax <sup>Q</sup>	Increased <b>pro growth signaling &amp; cell survival</b> & genomic instability.	Adult T cell leukemia <sup>Q</sup> / Lymphoma <sup>Q</sup>
Hep C	HCV core protein	Activates growth-promoting signal transduction pathways	Hepatocellular carcinoma <sup>Q</sup>
<b>Oncogenic DNA Viruses</b>			
HPV Low risk (6,11) High risk (16,18)	E6 <sup>Q</sup>	Degradation of <b>p53<sup>Q</sup></b> & stimulation of <b>TERT</b>	<b>Low risk:</b> Genital warts <sup>Q</sup> <b>High risk:</b> Squamous cell Ca cervix, Ca Oropharynx, anus, head/neck ( <b>laryngeal papilloma</b> ) <sup>Q</sup> , Esophagus
	E7 <sup>Q</sup>	Inactivates <b>RB<sup>Q</sup></b> & <b>CDKI</b> (p21 & p27)	
EBV (Epstein-Barr virus)	LMP1 <sup>Q</sup>	Activates <b>NF-kB</b> & <b>JAK/STAT</b> pathway	<b>Lymphoma:</b> Burkitt's, <sup>Q</sup> Hodgkin, T-cell & NK cell <b>Carcinoma:</b> Nasopharyngeal <sup>Q</sup> & gastric
	EBNA2	Activates <b>cyclin D</b>	
	vIL-10	B cell activation	
Hep B	HBx	Activates transcription factors	Hepatocellular carcinoma <sup>Q</sup>
<b>Bacteria</b>			
H.pylori	Cag-A	Growth factor stimulation	Gastric adenocarcinomas <sup>Q</sup> & gastric <b>MALToma</b> <sup>Q</sup>



## CLINICAL ASPECTS OF NEOPLASIA

### Cancer Cachexia

- **Progressive loss of body fat and lean body mass<sup>o</sup>** with profound weakness, anorexia & anemia, seen in cancer.
- **TNF<sup>o</sup>** is the **major contributor** to cachexia with advanced cancer.
- Equal loss of both **fat and lean muscle<sup>o</sup>**
- **Elevated basal metabolic rate**
- Evidence of **systemic inflammation** (e.g., an increase in acute phase reactants)

### Paraneoplastic Syndrome

Clinical syndrome	Major forms of underlying cancer
<b>Vascular and Hematologic changes</b>	
<i>Trousseau phenomenon<sup>o</sup></i>	Pancreatic <sup>o</sup> , Bronchogenic <sup>o</sup> & Colon Ca <sup>o</sup>
<i>DIC</i>	APML, prostatic carcinoma
<i>Nonbacterial thrombotic endocarditis</i>	Advanced cancers <sup>o</sup>
<i>Red cell aplasia<sup>o</sup></i>	Thymic neoplasms <sup>o</sup>
<b>Nerve and Muscle syndrome</b>	
<i>Myasthenia</i>	Bronchogenic Ca <sup>o</sup> , thymic neoplasms
<b>Dermatologic, soft tissue &amp; osseous changes</b>	
<i>Acanthosis nigricans (EGF)</i>	Gastric <sup>o</sup> , lung & uterine carcinoma
<i>Dermatomyositis</i>	Bronchogenic & Breast carcinoma <sup>o</sup>
<i>Clubbing ± Hypertrophic osteoarthropathy</i>	Bronchogenic Ca, Thymic neoplasms <sup>o</sup>
<b>Endocrinopathies</b>	
<i>Cushing syndrome (ACTH)</i>	Pancreatic, Small-cell Ca lung <sup>o</sup> , Neural tumors
<i>Syndrome of inappropriate antidiuretic hormone secretion (SIADH) (ADH)</i>	Small-cell Ca lung, Pancreas, Thymoma <sup>o</sup> , Mesothelioma, Bronchial adenoma, Carcinoid <sup>o</sup> , Ewing's sarcoma <sup>o</sup>
<i>Hypercalcemia (Pthrp)</i>	Squamous cell Ca lung, Breast Ca, Renal Ca
<i>Hypoglycemia (Insulin)</i>	Ovarian carcinoma, fibrosarcoma <sup>o</sup>
<i>Polycythemia (Crythropoetin)</i>	RCC, HCC, cerebellar hemangioma

### Tumor Lysis Syndrome (TLS)

- Caused by the **destruction** of a large number of **rapidly proliferating neoplastic cells.<sup>o</sup>**
- **Hyperuricemia<sup>o</sup>, hyperkalemia<sup>o</sup> (life threatening), hyperphosphatemia<sup>o</sup>, and hypocalcemia<sup>o</sup> ± Acidosis** seen.
- Hyperphosphatemia<sup>o</sup> (due to release of intracellular phosphate) produces a decrease in serum calcium.
- Deposition of **calcium phosphate in the kidney** and **hyperphosphatemia** may cause **acute renal failure**.
- Often seen during treatment of **Burkitt's lymphoma<sup>o</sup>, ALL, chronic leukemias<sup>o</sup>** & rarely, solid tumors

### Paraneoplastic Syndromes

- Signs and symptoms that **cannot be** readily explained by the **anatomic distribution** of the tumor or by the **elaboration of hormones** indigenous to the tissue from which the tumor arose.
- Seen in 10% of persons with cancer

#### Paraneoplastic Syndromes are Important to Recognize, for Several Reasons:

- Can be the **earliest** manifestation of an occult neoplasm.
- Can cause significant **clinical problems** and may even be **lethal**.
- May **mimic metastatic disease** and therefore confound treatment.



#### High Yield Facts

- **Cushing's syndrome** is the **most common** paraneoplastic endocrinopathy<sup>o</sup>
- **Hypercalcemia** is the **most common** paraneoplastic syndrome<sup>o</sup>
- Cancer-associated hypercalcemia is due to production of parathyroid hormone-related protein (**PTHrP**)<sup>o</sup>
- **Most common tumors in men** arise in the prostate<sup>o</sup> > lung, and colon/rectum.<sup>o</sup>
- In women, cancers of the breast<sup>o</sup>, lung, and colon/rectum are the most frequent
- Sacrococcygeal teratoma (SCT)<sup>o</sup> is the most frequent tumor in the neonatal period.



## LABORATORY DIAGNOSIS OF CANCERS

### Tissue Specimen

#### Histologic and Cytologic Methods

Sampling may be done by:

- Excision or biopsy, Needle aspiration, and Cytologic smears

#### Exfoliative Cytology


- Cells are collected after they have been either **spontaneously shed** by the body ("spontaneous exfoliation") or **manually scraped/brushed off** of a surface in the body ("mechanical exfoliation")
- Used in diagnosis of **Carcinoma stomach, bronchus, cervix** (PAP smear)

#### Immunohistochemistry

Using **specific antibodies**

- Categorization** of undifferentiated malignant tumors
- Determination of **site of origin of metastatic** tumors.
- Detection of molecules that have **prognostic or therapeutic significance**.

Marker	Tumors
Cytokeratin (CK)	Carcinoma (Squamous cell Ca, Adenocarcinoma)
Vimentin	Sarcoma <sup>Q</sup>
Neurofilament	Neural tumors
GFAP	Gliomas e.g., Astrocytoma <sup>Q</sup>
Desmin	Muscle tumors e.g., Rhabdomyosarcoma <sup>Q</sup>
S-100	Melanoma, <sup>Q</sup> Neuroendocrine Tumor, Schwannoma, Histiocytoma LCH <sup>Q</sup>
HMB 45 <sup>Q</sup>	Melanoma <sup>Q</sup>
Leucocyte common antigen (CD 45)	Lymphoma <sup>Q</sup>



IHC

### Tumor Markers

**Biomarker** found in the blood, urine, or body tissues that can be **elevated in cancer**, among other tissue types. Uses of tumor markers: (**not used for confirmatory diagnosis**)

- Screening**<sup>Q</sup> for common cancers
- Monitoring of cancer**<sup>Q</sup> survivors after treatment
- May help in **diagnosis**<sup>Q</sup> of specific tumor types

#### Important Tumor Markers

Tumor Markers	Tumor Types
<b>Glycoproteins</b>	
• CA-125 <sup>Q</sup>	Ovarian <sup>Q</sup> cancer
• CA-19-9 <sup>Q</sup>	Colon cancer, pancreatic cancer
• CA-15-3	Breast <sup>Q</sup> cancer
• CA 72-4	Gastric <sup>Q</sup> carcinoma
<b>Enzymes</b>	
• Prostatic acid phosphatase (PAP)	Prostate cancer <sup>Q</sup>
• Neuron-specific enolase (NSE)	Small-cell cancer of lung, Neuroblastoma <sup>Q</sup>
• Alkaline phosphatase (ALP)	Osteosarcoma <sup>Q</sup>
• Lactate Dehydrogenase (LDH)	Prostate Ca <sup>Q</sup> , testicular tumors, Lymphoma, Ewing's sarcoma <sup>Q</sup>
• Tyrosinase	Melanoma <sup>Q</sup>
• Gastrin	Pancreatic neuroendocrine tumor (Gastrinoma, ZES) <sup>Q</sup>
<b>Specific proteins</b>	
• Immunoglobulins (Ig)	Multiple myeloma and other gammopathies
• Beta 2 microglobulin <sup>Q</sup> ( $\beta_2M$ )	Multiple myeloma <sup>Q</sup>
• Prostate-specific antigen (PSA)	Prostate cancer
<b>Hormones</b>	
• Human chorionic gonadotropin (HCG)	Germ cell tumor, non-seminomatous <sup>Q</sup> testicular tumors
• Calcitonin <sup>Q</sup>	Medullary carcinoma of thyroid <sup>Q</sup>
• Catecholamine metabolites	Pheochromocytoma <sup>Q</sup>
<b>Oncofetal antigens</b>	
• $\alpha$ - Fetoprotein (AFP)	Liver cancers <sup>Q</sup> , non-seminomatous germ cell tumors of testis
• Carcinoembryonic antigen (CEA) <sup>Q</sup>	Colon, pancreas, lung, stomach & breast Ca



## Molecular Diagnosis (Discussed in Genetics Chapter)


- **PCR (Polymerase Chain Reaction)**
  - Used to produce **large amounts of target DNA fragment**, provided that the **DNA sequence of that region is known<sup>Q</sup>**
- **Array CGH (Comparative Genomic Hybridization)<sup>Q</sup>**
- **SNP (Single Nucleotide Polymorphism) Array**
- **Next gen Sequencing:** term used to describe several **newer DNA sequencing technologies<sup>Q</sup>** that are capable of sequencing entire human genome in few hours.<sup>Q</sup>

## Others

- **Flow Cytometry (Refer to Chapter 23)**
  - **Rapid & quantitative** measure of cellular antigens expressed by “liquid” tumors,
  - Advantage of flow cytometry over IHC is that **simultaneous identification of multiple antigens<sup>Q</sup>** on individual cells possible
- **Circulating Tumor Cells**
  - Detection, quantification, and **characterization of rare solid tumor cells<sup>Q</sup>** (e.g., carcinoma, melanoma) circulating in the blood

### Latest Update

#### Gompertzian growth



*Gompertzian growth.* A  $10^9$ , cancer is diagnosed; however at this stage cells are not in the cell cycle anymore, so they are not as responsive to treatment.  $10^{12}$  levels are not compatible with life (death).





R10<sup>th</sup>

## Latest Updates

CARs have extracellular domains consisting of antibodies that bind tumor antigens and intracellular domains that delivered signals that activate CTLs following their engagement with antigen on the surface of tumor cells.

Tisagenlecleucel (Kymriah<sup>TM</sup>) is FDA approved for the treatment of patients up to 25 years of age with B-cell precursor acute lymphoblastic leukemia (ALL) that is refractory or in second or later relapse. It is a CD19-directed genetically modified autologous T cell immunotherapy.

R10<sup>th</sup>

## Latest Updates

## Neoplasia

- Mc gene mutated is P53 (>75%)
- Temporary cell cycle arrest is caused by P53 by inducing p21 that blocks cyclin D 1 and stops the cell cycle
- Mutated Isocitrate dehydrogenase (IDH) produces 2 hydroxy-glutarate which is called **oncometabolite**. Thus enzymes of krebs cycle play a role in **oncometabolism**. Isocitrate dehydrogenase 1 and 2 (IDH1 and IDH2) are key metabolic enzymes, converting isocitrate to  $\alpha$ -ketoglutarate ( $\alpha$ KG). IDH1 and IDH2

mutations have been identified in multiple tumor types, including gliomas, cholangiocarcinoma and myeloid malignancies such as acute myeloid leukemia (AML) and myelodysplastic syndromes (MDS).

## Immunotherapy

- Personalised vaccines—against tumor antigens
- Adoptive immunotherapy—developing cell having **chimeric antigen receptors** have 2 domains—extracellular domain which has antibody that bind to tumor antigen and intracellular domain has a signal that activate cytotoxic T cells.



## NEXT Pattern Questions



1. The overall survival is increased by screening procedure in which of the following cancer?

- a. Prostate cancer      b. Lung cancer      c. Colon cancer      d. Ovarian cancer

Ans. (c) **Colon cancer**

(Ref: Robbins Basic Pathology 10th ed/pg 235)

"Colorectal cancer is an ideal candidate for screening strategies".

Because:

- It is common
- Has precursor lesion
- It is slow growing
- Testing is available

So, its detection by screening will increase survival.

**Tumors in which screening is done are:**

Cancer	Technique
Ovarian cancer	Ca-125, transvaginal sonography
Cervical cancer	Pap smear
Breast cancer	Mammography
Prostate cancer	Digital rectal examination, prostate specific antigen
Lung	Chest X-ray, sputum cytology



2. All of the following are features of malignant transformation by cultured cells except:

- a. Increased cell density  
b. Increased requirement for growth factors  
c. Alterations of cytoskeletal structures  
d. Loss of anchorage


Ans. (b) **Increased requirement for growth factors**

(Ref: Robbins Basic Pathology 10th ed/pg 205)


**Changes shown by cultured cells which suggest the malignant transformation are:**

- **Alterations of morphology:** Transformed cells often have a much rounder shape than control cells.
- **Increased cell density (loss of contact inhibition of growth):** Transformed cells often form multilayer, while control cells usually form a *monolayer*.
- **Loss of anchorage dependence:** Transformed cells can grow without attachment to the surface of the culture dish and will often grow in agar.
- **Loss of contact inhibition of movement:** Transformed cells grow over one another, while normal cells stop moving when they come into contact with each other.
- **A variety of biochemical changes,** including an increased rate of glycolysis, alterations of the cell surface (e.g., changes in the composition of glycoproteins or glycosphingolipids), and secretion of certain proteases. Alterations of cytoskeletal structures, such as actin filaments.

Diminished requirement for growth factors and, often, increased secretion of certain growth factors into the surrounding medium



3. A 50-year-old man experiences an episode of hemoptysis. On physical examination, he has puffiness of the face, pedal edema, and systolic hypertension. A chest radiograph shows an irregular perihilar 5 cm mass of the right lung. Laboratory studies show normal serum electrolytes. A transbronchial biopsy is performed, and the microscopic findings are shown in the figure. A bone scan shows no metastases. Immunohistochemical staining of the tumor cells is most likely to be positive for which of the following?



- Antidiuretic hormone
- Corticotropin
- Erythropoietin
- Parathyroid hormone-related peptide

Ans. (b) Corticotropin


(Ref: Robbins Basic Pathology 10th ed/pg 228)

The small cells have scant cytoplasm but marked hyperchromatism, consistent with small cell carcinoma. This patient has Cushing syndrome resulting from ectopic corticotropin production by the tumor, a form of paraneoplastic syndrome common to small cell carcinomas of the lung. Such small cell carcinomas are of neuroendocrine derivation. A syndrome of inappropriate antidiuretic hormone (SIADH) secretion from small cell carcinomas is also common, but leads to hyponatremia as well as edema. Erythropoietin production with polycythemia is more likely to be associated with a renal cell carcinoma. Insulin and gastrin production are most often seen in islet cell tumors of the pancreas. Hypercalcemia from a parathyroid hormone-related peptide (PTHrP) is more typically associated with pulmonary squamous cell carcinomas.




## Image-Based Questions

1. The given biopsy from intestine suggests:




- Squamous cell Ca
- Adenocarcinoma
- Adenoma
- Colorectal Ca


2. Biopsy from a lesion of breast mass is suggestive of:



- Lobular Ca
- Medullary Ca
- Fibroadenoma
- Carcinoma in situ




3. Given below is the biopsy from cervix of 45/F who presented with cervical discharge and occasional bleeding. Compare to the normal biopsy of left side (green circle), what is your interpretation of lesion on the right (red circle)?



- a. Carcinoma in situ
- b. Cervical anaplasia
- c. Cervical dysplasia
- d. CMV infection

5. The given figure shows:



- a. Aerobic glycolysis-Warburg effect
- b. Anaerobic glycolysis-Warburg effect
- c. TCA Cycle in tumor cells
- d. Angiogenesis in tumor cells

4. The given figure shows a very important component in chromosome of tumor cells for their replication potential. Identify?




- a. Angiogenic gene
- b. Telomerase
- c. Telomere
- d. Chromosome lethality

6. A 31-year-old woman has had dull, constant abdominal pain for 6 months. CT scan of the pelvis shows a 7 cm circumscribed mass that involves the right ovary and contains irregular calcifications. The right fallopian tube and ovary are surgically excised. The gross appearance of the ovary, which has been opened, is shown in the figure. What is the most likely diagnosis?




- a. Mucinous cystadenoma
- b. Choriocarcinoma
- c. Dysgerminoma
- d. Mature cystic teratoma





## Answers of Image-Based Questions

1. **Ans. (c) Adenoma**
  - The biopsy from intestine shows benign proliferation of glandular tissue without any infiltration into deeper layers.
2. **Ans. (d) Carcinoma in situ**
  - The tissue biopsy shows malignant clonal proliferation of epithelial cells limited to ducts and lobules by the basement membrane
3. **Ans. (c) Cervical dysplasia**
  - The one on the right with red circle shows abnormal proliferation, architecture and arrangement of cells compared to normal on left side.
4. **Ans. (c) Telomere**
  - A telomere is a region of repetitive nucleotide sequences (TTAGGG) at each end of a chromosome, which protects the end of the chromosome from deterioration or from fusion with neighboring chromosomes.
5. **Ans. (a) Aerobic glycolysis-Warburg effect**
  - Cancer cells preferred to aerobic glycolysis than to oxidative phosphorylation to produce lactate .
6. **Ans. (d) Mature cystic teratoma**
  - Characteristically they are unilocular cysts containing hair and sebaceous material along with hair shafts. Within the wall, tooth structures and areas of calcification are seen.



## Multiple Choice Questions


### TYPES OF NEOPLASMS

- Which of the following are features of anaplasia?** (PGI May 18)
  - Metaplasia
  - Pleomorphism
  - Loss of cell polarity
  - Abnormal nuclear morphology
  - Abnormal mitosis
- Lack of differentiation is called?** (Recent exam 2018)
  - Anaplasia
  - Metaplasia
  - Dysplasia
  - Carcinoma
- Which of the following is the most common malignant tumor in adult males in India?** (Recent Question 2016-17)
  - Lung cancer
  - Oropharyngeal carcinoma
  - Gastric carcinoma
  - Colorectal carcinoma
- Two most common cancer in Indian woman is:** (Recent Question 2016-17)
  - Carcinoma breast
  - Carcinoma cervix
  - Carcinoma colon
  - Carcinoma stomach
- C-KIT mutations are seen in:** (Recent Question 2015)
  - Gastrointestinal stromal tumors
  - Ovarian cancers
  - Neuroblastoma
  - Small-cell carcinoma of lung
- Most common cancer in the world** (Recent Question 2015)
  - Lung
  - Breast
  - Prostate
  - Cervix
- Most common cancer in females** (Recent Question 2015)
  - Lung
  - Breast
  - Stomach
  - Cervix
- Most common cancer in males** (Recent Question 2015)
  - Lung
  - Prostate
  - Stomach
  - Colorectum
- Most common cause of cancer death** (Recent Question 2015)
  - Breast
  - Liver
  - Lung
  - Brain
- Hamartoma is:** (Recent Question 2015)
  - Malignant tumor
  - Metastatic tissue
  - Development malformation
  - Hemorrhage in vessel
- Choristoma is:** (Recent Question 2015)
  - Dilated vascular malformation
  - Malignant stroma of stem cells
  - Normal tissue at abnormal site in the body
  - Benign tumor in which normal elements become abnormally overgrown
- Choristoma is a:** (AP PGMEET 2015)
  - Heterotopic (ectopic) rest of cells
  - Example of hamartoma
  - Benign tumor of trophoblastic cell
  - Benign tumor of cartilaginous tissue

- Next to metastasis, which is the most reliable feature to differentiate benign from malignant tumors?** (Recent Question 2015)
  - Anaplasia
  - Dysplasia
  - Local invasion
  - Loss of polarity
- Number of cancer cells present in the smallest clinically detectable mass** (Recent Question 2015)
  - $10^3$  cell
  - $10^6$  cell
  - $10^9$  cell
  - $10^{12}$  cell
- Overgrowth of a bile duct at localized region is?** (Recent Question 2013)
  - Hamartoma
  - Malignant tumor
  - Choristoma
  - Polyp
- Example of autonomous hyperplasia?** (JIPMER 2012)
  - Choristoma
  - Hamartoma
  - Fibromatosis
  - Endometrial hyperplasia
- Features(s) of hamartoma is/are:** (PGI Nov 2011)
  - Benign
  - Malignant
  - Malformation
  - Mostly symptomatic
  - Neoplasms
- High risk of malignancy is seen in?** (DNB June 11)
  - Simple hyperplasia with atypia
  - Simple hyperplasia without atypia
  - Complex hyperplasia with atypia
  - Complex hyperplasia without atypia
  - Intraductal carcinoma in situ

### TUMOR SUPPRESSOR GENES

- A 75-year-old male, known smoker presented to pulmonology department with history of cough. Biopsy was taken which showed the following. What is the change shown?** (Recent Pattern Question 2020)



- Dysplasia
  - Metaplasia
  - Hyperplasia
  - Atrophy
- If DNA is damaged in the cell cycle, which gene causes cell cycle arrest?** (AIIMS May 2019)
    - Rb
    - MYC
    - p53
    - K-RAS
  - Arrange the sequence of the event of the cell cycles:** (AIIMS May 2019)
    - Cyclin D-CDK4
    - Cyclin A-CDK1
    - Cyclin B-CDK2
    - Cyclin E-CDK2



**22. Proto-oncogenes to oncogenes transformation occurs by:** (AIIMS Nov 2019)

1. Point mutation
2. Promoter insertion
3. Amplification
4. Enhancer insertion
- a. 1 and 2 are correct
- b. 1 and 3 are correct
- c. 1, 3 and 4 are correct
- d. All are correct

**23. True about p53 Gene:** (PGI May 2019)

- a. Most common gene mutation found in human cancers
- b. Causes cell cycle arrest at G1/S check point
- c. Promote transcription of cell cycle inhibitors
- d. Known as guardian of genome
- e. Regulate cellular senescence

**24. Characteristics of proto-oncogenes?** (PGI Nov 2018)

- a. One mutation is enough for causing tumors
- b. Two mutations are needed for causing tumors
- c. Mutations of proto oncogenes is in somatic cells
- d. Mutations can be transferred through germ line

**25. Which of the following mutation is seen in Cowden syndrome?** (Recent Question 2018)

- a. PTEN mutation
- b. STK11 mutation
- c. PTCH mutation
- d. SMAD4 mutation

**26. Which is not a tumour suppressor gene?** (PGI May 2017)

- a. p53
- b. CD 95
- c. RAS
- d. PTEN
- e. Stk 7/Stk11

**27. On which cell cycle checkpoints BRCA-2 acts?** (PGI Nov 2017)

- a. G2-M
- b. G1-M
- c. M phase
- d. G1 phase
- e. S phase

**28. Which of the following Dyads are correctly matched?**

- a. RB1-retinoblastoma
- b. PTEN- Melanoma
- c. BRCA2-Breast cancer
- d. n-MYC-Neuroblastoma
- e. WT1 = Wilms' tumor

**29. If the RB gene phosphorylation is defective which of the following will happen?** (AIIMS May 2017)

- a. Cell cycle will stop at G2
- b. Cell cycle will stop at G1
- c. There will be no effect on cell cycle as RB gene phosphorylation is not needed
- d. The cell cycle progresses and cell divides

**30. All are true regarding Retinoblastoma except?** (JIPMER 2017)

- a. Play major role in cell cycle regulation
- b. Require deletion of both Rb genes
- c. Autosomal Dominant
- d. Located on chr 13p14

**31. Which of the following statement is true about p53 gene?** (PGI Nov 2016)

- a. Has tyrosine kinase activity
- b. Has pro-apoptotic activity present
- c. A tumor suppressor protein
- d. Has anti-apoptotic activity

**32. Homozygous loss of the VHL tumor suppressor protein;** (JIPMER 2016)

- a. Clear cell carcinomas
- b. Papillary renal cell carcinomas
- c. Chromophobe
- d. Belini duct Ca


**33. Most common gene involved in endometrial ca is?** (JIPMER 2016)

- a. PTEN
- b. BRAF mutation
- c. KRAS
- d. Mismatch repair genes

**34. BRAF mutation is seen in?** (Recent Question 2016-17)

- a. LCH
- b. Colon Ca
- c. Hairy cell leukemia
- d. AML M7

**35. There are different check points in cell growth and regulation. Which one is the primary point for regulation of cell growth?** (AIIMS Nov 2015)



- a. End of G1
- b. Start of G2
- c. End of S
- d. End of M

**36. K-RAS protooncogene is associated with** (Recent Question 2015)

- a. Colon cancer
- b. Breast cancer
- c. Bladder cancer
- d. Melanoma

**37. RET protooncogene is associated with** (Recent Question 2015)

- a. MEN1
- b. Medullary carcinoma of thyroid
- c. Small cell carcinoma of lung
- d. Melanoma

**38. The following protein is called "Governor of proliferation"** (Recent Question 2015)

- a. RB
- b. TP53
- c. APC
- d. Patched

**39. E6 protein of high risk human papilloma viruses bind to the following protein and there by promotes carcinogenesis** (Recent Question 2015)

- a. RB
- b. TP53
- c. APC
- d. MDM2

**40. The following protein is called "Gatekeeper of colonic neoplasia"** (Recent Question 2015)

- a. APC
- b. STK11
- c. SMAD2
- d. PTEN

**41. Hall mark mechanism of tumor suppressor gene inactivation** (Recent Question 2015)

- a. Loss of heterozygosity
- b. DNA hypermethylation
- c. DNA hypomethylation
- d. Histone deacetylation

**42. Protooncogene involved in GIST** (Recent Question 2015)

- a. KIT
- b. RAS
- c. RET
- d. MYC



- 43. Tumor suppressor gene mutated in familial gastric cancer** (Recent Question 2015)  
a. APC b. CDKN2A  
c. E-cadherin d. PTEN
- 44. All are associated with BRCA mutation except** (Recent Question 2015)  
a. Ovarian carcinoma  
b. Prostate carcinoma  
c. Endometrial carcinoma  
d. Papillary serous cancer of peritoneum
- 45. SIS proto-oncogene over-expression is seen in:** (Recent Question 2015)  
a. Astrocytoma b. Breast carcinoma  
c. Melanoma d. Gastric carcinoma
- 46. Tumor suppressor gene associated with pancreatic carcinoma** (Recent Question 2015)  
a. p53 b. P16/INK4a  
c. PTEN d. PTCH1 and PTCH
- 47. BRCA 1 and BRCA 2 genes are located on chromosomes** (Recent Question 2015)  
a. 13 and 17 b. 17 and 22  
c. 17 and 13 d. 13 and 22
- 48. The following is not a cell cycle inhibitor** (Recent Question 2015)  
a. p21  
b. p27  
c. p16/INK4a  
d. Cyclin D-CDK4 complex
- 49. The following is not one of the 4 key cell cycle regulators which are dysregulated in vast majority of human cancers** (Recent Question 2015)  
a. p16/INK4a b. Cyclin D  
c. p21 d. RB
- 50. The cell cycle check point which is important in cells exposed to ionizing radiation** (Recent Question 2015)  
a. G0-G1 check point b. G1-S check point  
c. G2-M check point d. S-G2 check point
- 51. Which of the following is a Tumor suppressor gene?**  
a. RB b. MYC (JIPMER 2015)  
c. RAS d. RET
- 52. AKT-1 gene mutation is associated with?** (Recent Question 2015)  
a. Stomach b. Breast  
c. Ovary d. Pancreas
- 53. CDK4 positive tumors are?** (PGI Nov 2015)  
a. Melanomas b. Sarcomas  
c. Glioblastomas d. Lobular Ca breast  
e. Prostate Ca
- 54. Gain of function in RAS gene mutation is equal to loss of function mutation in which of the following?** (Recent Question 2015)  
a. Rb b. bcr-tyrosine kinase  
c. bcl-2 d. c-myc  
e. GAP protein
- 55. Cyclin D dependent proteins are?** (PGI May 2014)  
a. CDK-1 b. CDK-2  
c. CDK-3 d. CDK-4  
e. CDK-6
- 56. Cells are most radio-resistant in** (AP PGME 14)  
a. S phase b. M phase  
c. G2 phase d. G1 phase
- 57. MYC gene is:** (Recent Question 2013)  
a. Protein kinase inhibitor b. Growth factor inhibitor  
c. GTPase d. Transcription activator
- 58. The most radiosensitive phase of cell cycle is:** (MH 16)  
a. G1 b. G2  
c. S d. M
- 59. Chromosome duplication takes place in:** (Recent Question 2014)  
a. M phase b. G0 phase Pattern  
c. S phase d. G2 phase
- 60. True about protooncogene:** (PGI May 2013)  
a. Regulate cell growth and gene expression  
b. Found in normal cells  
c. Induced by virus  
d. Inactivated by virus  
e. May convert to oncogene
- 61. Cyclin dependent kinase-2 (CDK-2) acts via:** (PGI May 12)  
a. Cyclin A b. Cyclin B  
c. Cyclin C d. Cyclin D  
e. Cyclin E
- 62. Protein structure on chromatids where the spindle fibers attach during cell division is called?** (DNB Aug 12 Pattern)  
a. Nucleolus b. Satellite  
c. Kinetochore d. Centromere
- 63. N-MYC amplification is associated with which tumor?** (COMEDK 11)  
a. Burkitt lymphoma  
b. Squamous cell carcinoma lung  
c. Astrocytoma  
d. Neuroblastoma
- 64. Which among the following pairs of Oncogenes is activated by Translocation?** (DNB Dec 10)  
a. SIS and HST-1 b. HGF and L-MYC  
c. TGF and CDK4 d. ABL and C-MYC
- 65. Which of the following genes is growth promoting marker for oncogenesis?** (MH 16)  
a. Rb gene b. RAS gene  
c. p53 gene d. BRCA 1 gene
- 66. Tumor suppressor gene is not involved in:** (Recent Question 2014)  
a. Breast cancers  
b. Neurofibromatosis  
c. Multiple endocrine neoplasia  
d. Retinoblastoma
- 67. Cancers are usually associated with?** (AIIMS Nov 2014)  
a. Hypomethylation of oncogenes  
b. Methylation of tumor suppressor genes  
c. Loss of heterozygosity  
d. Mutation in introns
- 68. RET gene mutation is associated with?** (Recent Question 2013)  
a. Pheochromocytoma  
b. Medullary carcinoma thyroid  
c. Lymphoma  
d. Renal cell carcinoma





**69. Which of the following tumor suppressor gene mutation occurs in endometrial carcinoma?**

(Recent Question 2013)

- a. P53
- b. Rb
- c. PTEN
- d. APC

**70. E-cadherin gene deficiency is seen in:**

(Recent Question 2013)

- a. Gastric Ca
- b. Intestinal Ca
- c. Thyroid Ca
- d. Pancreatic Ca

**71. Li Fraumeni syndrome is due to mutation of which gene:**

(Recent Question 2013)

- a. p21
- b. p53
- c. p41
- d. p43

**72. Gene for Wilms' tumor is located on:**

(Recent Question 2013/JIPMER 12)

- a. Chromosome 1
- b. Chromosome 10
- c. Chromosome 11
- d. Chromosome 12

**73. True statements about P53. gene are all except:**

(AI 08, AIIMS Nov 10, DNB June 10)

- a. Arrests cell cycle at G1 Phase
- b. Product is 53 KD protein
- c. Located on chromosome 17
- d. Wild form is associated with increased risk of childhood tumors

**74. BRCA-1 Gene is located on:** (AIIMS Nov 08)(MH 2015)

- a. Chromosome 13
- b. Chromosome 11
- c. Chromosome 17
- d. Chromosome 22

#### OTHER HALLMARKS OF CANCER

**75. Gene promoting metastasis:** (Recent Question 2015)

- a. SNAIL
- b. TWIST
- c. Both
- d. None

**76. Basement membrane degradation is mediated by**

(Recent Question 2015)

- a. Oxidase
- b. Elastase
- c. Metalloproteinase
- d. Myeloperoxidase

**77. Excessive fibrosis in a tumor is called:**

(Recent Question 2014)

- a. Anaplasia
- b. Metaplasia
- c. Desmoplasia
- d. Dysplasia

**78. Dysplasia is characterized by all except:**

(Recent Question 2013)

- a. High nuclear to cytoplasmic ratio
- b. Loss of architecture
- c. Pleomorphism
- d. Invasion

**79. Feature of dysplasia are:**

(PGI May 2011)

- a. Prominent nucleus
- b. Nuclear enlargement
- c. Nuclear hyperchromia
- d. Coarsening and clumping of chromosome
- e. Increased apoptosis

**80. All of the following are angiogenic factors EXCEPT:**

(DNB Dec 10)

- a. VEGF
- b. Platelet activating factor
- c. IFN $\alpha$
- d. TGF- $\beta$

#### CARCINOGENESIS

**81. Cancers associated with viruses?** (PGI May 2017)

- a. Hepatocellular cancer
- b. Kaposi sarcoma
- c. Nasopharyngeal cancer
- d. Small cell Ca lung
- e. Prostatic Ca

**82. Human papilloma virus is/are associated with all except:** (Recent Question 2016-17)

- a. Oropharyngeal tumors
- b. Carcinoma nasopharynx
- c. Carcinoma anal canal
- d. Carcinoma pancreas
- e. Carcinoma cervix

**83. Smoking increases risk of all the following cancers except** (Recent Question 2015)

- a. Prostate
- b. Oral cavity
- c. Bladder
- d. Pancreas

**84. Vinyl chloride is associated with** (Recent Question 2015)

- a. Hepatoblastoma
- b. Liver angiosarcoma
- c. Hepatocellular carcinoma
- d. Hemangioblastomas

**85. Cancer which is more prevalent in less developed countries than developed countries**

(Recent Question 2015)

- a. Lung
- b. Breast
- c. Stomach
- d. Esophagus

**86. The following is not an initiator of carcinogenesis in tobacco smoke** (Recent Question 2015)

- a. Nicotine
- b. Tar
- c. Polycyclic aromatic hydrocarbons
- d. Nitrosamine

**87. The initial hematopoietic change after total body irradiation** (Recent Question 2015)

- a. Neutropenia
- b. Anemia
- c. Lymphopenia
- d. Thrombocytopenia

**88. Skin cancer is caused by exposure to** (MH PG 2014)

- a. Asbestos
- b. Arsenic
- c. Nitrosamine
- d. Vinyl chloride

**89. Least affected by radiation is:** (Recent Question 2014)

- a. GIT
- b. Bone marrow
- c. Cartilage
- d. Lymphocytes

**90. Kaposi sarcoma is seen with:** (Recent Question 2014)

- a. HCV
- b. HPV
- c. HSV
- d. HHV-8

**91. Which of the following human papilloma viruses (HPV) is a low risk oncogenic virus?** (AP PGME 14)

- a. 11
- b. 16
- c. 18
- d. 31

**92. Human Papilloma Virus does not cause:**

(PGI May 2013)

- a. Oropharyngeal carcinoma
- b. Cervical carcinoma
- c. Esophageal carcinoma
- d. Cutaneous carcinoma
- e. Burkitt's lymphoma

**93. Pathogenic mechanism of HPV in cervical Ca**

(JIPMER 2013)

- a. Down-regulation of P16INK4
- b. Degradation of Cyclin D1
- c. Instability of E6 and E7
- d. Up-regulation of BCL-2



- 94. UV radiation:** (Recent Question 2013)  
 a. Prevents formation of Pyrimidine dimers  
 b. Stimulates formation of Pyrimidine dimers  
 c. Purine dimers  
 d. Single gene deletion

- 95. The least radio sensitive tissue is:** (AI 10)  
 a. Nervous tissue                      b. Bone  
 c. Kidney                                d. Thyroid

### GENETIC BASIS OF CANCERS

- 96. All of the following are hereditary tumors except?** (Recent Question 2016-17)  
 a. Retinoblastoma                      b. Ewings Sa  
 c. Ca breast                              d. Nasopharyngeal Ca  
 e. Ca pancreas

- 97. Tumors associated with AIDS?** (PGI Nov 2016)  
 a. DLBCL                                b. Ca breast  
 c. Ca pancreas                          d. Kaposi Sarcoma  
 e. Ca lung

- 98. Chromophthis mechanism of carcinogenesis has been found to be associated with which of the following tumors?** (Recent Question 2016-17)  
 a. Sarcomas                              b. Osteosarcoma  
 c. RCC                                    d. ALL

- 99. True statement of Ataxia telangiectasia is?** (PGI Nov 2016)  
 a. Seizure  
 b. IgA is low or absent  
 c. Increased risk of Leukemia  
 d. Hypoplasia of thymus  
 e. Adaptive immune system is normal

- 100. Which of the following marker favours diagnosis of preinvasive and invasive cervical cancer:** (Recent Question 2016-17)  
 a. Ki67  
 b. Oncoprotein E6  
 c. p16INK4, cyclin E, and Ki-67  
 d. Oncoprotein E8

- 101. Which of the following marker/mutation is/are seen in papillary carcinoma of thyroid:** (Recent Question 2016-17)  
 a. Synaptophysin                      b. RET/PTC  
 c. P53                                      d. NTRK1  
 e. RAS

- 102. Chromothrips is false statement** (Recent Question 2015)  
 a. Include simple deletions, inversions and translocations in chromosomes  
 b. Found in higher frequencies in osteosarcomas and bone cancers  
 c. Hundreds of chromosome breaks occur across single or several chromosomes  
 d. DNA is repaired in a haphazard manner

- 103. BRAF mutation is seen in?** (Recent Question 2015)  
 a. Melanoma  
 b. Squamous Cell Ca  
 c. Papillary Ca thyroid  
 d. Retinal hemangioblastoma

- 104. Most common translocation in Ewing's sarcoma is:** (Recent Question 2015)  
 a. t(11;22)                                b. t(9;22)  
 c. t(8;14)                                d. t(x;11)

### PARANEOPLASTIC SYNDROMES

- 105. Paraneoplastic syndrome(s) associated with lymphoma?** (PGI Nov 2017)  
 a. SIADH                                b. Hypercalcemia  
 c. Cushing's syndrome              d. Acanthosis nigricans

- 106. All are signs of paraneoplastic syndrome in skin except?** (PGI Nov 2016)  
 a. Tripe palm  
 b. Acanthosis nigricans  
 c. Superficial thrombophlebitis  
 d. Dermatomyositis  
 e. Osler's node

- 107. Malignancy associated with hypercalcemia** (PGI Nov 2016)  
 a. Breast cancer                        b. Prostate cancer  
 c. Small cell lung cancer              d. Non-small lung cancer

- 108. Migratory thrombophlebitis is seen in all except?** (JIPMER 2012, AI 08)  
 a. Lung Ca                                b. Prostate Ca  
 c. Colon Ca                                d. Pancreatic Ca

### LABORATORY DIAGNOSIS OF CANCER

- 109. Tumour causing phosphaturia and osteomalacia is?** (AIIMS Nov 18)  
 a. Peripheral nerve sheath  
 b. Meningioma  
 c. Fibrosarcoma  
 d. Osteosarcoma

- 110. Sentinel lymph node biopsy is most significant for ?** (AIIMS Nov 18)  
 a. Ca Vulva                                b. Ca Vagina  
 c. Ca Endometrium                      d. Ca Cervix

- 111. Nobel Prize for 2018 was awarded for what contribution?** (AIIMS Nov 18)  
 a. Negative immune regulation in treatment of cancer  
 b. CAS9-CRISPER  
 c. DNA repair syndromes  
 d. Stem cell transplant

- 112. Gastrin is a marker of which carcinoma?** (JIPMER 18)  
 a. Medullary carcinoma of thyroid  
 b. GIST  
 c. Gastric carcinoma  
 d. Pancreatic carcinoma/neuroectodermal tumour

- 113. Markers of carcinoma pancreas ?** (PGI Nov 2018)  
 a. Ca19-9                                b. Ca 125  
 c. CEA                                      d. AFP

- 114. Sarcoma on paraffin mount shows what markers?** (PGI Nov 2018)  
 a. Desmin                                b. Vimentin  
 c. PAX8                                    d. WT1  
 e. HMB 45



115. A 5-year-old child was presenting with proptosis. Microscopic examination has revealed round cell tumor and positive for desmin immunohistochemical marker. Most likely diagnosis is? (AIIMS May 18)
  - a. Leukemia
  - b. Embryonal rhabdomyosarcoma
  - c. Lymphoma
  - d. Primitive Neuroectodermal Tumor (PNET)
116. Biopsy from an eight-year-old child with leg swelling was showing small round blue tumor cells consistent with diagnosis of Ewing's sarcoma. What will be the best method to detect translocation t (11;22) in this malignancy? (AIIMS May 18)
  - a. Conventional karyotyping
  - b. Next generation sequencing
  - c. FISH
  - d. PCR
117. Which of the following immunohistochemical marker is positive in neuroendocrine tumor? (Recent exam 2018)
  - a. Cytokeratin
  - b. Synaptophysin
  - c. Calretinin
  - d. GFAP
118. Which of the following neoplasms shows ALK positivity? (AIIMS May 2017)
  - a. Ewing sarcoma
  - b. Inflammatory myofibroblastic tumor
  - c. Synovial sarcoma
  - d. Fibromatosis
119. Fixative agent for PAP smear? (AIIMS Nov 2017)
  - a. Norma saline
  - b. Formalin
  - c. 95% ethanol
  - d. Air drying
120. Not seen in tumour lysis syndrome? (AIIMS Nov 2017)
  - a. Hyperphosphatemia
  - b. Hyperuricaemia
  - c. Hypercalcemia
  - d. Hyperkalaemia
121. In CA breast, based on Which stage/grade of IHC staining, FISH for gene amplification will be done? (AIIMS Nov 2017)
  - a. Her2 neu 3+
  - b. Her2 Nu 2+
  - c. Her2 neu +
  - d. Will be done irrespective of above
122. ALK1 mutation seen mc with which hereditary hemorrhagic telangiectasia? (JIPMER 2017)
  - a. Type 1
  - b. Type 2
  - c. Type 3
  - d. Type 4
123. Carbohydrate marker of recurrent breast cancer is? (JIPMER 2017)
  - a. CA 15.3
  - b. CA 125
  - c. CEA
  - d. CA 19.9
124. For detection of carcinoma lip, stain used is? (AIIMS Nov 2016)
  - a. Giemsa
  - b. Crystal violet
  - c. Toulidine blue
  - d. Hematoxylin and eosin
125. AFP and CEA both rise in (Recent Question 2016-17)
  - a. Testicular ca
  - b. Hepatic ca
  - c. RCC
  - d. Germ cell tumor of ovary
126. A surgeon suspecting testicular carcinoma in a patient asks the intern to send the sample for histopathology, what is the fluid in which the intern should send the sample to the pathologist? (AIIMS May 2016)
  - a. Bouvin solution
  - b. 10% formalin
  - c. 95% ethanol
  - d. Alcohol
127. High level of  $\beta$  hCG is/are seen in all except: (Recent Question 2016-17)
  - a. Down syndrome
  - b. Neural tube defect
  - c. Germ cell tumor
  - d. Gestational trophoblastic disease
  - e. Multiple pregnancy
128. Neuroendocrine cell tumor markers are: (Recent Question 2016-17)
  - a. Chromogranin A
  - b. CD56
  - c. Neuron-specific enolase
  - d. Synaptophysin
  - e. Cytokeratin 7
129. Not true about cancer cachexia (Recent Question 2015)
  - a. Equal loss of both fat and lean muscle
  - b. Elevated basal metabolic rate
  - c. No evidence of systemic inflammation
  - d. TNF $\alpha$  (cachetin) plays an important role
130. Grading of cancers is does not depend on (Recent Question 2015)
  - a. Degree of differentiation
  - b. Number of mitoses
  - c. Architectural features
  - d. Size of primary lesion
131. Tumor markers and tumors -find the wrong match (Recent Question 2015)
  - a. Desmin-Carcinoma
  - b. Vimentin-sarcoma
  - c. Leucocyte specific antigen-lymphoma
  - d. S100-melanoma
132. Acid phosphatase is a tumor marker for (Recent Question 2015)
  - a. Pancreatic carcinoma
  - b. Prostatic carcinoma
  - c. Papillary carcinoma of thyroid
  - d. Renal cell carcinoma
133. Tumor marker for non-seminomatous testicular tumors (Recent Question 2015)
  - a. CEA
  - b. CA-125
  - c. CA-19-9
  - d. Alpha-fetoprotein
134. HMB-45 is positive in (Recent Question 2015)
  - a. Small cell carcinoma of lung
  - b. Adenocarcinoma of lung
  - c. Squamous cell carcinoma of lung
  - d. Angiomyolipoma of lung
135. CA-125 is used is the monitoring and treatment of (Recent Question 2015)
  - a. Ovarian cancer
  - b. Colon cancer
  - c. Breast cancer
  - d. Pancreatic cancer
136. An undifferentiated malignant tumor on immuno-histochemical stain shows cytoplasmic positivity of most of the tumor cells for cytokeratin. The most probable diagnosis of the tumor is: (MH 16)
  - a. Lymphoma
  - b. Carcinoma
  - c. Sarcoma
  - d. Malignant melanoma



- 137. Hypercalcemia as a paraneoplastic syndrome in lymphomas is due to elaboration of** (Recent Question 2015)
- PTHrP
  - PTH
  - 1,25dihydroxy vitamin D
  - PGE2
- 138. Molecular profiling of cancer cells is obtained by** (Recent Question 2015)
- Flow cytometry
  - Immunohistochemistry
  - DNA microarray analysis
  - PCR
- 139. Antibody not associated with paraneoplastic cerebellar degeneration** (Recent Question 2015)
- Anti-Yo
  - Anti-Hu
  - Anti-Ri
  - Anti-Tr
- 140. Alpha feto protein is/are increased in:** (PGI May 2015)
- Yolk sac tumor
  - Seminoma
  - Dysgerminoma
  - Non-seminoma
  - Hepatocellular carcinoma
- 141. Hypoglycemia occur as a paraneoplastic syndrome in:** (Recent Question 2015)
- Bronchogenic carcinoma
  - CA pancreas
  - Fibrosarcoma
  - RCC
- 142. Test for carcinogenicity** (Recent Question 2015)
- Kveim's test
  - Ame's test
  - Schilling's test
  - Schick test
- 143. Bortezomib, a proteasome inhibitor is used in the treatment of** (Recent Question 2015)
- AML M3
  - CML
  - Multiple myeloma
  - GIST
- 144. Chemical carcinogen associated with cancer of renal pelvis** (Recent Question 2015)
- Polycyclic hydrocarbons
  - Cyclophosphamide
  - Asbestos
  - Cadmium
- 145. Paraneoplastic neurologic syndrome not associated with antibodies** (Recent Question 2015)
- Lambert-Eaton myasthenic syndrome
  - Stiff pearson syndrome
  - Limbic encephalopathy
  - Necrotizing myelopathy
- 146. The hallmark change in cell physiology that determines malignant phenotype** (Recent Question 2015)
- Self sufficiency in growth signals
  - Sustained angiogenesis
  - Resistance to apoptosis
  - Aerobic glycolysis
- 147. Keratinization and pearl formation is seen in?** (Recent Question 2015)
- Squamous cell Ca
  - Adenocarcinoma
  - Small cell Ca
  - Basal cell Ca
- 148. Which of the following is not correctly matched?** (AIIMS Nov 14)
- Melanoma - s100
  - Carcinoma - desmin
  - Sarcoma - vimentin
  - Lymphoma -leucocyte common antigen
- 149. Calcitonin is a marker of:** (Recent Question 2014)
- Prostate cancer
  - Medullary carcinoma of thyroid
  - Pheochromocytoma
  - Pancreatic cancer
- 150. MIC-2 is a marker of?** (DNB Aug 12 Pattern)
- Ewing sarcoma
  - Chronic lymphocytic leukemia
  - Mantle cell lymphoma
  - All of these
- 151. Not a marker for muscle tumor:** (Recent Question 2014)
- Desmin
  - Actin
  - Neurofilament
  - Intermediate filament
- 152. The following is not a marker of melanoma:** (JIPMER 2014)
- S-100
  - MITF
  - CK-20
  - Vimentin
- 153. Immunohistochemical marker for Rhabdomyosarcoma is:** (JIPMER 2014)
- Desmin
  - Vimentin
  - Cytokeratin
  - Neurofilament
- 154. SS18 -SSX1 gene is associated with:** (Recent Question 2013)
- Liposarcoma
  - Rhabdomyosarcoma
  - Synovial sarcoma
  - Ewing's sarcoma
- 155. Maternal serum AFP raised in:** (PGI May 2013)
- Gestational Trophoblastic disease
  - Down syndrome
  - Omphalocele
  - Sacroccygealtetatomia
  - Neural tube defect
- 156. True about Carcino embryonic antigen (CEA) is/are:** (PGI May 2013)
- Used for monitoring of recurrence of colon cancer
  - Specific for colon cancer
  - Increased in smokers
  - Increased in colon cancer
  - Increased in Renal Carcinoma
- 157. True about CA-125 is/are:** (PGI May 2013)
- It is a Glycoprotein
  - It is a specific marker
  - It is Increased in colon carcinoma
  - Normal range in pre-menopausal females is 200 U/ml
  - May be elevated in Pelvic inflammatory disease
- 158. Which amongst the following a marker of Carcinoma:** (AI 12)
- Cytokeratin
  - Vimentin
  - Calretinin
  - CD45
- 159. Conventional cytogenetics are difficult in solid tumors especially in case of carcinoma cervix because of:** (AIIMS Nov 10)
- High mitotic rate
  - Bacterial contamination of the specimen
  - Good metaphase activity
  - Inadequate biopsy specimen





**160. Increased LDH helps in diagnosis of?**

(DNB Aug 12)

- Prostate carcinoma
- Hepatocellular carcinoma
- Pancreatic carcinoma
- Renal cell carcinoma

**161. AFP is elevated in:**

(PGI May 2011)

- HCC
- Hepatoblastoma
- Infant hemangioendothelioma
- Amebic liver abscess
- Embryonic sarcoma

**162. Ames test in neoplasia is a test for:**

(AP 2013)

- Teratogenicity
- Mutagenicity
- Carcinogenicity
- Clonality

**163. Elevated CA -125 are seen in:**

(PGI Nov 10)

- Abdominal TB
- Ca cervix
- Endometriosis
- Ovarian Ca
- Endometrial Ca

**164. A 67-year male smoker presents with hemoptysis and cough. Bronchoscopic biopsy revealed undifferentiated tumor. The immunohistochemical marker that will be most helpful is:**

(DNB June 10, AIIMS Nov 09)

- Calretinin
- Vimentin
- Cytokeratin
- TTF1

**165. Which of the following is not a tumor marker:**

(DNB Dec 10)

- CEA
- Tyrosinase
- Human leucocyte antigen
- AFP



## Answers with Explanations

**1. (b, c, d, e) b. Pleomorphism c. Loss of cell polarity d. Abnormal nuclear morphology; e. Abnormal mitosis**

**2. Ans. (a) Anaplasia** (Ref: Robbins 9th ed p 269)

Malignant neoplasms that are composed of poorly differentiated cells are said to be anaplastic. Lack of differentiation, or anaplasia, is considered a hallmark of malignancy. The term anaplasia means "to form backward," implying a reversal of differentiation to a more primitive level.

**3. Ans. (b) Oropharyngeal carcinoma**

(Ref: Harrison's 19th ed/pg 467; table 99.1)

- Most common cancer all over the world (overall, considering both sexes together): **Lung cancer**
- Most common cancer all over the world: **Prostate cancer**
- Most common cancer (females) all over the world: **Breast cancer**
- Most common cause of cancer death in the world: **Lung cancer**
- 2<sup>nd</sup> most common cancer worldwide (overall): **Breast cancer**
- Causes of cancer death: lung > stomach > liver > colorectal > breast

**4. Ans. (a) Carcinoma breast, b. Carcinoma cervix**

(Ref: Park 23rd/ 382-83; Harshmohan 7th/198)

**5. Ans. (a) Gastrointestinal stromal tumors**

(Ref: Robbins 9th/pg 298; 8th/pg 295)

**c-Kit**

- It is a proto oncogene

- It is also known as Mast/stem cell growth factor receptor (SCFR), tyrosine-protein kinase Kit or CD117
- It is a receptor tyrosine kinase protein that in humans is encoded by the KIT gene.
- Activating mutations in this gene are associated with gastrointestinal stromal tumors, testicular seminoma, mast cell disease, melanoma, acute myeloid leukemia
- Inactivating mutations are associated with the genetic defect piebaldism.

**6. Ans. (a) Lung**

(Ref: Harrison's 19th ed/pg 467; table 99.1)

**7. Ans. (b) Breast**

(Ref: Harrison's 19th ed/pg 467; table 99.1)

**8. Ans. (b) Prostate**

(Ref: Harrison's 19th ed/pg 467; table 99.1)

**9. Ans. (c) Lung**

(Ref: Harrison's 19th ed/pg 467; table 99.1)

**10. Ans. (c) Development malformation**

(Ref: R 9th/pg 267)

**Hamartoma:**

- It is a **disorganized but benign-appearing mass** composed of cells **indigenous to that particular site**.
- Previously it was considered a **developmental malformation** but, hamartomas have been found to have **clonal recurrent translocations** involving genes encoding certain chromatin proteins



- So, hamartomas are **considered neoplasms now**.
- E.g., Pulmonary chondroid hamartoma contains islands of disorganized, but histologically normal cartilage, bronchi, and vessels.

**11. Ans. (c) Normal tissue at abnormal site in the body**

(Ref: Robbins 9th/pg 267; 8th/pg 262)

**12. Ans. (d) Benign tumor of cartilaginous tissue**

(Ref: Robbins 9th/pg 267; 8th/pg 262)

**13. Ans. (c) Local invasion**

(Ref: Robbins 9th/pg 268; 8th/pg 263)

**14. Ans. (c) 10<sup>9</sup> cell**

(Ref: Robbins 9th/pg 267; 8th/pg 268)

**15. Ans. (a) Hamartoma** (Ref: Robbins 9th/pg 267)

**Von Meyenburg Complexes (bile duct hamartomas)** are small clusters of dilated bile ducts embedded in a fibrous, sometimes hyalinized, stroma located close to or within portal tracts.

**16. Ans. (c) Fibromatosis**

(Ref: Underwood's Pathology, by Simon Cross: Chapter 4)

**Fibromatoses** are apparently **autonomous proliferation of myofibroblasts**, occasionally forming **tumor like** masses. Eg Palmar Fibromatoses (Dupuytren's contractures), Desmoid tumor, Retroperitoneal fibromatosis and Peyronie's disease of penis.

**17. Ans. (c, e) c. Malformation; e. Neoplasms**

(Ref: Robbins 9th/pg 267; 8th/pg 262; Refer Ans 6)

**18. Ans. (c, e) c. Complex hyperplasia with atypia; e. Intraductal Carcinoma in situ**

(Ref: Robbins 9th/pg 270)

Discussing the options one by one:

Lesions	Relative Risk of malignancy	Description
<b>A. Simple hyperplasia with atypia</b>	1	Proliferation of ductal epithelium and/or stroma <b>without cytologic or architectural features suggestive of carcinoma in situ</b>
<b>B. Simple hyperplasia without atypia</b>	≤1	Epithelial hyperplasia i.e. presence of <b>&gt; 2 cell layers</b>

Contd...

Lesions	Relative Risk of malignancy	Description
<b>C. Complex hyperplasia with atypia</b>	4-5	Cellular proliferation <b>resembling carcinoma in situ</b> but lacking sufficient features for diagnosis as carcinoma
<b>D. Complex hyperplasia without atypia</b>	1.5-2	Individual cells may be <b>enlarged</b> , but, as in simple hyperplasia, the internal makeup of the cells is considered to be <b>normal</b>
<b>E. Intraductal Carcinoma in-situ</b>	8 -10	<b>Dysplastic changes are marked</b> and involve the <b>full thickness of the epithelium</b> , but <b>does not penetrate the basement membrane</b>

**19. Ans. (b) Metaplasia** (Ref: Robbins 9th ed/pg 270)

**20. Ans. (c) p53** (Ref: Robbins 9th ed/pg 294)

**21. Ans. (A→D→B→C)** (Ref: Robbins 9th ed/pg 25)

**22. Ans. (b) 1 and 3 are correct** (Ref: Robbins 9th ed)

**23. Ans. (a, b, c, d, e) a. Most common gene mutation found in human cancers; b. Causes cell cycle arrest at G1/S check point; c. Promote transcription of cell cycle inhibitors; d. Known as guardian of genome; e. Regulate cellular senescence** (Ref: Robbins 9th ed/pg 294)

**24. (a, c, d) a. One mutation is enough for causing tumors; c. Mutations of proto oncogenes is in somatic cells; d. Mutations can be transferred through germ line**

Oncogenes are produced from proto-oncogene by point mutations (single mutations) in both somatic cells or germ cells.

**25. Ans. (a) PTEN mutation**

(Ref: Robbins 9th ed p 298)

PTEN (phosphatase and tensin homologue) is a membrane-associated phosphatase encoded by a gene on chromosome 10q23 that is mutated in Cowden syndrome, an autosomal dominant disorder marked by frequent benign growths, such as skin appendage tumors, GI, and CNS growths; breast, endometrial, and thyroid carcinoma

**26. Ans. (b, c) b. CD 95; c. RAS**

**27. Ans. (a, e) a. G2-M; b. S phase** (Ref: R 315)

BRCA1 deficiency causes abnormalities in the S-phase checkpoint, the G<sub>2</sub>/M checkpoint, the spindle checkpoint and centrosome duplication. Defects in this pathway leads to the activation of the salvage nonhomologous end joining pathway, formation of dicentric chromosomes, bridge-fusion-breakage cycles, and massive aneuploidy.



28. **Ans. (a, c, d, e) a. RB1-retinoblastoma; c. BRCA2-Breast cancer; d. n-MYC-Neuroblastoma; e. WT1 = Wilms' tumor** (Ref: Robbins 291)

PTEN (phosphatase and tensin homologue) mutated in Cowden syndrome, an autosomal dominant disorder marked by frequent benign growths, such as skin appendage tumors, GI, and CNS growths; breast, endometrial, and thyroid carcinoma.

29. **Ans. (b) Cell cycle will stop at G1** (Ref: R9/ 290-292)

- When hypophosphorylated, RB exerts antiproliferative effects by binding and inhibiting E2F transcription factors that regulate genes required for cells to pass through the G1-S phase cell cycle checkpoint. Normal growth factor signalling leads to RB hyperphosphorylation and inactivation, thus promoting cell cycle progression. Thus defective phosphorylation will result in RB gene stopping the cell cycle at G1S transition

30. **Ans. (d) Located on chr 13p14**

(Ref: Rb gene is located on Chr 13q14)

31. **Ans. (a, b, c) a. Has tyrosine kinase activity, b. Has pro-apoptotic activity present, c. A tumor suppressor protein**

(Ref: Robbins 9th/294)

32. **Ans. (a) Clear cell carcinomas**

(Ref: Robbins 9th/pg 721; 8th/pg 730)

33. **Ans. (a) PTEN** (Ref: Robbins 9th/pg 721; 8th/pg 730)

34. **Ans. (c) Hairy cell leukemia** (Ref: Robbins 9th/1014)

35. **Ans. (a) End of G1** (Ref: Robbins 9th/pg 25; 8th/pg 285)

G1-S is the primary point for regulation of cell growth

36. **Ans. (a) Colon cancer**

(Ref: Robbins 9th/pg 721; 8th/pg 730)

37. **Ans. (b) Medullary carcinoma of thyroid**

(Ref: Robbins 9th/pg 721; 8th/pg 730)

38. **Ans. (a) RB** (Ref: Robbins 9th/pg 294; 8th/pg 291)

RB gene (Chr 13q14) is the 'Governor of proliferation' and Key **negative regulator** of G1/S cell cycle transition

39. **Ans. (b) TP53** (Ref: Robbins 9th/pg 294; 8th/pg 291)

- HPV's genome consists of an early (E) gene region, a late (L) gene region, and a noncoding region that contains regulatory elements.
- E1, E2, E5, E6, and E7 proteins are expressed early in the growth cycle and are necessary for viral replication and cellular transformation.

- E6 and E7 proteins cause malignant transformation by targeting the human cell cycle regulatory molecules p53 and Rb (retinoblastoma protein), respectively, for degradation.

40. **Ans. (a) APC** (Ref: Robbins 9th/pg 288; 8th/pg 284)

#### Adenomatous polyposis coli (APC) gene

- On Chr 5q21<sup>Q</sup>
- Gate-keeper of Colonic Neoplasia<sup>Q</sup>
- Component of the WNT signaling pathway<sup>Q</sup>
- Controls cell fate, adhesion, and cell polarity during embryonic development.
- APC **controls** oncogenic effects of  $\beta$ -catenin so prevents carcinomas
- Germline loss-of-function mutation  $\rightarrow$  **Familial adenomatous polyposis colon Cancer**; (AD)<sup>Q</sup>
- Other tumors induced: **Hepatoblastomas, Hepatocellular carcinomas**<sup>Q</sup>

41. **Ans. (a) Loss of heterozygosity** (Ref: Robbins 9th/pg 288)

42. **Ans. (a) KIT** (Ref: Robbins 9th/pg 288; 8th/pg 285)

43. **Ans. (c) E-cadherin** (Ref: Robbins 9th/pg 288; 8th/pg 285)

#### E-cadherin

- Loss-of-contact inhibition, by mutation of the E-cadherin/ $\beta$ -catenin axis is a key characteristic of carcinomas.
- Loss of E-cadherin contributes to the malignant phenotype by allowing easy disaggregation of cells, which can then invade locally or metastasize.
- Reduced cell surface expression of E-cadherin has been seen in Ca **esophagus, colon, breast, ovary, and prostate**.
- Germline loss- of-function mutations of the E-cadherin gene, known as CDH1, cause **familial gastric carcinoma**
- Some of the sporadic gastric carcinomas are also associated with loss of E-cadherin expression.

44. **Ans. (c) Endometrial carcinoma**

(Ref: Robbins 9th/pg 288)

45. **Ans. (a) Astrocytoma**

(Ref: Robbins 9th/pg 288; 8th/pg 285)

46. **Ans. (b) P16/INK4a**

(Ref: Robbins 9th/pg 288; 8th/pg 285)

Germline mutations of *p16* (*CDKN2A*) are present in 25% of melanoma-prone kindreds, Somatic acquired deletion or inactivation of *p16* is seen in:

- 75% of pancreatic carcinomas, glioblastomas, esophageal cancers, and non-small-cell lung carcinomas.



47. Ans. (c) **17 and 13** (Ref: Robbins 9th/pg 288; 8th/pg 285)

48. Ans. (d) **Cyclin D-CDK4 complex** (Ref: R 9th/pg 25)

49. Ans. (c) **p21** (Ref: Robbins 9th/pg 288; 8th/pg 285)

**Loss of normal cell cycle control is central to malignant transformation and that at least one of four key regulators of the cell cycle (p16/INK4a, cyclin D, CDK4, RB) is dysregulated in the vast majority of human cancers.**

50. Ans. (c) **G2-M check point** (Ref: Robbins 9th/pg 25)

Cells exposed to **ionizing radiation** → Cell cycle **arrested** in G<sub>2</sub> and **repair mechanisms activated**. Thus G2-M check point is important in cells exposed to ionizing radiation

51. Ans. (a) **RB** (Ref: Robbins 9th/pg 292; 8th/pg 288)

52. Ans. (b) **Breast** (Ref: Robbins 9th/pg 292; 8th/pg 288)

PTEN acts as a tumor suppressor by serving as a brake on the PI3K/AKT arm of the receptor tyrosine kinase pathway PTEN (phosphatase and *tensin* homologue) is a membrane-associated phosphatase, that is encoded by a gene on chromosome 10q23

It is mutated in Cowden syndrome, an autosomal dominant disorder marked by skin appendage tumors, and an increased incidence of epithelial cancers, particularly of the breast, endometrium, and thyroid

53. Ans. (a, b, c); **a. Melanomas; b. Sarcomas; c. Glioblastomas**

(Ref: Robbins 9th/pg 292; 8th/pg 288)

CDK4 activation by amplification or point mutation is seen in Glioblastoma, melanoma, sarcoma

54. Ans. (e) **GAP protein**

(Ref: Robbins 9th/pg 292; 8th/pg 288)

- RAS has an intrinsic GTPase activity that is accelerated by *GTPase-activating proteins* (GAPs), which bind to the active RAS and augment its GTPase activity by more than 1000-fold, thereby terminating signal transduction.
- Thus, GAPs prevent uncontrolled RAS activity.
- Gain-of-function mutations in RAS proteins and loss-of-function mutations in GAPs lead to unchecked proliferation of cells.

55. Ans. (d) **CDK-4; e. CDK-6** (Ref: Robbins 9th/pg 25)

56. Ans. (a) **S phase**

(Ref: Robbins 9th/pg 25; 8th/pg 285; *Cancer, By David Morris, 2003, Chapter 5*)

Cells are **most radio-resistant in S phase** and **most radio-sensitive in M > G<sub>2</sub> phase**.

57. Ans. (d) **Transcription activator**

(Ref: Robbins 9th/pg 284)

## MYC gene

- MYC is a **transcription factor** that acts to **reprogram somatic cells into pluripotent stem cells**.
- MYC translocations are seen in **Burkitt lymphoma** and is amplified in some **breast, colon, lung Carcinomas**.
- NMYC and LMYC genes are also amplified in **neuroblastomas** and **small cell cancers of lung**, respectively
- Notch signaling (T cell ALL), Wnt signaling (colon Ca) and Hedgehog signaling (medulloblastoma) pathways transform cells in part through upregulation of MYC gene.

58. Ans. (d) **M** (Ref: Robbins 9th/pg 25; 8th/pg 285)

59. Ans. (c) **S phase** (Ref: Robbins 9th/pg 25; 8th/pg 285)

**S is the synthetic phase** where **DNA replication** takes place

60. Ans. (a, b, c, e); **a. Regulate cell growth and gene expression; b. Found in normal cells; c. Induced by virus; e. May convert to oncogene** (Ref: Robbins 9th/pg)

## Proto-Oncogenes

- **Unmutated counterparts** of genes found **normally in a cell** that promote **autonomous cell growth** in cancer
- **Function:** Regulate **Cell growth<sup>o</sup>, proliferation, inhibition of apoptosis<sup>o</sup> and nuclear transcription<sup>o</sup>**
- **Chromosomal translocation is the most common mechanism for activation of proto-oncogenes<sup>o</sup>**
- **Proto-oncogenes** can also be **activated** by virus to **oncogenes**

61. Ans. (a, e); **a. Cyclin A; e. Cyclin E**

(Ref: Robbins 9th/pg 25; 8th/pg 285; Refer to Ans 29 Above)

**CDK2** forms a complex with **cyclin E** in late G<sub>1</sub>, which is involved in **G<sub>1</sub>/S transition**.

**CDK2** also forms a complex with **cyclin A** at the S phase that facilitates **G<sub>2</sub>/M transition**.

62. Ans. (c) **Kinetochore**

(Ref: Emery's elements of Medical Genetics, 14th ed/pg 17)


- **Kinetochore** is a **protein structure on chromatids** where the spindle fibers attach during cell division to pull sister chromatids apart.
- **"Satellites:" Highly repetitive and abundant DNA sequences with sequence homogeneity**, that are easily separable from the main mass of DNA. **This satellite DNA includes tandem arrays—many copies, one right after another**
- **Centromere** region forms the **"pinched waist"** of metaphase chromosomes, and is the site to which the spindle fibers attach, to separate daughter chromatids in mitosis.

63. Ans. (d) **Neuroblastoma** (Ref: Robbins 9th/pg 288)

**N-MYC** causes, **Neuroblastoma, small-cell Ca lung<sup>o</sup>**

64. Ans. (d) **ABL and C-MYC** (Ref: Robbins 9th/pg 288)





### Discussing the options one by one:

Oncogene	Method of activation
A. SIS and HST-I	Overexpression
B. HGF and L-MYC	Overexpression
C. TGF and CDK4	Overexpression
D. ABL and C-MYC	Translocation
E. RAS and BRAF	Point mutation

65. Ans. (b) **Ras gene**

(Ref: Robbins 9th/pg 288; 8th/pg 284)

66. Ans. (c) **Multiple endocrine neoplasia**

(Ref: R 9th/pg 291)

**Multiple endocrine neoplasia (MEN) is caused by RET which is a Proto-oncogene**

67. Ans. (b) **Methylation of tumor suppressor genes**

(Ref: Robbins 9th/pg 319)

#### Epigenetic alterations in cancers:

- **Abnormal DNA methylation:** hypomethylation or hypermethylation
- **Silencing of tumor suppressor genes by local hypermethylation of DNA is the most common mechanism of cancers**
- Changes in histones near genes that **influence cellular behavior can also predispose to Cancers**

68. Ans. (b) **Medullary carcinoma thyroid**

(Ref: R 9th/pg 284)

#### The RET proto-oncogene

- A **receptor tyrosine kinase** that undergoes oncogenic conversion by **mutation and gene rearrangements**.
- RET protein is a **receptor for glial cell line-derived neurotrophic factor** and structurally related proteins that promote cell survival during neural development.
- Normally **expressed in neuroendocrine cells, such as parafollicular C cells of thyroid, adrenal medulla and parathyroid cell precursors**.
- In MEN-2A, mutations in RET extracellular domain → Medullary thyroid Ca, Adrenal and parathyroid tumors.
- In MEN-2B, mutations in cytoplasmic domain → Thyroid and adrenal tumors without parathyroid involvement
- In **familial cases, mutation is in germline** while in **sporadic cases, somatic rearrangements** are seen

69. Ans. (c) **PTEN** (Ref: Robbins 9th/pg 291; 8th/pg 287)

#### PTEN (Phosphatase and Tensin homologue)

- PTEN is a membrane-associated phosphatase encoded by a gene on **chromosome 10q23**
- Mutated in **Cowden syndrome** (an **autosomal dominant** disorder with **tumors of skin appendages** and an increased incidence of epithelial **cancers of breast, endometrium, and thyroid**).
- It acts as a **tumor suppressor** by serving as a brake on the pro-survival/pro-growth **PI3K/AKT pathway**.

- By phosphorylating a number of substrates, including BAD and MDM2, AKT **enhances cell survival**.
- **PI3K/AKT pathway is the most commonly mutated pathway in human cancers.**

70. Ans. (a) **Gastric Ca** (Ref: Robbins 9th/pg 291; 8th/pg 287)

#### E-cadherin

- Loss-of-contact inhibition, by mutation of the E-cadherin/ $\beta$ -catenin axis is a key characteristic of carcinomas.
- Loss of E-cadherin contributes to the malignant phenotype by allowing easy disaggregation of cells, which can then invade locally or metastasize.
- Reduced cell surface expression of E-cadherin has been seen in Ca **esophagus, colon, breast, ovary, and prostate**.
- Germline loss- of-function mutations of the E-cadherin gene, known as CDH1, cause **familial gastric carcinoma**
- Some of the sporadic gastric carcinomas are also associated with loss of E-cadherin expression.

71. Ans. (b) **p53**

(Ref: Robbins 9th/pg 296; 8th/pg 290)

72. Ans. (c) **Chromosome 11**

(Ref: Robbins 9th/pg 298)

73. Ans. (d) **Wild form is associated with increased risk of child-hood tumors** (Ref: Robbins 9th/pg 294)

74. Ans. (c) **Chromosome 17** (Ref: Robbins 9th/pg 291)

- BRCA1 (chromosome 17q)
- BRCA2 (chromosome 13q)
- Wilms' tumor gene or WT1 gene (chromosome 11)
- NF2 gene (chromosome 22)

75. Ans. (c) **Both** (Ref: Robbins 9th/pg 721)

Genes that promote epithelial-mesenchymal transitions, like *TWIST* and *SNAIL*, may be important metastasis genes in epithelial tumors

76. Ans. (c) **Metalloproteinase** (Ref: Robbins 9th/pg 721)

**Degradation of ECM (basement membrane)** is carried out by **Metalloproteinases** (MMPs type **2 and 9**) also known as **Type IV collagenase<sup>o</sup>**, cathepsin D, and urokinase plasminogen activator

77. Ans. (c) **Desmoplasia**

(Ref: Robbins 9th/pg 266; 8th/pg 260)

#### Basic components of tumors:

- (1) **Neoplastic cells:** constitute the tumor parenchyma
  - (2) **Reactive stroma:** made up of connective tissue, blood vessels, cells of the immune system
- **Desmoplasia**-abundant **collagenous stroma** (fibrosis) in a tumor, stimulated by **parenchymal cells<sup>o</sup>**

78. Ans. (d) **Invasion**

(Ref: Robbins 9th/pg 271; 8th/pg 265)



## Dysplasia

- **Dysplasia** literally means “disordered growth”<sup>Q</sup>
- Epithelial dysplasia is a **premalignant lesion**: increased risk of cancer

## Features of Dysplasia:

- **Pleomorphism**<sup>Q</sup> —variation in size and shape
- Abnormal **nuclear** morphology-**high N:C ratio (nuclear enlargement), hyperchromatic**<sup>Q</sup> nuclei
- Increased but typical **mitotic** figures<sup>Q</sup>
- **Loss of polarity**- disturbed orientation of cells

79. Ans. (a, b, c); a. **Prominent nucleus**; b. **Nuclear enlargement**; c. **Nuclear hyperchromia**

(Ref: R 9th/pg 271)

80. Ans. (c) **IFN a** (Ref: Robbins 9th/pg 306; 8th/pg 298)

## Angiogenesis

- Tumor cannot enlarge beyond **1 to 2 mm**<sup>Q</sup> in diameter unless it has the capacity to induce angiogenesis.
- **Angiogenesis** is an important requirement for tumors to undergo metastasis

81. Ans. (a, b, c) a. **Hepatocellular cancer**; b. **Kaposi sarcoma**; c. **Nasopharyngeal cancer**

82. (b) **Carcinoma nasopharynx**, d. **Carcinoma pancreas**

(Ref: Robbins 9th/ 326-27)]

83. Ans. (a) **Prostate** (Ref: Robbins 9th/pg 276)

- 90% of lung cancers occur in smokers.
- Smoking is also associated with an increased risk of cancers of the oral cavity, larynx, esophagus, stomach, bladder, and kidney, as well as some forms of leukemia.
- Cessation of smoking reduces the risk of lung cancer.

84. Ans. (b) **Liver Angiosarcoma** (Ref: Robbins 9th/pg 428)

85. Ans. (d) **Esophagus** (Ref: Robbins 9th/pg 428; 8th/pg 423)

- Liver (2-fold), cervical (2-fold), and esophageal (2- to 3-fold) cancers are more common in less developed countries.
- Stomach cancer incidence is similar in more and less developed countries but is much more common in Asia than North America or Africa.

86. Ans. (a) **Nicotine** (Ref: Robbins 9th/pg 428; 8th/pg 423)

87. Ans. (c) **Lymphopenia**

(Ref: Robbins 9th/pg 428; 8th/pg 423)

## Radio-sensitive tissues

- Bone marrow cells and all stem cells
- Lymphocytes
- Immune response cells
- Mucosa lining of small intestines
- Breast tissue
- Gonads
- Sebaceous (fat) glands of skin

88. Ans. (b) **Arsenic** (Ref: Robbins 9th/pg 721; 8th/pg 730)

89. Ans. (c) **Cartilage** (Ref: Robbins 9th/pg 428; 8th/pg 423)

The main types of ionizing radiation are:

- X-rays and gamma rays (electromagnetic waves of very high frequencies),
- High-energy neutrons
- Alpha particles (composed of two protons and two neutrons)
- Beta particles (essentially electrons).

**In the human body, the tissue most resistant to the effect of radiation is Cartilage > Bone**

The body tissues can be divided into:

Radio-sensitive	Radio-resistant
<ul style="list-style-type: none"> <li>• Bone marrow cells and all stem cells</li> <li>• Lymphocytes</li> <li>• Immune response cells</li> <li>• Mucosa lining of small intestines</li> <li>• Breast tissue</li> <li>• Gonads</li> <li>• Sebaceous (fat) glands of skin</li> </ul>	<ul style="list-style-type: none"> <li>• Brain and its Neurons</li> <li>• Kidney</li> <li>• Liver</li> <li>• Heart, large arteries and veins</li> <li>• Mature blood cells</li> <li>• Muscle cells</li> <li>• Cartilage</li> </ul>

90. Ans. (d) **HHV-8** (Ref: Robbins 9th/pg 325; Refer to pretexts)

91. Ans. (a) **11** (Ref: Robbins 9th/pg 325; 8th/pg 314)

**HPV 6 and 11** are **low risk** oncogenic viruses, while HPV 16 and 18 are high risk oncogenic viruses.

92. Ans. (e) **Burkitt's lymphoma** (Ref: Robbins 9th/pg 325)

HPV causes Genital warts,<sup>Q</sup> Squamous cell Ca of cervix, anogenital region, head/neck<sup>Q</sup>  
While Burkitt's lymphoma is caused by EBV

93. Ans. (c) **Instability of E6 and E7**

(Ref: Robbins 9th/pg 325)

94. Ans. (b) **Stimulates formation of Pyrimidine dimers**

(Ref: Robbins 9th/pg 428; 8th/pg 423)

Ultraviolet B Rays	Ionizing Radiation
<ul style="list-style-type: none"> <li>• Leads to <b>formation of pyrimidine dimers in DNA</b><sup>Q</sup></li> <li>• Produces skin cancers like Squamous cell Ca, Basal cell Ca, and melanoma of skin</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Particulate</b> radiation (<math>\alpha</math> and <math>\beta</math> particles, protons, neutrons) are all carcinogenic</li> <li>• <b>Electromagnetic</b> (x-rays, <math>\gamma</math> rays)</li> <li>• X-ray causes DNA mutation by <b>Pyrimidine dimer breakdown</b></li> </ul>

95. Ans. (b) **Bone** (Ref: Robbins 9th/pg 428; 8th/pg 423)

96. Ans. (b) **Ewings Sa, e. Ca pancreas**

(Ref: Annexure 5; Harrisons 19th/ pg 595)



**97. Ans. (a) DLBCL; d. Kaposi Sarcoma**

(Ref: Robbins 9th/pg 316-318)

Most common mutations in type I endometrioid carcinomas act to increase signaling through the PI3K/AKT pathway, which is a hallmark of this particular tumor type.

**Other genes involving PI3K/AKT pathway are:**

- PTEN 30% to 80%
- PIK3CA, 40% of endometrioid carcinomas.
- KRAS, 25% of cases.
- ARID1A, 25-33% onethird of tumors.
- DNA mismatch repair genes 20%

**98. Ans. (b) Osteosarcoma** (Ref: Robbins 9th/318)

Chromothripsis (A process in which in which a chromosome is "shattered" and then reassembled in a haphazard way.) has been observed in all 1-2% of cancers and 25% of osteosarcomas and then gliomas.

**99. Ans. (b, c, d); b. IgA is low or absent, c. Increased risk of Leukemia, d. Hypoplasia of thymus**

(Ref: Harrison 19/2630)

**Ataxia telangiectasia:**

Immunodeficiency	Thymic hypoplasiaQ (most consistent defect) with cellular and humoral (IgA and IgG2) <sup>Q</sup> immunodeficiency
Tumors seen	Lymphomas, <sup>Q</sup> Hodgkin's disease, <sup>Q</sup> T cell ALL and Breast cancer
Neuropathologic changes	<ul style="list-style-type: none"> <li>• Loss of Purkinje, granule, and basket cells in the cerebellar cortex<sup>Q</sup> (most striking change) and deep cerebellar nuclei.</li> </ul>

**100. Ans. (a, b, c) a. Ki67, b. Oncoprotein E6, c. p16INK4, cyclin E, and Ki-67**

(Ref: Robbins 9th/1002-04; Harrison 19th/595; Harshmohan 7th/716)

**101. Ans. (b) RET/PTC, d. NTRK1, e. RAS**

(Ref: Robbins 9th/1095; Harrison 19th/2305; L and B 26th/764)

**102. Ans. (a) Include simple deletions, inversions and translocations in chromosomes**

(Ref: Robbins 9th/pg 428)

**103. Ans. (a) Melanoma** (Ref: Robbins 9th/pg 286; 8th/pg 283)

**BRAF is a serine/threonine protein kinase**, and is a member of the RAF family.

**BRAF Mutations** have been detected in: **Hairy cell leukemias, melanomas, benign nevi, colon carcinomas and dendritic cell tumors.**

**104. Ans. (a) t (11;22)** (Ref: Robbins 9th/pg 317; 8th/pg 305)

**105. Ans. (b) Hypercalcemia**

(Ref: Robbins 331)

Hypercalcemia is seen with Adult T-cell leukemia/lymphoma; Acanthosis nigricans is seen with Gastric carcinoma, Lung carcinoma, Uterine carcinoma

**106. Ans. (a, e) a. Tripe palm, e. Oslers node**

**Dermatologic Disorders as paraneoplastic syndromes:**

**Acanthosis nigricans**

- Gastric carcinoma
- Lung carcinoma
- Uterine carcinoma

**Dermatomyositis**

- Bronchogenic carcinoma
- Breast carcinoma

**107. Ans. (d) Nonsmall lung cancer** (Ref: Robbins 9th/719)

Note that tumors that produce ACTH and ADH are predominantly small cell carcinomas, whereas those that produce hypercalcemia are mostly squamous cell carcinomas.

**108. Ans. (b) Prostate Ca**

(Ref: Robbins 9th/pg 332; 8th/pg 321)

**Migratory thrombophlebitis**

- Also called 'Trousseau sign'
- Episodes of thrombophlebitis which are **recurrent** and appear in **different locations** over time
- It is seen in: **Pancreatic Ca, Bronchogenic Ca and Colon cancer**
- Not seen in Prostate Cancer

**109. Ans. (c) Fibrosarcoma**

Acquired hypophosphataemic osteomalacia is a rare tumour-associated disorder, first recognized in 1947. It is characterized by hypophosphataemia, phosphaturia, normocalcemia and osteomalacia in the absence of a nutritional or drug history suggestive of vitamin D deficiency or generalized renal tubular defects. Patients typically present with bone pain and proximal muscle weakness. Tumours associated with this disorder are generally of mesenchymal origin and benign. This condition has occasionally been associated with haemangiopericytoma, fibrous dysplasia, osteosarcoma, chondroblastoma, chondromyxoid fibroma, malignant fibrous histiocytoma, giant cell tumour, haemangioma, paraganglioma, prostate cancer and oat cell carcinoma of the lung

**110. Ans. (a) Ca Vulva**

The initial spread of vulval ca spreads to Inguinal, pelvic, ileal and periaortic lymph node. This allows the surgeon to assess the lymph node status by sentinel lymph node biopsy. Rest of the options mainly shows invasive property where they spread via hemtolympoid pathway.

**111. Ans. (a) Negative immune regulation in treatment of cancer**

The 2018 Nobel Prize in Physiology or Medicine was awarded to James P. Allison and Tasuku Honjo "for their discovery of cancer therapy by inhibition of negative



immune regulation". Their pioneering work on the CTLA4 and PD1 immune checkpoints revealed that these pathways act as so-called 'brakes' on the immune system, and showed that inhibition of these checkpoint pathways allow T cells to more effectively eradicate cancer cells. This research laid the foundation for the clinical development of immune checkpoint inhibitors, which have dramatically improved outcomes for many people with cancer.

112. Ans. (d) **Pancreatic carcinoma/neuroectodermal tumour**

113. Ans. (a, b) **a. Ca19-9; b. Ca 125**

114. Ans. (a) **Desmin**

115. Ans. (b) **Embryonal rhabdomyosarcoma**

116. Ans. (c) **FISH**

117. Ans. (b) **Synaptophysin** (Ref: Robbins 9th ed p 717)

The classic list of "NE markers" includes Synaptophysin (SYN), Chromogranin A (CHR), Neuron Specific Enolase (NSE), and CD56 (NCAM or Neural Cell Adhesion Molecule) and CD57 (Leu-7). We apply these marker by IHC method to identify the differentiation.

118. Ans. (b) **Inflammatory myofibroblastictumor**

(Ref: Robbins Basic Pathology, Chapter Neoplasia)

- Anaplastic lymphoma kinase (ALK) - Location: 2p23. It's also seen in Adenocarcinoma lung, Neuroblastoma, Inflammatory myofibroblastic tumor, Diffuse large B cell lymphoma. It's negative in Fibromatosis, one of the important thing to differentiate from difficult cases of inflammatory myofibroblastic tumor

119. Ans. (c) **95% ethanol**

120. Ans. (c) **Hypercalcemia**

121. Ans. (b) **Her2 neu 2+**

122. Ans. (a) **Type 1**

**ALK (2p23) gene mutation is seen in** (hereditary hemorrhagic telangiectasia-1, Adenocarcinoma lung, Neuroblastoma, anaplastic large B cell lymphoma)

123. Ans. (a) **CA 15.3**

124. Ans. (c) **Toluidine blue**

(Ref: Early Diagnosis and Treatment of Cancer Series: Head and Neck Cancers, Wayne Koch Pg. 54)

- Toluidine blue stain is used as a marker to differentiate lesions at high risk of progression in order to improve early diagnosis of oropharyngeal carcinomas.

- Toluidine blue, an acidophilic metachromatic dye of thiazine group selectively stains acidic tissue components (sulfates, carboxylates and phosphate radicals), thus staining DNA and RNA.
- Toluidine blue has been established as a diagnostic adjunct in detecting oral lesions related to invasive carcinomas, carcinoma in situ or early asymptomatic oral carcinomas.

125. Ans. (b) **Hepatic ca** (Ref: Robbins 9th /338, Henry 1392)

Carcinoembryonic antigen (CEA) is raised in **ca colon, lung, liver, pancreas, stomach, and breast**, and alpha-fetoprotein (AFP), which is produced by **hepatocellular carcinomas, yolk sac remnants in the gonads, and occasionally teratocarcinomas and embryonal cell carcinomas**.

126. Ans. (b) **10% formalin**

(Ref: Annexure 4; Bancrofts Stains)

This is a tricky Question!

- When you first see the term analysis for sperm, the answer is bouin's fluid. But what the examiner wants to know is whether you know that for histological diagnosis, the fixative is 10% formalin.

127. Ans. (b) **Neural tube defect** (Ref: Robbins (SEA) 9th/337)

- Beta hCG increased in: Gastric and pancreatic Ca, hepatoma, Ovarian Ca, germ cell tumor of testis, trophoblastic tumors, choriocarcinoma, and testicular tumors.

128. Ans. (a, b, c, d) **a. Chromogranin A, b. CD56, c. Neuron-specific enolase, d. Synaptophysin**

(Ref: Harrison 19th/558, 2337, 120e-2t)

129. Ans. (c) **No evidence of systemic inflammation**

(Ref: Robbins 9th/pg 331; 8th/pg 321)

### Cancer Cachexia:

**Progressive loss of body fat and lean body mass<sup>o</sup>** with profound weakness, anorexia and anemia, seen in cancer. **TNF<sup>o</sup>** is the **major contributor** to cachexia with advanced cancer.

Equal loss of both **fat and lean muscle<sup>o</sup>**

**Elevated basal metabolic rate**

Evidence of **systemic inflammation** (e.g., an increase in acute phase reactants)

130. Ans. (d) **Size of primary lesion** (Ref: Robbins 9th/pg 11)

**Grading of a tumor is based on its histopathological appearance, that includes** Degree of differentiation, Number of mitoses, architectural features **and not on the size of primary lesion**

131. Ans. (a) **Desmin-Carcinoma** (Ref: Robbins 9th/pg 11)

**Desmin** is an IHC marker for **muscle tumors** and not Carcinoma





132. Ans. (b) **Prostatic carcinoma** (Ref: Robbins 9th/pg 337)

133. Ans. (d) **Alpha-fetoprotein** (Ref: Robbins 9th/pg 337)

134. Ans. (d) **Angiomyolipoma of lung** (Ref: R 9th/pg 337)

Lymphangiomyomatosis and **angiomyolipoma**: closely related entities characterized by hamartomatous proliferation of **HMB-45-positive** smooth muscle

135. Ans. (a) **Ovarian cancer** (Ref: Robbins 9th/pg 337)

136. Ans. (b) **Carcinoma** (Ref: Robbins 9th/pg 11; 8th/pg 335)

137. Ans. (c) **1,25dihydroxy vitamin D** (Ref: R 9th/pg 337)

138. Ans. (c) **DNA microarray analysis** (Ref: R 9th/pg 337)

Microarray can study several genes at a time. Hence it can be used in molecular profiling of cancer cells.

139. Ans. (b) **Anti-Hu** (Ref: Robbins 9th/pg 337; 8th/pg 327)

**Antibodies to intracellular antigens, syndromes, and associated cancers**

Antibody	Associated Neurologic Syndrome(s)	Tumors
Anti-Hu (ANNA1)	encephalomyelitis, subacute sensory neuronopathy	SCLC
Anti-Yo (PCA1)	Cerebellar degeneration	Ovary, breast
Anti-Ri (ANNA2)	Cerebellar degeneration, opsoclonus, brainstem encephalitis	Breast, gynecologic, SCLC
Anti-Tr	Cerebellar degeneration	Hodgkin's lymphoma

140. Ans. (a, d, e); a. **Yolk sac tumor**; d. **Non-seminoma**; e. **Hepatocellular carcinoma** (Ref: Robbins 9th/pg 337)

141. Ans. (c) **Fibrosarcoma** (Ref: Robbins 9th/pg 337)

Hypoglycemia is seen as a paraneoplastic syndrome in fibrosarcoma and osteosarcoma

142. Ans. (b) **Ames test** (Ref: Robbins 9th/pg 337; 8th/pg 327)

- The **Ames test** is a **biological assay** to **assess the mutagenic potential of chemical** compounds.
- Ames test **uses bacteria** (*Salmonella typhimurium* that carry mutations in genes involved in histidine synthesis) to test whether a given chemical can cause mutations in the DNA of test organism.
- A **positive test indicates** that the **chemical might act as a carcinogen**

It is a **quick and convenient** assay to estimate the carcinogenic potential of a compound because standard carcinogen assays on mice are time-consuming and expensive

143. Ans. (c) **Multiple myeloma** (Ref: Robbins 9th/pg 337)

Bortezomib, a proteasome inhibitor is used in the treatment of Multiple myeloma

144. Ans. (a) **Polycyclic hydrocarbons** (Ref: R 9th/pg 337)

145. Ans. (d) **Necrotizing myelopathy** (Ref: R 9th/pg 337)

146. Ans. (d) **Aerobic glycolysis** (Ref: Robbins 9th/pg 337)

- **Cancer cells** tend to **convert most glucose to lactate**<sup>o</sup> even in presence of **ample oxygen** (**aerobic glycolysis**)<sup>o</sup>

147. Ans. (a) **Squamous cell Ca** (Ref: Robbins 9th/pg 1155)

**Malignant squamous epithelium with prominent central keratin pearls** are seen in **Squamous cell Ca**

148. Ans. (b) **Carcinoma - desmin**

(Ref: Robbins 9th/pg 11; 8th/pg 35; Refer table in Ans 158)  
*Desmin is an IHC marker for muscle tumors and not Carcinoma*

149. Ans. (b) **Medullary carcinoma of thyroid**

(Ref: Robbins 9th/pg 337; 8th/pg 327)

**Calcitonin** is a marker of **Medullary carcinoma of thyroid**; Hormones acting as tumor markers are-

150. Ans. (a) **Ewing sarcoma**

(Ref: Robbins 9th/pg 1203)

**Mic2** is a **tumor marker** for **Ewing's sarcoma**

**CD99** is a **product of the MIC2 gene** located on X and Y chromosomes.

It is seen in **Ewing's sarcoma/primitive neuroectodermal tumor (PNET)** and **lymphoblastic tumors**.

151. Ans. (c) **Neurofilament**

(Ref: Robbins 9th/pg 11; 8th/pg 35)

Markers for **muscle tumor** are: **Desmin, Actin, Intermediate filament, MYOD, myogenin**

152. Ans. (c) **CK-20**

(Ref: Robbins 9th/pg 11; 8th/pg 35)

The markers of **Melanoma** are **S-100, MITF (microphthalmia-associated transcription factor)** and **Vimentin**

153. Ans. (a) **Desmin**

(Ref: Robbins 9th/pg 11; 8th/pg 35)

154. Ans. (c) **Synovial sarcoma**

(Ref: Robbins 9th/pg 1223)

### Synovial Sarcoma

- Immunohistochemistry yields positive reactions for **keratin** and **epithelial membrane antigen**;



- Most synovial sarcomas show a characteristic chromosomal translocation **t(x;18)(p11;q11)** producing **SS18-SSX1, SSX2, or SSX4** fusion genes that encode chimeric transcription factors

**155. Ans. (c, d, e); c. Omphalocele; d. Sacrococcygeal teratoma; e. Neural tube defect**

(Ref: Henry's 22nd/ pg 1392; Robbins 9th/pg 337; 8th/pg 327)

#### Alpha-fetoprotein

Structurally **related to Albumin**<sup>Q</sup>

In the fetus, AFP is **synthesized by yolk sac<sup>Q</sup> and fetal Hepatocytes<sup>Q</sup>**

**Maternal serum AFP** raised in **Omphalocele<sup>Q</sup>, Sacrococcygeal teratoma<sup>Q</sup>, Neural tube defect<sup>Q</sup>**

AFP can be used for **prenatal screening** for:

- **Neural tube defects<sup>Q</sup>** (anencephaly and spina bifida)
- **Down syndrome<sup>Q</sup>** (as part of triple test)
- **Omphalocele** and **gastroschisis** (Acetylcholinesterase levels may also be increased)
- **Sacroccoccygeal teratoma<sup>Q</sup>**

**Increased levels in Liver cancers<sup>Q</sup>, non-seminomatous<sup>Q</sup> germ cell tumors of testis and non-neoplastic lesions like Amebic liver abscess<sup>Q</sup> and Hepatitis<sup>Q</sup> Sacrococcygeal teratoma (SCT) is the most frequent tumor in the neonatal period.**

**156. Ans. (a, c, d); a. Used for monitoring of recurrence of colon cancer; c. Increased in smokers; d. Increased in colon cancer**

(Ref: Henry's 22nd/ pg 1392; Robbins 9th/pg 337; 8th/pg 327)

#### Carcino embryonic antigen (CEA)

- **Tumor marker for Carcinomas of colon, pancreas, lung, stomach and breast.**
- Can be used for **monitoring of recurrence of colon cancer**, but it is **not specific for colon cancer**.
- As **liver metabolizes CEA**, **liver damage** impairs CEA clearance and leads to **increased levels** in blood
- Also increased in **smokers, Alcoholic Cirrhosis, Ulcerative Colitis, Pancreatitis, hepatitis**

**157. Ans. (a, c, e); a. It is a Glycoprotein; c. It is Increased in colon carcinoma; e. May be elevated in Pelvic inflammatory disease**

(Ref: Robbins 9th/pg 337; 8th/pg 327; Ref: Henry's 22nd/ pg 1393)

#### CA-125:

- **Normal range** in pre-menopausal females is **0-35 U/mL**
- **Increased in Ovarian Ca:** non-mucinous epithelial (> 80%), serous, endometrioid and clear cell types
- Other Cancers with increased CA-125 levels are **Non-Hodgkin Lymphoma, Lung cancer, Endometrial Ca**
- Non-malignant diseases with increased CA-125: **Liver cirrhosis, Pelvic inflammatory disease including endometriosis and advanced abdominal or pelvic tuberculosis**

**158. Ans. (a) Cytokeratin** (Ref: Robbins 9th/pg 11)

**159. Ans. (b) Bacterial contamination of the specimen**

(Ref: Halder A, Halder S and Fauzdar A. A preliminary investigation of genomic screening in cervical carcinoma by comparative genomic hybridization. Indian J Med Res 122, November 2005, pp 434-446)

This question is a direct pick up from the above publication by our AIIMS faculty!

**"Conventional cytogenetics are difficult in solid tumors especially in case of carcinoma cervix as most biopsy sample is contaminated/infected with microorganisms and quality of metaphase preparation is poor."**

**160. Ans. (a) Prostate carcinoma**

(Ref: Robbins 9th/pg 337)

**161. Ans. (a, b, c, d); a. HCC; b. Hepatoblastoma; c. Infant hemangioendothelioma; d. Amebic liver abscess**

(Ref: Robbins 9th/pg 337; 8th/pg 327)

**162. Ans. (b) Mutagenicity**

(Ref: Robbins 9th/pg 337; 8th/pg 327)

**163. Ans. (a, b, c, d, e); a. Abdominal TB; b. Ca cervix; c. Endometriosis; d. Ovarian ca; e. Endometrial ca**

(Ref: Robbins 9th/pg 337; 8th/pg 327)

**164. Ans. (c) Cytokeratin**

(Ref: Robbins 9th/pg 337; 8th/pg 327)

An elderly smoker has a high risk of Small cell Ca and Squamous cell Ca lung; Of the given options, cytokeratin is a marker of squamous cell Carcinoma; Discussing the options one by one,

Marker	Tumor
A. Calretinin	Malignant Mesothelioma
B. Vimentin	Mesenchymal tumors
C. Cytokeratin	Epithelial carcinomas (Squamous cell Ca and Adenocarcinoma)
D. TTF1	Adenocarcinoma

**165. Ans. (c) Human leucocyte antigen** (Ref: R 9th/pg 337)

CEA, Tyrosinase and AFP are tumor markers

- **Tyrosinase** has been demonstrated to be a **sensitive marker for melanoma**
- Human leucocyte antigen is not a tumor marker.

[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines, typical of notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Diseases of Infancy and Childhood

## Key Points

- » **Agenesis is complete absence of an organ and its primodium**
- » **Aplasia is absence of an organ due to failure of growth of existing primodium**
- » Cystic fibrosis results from abnormal function of an **epithelial chloride channel protein (CFTR) gene on chr7q31.2**
- » **Increased sweat chloride (>70 mEq/L) concentration on two or more occasions suggests Cystic fibrosis.**
- » **Hemangiomas are the most common tumors of infancy**
- » **The most frequent childhood cancers is Leukemia (B-ALL)**
- » **Most common site of neuroblastoma is adrenal medulla**
- » **Wilms' tumor is most common primary renal tumor of childhood**
- » **WAGR syndrome, Denys-Drash and Beckwith-Wiedemann syndrome (BWS) are associated with Wilms' tumor**
- » **Anaplasia in Wilms' tumor signifies presence of TP53 mutations, emergence of resistance to chemotherapy with increased risk of recurrence and death-adverse prognosis**

## Key Recent Updates

- » Wilms' tumor being mixed tumors has Desmin, Vimentin and Cytokeratin positive
- » Thyroid transcription factor-1 (TTF-1) is a sensitive marker for pulmonary and thyroid adenocarcinomas
- » Most sensitive and specific IHC marker for Neuroblastoma is PHOXB2.





## CONGENITAL ANOMALIES

### Organ-Specific Disorders of Development

- **Agenesis:**
  - Complete absence of an organ<sup>Q</sup> and its associated primodium. E.g., renal agenesis<sup>Q</sup>
- **Aplasia:**
  - Absence of organ<sup>Q</sup> due to failure of growth of existing primodium. E.g., thymic aplasia<sup>Q</sup>, found in DiGeorge syndrome
- **Atresia:**
  - Absence of an opening<sup>Q</sup> usually of a hollow viscera such as trachea, intestine. E.g., esophageal atresia
- **Hypoplasia:**
  - Incomplete development or decreased size<sup>Q</sup> of an organ with decreased number of cells



#### High Yield Facts

- **Hereditary disorders:** Derived from parents and are transmitted in the germ line through the generations and therefore, are familial.<sup>Q</sup>
- **Congenital** means “born with”<sup>Q</sup>
- **Some congenital diseases are not genetic;** e.g. Congenital syphilis<sup>Q</sup>
- **Malformations:** Due to intrinsically abnormal developmental process (multifactorial); e.g. Anencephaly, congenital heart defects<sup>Q</sup>

### Fetal Alcohol Syndrome

Due to teratogenic effect of alcohol

#### Prenatal and postnatal growth retardation, psychomotor disturbances

Facial anomalies	Microcephaly, short palpebral fissures, and maxillary hypoplasia
Heart defects	Atrial septal defect



#### High Yield Facts

- **Disruptions:** Secondary destruction (extrinsic disturbance in morphogenesis)<sup>Q</sup> of an organ that was previously normal in development. Not heritable and no risk of recurrence. E.g. Amniotic bands<sup>Q</sup>
- **Deformations:** Structural abnormalities (extrinsic disturbance in morphogenesis) due to compression of growing fetus by abnormal biomechanical forces like bicornuate uterus. E.g., Clubfeet<sup>Q</sup>. Most common underlying factor responsible for deformation is uterine constraint<sup>Q</sup>
- **Sequence:** Cascade of anomalies triggered by one initiating aberration. E.g. Potter sequence (oligohydramnios sequence): Renal agenesis → Oligohydramnios → flattened facies, abnormalities of the hands and feet and pulmonary hypoplasia.
- **Malformation syndrome:** Constellation of congenital anomalies that *cannot* be explained on the basis of a single, localized, initiating defect. Usually caused by a specific chromosomal abnormality. E.g. Down's syndrome<sup>Q</sup>

## CYSTIC FIBROSIS (MUCOVISCIDOSIS)

- Most common lethal genetic disease<sup>Q</sup> that affects Caucasian populations
- Autosomal recessive, inherited disorder of ion transport channel.
- **Reproductive tracts:** Male infertility<sup>Q</sup>

#### Chiefly Affects

- Fluid secretion from exocrine glands<sup>Q</sup>
- Epithelial lining of the respiratory tract<sup>Q</sup>: Chr lung disease due to recurrent infections,
- Gastrointestinal: Pancreatic insufficiency, steatorrhea, malnutrition, hepatic cirrhosis, intestinal obstruction



#### High Yield Facts

##### Cystic fibrosis

- *Haemophilus influenzae*<sup>Q</sup> and *S. aureus*<sup>Q</sup> are the first organisms recovered from lung secretions
- Antibiotic-resistant and mucoid *P. aeruginosa*<sup>Q</sup> colonizes next.
- *Burkholderia cepacia* is pathognomonic organism



## Primary Defect


- Abnormal function of an **epithelial chloride channel protein**<sup>o</sup> encoded by the **Cystic Fibrosis Transmembrane conductance Regulator (CFTR)** gene on chr7q31.2<sup>o</sup>

## Other Defects Lie in

- Outwardly rectified **chloride channels**<sup>o</sup>

- Inwardly rectified **potassium channels (Kir 6.1)**<sup>o</sup>
- Epithelial sodium channel (ENaC)**<sup>o</sup>

Of these, the **interaction of CFTR with ENaC**<sup>o</sup> is the most important step



## Mechanism: Effect of CFTR Gene Mutation

### ENaC (Epithelial sodium channel)

Site	Normal Function	In cystic fibrosis	Effect seen in CF
Apical exocrine glands of airway and GIT	Na uptake from luminal fluid	Increased Na and H <sub>2</sub> O uptake from lumen	Thick, viscid secretions <sup>o</sup>
Sweat ducts	Absorb luminal Na <sup>+</sup>	Decreased reabsorption of Na and Cl	Salty sweat (high Na and Cl) <sup>o</sup> (Hallmark of CF)

- Most common mutation: **class II mutation (ΔF508)**<sup>o</sup>
- Leads to **defective processing**<sup>o</sup> of the protein from the E.R to the Golgi apparatus

## Systemic Involvement

### Respiratory Tract

- Upper respiratory tract disease** is almost **universal** in patients with CF<sup>o</sup>
- Chronic cough**<sup>o</sup> and **sputum**<sup>o</sup> production
- Bronchiectasis**<sup>o</sup>, **atelectasis**<sup>o</sup>, infiltrates, hyperinflation, **Nasal polyps**<sup>o</sup>

### Genitourinary System

- Azoospermia and infertility**<sup>o</sup>
- Congenital **bilateral absence or obliteration**<sup>o</sup> of the **vas deferens**<sup>o</sup>, due to defective liquid secretion may be seen

### Gastrointestinal and Nutritional Abnormalities

- Intestinal: Meconium ileus**<sup>o</sup>, **distal intestinal obstruction syndrome (DIOS)**<sup>o</sup>, rectal prolapse.
- Pancreatic: Pancreatic exocrine insufficiency**<sup>o</sup>, recurrent acute/chronic **pancreatitis**<sup>o</sup>

- Hepatic: Focal biliary cirrhosis**, or multilobular cirrhosis, prolonged **neonatal jaundice**.
- Nutritional: Failure to thrive (protein-calorie malnutrition)**, hypoproteinemia, edema, fat-soluble vitamin deficiency
- Salt-loss syndromes**: acute salt depletion, chronic **metabolic alkalosis**<sup>o</sup>

### Diagnosis

Criteria for Diagnosis of Cystic Fibrosis

- One or more characteristic **phenotypic features**<sup>o</sup>
- OR** a **history**<sup>o</sup> of cystic fibrosis in a sibling
- OR** a positive **newborn screening test** result (**Serum Immuno-reactive trypsinogen**)

### AND

- Increased sweat chloride (>70 meq/L)**<sup>o</sup> concentration on **two or more occasions**
- OR** identification of **two cystic fibrosis mutations**<sup>o</sup>
- OR** demonstration of **abnormal epithelial an ion transport (transepithelial potential difference is raised)**<sup>o</sup>

Sequencing the CFTR gene is the **gold standard** for diagnosis of cystic fibrosis<sup>o</sup>




## MALIGNANT TUMORS

### Mnemonic

Small Round Blue Cell Tumors<sup>Q</sup>  
Tumors with similar histology of small round cell include:

#### "Low NEW MRP"

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li>• Lymphoma</li> <li>• Ewing's Sarcoma</li> <li>• Medulloblastoma</li> <li>• Primitive Neuroectodermal tumor</li> </ul> | <ul style="list-style-type: none"> <li>• Neuroblastoma</li> <li>• Wilms' Tumor</li> <li>• Retinoblastoma</li> </ul> |
|---|---|



Small round blue cell tumor



### High Yield Facts

The most frequent childhood cancers are:  
(in order of decreasing frequency)<sup>Q</sup>

Leukemia (Most common is ALL)<sup>Q</sup> > Neuroblastoma > Wilms' tumor > Hepatoblastoma > Retinoblastoma

- Hemangiomas<sup>Q</sup> are the most common tumors of infancy
- Characteristic chromosomal translocation,  $t(12;15)^Q$ , has been described in congenital-infantile fibrosarcomas, (ETV6-NTRK3 fusion transcript)
- Sacrococcygeal teratomas are the most common teratomas of childhood<sup>Q</sup>

## NEUROBLASTIC TUMORS

Tumors of the sympathetic ganglia and adrenal medulla that are derived from primordial neural crest cells.


### Neuroblastoma

#### Sites of Tumor

- Adrenal medulla<sup>Q</sup> (MC site), along the sympathetic chain in the paravertebral region<sup>Q</sup> of the abdomen (25%) and posterior mediastinum<sup>Q</sup> (15%) > pelvis<sup>Q</sup>, neck, and brain (cerebral neuroblastomas)



Neuroblastoma (Adrenal)



Homer Wright rosettes in neuroblastoma

### Morphology

- **Gross:** Often sharply demarcated by a fibrous pseudo-capsule
- **Microscopic:** Foci of punctate intratumoral calcification
- **Homer-Wright pseudorosettes<sup>Q</sup>** with central space filled with neuropil (eosinophilic fibrillar material)<sup>Q</sup>
- **Accompanied with:** Primitive neuroblasts (ganglioneuroblastoma)<sup>Q</sup>
- Stains +ve with Neuron-specific enolase (NSE)
- **E.M:** central dense cores (containing catecholamines) surrounded by a peripheral halo.


### Metastasis

- Hematogenous<sup>Q</sup> and lymphatic<sup>Q</sup> route to liver, lungs, bones, and bone marrow. Proptosis and ecchymosis (spread to the periorbital region)<sup>Q</sup> is common.
- Disseminated neuroblastomas may present with multiple cutaneous metastases ("blueberry muffin baby")<sup>Q</sup>





- **Stage 4S("S"=special): Localized primary tumor** (as defined for stages 1,2A, or 2B) with dissemination limited to **skin, liver, and/or bone marrow**; **stage 4S is limited to infants younger than 1 year**



## Prognostic Factors in Neuroblastoma

Variable	Favorable	Unfavorable
Stage	Stage 1, 2A, 2B, 4S <sup>Q</sup>	Stage 3, 4
Age	<18 months <sup>Q</sup>	>18 months
Schwannianstroma	Present	Absent
Mitosis-karyorrhexis index	< 200/5000 cells	> 200/5000 cells
DNA ploidy	Hyperdiploid <sup>Q</sup>	Near-diploid
N/myc	Not amplified	Amplified
Chromosome 11q loss	Absent	Present
TRKA expression	Present <sup>Q</sup>	Absent
TRKB expression	Absent	Present



## High Yield Facts

### Neuroblastoma

- Most common **extracranial solid tumor** of **childhood**<sup>Q</sup>
- Most common **abdominal tumor** of **childhood**<sup>Q</sup>
- Most **frequently diagnosed** tumor of **infancy (<1 year of age)**<sup>Q</sup>
- Median age at diagnosis is **18 months**<sup>Q</sup>
- **Mature ganglion cells (ganglioneuroma;** accompanied by the appearance of Schwann cells) on histology → signifies favorable outcome<sup>Q</sup>
- 98% cases are sporadic<sup>Q</sup>
- 2% cases are **familial**<sup>Q</sup> (Germline mutations in the **anaplastic lymphoma kinase (ALK gene)**<sup>Q</sup>)
- 90% of neuroblastomas, regardless of location, produce catecholamines: **vanillylmandelic acid (VMA)** and **homovanillic acid**

## WILMS' TUMOR

- Most common **primary renal tumor of childhood**<sup>Q</sup>
- Peak incidence for Wilms' tumor is between **2 and 5 years**<sup>Q</sup>

### Can be present as:

- **Synchronous:** Both kidneys involved **simultaneously**<sup>Q</sup>
- **Metachronous:** Kidney affected **one after the other**<sup>Q</sup>

### Groups of congenital malformations with increased risk of Wilm's tumor:

WAGR syndrome (33% risk) <sup>Q</sup>	Denys-Drash syndrome (90% risk, Max) <sup>Q</sup>	Beckwith-Wiedemann syndrome (BWS)
<ul style="list-style-type: none"> <li>• <b>WT1 gene: Chr 11p13</b><sup>Q</sup></li> <li>• Wilms' tumor, Aniridia, Genital anomalies, and mental Retardation</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Gonadal dysgenesis</b><sup>Q</sup> (male pseudohermaphroditism)</li> <li>• Early-onset nephropathy (<b>diffuse mesangial sclerosis</b>)<sup>Q</sup></li> <li>• Increased risk of <b>gonadoblastomas</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• <b>WT2 gene: 11p15.5</b></li> <li>• Organomegaly: <b>Macroglossia</b><sup>Q</sup>, <b>hemihypertrophy</b><sup>Q</sup>, <b>omphalocele</b>, (adrenal <b>cytomegaly</b>)<sup>Q</sup></li> <li>• <b>Genomic imprinting</b><sup>Q</sup></li> </ul>

*β-catenin mutations is seen in 10% of sporadic Wilms' tumor.*

**Nephrogenic rests**<sup>Q</sup>: Putative **precursor lesions** of Wilms' tumors.

- **100%**<sup>Q</sup> in cases of **bilateral Wilms' tumor**
- **Increased risk** of developing Wilms' tumors in the **contralateral kidney**<sup>Q</sup>

## Morphology




### Triphasic combination:

- **Blastemal**-small blue tumor cells
- **Stromal**-fibrocytic or myxoid
- **Epithelial cell** -tubules or glomeruli

## Anaplasia

- **Characteristic: Large, hyperchromatic, pleomorphic nuclei and abnormal mitoses.**
  - Loss of 11q and 16q, and gain in 1q
- **Signifies:** Presence of **TP53 mutations**<sup>Q</sup> and the **emergence of resistance**<sup>Q</sup> to chemotherapy
- **Increased risk** of recurrence and death-**adverse prognosis**<sup>Q</sup>






Wilms' Tumor




## Image-Based Questions

1. 5 m Infant present with Abdominal mass in AIIMS Peds OPD along with dark pigmentation around eyes. CT scan suggested abdominal mass with punctate calcification. Biopsy of the same was done which has been shown below. What is your diagnosis?



- Wilms' tumor with nephrogenic nests
- Neuroblastoma with rosettes
- Renal cell Ca with Papillary pattern
- Hepatoblastoma with rosettes

2. 2/male presented with abdominal mass. On CT scan intra-renal mass was seen, the biopsy of which has been shown below. What is your diagnosis?



- Wilms' tumor with triphasic pattern
- Neuroblastoma with rosettes
- Renal cell Ca with Papillary pattern
- Hepatoblastoma with rosettes



## Answers of Image-Based Questions

- Ans. (b) **Neuroblastoma with rosettes**
  - Abdominal mass along with dark pigmentation around eyes (Raccoon eyes) with CT scan showing punctate calcification. Biopsy from the same shows Homer-Wright rosette.
- Ans. (a) **Wilms' tumor showing intrarenal mass with triphasic pattern in biopsy.**




## Multiple Choice Questions

1. **Feature(s) of Familial Mediterranean fever:**  
(PGI May 2019)
  - a. Caused by mutations of the MEFV gene
  - b. First fever attack occur only after 20 years of age
  - c. Presentation include episodic bouts of acute peritonitis
  - d. Amyloidosis occur as a complication
  - e. Colchicine is used for treatment
2. **Which of the following is true about Wilms' tumor?**  
(PGI Nov 2018)
  - a. Associated with aniridia
  - b. Small unipolar cysts
  - c. Presents in neonates
  - d. Has pseudocapsule
3. **Which are primitive neuroectodermal tumors?**  
(PGI Nov 2018)
  - a. Medulloblastoma
  - b. Craniopharyngioma
  - c. Meningioma
  - d. Rhabdomyosarcoma
4. **Wilms' tumor has the following markers positive except?**  
(JIPMER 2017)
  - a. Desmin
  - b. Vimentin
  - c. TTF-1
  - d. Cytokeratin
5. **All are good prognostic factors in neuroblastoma except:**  
(Recent Question 2015)
  - a. Stage 4S
  - b. Trk A expression
  - c. Trk B expression
  - d. Age <18 months
6. **For diagnosing cystic fibrosis sweat chloride level should be more than:**  
(Recent Question 2015)
  - a. 20 mEq/L
  - b. 40 mEq/L
  - c. 60 mEq/L
  - d. 70 mEq/L
7. **Most common cancer in children less than 10 years**  
(Recent Question 2015)
  - a. Leukemia
  - b. Neuroblastoma
  - c. Brain tumor
  - d. Wilms' tumor
8. **False statement about cystic fibrosis**  
(Recent Question 2015)
  - a. Sweat chloride level >70 mEq/L is diagnostic
  - b. ENAC activity of the sweat ducts increases
  - c. Mycobacterium tuberculosis infection is rare
  - d. Distal intestinal obstruction syndrome occurs in children
9. **The following is not a category A agent of bioterrorism**  
(Recent Question 2015)
  - a. Anthrax
  - b. Botulism
  - c. Plague
  - d. Brucellosis
10. **Most common site of metastases in neuroblastoma**  
(Recent Question 2015)
  - a. Lung
  - b. Skull
  - c. Liver
  - d. Vertebrae
11. **All are true about Wilms' tumor except**  
(Recent Question 2015)
  - a. Triphasic morphology
  - b. MC renal malignancy in children
  - c. Associated with cysts in liver
  - d. Does not respond to chemotherapy and radiotherapy
12. **Find the false statement about cystic fibrosis**  
(Recent Question 2015)
  - a. A normal sweat chloride test does not exclude the diagnosis
  - b. Inhaled recombinant human deoxyribonuclease reduces the risk of acute exacerbations
  - c. Ivacaftor, a potentiator of the CFTR channel is used to for patients with ( $\Delta F508$ ) mutation
  - d. All the above
13. **The following is not one of the core prognostic factor of neuroblastoma:**  
(Recent Question 2015)
  - a. Age at diagnosis
  - b. Morphology
  - c. Amplification of MYCN gene
  - d. TRKA expression
14. **The genetic-abnormality associated with Beckwith-wiedeman syndrome**  
(Recent Question 2015)
  - a. Negative missense mutation
  - b. Genomic imprinting
  - c. Deletion
  - d. Balanced translocation
15. **Most common tumor of infancy:**  
(Recent Question 2015)
  - a. Hemangioma
  - b. Brain tumor
  - c. Leukemia
  - d. Neuroblastoma
16. **Passive smoke inhalation in non-smokers can be estimated by measuring the blood levels of**  
(Recent Question 2015)
  - a. Cotinine
  - b. Nicotine
  - c. Hydrocarbons
  - d. Carbon monoxide
17. **The following tumor is not common in the first decade**  
(Recent Question 2015)
  - a. Ameloblastoma
  - b. Retinoblastoma
  - c. Neuroblastoma
  - d. Rhabdomyosarcoma
18. **The following are small round blue cell tumors A/E**  
(Recent Question 2015)
  - a. Lymphomas
  - b. Osteosarcoma
  - c. Neuroblastoma
  - d. Rhabdomyosarcoma
19. **True regarding stge IV-S of neuroblastoma A/E**  
(Recent Question 2015)
  - a. Limited to infants < 1 year
  - b. Primary localize tumor
  - c. Dissemination to bone
  - d. Good prognosis
20. **A mother brings her 10 months old baby with the complaint that the swat is very salty. Past history revealed revealed meconium ileus in the new born period. Diagnosis**  
(Recent Question 2015)
  - a. Hirschprung's disease
  - b. Hyaline membrane disease
  - c. Necrotizing enterocolitis
  - d. Cystic fibrosis
21. **Most common cause of non-immune hydrops**  
(Recent Question 2015)
  - a. Chromosomal abnormalities
  - b. Fetal anemia
  - c. Intrauterine infections
  - d. Cardiovascular causes
22. **Cystic fibrosis gene is located in chromosome**  
(PGI Nov 2015/Recent Question 2015)
  - a. 7p
  - b. 7q
  - c. 13p
  - d. 22q



23. **Most common cause of death in Cystic fibrosis**  
(Recent Question 2015)
  - a. Lower respiratory tract infections
  - b. Cardiovascular defects
  - c. Pancreatitis
  - d. Malnutrition and malabsorption
24. **The following tumor is not common in the first decade**  
(Recent Question 2015)
  - a. Ameloblastoma
  - b. Retinoblastoma
  - c. Neuroblastoma
  - d. Rhabdomyosarcoma
25. **The following are small round blue cell tumors A/E**  
(Recent Question 2015)
  - a. Lymphoma
  - b. Osteosarcoma
  - c. Neuroblastoma
  - d. Rhabdomyosarcoma
26. **Stippled calcification is seen in** (Recent Question 2015)
  - a. Wilms' tumor
  - b. Pheochromocytoma
  - c. Teratoma
  - d. Neuroblastoma
27. **Most common abdominal mass in children**  
(Recent Question 2015)
  - a. Hydronephrosis
  - b. Wilms' tumor
  - c. Neuroblastoma
  - d. Rhabdomyosarcoma
28. **Tumor with triphasic combination of blastmal, stromal and epithelial cell types**  
(Recent Question 2015)
  - a. Wilms' tumor
  - b. Teratoma
  - c. Melanoma
  - d. Neuroblastoma
29. **The following factor is associated with good prognosis in neuroblastoma**  
(Recent Question 2015)
  - a. TRLB expression
  - b. TRKA expression
  - c. MRP expression
  - d. Telomerase expression
30. **A mother brings her 10-month-old baby with the complaint that the sweat is very salty-past history revealed meconium ileus in the new born period. What is your diagnosis ?**  
(Recent Question 2015)
  - a. Hirschsprung's disease
  - b. Hyaline membrane disease
  - c. Necrotizing enterocolitis
  - d. Cystic fibrosis
31. **Which is not associated with bilateral renal agenesis?**  
(Recent Question 2015)
  - a. Potter facies
  - b. Renal agenesis
  - c. Polyhydramnios
  - d. Oligohydramnios
32. **Most common malignancy in children?**  
(Recent Question 2015)
  - a. Leukemia
  - b. Brain tumors
  - c. Neuroblastoma
  - d. Retinoblastoma
33. **WAGR syndrome includes all except?**  
(Recent Question 2015)
  - a. Wilms' tumor
  - b. Aniridia
  - c. Growth retardation
  - d. Mental retardation
34. **Most common cause of extracranial solid tumor in children:**  
(Recent Question 2015)
  - a. Neuroblastoma
  - b. Wilms' tumor
  - c. Thymoma
  - d. Osteosarcoma
35. **Most important prognostic factor of Wilms' tumor:**  
(Recent Question 2015)
  - a. Histopathology
  - b. Ploidy of cells
  - c. Age < 1 yr
  - d. Mutation of 12p gene
36. **Small round cell tumors include:**(Recent Question 2014)
  - a. Wilms' tumor
  - b. Retinoblastoma
  - c. Rhabdomyosarcoma
  - d. All
37. **Sweat chloride in cystic fibrosis:**(Recent Question 2014)
  - a. Decreased
  - b. Increased
  - c. No change
  - d. May increase or decrease
38. **All are good prognostic factors for neuroblastoma except -**  
(Recent Question 2014)
  - a. Trk-A expression absent
  - b. Absence of 1 p loss
  - c. Absence of 17 p gain
  - d. Absence of 11q loss
39. **Not a childhood tumor is?** (Recent Question 2014)
  - a. Neuroblastoma
  - b. Wilms' tumor
  - c. Small cell carcinoma
  - d. Retinoblastoma
40. **Cystic fibrosis causes all except?**(Recent Question 2014)
  - a. Decreased chloride in sweat
  - b. Infertility
  - c. Increased infection
  - d. Pancreas involvement
41. **Children with germline retinoblastoma are more likely to develop other primary malignancies in their later lifetime course. Which of the following can occur in such patients?**  
(AIIMS Nov 13)
  - a. Osteosarcoma of lower limbs and soft tissue sarcoma
  - b. Thyroid carcinoma
  - c. Seminoma
  - d. Squamous cell carcinoma
42. **Bilateral proptosis is characteristically present in**  
(AIIMS Nov 13)
  - a. Retinoblastoma
  - b. Rhabdomyosarcoma
  - c. Neuroblastoma
  - d. PNET
43. **Which histological finding in resected kidney indicated Bilateral Wilms' tumor?**  
(JIPMER 2013)
  - a. Blastemal component
  - b. Nephrogenic rests
  - c. Skeletal muscle differentiation
  - d. Abnormal mitotic figures
44. **In cystic fibrosis mutation occurs at?**  
(DNB Aug 12 Pattern)
  - a. One gene
  - b. Two gene
  - c. Three gene
  - d. Four gene
45. **All are true about cystic fibrosis except?** (JIPMER 2012)
  - a. Recurrent respiratory infections
  - b. Majority of males are infertile
  - c. Fasting hyperglycemia is a feature of early disease
  - d. Sweat chloride is >70 meq/l
46. **Which of the following congenital lesion is a deformity?**  
(PGI Nov 2011)
  - a. Potter sequence
  - b. CTEV
  - c. Congenital heart disease
  - d. Cleft lip
  - e. Imperforate anus
47. **Anaplasia in Wilms' tumor is evident by** (PGI Nov 10)
  - a. Increased mitosis
  - b. Pleomorphic nuclei
  - c. Large nucleus
  - d. p53 mutation
  - e. Increased response to chemotherapy
48. **True about nephrogenic rest:** (PGI Nov 10)
  - a. Associated with Wilms' tumor
  - b. Increased risk of Wilms' tumor in contralateral kidney
  - c. No association with Wilms' tumor
  - d. Abnormal embryonal renal tissue
  - e. High risk of Neuroblastoma





## Answers with Explanations

1. **Ans. (a, c, d, e); a. Caused by mutations of the MEFV gene; c. Presentation include episodic bouts of acute peritonitis; d. Amyloidosis occur as a complication; e. Colchicine is used for treatment** (Ref: R 9th/pg 259-260)
2. **Ans. (a, d) a. Associated with aniridia; d. Has pseudocapsule**  
Wilms' tumor is associated with WAGR syndrome which includes Aniridia, grossly have multilocular cysts, mostly well circumscribed and surrounded by a pseudocapsule. It is a pediatric tumor and can present in neonates.
3. **Ans. (a) Medulloblastoma**
4. **Ans. (c) TTF-1**  
Wilms' tumor being mixed tumor has Desmin, Vimentin and Cytokeratin positive. Thyroid transcription factor-1 (TTF-1) is a sensitive marker for pulmonary and thyroid adenocarcinomas
5. **Ans. (c) Trk B expression** (Ref: Robbins 9th/pg 475-479)
6. **Ans. (d) 70 mEq/L** (Ref: Robbins 9th/pg 466-470)
7. **Ans. (a) Leukemia** (Ref: Robbins 9th/pg 473-475)
8. **Ans. (b) ENAC activity of the sweat ducts increases**  
(Ref: Robbins 9th/pg 466-470; 8th 465-470)  
**In cystic fibrosis, there is decreased reabsorption of Na & Cl from sweat glands.**
9. **Ans. (d) Brucellosis** (Ref: Harrison 19th ed)  
**Category A** agents are the highest-priority pathogens. Greatest risk to national security because they (1) be easily disseminated (2) result in high mortality rates (3) might cause public panic (4) require special action for public health. **Category B** agents are the second highest priority pathogens and include those that are moderately easy to disseminate. **Category C** agents are the third highest priority.
10. **Ans. (b) Skull** (Ref: Robbins 9th/pg 473-479)
11. **Ans. (d) Does not respond to chemotherapy and radiotherapy** (Ref: Robbins 9th/pg 479-481)
12. **Ans. (c) Ivacaftor, a potentiator of the CFTR channel is used to for patients with ( $\Delta F508$ ) mutation**  
(Ref: Robbins 9th/pg 466-470; 8th/pg 465-470)
13. **Ans. (d) TRKA expression** (Ref: Robbins 9th/pg 479-481)
14. **Ans. (b) Genomic imprinting** (Ref: R 9th/pg 479-481)
15. **Ans. (a) Hemangioma** (Ref: Robbins 9th/pg 475)
16. **Ans. (a) Cotinine** (Ref: Robbins 9th/pg 475)
17. **Ans. (a) Ameloblastoma** (Ref: Robbins 9th/pg 474-475)
18. **Ans. (b) Osteosarcoma** (Ref: Robbins 9th/pg 474-475)
19. **Ans. (c) Dissemination to bone** (Ref: R 9th/pg 475-479)
20. **Ans. (d) Cystic fibrosis** (Ref: Robbins 9th/pg 466-470)
21. **Ans. (d) Cardiovascular causes** (Ref: R 9th/pg 475)
22. **Ans. (b) 7q** (Ref: Robbins 9th/pg 466-470)
23. **Ans. (a) Lower respiratory tract infections**  
(Ref: Robbins 9th/pg 466-470; 8th/pg 475)
24. **Ans. (a) Ameloblastoma** (Ref: Robbins 9th/pg 475-479)
25. **Ans. (b) Osteosarcoma** (Ref: Robbins 9th/pg 475-479)
26. **Ans. (d) Neuroblastoma** (Ref: Robbins 9th/pg 475-479)
27. **Ans. (c) Neuroblastoma** (Ref: Robbins 9th/pg 475-479)
28. **Ans. (a) Wilms' tumor** (Ref: Robbins 9th/pg 479-481)
29. **Ans. (b) TRKA expression** (Ref: Robbins 9th/pg 475-479)
30. **Ans. (d) Cystic fibrosis** (Ref: Robbins 9th/pg 466-470)
31. **Ans. (c) Polyhydramnios**  
(Ref: Robbins 9th/pg 452; 8th/pg 448N19: 1827)  
Potter sequence (oligohydramnios sequence)
  - Bilateral Renal agenesis  $\rightarrow$  Oligohydramnios  $\rightarrow$  flattened facies, abnormalities of hands and feet and pulmonary hypoplasia.
  - Neonates with bilateral renal agenesis die of pulmonary insufficiency from pulmonary hypoplasia rather than renal failure.
32. **Ans. (a) Leukemia** (Ref: Robbins 9th/pg 473-475)  
**The most frequent childhood cancers are: (in order of decreasing frequency)<sup>Q</sup>**
  - Leukemia (Most common is ALL),<sup>Q</sup> Neuroblastoma,<sup>Q</sup> Wilms' tumor, Hepatoblastoma, Retinoblastoma
33. **Ans. (c) Growth retardation** (Ref: R 9th/pg 479-481)
34. **Ans. (a) Neuroblastoma** (Ref: Robbins 9th/pg 475-479)  
**Neuroblastoma** is the:
  - Most common **extracranial solid tumor** of childhood<sup>Q</sup>
  - Most common **abdominal tumor** of childhood<sup>Q</sup>
  - Most **frequently diagnosed** tumor of infancy (<1 year of age)<sup>Q</sup>





### 35. Ans. (a) **Histopathology**

(Ref: Robbins 9th/pg 479-481; 8th/pg 479-481N19: 1759)

- **Histology** plays a major role in **risk stratification of Wilms' tumor**.
- **Absence of anaplasia** is considered a favorable histologic finding.
- Other prognostic factors for Wilms' tumor are: **age, stage, tumor weight, and loss of heterozygosity** at chromosomes 1p & 16q.

### 36. Ans. (d) **All** (Ref: Robbins 9th/pg 475; 8th/pg 475)

**Small round blue cell tumors** are Tumors with **similar histology of small round cell**.

#### "Low NEW MRP"

- |  |   |
|--|---|
| <ul style="list-style-type: none"> <li>• Lymphoma</li> <li>• EwingSarcoma</li> <li>• Medulloblastoma</li> <li>• Primitive Neuroectodermal Tumor</li> </ul> | <ul style="list-style-type: none"> <li>• Neuroblastoma</li> <li>• Wilms' Tumor</li> <li>• Retinoblastoma</li> </ul> |
|--|---|

### 37. Ans. (b) **Increased** (Ref: Robbins 9th/pg 466-470)

### 38. Ans. (a) **Trk-A expression absent** (Ref: R 9th/pg 475-479)

### 39. Ans. (c) **Small cell carcinoma** (Ref: Robbins 9th/pg 475)

Small cell Carcinoma of lungs is a tumor of adults.

### 40. Ans. (a) **Decreased chloride in sweat**

(Ref: R 9th/pg 466-470)

### 41. Ans. (a) **Osteosarcoma of lower limbs and soft tissue sarcoma** (Ref: Robbins 9th/pg 475, 1339; 8th/pg 475, 1365)

Clinical features of Neuroblastoma: reflect the tumor site and Extent of disease

Extent of disease	Clinical features
<b>Localized disease</b>	<b>Asymptomatic mass or as mass-related symptoms</b> ; E.g. spinal cord compression, bowel obstruction and superior vena cava syndrome.
<b>Metastatic disease</b>	Fever, irritability, failure to thrive, bone pain, cytopenias, bluish subcutaneous nodules, <b>bilateral orbital proptosis, and periorbitalecchymoses (raccoon eyes)</b>
<b>Neurologic involvement</b>	Horner syndrome, nerve root compression
<b>Paraneoplastic syndrome</b>	Opsoclonus-myoclonus-ataxia syndrome
<b>If catecholamines produced</b>	Increased sweating, hypertension, secretory diarrhea (due to release of VIP)
<b>Extensive tumors</b>	Tumor lysis syndrome and DIC
<b>Infants with stage 4S</b>	Subcutaneous tumor nodules, massive liver involvement, limited bone marrow disease & a small primary tumor without bone involvement or other metastases.

### 42. Ans. (a) **Retinoblastoma**

(Ref: Robbins 9th/pg 475-479; 8th/pg 475-478)

Children with **germline retinoblastoma** are more likely to develop other primary malignancies like **Osteosarcoma of lower limbs** and **soft tissue sarcoma**, in their later lifetime course.

### 43. Ans. (b) **Nephrogenic rests** (Ref: Robbins 9th/pg 479-481)

Nephrogenic rests

- They are putative **precursor lesions of Wilms' tumors**
- Patients with presence of **nephrogenic rests in the resected specimen, are at an increased risk of developing Wilms' tumors in the contralateral kidney** and require frequent and regular surveillance.

### 44. Ans. (a) **One gene** (Ref: Robbins 9th/pg 466-470)

**Cystic Fibrosis** an **Autosomal recessive disease**, caused by mutation in **Cystic Fibrosis Transmembrane conductance Regulator (CFTR) gene on chr7q31.2**, which leads to abnormal function of **ion transport channel**.

### 45. Ans. (c) **Fasting hyperglycemia is a feature of early disease**

(Ref: Refer Ans 7 & 10; Nelson 19th/pg 1996; Robbins 9th/pg 466-470; 8th/pg 465-470)

Cystic fibrosis-related diabetes (CFRD)

- **No diabetes is seen in CF patients younger than 10 yr while 40-50% show diabetes at ≥ 20 yr age;**
- **So diabetes is a late feature of CF, seen only in individuals who survive to adolescence & beyond.**
- Patients with CFRD have features of both **T1DM and T2DM**.
- There is pancreatic damage leading to slowly progressive insulin deficiency, along with insulin resistance

### 46. Ans. (b) **CTEV** (Ref: Robbins 9th/pg 452; 8th/pg 448)

**Deformations or deformities are structural abnormalities (extrinsic disturbance in morphogenesis)** due to **compression of growing fetus by abnormal biomechanical forces** like bicornuate uterus. Eg: **Clubfeet**<sup>o</sup>

### 47. Ans. (a, b, c, d) **a. Increased mitosis; b. Pleomorphic nuclei; c. Large nucleus; d. p53 mutation**

(Ref: Robbins 9th/pg 481)

**Anaplasia in Wilms' tumor** (seen in **5%** of cases) is defined as the presence of cells with:

- Large, hyperchromatic, **pleomorphic nuclei**
- **Abnormal mitoses**

Anaplasia also correlates with the **presence of TP53 mutations; loss of p53 function** makes anaplastic cells relatively **unresponsive to cytotoxic chemotherapy**.

### 48. Ans. (a, b, d) **a. Associated with Wilms' tumor; b. Increased risk of Wilms' tumor in contralateral kidney; d. Abnormal embryonal renal tissue**

(Ref: R 9th/pg 479-481)

# White Blood Cells and its Disorders

## Key Points

- » Hematopoiesis starts between 3-4<sup>th</sup> wk of Intrauterine life in yolk sac
- » Leukopenia is abnormally **low WBC (TLC <4000/uL)** while **Leukocytosis is increase in the count of WBCs (>11,000/uL)**
- » **Leukemia** refers to hematological neoplasms with involvement of **bone marrow & peripheral blood** **Lymphoma** refers to **discrete tissue masses<sup>o</sup>** usually involving Lymph node, Spleen and Liver
- » ALL is the **most common cancer of children.**
- » **The most common leukemia** of adults in the **Western world is CLL.**
- » The most common site for extranodal lymphoma is stomach
- » **Follicular lymphoma is the most common** form of **indolent (low grade) NHL** in the West.
- » **DLBCL, is the most common** form of **NHL in India**
- » Burkitts lymphoma shows **"starry sky" pattern**
- » Diagnostic Hallmark of Hodgkins lymphoma are Reed-Sternberg cells
- » Cut off for blast counts in AML is <20% if AML is associated with cytogenetic abnormalities like t(15;17), t(8;21), inv(16)
- » **BCR-ABL gene<sup>o</sup>** (210 kDa in size) is hallmark of CML

## Key Recent Updates

- » CSF3R mutation is seen in CNL
- » Provisional response to TKI is added in accelerated phase of CML.



## HEMATOPOIESIS

Formation of blood components during embryonic stage and throughout adult life

- **Definitive hematopoiesis:**
  - Forms multipotent hematopoietic cells (HSCs) at 4th week of intrauterine life around Aorta, gonads and mesonephros
- **Sites at different ages:**


Age group	Site of Hematopoiesis
Embryo <sup>Q</sup>	Till the 3 <sup>rd</sup> wk in Yolk sac; <sup>Q</sup> Up to 3 <sup>rd</sup> month in Liver <sup>Q</sup>
Fetus	4 <sup>th</sup> month onwards: Bone marrow <sup>Q</sup>
Birth	Bone marrow <sup>Q</sup>
Child	Bone marrow: <b>throughout the skeleton</b> <sup>Q</sup>
Adult	Bone marrow: <b>Flat bone</b> (Vertebra, ribs, sternum, pelvis) <sup>Q</sup> & proximal epiphysis of humerus & femur

### Properties of Hematopoietic Stem Cells

- **Pluripotency**<sup>Q</sup> - Ability of a single HSC to generate all mature blood cells
- Capacity for **self-renewal**<sup>Q</sup>
- **Not** seen usually in **peripheral blood**<sup>Q</sup>
- Under conditions of **stress**, e.g. severe anemia or acute inflammation, HSCs are **mobilized from bone marrow** and appear in the **peripheral blood**.<sup>Q</sup>

### Morphology of Bone Marrow

Normal Myeloid: Erythroid ratio = 3-4:1 <sup>Q</sup>	Light Microscopy
Normal ratio of marrow cells: fat cells=1:1 <sup>Q</sup>	<ul style="list-style-type: none"> <li>• Thin-walled <b>sinusoids</b><sup>Q</sup> lined by <b>single layer of endothelial cells</b></li> <li>• Clusters of <b>hematopoietic &amp; fat cells</b> within the interstitium</li> <li>• <b>Megakaryocytes</b> lie <b>next to sinusoids</b><sup>Q</sup> where they release platelets</li> <li>• Red cell precursors (<b>Erythroblasts</b>) surround macrophages (so-called <b>nurse cells</b>)<sup>Q</sup></li> </ul>
Normal Cellularity (%) = <b>100 – Age</b> <sup>Q</sup> <ul style="list-style-type: none"> <li>• Decreases with age</li> <li>• At 10 yrs = 100-10 = 90% cellularity</li> <li>• At 30 yrs = 100-30 = 70% cellularity</li> </ul>	



### High Yield Facts

- **Agranulocytosis:** Clinically significant reduction in **neutrophils** making one susceptible to bacterial & fungal infections.<sup>Q</sup>
- **Drugs** are the most common cause of **agranulocytosis**<sup>Q</sup>
- Serious infection increases when ANC < 500/mm<sup>3</sup>

## DISORDERS OF WHITE BLOOD CELLS

- Quantitative defects
  - Leucopenia
  - Leucocytosis
- Qualitative defects

### Leukopenia

- **Definition:**
  - An abnormally **low white cell count** (leukopenia; TLC <4000/ $\mu$ L)<sup>Q</sup>



## Lymphopenia

- **Definition**
  - Reduction in number of **lymphocytes** in blood
- **Etiology:**
  - **Congenital immunodeficiency diseases, e.g. SCID<sup>Q</sup>**
  - **Human immunodeficiency virus (HIV) infection<sup>Q</sup>**
  - **Glucocorticoids<sup>Q</sup>** or cytotoxic drugs
  - Autoimmune disorders
  - Malnutrition
  - **Acute viral infections<sup>Q</sup>**

## Neutropenia

- **Definition**
  - Reduction in the number of neutrophils (**<1500/ $\mu$ L**) in the blood<sup>Q</sup>
- **Etiology**
  - A. Drug Induced Neutropenia**
    - **Anti-bacterials** – **Chloramphenicol<sup>Q</sup>**, Cotrimoxazole, Ciprofloxacin, Nitrofurantoin

- **Anti-inflammatory- Ibuprofen<sup>Q</sup>**
- **Antithyroid-** Carbimazole, Propylthiouracil
- **Anticonvulsants-** Valproate, Phenytoin
- B. Inadequate or ineffective granulopoiesis:<sup>Q</sup>**
  - **Aplastic anemia<sup>Q</sup>**
  - Infiltrative marrow disorders (e.g., tumors, granulomatous disease)
  - Infections: **Viral- Parvo B19<sup>Q</sup>, HIV, EBV**
  - Ineffective hematopoiesis: **Megaloblastic anemias<sup>Q</sup>** & Myelodysplastic syndromes
  - **Kostmann syndrome<sup>Q</sup>:** Autosomal recessive congenital neutropenia
  - Cyclic Neutropenia
- C. Accelerated destruction or sequestration of neutrophils**
  - Immunological injury to neutrophils, e.g. **SLE<sup>Q</sup>**
  - **Splenomegaly<sup>Q</sup>**
  - Increased peripheral utilization: **bacterial, fungal, or rickettsial infections**

## Leukocytosis

Increase in the number of WBCs (**>11,000/ $\mu$ L**)<sup>Q</sup>

### Causes of Leukocytosis<sup>Q</sup>

Type of Leukocytosis	Causes
<b>Neutrophilia (&gt;75%)</b>	<ul style="list-style-type: none"> <li>• <b>Infection (bacterial)<sup>Q</sup> &amp; Inflammation</b> including <b>MI<sup>Q</sup></b></li> <li>• Acute <b>stress</b> states (burns, post-surgery)</li> <li>• <b>Myeloproliferative</b> disorders: CML, <b>Polycythemia vera</b></li> <li>• Others: <b>steroid<sup>Q</sup></b> therapy, <b>Renal failure<sup>Q</sup></b></li> </ul>
<b>Eosinophilia (&gt;400/<math>\mu</math>L)</b>	<ul style="list-style-type: none"> <li>• <b>Allergies:</b> <b>Asthma<sup>Q</sup>, hay fever<sup>Q</sup>, urticaria</b></li> <li>• <b>Skin diseases:</b> Eczema, <b>dermatitis herpetiformis<sup>Q</sup></b></li> <li>• <b>Parasitic:</b> Ascariasis, <b>Hookworm<sup>Q</sup></b>, Filariasis, Trichinosis</li> <li>• Others: <b>Tropical eosinophilia<sup>Q</sup>, Hypereosinophilic syndrome, Hodgkin's disease</b></li> </ul>
<b>Basophilia (&gt;1%)</b>	CML <sup>Q</sup> , PCV, Ulcerative colitis <sup>Q</sup> , Mastocytosis <sup>Q</sup> , Myxedema
<b>Monocytosis</b>	<ul style="list-style-type: none"> <li>• <b>Infections:</b> TB<sup>Q</sup>, Kala azar, malaria, Syphilis</li> <li>• <b>Malignancies:</b> <b>AML-M4/5<sup>Q</sup></b>, CMML, Hodgkin's</li> <li>• <b>Inflammatory</b> diseases: Ulcerative colitis<sup>Q</sup>, Crohn's, SLE, Sarcoidosis</li> </ul>
<b>Lymphocytosis</b>	<ul style="list-style-type: none"> <li>• <b>Bacterial:</b> TB, <b>brucellosis</b>, Syphilis, <b>pertussis</b>, <b>Diphtheria<sup>Q</sup></b></li> <li>• <b>Viral infections:</b> <b>Infectious mononucleosis<sup>Q</sup></b>, Mumps, <b>Malignancies:</b> <b>CLL<sup>Q</sup>, NHL, Hairy Cell Leukemia<sup>Q</sup></b></li> </ul>

### Qualitative defects in WBCs

Anomaly	Inheritance	Characteristic	Other features
<b>May Hegglin anomaly</b>	AD	<b>Basophilic inclusions</b> in WBCs	<b>Giant platelets &amp; thrombocytopenia</b>
<b>Alder-Reilly anomaly</b>	AR	<b>Lilac inclusions</b> in Neutrophils	Stains with <b>Toluidine Blue</b>
<b>Pegler Huet anomaly</b>	AD	<b>Hypo-segmented</b> Neutrophils	<b>Distinct from Pseudo Pegler-Huet anomaly</b>
<b>Döhle bodies</b>		Patches of dilated endoplasmic reticulum that appear as sky-blue cytoplasmic "puddles."	
<b>Toxic granules</b>		Which are coarser and darker than the normal neutrophilic granules, represent abnormal azurophilic (primary) granules.	






## High Yield Facts

- Infection of **B lymphocytes** by EBV occurs in **Infectious Mononucleosis**
- Atypical lymphocytes that are characteristic of infectious mononucleosis are **CD8+ T lymphocytes** called **DOWNY cells**, that develop in response to the infected B lymphocytes.
- Reed Sternberg like cell are seen in **Infectious mononucleosis**, **Adult T cell lymphoma**, **Diffuse large B cell lymphoma**
- **Kikuchi disease** or 'histiocytic necrotizing lymphadenitis' is benign, recurrent necrotizing lymphadenitis
- Eosinophilic abscess in lymph node is characteristically seen in - **Kimura's disease**
- **Typhoid** (Enteric fever) presents with **lymphopenia** and not **leukocytosis**


## NEOPLASTIC PROLIFERATIONS OF WHITE CELLS

- **Leukemia:** Hematological Neoplasms with involvement of **bone marrow and peripheral blood**<sup>Q</sup>
- **Lymphoma:** Hematological Neoplasms where proliferations arise as **discrete tissue masses**<sup>Q</sup> usually involving Lymph node, Spleen, Liver<sup>Q</sup>



## Lymphoid Neoplasms

### World Health Organization (WHO) 2008 Classification of Lymphoid Neoplasms



## High Yield Facts

- **CD19** is the earliest recognizable marker of B cells & is lost when B cell becomes a plasma cell
- **CD 34** is the surface glycoproteins that is most often expressed in human hematopoietic stem cell
- Outside the hematopoietic system, **CD34** is expressed on **endothelial cells**.
- **CD 45** is found in **all hematopoietic cells except erythrocytes**
- **CD 46** is a receptor of pathogen like HHV-6, Streptococci
- **PAX9** – Marker of B cells

### Acute Lymphoblastic Leukemia (ALL)

- **Definition:**
  - Neoplasms of **immature B (pre-B) or T (pre-T) cells** which are referred to as **lymphoblasts**
- **Epidemiology:**
  - ALL is the **most common cancer of children**<sup>Q</sup>, Peak Incidence: **3<sup>rd</sup> yr**<sup>Q</sup>
  - **Hispanics** have the highest incidence of any ethnic group.




- **Pathogenesis:**
  - T-ALLs have **gain-of-function** mutations in **NOTCH1**<sup>Q</sup>
  - B-ALLs have **loss-of-function** mutations **PAX5**<sup>Q</sup>, **E2A**<sup>Q</sup> & **EBF**<sup>Q</sup>, or **t(12;21)**<sup>Q</sup> involving the genes **ETV6** and **RUNX1**, 2 genes that are needed in very early hematopoietic precursor.
- **Clinical features:**
  - Abrupt **stormy onset**
  - Symptoms related to depression of marrow function:
    - **Fatigue** due to anemia;
    - **Fever** due to neutropenia; and
    - **Bleeding** due to thrombocytopenia
  - **Marrow expansion** and infiltration of the **sub-periosteum: sternal tenderness**<sup>Q</sup>
- Generalized lymphadenopathy, Hepatosplenomegaly; **testicular enlargement**
- **T-ALL:** Mediastinal mass (**Superior Mediastinal Syndrome**)<sup>Q</sup>
- **CNS features:** headache, vomiting, and nerve palsies
- **Morphology:**
  - Hypercellular Bone Marrow; > **20% lymphoblasts**<sup>Q</sup>
  - Compared with myeloblasts, **lymphoblasts** have **more condensed chromatin**, less conspicuous nucleoli & scanty agranular cytoplasm
  - Cytochemistry: Myeloperoxidase (**MPO**) **-ve**, Sudan Black B (**SBB**): **-ve**<sup>Q</sup>
  - **Diagnosis of choice**
  - Flow cytometry

## Classification of ALL

### FAB (French American British) Classification

ALL-subtype	L1	L2	L3 (Mature B-cells)Q
<b>Morphology of Blasts</b>	<ul style="list-style-type: none"> <li>• Small Homogenous Blasts</li> <li>• Little Cytoplasm</li> <li>• Regular Nucleus,</li> <li>• Small indistinct nucleoli</li> </ul>	<ul style="list-style-type: none"> <li>• Large heterogeneous blasts</li> <li>• One or more nucleoli</li> </ul>	<ul style="list-style-type: none"> <li>• Large homogenous blasts</li> <li>• Abundant basophilic cytoplasm</li> <li>• Prominent <b>cytoplasmic vacuolation</b><sup>Q</sup></li> <li>• Resemble <b>Burkitt</b><sup>Q</sup> lymphoma</li> </ul>
<b>Age group</b>	Children	Adults	Adults
<b>Prognosis</b>	Good <sup>Q</sup>	Intermediate <sup>Q</sup>	Poor <sup>Q</sup>
<b>Cytochemistry</b>	PAS +	PAS +	PAS-, SBB+ <sup>Q</sup>



### High Yield Facts

- **Mature B-cell ALL** is an uncommon type of **ALL**<sup>Q</sup> (1-2% of ALL cases) in children.
- Both **B-cell ALL** and **Burkitt lymphoma** are characterized by **FAB L3**<sup>Q</sup> morphology,
- **Mature B-cell ALL** is associated with **t(8;14)** & overexpression of the **c-myc** oncogene
- **T-ALL** commonly presents with Mediastinal mass (**Superior Mediastinal Syndrome**)<sup>Q</sup>
- **T-ALL** are **aggressive lymphomas**.<sup>Q</sup>
- In **T-ALL**, cells are positive for markers of blasts like- **Tdt**,<sup>Q</sup> **CD34** & **T cell markers CD1, CD2, CD5, CD7**<sup>Q</sup>
- **Response to treatment is the best prognostic marker in ALL**

### WHO 2018: Classification of ALL

- B-Lymphoblastic leukemia, Not otherwise specified (NOS)
- B-Lymphoblastic leukemia with recurrent cytogenetic abnormality

1. t(12;21)ETV6-RUNX1	3. t(9;22) BCR-ABL1	5. B cell ALL with hyperdiploidy	7. BCR-ABL 1 like ALL (Provisional entity in 2018 WHO)
2. t(v,11)(KMT2A-MLL)	4. t(5;14) IgH-IL3	6. B cell ALL t(1, 19)-TCF3 -PBX1	8. iAMP 21 ALL.

- T-Lymphoblastic leukemia




## Prognostic Factors in Acute Lymphoblastic Leukemia

Determinants	Favorable	Unfavorable
WBC/uL	<10,000	>2,00,000 <sup>a</sup>
Age	2–9 yr	<1 y, >10 y <sup>a</sup>
Gender	Female	Male <sup>a</sup>
Ethnicity	White	Black <sup>a</sup>
L. node, liver, spleen enlargement	Absent	Massive
Testicular enlargement	Absent <sup>a</sup>	Present <sup>a</sup>
Central nervous system leukemia	Absent	Present <sup>a</sup>
FAB morphologic features	L1 <sup>a</sup> Early pre-B-cell ALL	L2 <sup>a</sup> Pre-B-cell ALL Mature B-cell
Ploidy	Hyperdiploidy	Hypodiploidy<45
Cytogenetic markers	Trisomy 4, 10, 17 <sup>a</sup> t(12;21)	t(9;22) t(4;11)
Remission states	< 14 days	> 14 days

## Acute Myeloid Leukemias (AML)

- **Definition:**
  - Neoplasms of myeloid cells which are referred to as myeloblasts<sup>o</sup>
- **Clinical features:**
  - Same as ALL but in addition certain types of AML show:
    - Chloromas (AML M2 > 5 > 4)<sup>o</sup>
    - Gingival hyperplasia (AML M5 > 4)<sup>o</sup>
    - DIC (AML M3)<sup>o</sup>
- **Predisposing Conditions:**

Genetic factors	Congenital bone marrow failure syndromes	Drugs
• Down's syndrome	• Kostmann syndrome	• Benzene
• Fanconi's anemia	• Diamond – Blackfan anemia	• Alkylating agents
• Bloom's syndrome		• Epipodophylotoxins
• Neurofibromatosis type 1		• Ionizing radiation
• Klinefelter syndrome		
• Turner syndrome		



Pathogenesis of acute myeloid leukemia



## Major Subtypes of AML in the WHO Classification 2018

Class	Prognosis	FAB Subtype	Morphology/Comments
<b>I. AML WITH RECURRENT GENETIC ABERRATIONS</b>			
<b>a. AML with balanced translocations</b>			
AML with t(8;21) <sup>q</sup> RUNX <sub>1</sub> – RUNX <sub>1</sub> T <sub>1</sub>	Favorable	M2 <sup>q</sup>	Auer rods++; abnormal cytoplasmic granules
AML with inv (16) <sup>q</sup> CBFB-MYH II	Favorable	M4	abnormal eosinophilic precursors <sup>q</sup>
AML with t(15;17) PML-RARA	Intermediate	M3	Auer rods +++, high incidence of DIC <sup>q</sup>
AML with t(9;11) KMT <sub>2A</sub> -MLL	Poor	Variable	
AML with BCR-ABL <sub>1</sub> mutation			Responds to R <sub>x</sub>
<b>b. AML with gene mutations</b>			
AML with mutated NPM <sub>1</sub>	Favorable		
AML with Biallelic mutation of CEBPA	Favorable		
<b>II. AML WITH MDS-LIKE FEATURES</b>			
AML with MDS-like cytogenetic aberrations	Poor	Variable	Associated with 5q-, 7q-, Monosomy 5 and 7 <sup>q</sup>
<b>III. AML, THERAPY-RELATED</b>			
1. Post alkylating agents	Very poor	Variable	5–10 years after exposure • Unbalanced loss of chr. 5 & 7 & loss of p53
2. Post DNA topoisomerase II			1–5 years after exposure • Balanced chromosomal translocations
<b>IV. AML, NOT OTHERWISE SPECIFIED (previously FAB)</b>			
AML, minimally differentiated	Poor <sup>q</sup>	M0	MPO –VE
AML without maturation	Intermediate	M1	MPO +ve in >3% of blasts
AML with myelocytic maturation	Intermediate	M2	myelocytic maturation, Auer Rods ++
AML with myelomonocytic maturation	Intermediate	M4	Myelocytic and monocytic differentiation <b>MPO +, NSE +<sup>q</sup></b>
AML with monocytic maturation	Intermediate	M5	Monoblasts and pro-monocytes predominate, <b>Non-specific esterase (NSE)+<sup>q</sup></b>
AML with erythroid maturation	Poor <sup>q</sup>	M6	>80% erythroid precursors
AML with megakaryocytic maturation	Poor <sup>q</sup>	M7	>50% megakaryocytic blasts <b>Most common Acute Leukemia in Down syndrome</b>

### Diagnosis of AML

#### ■ Bone marrow Morphology:

- > 20% myeloid blasts in the bone marrow<sup>q</sup>
- Myeloblasts have **delicate nuclear chromatin**, 2-4 nucleoli, and moderate cytoplasm with or without Auer rods
- **Auer rods: Most reliable** morphological feature of AML<sup>q</sup>
  - Needle-like azurophilic fusiform inclusions in cytoplasm of myeloblasts
  - Stain +ve with **MPO and Sudan Black B<sup>q</sup>**
  - Seen in AML **M2, M3** (also seen in **CML blast crisis** and **MDS**)<sup>q</sup>
  - **Faggots:** bundles of Auer Rods in crisscross pattern<sup>q</sup>
- **Phi Body:** round or oval inclusions in blasts<sup>q</sup>

#### ■ Cytochemistry:

- Blasts stain positive for:
  - **Myeloid stain:** Myeloperoxidase (**MPO**), Sudan Black B (**SBB**)-M2/3

- **Monocytic stain:** Non-specific esterase (**NSE**) M5<sup>q</sup>
- Both **MPO/SBB & NSE:** M4<sup>q</sup>

#### ■ Diagnosis of choice

- Flow cytometry



### High Yield Facts

Cytochemical stains	Cells stained
<b>Myeloperoxidase (MPO)</b>	Myeloid <sup>q</sup>
<b>Sudan Black B (SBB)</b>	Myeloid <sup>q</sup>
<b>Periodic acid Schiff (PAS)</b>	Lymphoid (Block positivity) <sup>q</sup>
<b>Non-specific esterase (NSE)</b>	Monocytes <sup>q</sup> >> Myeloid
<b>Acid phosphatase</b>	T-lymphocyte <sup>q</sup>
<b>Tartrate resistant acid phosphatase (TRAP)</b>	Hairy cell leukemia <sup>q</sup>







## High Yield Facts


- **Pan B-** marker is **CD 19<sup>a</sup>**
- **Pan T-** marker is **CD3<sup>a</sup>**
- **Memory cells** have **CD45RO<sup>a</sup>**
- **CD 71:** Transferrin receptors
- **CD95<sup>a</sup>** is the major receptor for **apoptosis**
- **Cut off for blast counts in AML** is **< 20%** if AML is associated with cytogenetic abnormalities like **t(15;17), t(8;21), inv(16)<sup>a</sup>**
- AML causing **gum hypertrophy/infiltration** are **AML-M5, M4<sup>a</sup>**
- AML causing **extramedullary blast proliferations (Chloroms)** are **AML M2, M4, M5<sup>a</sup>**
- AML causing blast infiltrations in skin (**leukemia cutis**) are **AML M5, M4<sup>a</sup>**
- **Disseminated intravascular coagulation (DIC)<sup>a</sup>** can be seen in Acute promyelocytic leukemic (**APML, M3**)

## Differences between Myeloblast and lymphoblast in Acute Leukemia

Parametres	Myeloblast	Lymphoblast
Size	Larger (18-20 $\mu$ )	Smaller (10-18 $\mu$ )
Cytoplasm	Moderate and granular	Scant and agranular
N/C ratio	High	Very high
Nuclear chromatin	Fine and stippled	Coarser
Nucleoli	2-5 prominent	0-2 Inconspicuous
Auer rods	Present	Absent
Accompanying cells	Myelocytes, metamyelocytes, stab and neutrophils	Lymphocytes



## Origin of peripheral B cell neoplasms





## Peripheral B-Cell Neoplasms


### Chronic Lymphocytic Leukemia (CLL)/ Small Lymphocytic Lymphoma (SLL)

- **Clinical criteria:**
  - Absolute **clonal lymphocytes** >5000/uL<sup>Q</sup>
  - **MC leukemia** of adults in the **Western world**.
  - Median age at diagnosis is **60 years**; **M:F = 2 : 1**
- MC mutation-Del 13q.
- Cell of origin-Naive B cell.
- **Peripheral Smear:**
  - **Small round lymphocytes** with scant cytoplasm (**CONVENT** girl appearance)<sup>Q</sup>
  - Occasional cells have distorted outline called **smudge cells**<sup>Q</sup>
  - Rarely **Warm type AIHA** may develop showing Spherocytes<sup>Q</sup>
- **Morphology of Lymph nodes:**
  - **Diffusely effaced**<sup>Q</sup> by an infiltrate of predominantly small lymphocytes (6- 12µm) between which lies larger activated lymphocytes- **proliferation centers**<sup>Q</sup> (**pathognomonic for CLL/SLL**), which contain **mitotically active cells**.
  - Overall CLL has **low mitotic rate**<sup>Q</sup> exc in proliferative center
- **Diagnosis of choice**
  - **Immunophenotyping**
  - **Dim Surface Ig** (usually IgM or IgM and IgD)<sup>Q</sup>
  - **Pan B-cell markers** CD19 + and CD20+<sup>Q</sup>
  - **CD23+ and CD5+<sup>Q</sup>**


- **Poor prognosis markers:**
  - Rai (stage 3 /4) and Binnet (stage C)
  - High **β2 microglobulin**
  - **Diffuse marrow** involvement
  - Lymphocyte doubling time (<1 yr)
  - **High LDH**
  - Increased Serum **CD23**<sup>Q</sup>
  - **Lack of somatic IghVh hypermutation**<sup>Q</sup>
  - **ZAP-70 +ve**
  - **CD38 +ve**
  - Presence of **NOTCH1 mutations**<sup>Q</sup>
- **Transformations of CLL (poor prognosis):**
  - Diffuse large B-cell lymphoma - **Richter syndrome** (5% - 10%)<sup>Q</sup>
  - Large-cell transformation to **prolymphocytic leukemia (PLL)**<sup>Q</sup>
  - **Acute Leukemia**
  - 2<sup>nd</sup> malignancy- **Melanoma and CNS tumors**<sup>Q</sup>

### High Yield Facts

- Rai and Binnet staging system was used for CLL
- Almost never develops after radiation
- M.C genetic anomalies in CLL are **del 13q14.3<sup>Q</sup>**, 11q, and 17p, and trisomy 12q.<sup>Q</sup>
- Micro-RNAs: **miR-15a and miR-16-1** (tumor suppressor genes): **good prognosis in CLL**<sup>Q</sup>
- CLL with Somatically **hypermuted** Ig genes have **indolent course**<sup>Q</sup>
- CLL with **Unmutated** Ig genes (naive B-cell origin) have **aggressive course**<sup>Q</sup>



CLL with smudge cells



Lymph node biopsy showing effacement by small lymphocytes

### Mantle Cell Lymphoma

- **Definition:**
  - Tumor arising from **mantle zone**<sup>Q</sup> which surrounds germinal centers
- **Seen in:**
  - **M>F**; Most common age: **5<sup>th</sup>-6<sup>th</sup> decade**
- **Pathogenesis:**
  - **t(11;14)**<sup>Q</sup> → overexpression of **cyclin D1** → promotes **G1-to S-phase** progression during the cell cycle.
- **Presentation:**
  - Usual presentation: **Lymphadenopathy** (with occasional spill to peripheral blood)<sup>Q</sup>
  - Unusual presentation: **Lymphomatoid polyposis**<sup>Q</sup>- mucosal involvement of the **small bowel or colon** producing **polyp-like lesions**
- **Morphology of lymph node:**
  - A **homogeneous population of small lymphocytes** with deeply **clefted (cleaved) nuclear** contours




- **Immunophenotype:**
  - Express high levels of **cyclin D1, CD19, CD20, CD45** and **surface Ig<sup>Q</sup>**.
  - **CD5+ve<sup>Q</sup>** and **CD23-ve** which help to distinguish it from CLL/SLL.
  - **Most sensitive marker is SOX II**
- **Prognosis:**
  - Poor; Median survival of **3 - 4 years**.

### Follicular Lymphoma

**Most common form of indolent (low grade) NHL in the West.<sup>Q</sup>**

- **Clinical feature**
  - Presents with painless, generalized **lymphadenopathy**.
- **Pathogenesis**
  - Arises from **germinal center of B cells<sup>Q</sup>**
  - Hallmark translocation **t(14;18)<sup>Q</sup>**
  - Mutations in **MLL gene<sup>Q</sup>** (histone-methyl transferase) that regulates **gene expression (90%)**
- **Morphology**
  - **Lymph node:** Predominantly **nodular** or **nodular & diffuse** growth pattern with **centrocytes** (small cleaved cells) along with **centroblasts<sup>Q</sup>**
  - **Bone marrow:** **paratrabecular** lymphoid aggregates<sup>Q</sup>
- **Immunophenotype**
  - Resemble **germinal center B cells** **CD19, CD20, CD10, surface Ig, and BCL6<sup>Q</sup>**
  - **CD5 -ve<sup>Q</sup>**
  - **BCL2** is expressed in more than 90% of cases<sup>Q</sup>
- **Histologic transformation occurs to:**
  - Diffuse large B-cell lymphoma (**DLBCL**)
  - Burkitt's lymphoma (**BL**)



Follicular lymphoma (Lymph node Biopsy)

### Diffuse Large B-Cell Lymphoma (DLBCL)

**Most common form of NHL in India<sup>Q</sup>**

- **Epidemiology:**
  - **M>F**, Median age = **60 yrs<sup>Q</sup>**
- **Pathogenesis:**
  - Pathogenic event is dysregulation of **BCL6<sup>Q</sup>**
  - **10% - 20% have t(14;18)<sup>Q</sup>**
- **Morphology:**
  - Tumor cells have **large cell size** (4-5 times size of small lymphocyte) & **diffuse pattern** of growth. B4 is involved late
- **Immunophenotype:**
  - **CD19+ and CD20+, CD10+ and BCL6+, surface Ig+<sup>Q</sup>**
- **Special Subtypes:**

Immunodeficiency-associated large B-cell lymphoma	Primary effusion lymphoma
<ul style="list-style-type: none"> <li>• Severe T-cell immunodeficiency</li> <li>• (HIV, allogeneic bone marrow transplantation)</li> <li>• Co-infection with <b>EBV<sup>Q</sup></b></li> </ul>	<ul style="list-style-type: none"> <li>• Malignant pleural effusion or ascites in advanced <b>HIV<sup>Q</sup></b> infected patients</li> <li>• Co-infection with <b>KSHV/HHV-8<sup>Q</sup></b></li> <li>• IHC : <b>CD38, CD30+, CD20 -</b></li> </ul>

- **Prognosis:**
  - **Poor prognosis with aggressive course**

### Malt Lymphoma (MALToMa)

- **MALT lymphomas** express **B-cell antigens (CD19 and CD20)** & **monotypic surface Ig (IgM without IgD)**.
- **MALTomas** may be **CD43+** but lack other small B-cell lymphoma markers (CD5, CD10, CD23 & cyclin D1)
- **MALToma** of salivary glands in sjogrens & hashimoto thyroiditis show morphology of marginal zone lymphoma



### High Yield Facts

- **Nasal NK/T lymphoma** may present with facial swelling/ destruction, so called **lethal midline granuloma** or **polymorphic reticulosis**.
- **Most common ocular lymphoma** is **B-cell NHL**
- **The most site for extranodal lymphoma** is **Stomach**

### Burkitt's Lymphoma


<b>Pathogenesis</b>	<b>MYC gene<sup>Q</sup> (chr 8)</b> (transcriptional regulator)-characteristic but not specific		
<b>Hallmark translocations:</b>	<ul style="list-style-type: none"> <li>• <b>t(8;14) myc; IgH - most characteristic<sup>Q</sup></b></li> <li>• <b>t(2;8) myc ; Ig κ</b></li> <li>• <b>t(8;22) myc ; λ</b></li> </ul>		
<b>Subtypes/Varieties</b>	<b>African (endemic)</b>	<b>Sporadic (Non-endemic)</b>	<b>Immunodeficiency associated (HIV)</b>
<b>Site of involvement</b>	<ul style="list-style-type: none"> <li>• <b>Mandible (M.C)<sup>Q</sup></b></li> <li>• <b>Abdominal viscera</b>; E.g. kidneys, ovaries, adrenals</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Ileocecal region(M.C)<sup>Q</sup></b></li> <li>• <b>Peritoneum</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Lymph nodes</b></li> <li>• <b>Bone marrow</b></li> </ul>
<b>EBV infection</b>	<b>100%<sup>Q</sup></b>	<b>20-30%</b>	<b>25-40%</b>

Contd...






<b>Morphology:</b>	<b>Tumor:</b> High mitotic index <sup>Q</sup> , numerous apoptotic cells, combined with <b>benign macrophages</b> . <ul style="list-style-type: none"> <li>Macrophages have abundant clear cytoplasm: characteristic “<b>starry sky</b>” pattern.<sup>Q</sup></li> </ul> <b>Bone Marrow:</b> <ul style="list-style-type: none"> <li><b>Clumped nuclear chromatin</b> with distinct nucleoli, and royal blue cytoplasm containing clear cytoplasmic vacuoles.</li> </ul>
<b>Immunophenotype</b>	Surface IgM+, CD19+, CD20+, <b>CD10+</b> , and <b>BCL6+</b> <sup>Q</sup>
<b>Prognosis</b>	Burkitt lymphoma is <b>very aggressive</b> but <b>responds well</b> to intensive <b>chemotherapy</b>



Burkitt lymphoma with starry sky pattern




Cells showing cytoplasmic vacuoles

### Hairy Cell Leukemia

Chronic B-cell leukemia characterized by **hairy cells**, **pancytopenia** and **splenomegaly**<sup>Q</sup>

- **Epidemiology:**
  - Median age: **55 years**; M:F ratio of **4-5 : 1**.
- **Pathogenesis:**
  - 90% of cases with activating point mutations in the **serine/threonine kinase BRAF v600E**<sup>Q</sup> (also +ve in **melanoma**, **LCH** and **papillary lung Ca**)<sup>Q</sup>
- **Morphology:**
  - **Peripheral smear:** Pancytopenia with **monocytopenia**; ‘**Hairy**’ tumor cells- Tumor cells with Fine hair like projections, best recognized under the **phase-contrast microscope**.<sup>Q</sup>
  - On Electron microscopy: Hairy cells show **ribosomal-lamellar complexes**<sup>Q</sup>
  - Bone Marrow Aspirate: **Dry Tap** (due to **fibrosis in marrow**)<sup>Q</sup>
  - Bone Marrow Biopsy: Tumor nuclei surrounded by zone of clear cytoplasm giving rise to “**honeycomb**”<sup>Q</sup>, “**fried egg**”<sup>Q</sup>/ “**chicken wire mesh**”<sup>Q</sup> appearance
- **Immunophenotype:**
  - CD19+ and CD20+, **surface Ig +**
  - **Characteristic marker:** **CD103**<sup>Q</sup> along with **CD11c**, **CD25**

- **Cytochemical markers:**
  - Tartrate resistant acid phosphatase (**TRAP**), **DBA 44** and **Annexin A1**.<sup>Q</sup>
- **Clinical Features:**
  - Infiltration of the bone marrow, liver, and spleen- **massive splenomegaly**.<sup>Q</sup>
  - **Pancytopenia** resulting from marrow involvement and splenic sequestration.
  - **Atypical mycobacterial** infections due to monocytopenia.




Hairy cell leukemia






## High Yield Facts

Infections and Associations with Lymphoma	
Agent	Type of Lymphoma
<b>Epstein Barr virus (EBV)</b>	<ul style="list-style-type: none"> <li>Burkitt lymphoma (Africa)</li> <li>Post Transplant Lymphoproliferative Disorders</li> <li>AIDS-related lymphoma (central nervous system, others)</li> <li>Natural killer/T-cell nasal lymphoma</li> <li>Hodgkin lymphoma</li> </ul>
<b>Human T-lymphotropic virus I (HTLV-1)</b>	Adult T-cell leukemia/lymphoma
<b>Human herpes virus 8 (HHV8) or Kaposi sarcoma-associated herpes virus (KSHV)</b>	<ul style="list-style-type: none"> <li><b>Primary effusion lymphoma</b></li> <li><b>Plasmablastic lymphoma</b></li> </ul>
<b>Helicobacter pylori</b>	Gastric MALToma
<b>Hepatitis C virus</b>	Splenic marginal zone lymphoma; other B-cell lymphomas
<b>Campylobacter jejuni</b>	Immunoproliferative small intestinal disease
<b>Borrelia burgdorferi</b>	Primary cutaneous B-cell lymphoma
<b>Chlamydia psittaci</b>	Ocular adnexal lymphoma



Clover cells (Adult T cell leukemia/lymphoma)



Sezary cell (Peripheral T cell lymphoma)




## High Yield Facts

- Clover leaf/flower cells:** Adult T cell leukemia/lymphoma
- Sezary cells:** Peripheral T cell lymphoma
- Hallmark/Doughnut cells:** Anaplastic large cell lymphoma

## Plasma Cell Neoplasms and Related Disorders

### Multiple Myeloma

- Definition:** Malignant proliferation of plasma cells derived from a **single clone**.
- Classification and Diagnostic Criteria of Plasma cell neoplasms



#### Latest Update

##### Myeloma Defining Events (any 1 is sufficient for diagnosing multiple myeloma)

- $\geq 60\%$  clonal plasma cells
- Involved/uninvolved free light chain ratio ( $>100$ )
- $\geq$  focal lesions of  $> 5$  mm in size on (MRI)

#### Classification of Plasma Cell Neoplasia (WHO 2017)

MGUS	Smoldering Myeloma	Multiple Myeloma
IgG/A/M MGUS [All criteria must be met] <ul style="list-style-type: none"> <li>• Serum monoclonal protein (IgG or IgA or IgM <math>&lt;3</math> g/dL AND</li> <li>• Clonal BM plasma cells <math>&lt;10\%</math> AND</li> <li>• No myeloma defining events (see below)</li> </ul>	<ul style="list-style-type: none"> <li>• Serum monoclonal protein (IgG or IgA) <math>\geq 3</math> g/dL or</li> <li>• Urinary monoclonal protein <math>\geq 500</math> mg/24 h and/or</li> <li>• Clonal BM plasma cells <math>10\% - 60\%</math></li> </ul>	<ul style="list-style-type: none"> <li>• Clonal BM plasma cells of <math>\geq 10\%</math> or</li> <li>• Biopsy-proven bony or extramedullary plasmacytoma</li> </ul>
	AND	AND
	<ul style="list-style-type: none"> <li>• No myeloma defining events or amyloidosis (no CRAB and no SLiM) as details below</li> </ul>	<ul style="list-style-type: none"> <li>• 1 or more myeloma defining events as details below</li> <li><math>\geq 1</math> CRAB feature(s)</li> <li>OR</li> <li><math>\geq 1</math> SLiM feature(s)</li> </ul>

Myeloma defining events are evidence of end organ damage that can be attributed to the underlying plasma cell proliferative disorder, especially

**C:** Calcium elevation ( $>11$  mg/dL or  $>1$  mg/dL higher than ULN)

**R:** Renal insufficiency (creatinine clearance  $< 40$  mL/min or serum creatinine  $>2$  mg/dL)

**A:** Anemia (Hb  $<10$  g/dL or 2 g/dL  $<$  normal)

**B:** Bone disease ( $\geq 1$  lytic lesions on skeletal radiography, CT, or PET-CT).

OR, in the absence of **CRAB**, any one or more of the following biomarkers or malignancy, referred to here as the

**SLiM** criteria: **SLiM**: **S** =  $\geq 60\%$  clonal BM plasma cells; **Li** = Serum free Light chain ratio involved: uninvolved  $\geq 100$ ; **M** =  $>1$  focal lesions ( $\geq 5$  mm each) detected by MRI studies

## Lab Diagnosis of Multiple Myeloma

### Peripheral smear:


- Anemia: Normocytic, normochromic
- Rouleaux formation** with basophilic background staining
- Few plasma cells may be seen (**Plasma cells >20% or >2,000/uL** → **Plasma cell leukemia**)<sup>Q</sup>

### Bone marrow:


- Diagnostic hallmark:** Infiltration of marrow by **plasma cells**<sup>Q</sup>
- Plasmablasts** may be present
- Mott Cells/Grape cells**<sup>Q</sup>: Cells with small spherical inclusions of Immunoglobulins
- Flame cells/Thesaurocytes**<sup>Q</sup>: Orange red flame like peripheral rim
- Inclusion bodies in plasma cells:
  - Dutcher body**- Intranuclear<sup>Q</sup>
  - Russel body**-Intracytoplasmic<sup>Q</sup>

### Other Investigations:


- Serum electrophoresis:** Monoclonal band (**M spike**)<sup>Q</sup>
- Immunofixation electrophoresis:** Distinguishes **Ig class**
- Serum/Urine **free light chain assay:** **k/λ chains**
- Bence jones protein in urine**<sup>Q</sup> : Free light chains which precipitates at 55-60°C and disappear on heating to 95°C
- Immunophenotyping:** **CD 38 +ve, CD138 +ve, cIg +ve, CD19-ve**<sup>Q</sup>
- Cytogenetics:**
  - t(11;14)-diagnostic hallmark, good prognosis**<sup>Q</sup>
  - del 13q, t(4;14), t(14;16): Poor prognosis
- Imaging :**
  - X-ray (punched out lytic lesions:**<sup>Q</sup> skull,<sup>Q</sup> spine, ribs, pelvis);
- Serum  $\beta_2$  microglobulin: **<3.5mg/L** indicates **good prognosis**<sup>Q</sup>



Rouleux RBCs



Plasma cells



Mott cells

## High Yield Facts

- M spike : **IgG (most common)**<sup>Q</sup> >A>M>D>E
- Free light chains are called Bence-Jones proteins- **not detected by urine protein dipsticks**<sup>Q</sup>
- IL-6**<sup>Q</sup> helps in **survival and proliferation** of myeloma cell proliferation.
- Other growth factors for myeloma cells-**IL1  $\beta$ <sup>Q</sup> and VEGF<sup>Q</sup>**
- Most common** cytogenetic abnormality in myeloma is 13q->t (11,14)
- Osteolytic lesions**<sup>Q</sup> in bone are due to involvement of **RANK-L**<sup>Q</sup> (receptor activator of nuclear factor kappa B ligand) → **activates osteoclasts**
- POEMS**<sup>Q</sup>: Polyneuropathy, Organomegaly, Endocrinopathy, Multiple Myeloma & Skin Changes
- Alkaline Phosphatase levels in multiple myeloma is normal** and not raised, as there is no bone formation and only bone lysis.

- Mc mutation: **MYD88**

### Clinical features:

- Anemia, lymphadenopathy, Hepatosplenomegaly & hyperviscosity
- Immunophenotyping:**
  - CD 138 +, cy IgM+, CD19 +

## HODGKIN LYMPHOMA (HL)

### Characteristics:

- Arises in lymph nodes (**M.C cervical region**)<sup>Q</sup> & spreads to anatomically **contiguous lymphoid tissues**<sup>Q</sup>

### Clinical features:

- Pel Ebstein fever** (Intermittent fever every alternate week)<sup>Q</sup>
- Lymphadenopathy**; Affected lymph nodes become painful with alcohol ingestion<sup>Q</sup>

### Pathogenesis: Mc mutation: Rel transcription activators

- Activation of the **transcription factor NF- $\kappa$ B** is a common event in **classical HL**.
- Cytokines (e.g., **IL-5, IL-10, M-CSF**), chemokines (e.g., **eotaxin**), & other factors (e.g., **immunomodulatory factor galectin-1**) that are secreted by Reed-Sternberg cells

## Waldenstrom Macroglobulinemia

### Definition:

- Indolent **lymphoproliferative disorder** characterized by Lymphoplasmacytic cell proliferation in marrow with secretion of **IgM**






#### ■ **Morphology:**

- Reed-Sternberg cells (**R.S cells**) surrounded by **T lymphocytes** in a rosette-like manner<sup>Q</sup>
- **Diagnostic Hallmark: Reed-Sternberg cells:** (45 µm) binucleate cell or single nucleus with multiple nuclear lobes.<sup>Q</sup>

#### ■ **Poor prognostic markers:**

- Albumin <4.0 g/dL, Hemoglobin <10.5 g/dL, Male sex, 45 years of age or more, Stage IV disease, Leukocytosis at or above 15,000/mm<sup>3</sup>, Lymphocytopenia



#### **Description of different Subtypes of Hodgkin Lymphoma**

Subtype	Morphology	Immunophenotype	Association with EBV	Typical Clinical Features
<b>Nodular sclerosis</b>	Lacunar cells (clear space around nucleus) <sup>Q</sup> Fibrous strands <sup>Q</sup> & plasma cells <sup>Q</sup>	CD15+, CD30+;	usually EBV-	<b>MC in World<sup>Q</sup>; M=F;</b> usually stage I or II; frequent <b>mediastinal</b> involvement; <b>Good prognosis<sup>Q</sup></b>
<b>Mixed cellularity</b>	Mononuclear cells	CD15+, CD30+;	<b>70% EBV+</b>	<b>MC in India<sup>Q</sup>, stage III or IV;</b> M > F; biphasic incidence; <b>Good prognosis<sup>Q</sup></b>
<b>Lymphocyte rich</b>	Mononuclear cells	CD15+, CD30+;	<b>40% EBV+</b>	Uncommon; M > F, <b>Good prognosis<sup>Q</sup></b>
<b>Lymphocyte depletion</b>	Reticular variant:	CD15+, CD30+;	<b>90% EBV+ (Maximum)<sup>Q</sup></b>	Uncommon; M>F; <b>HIV infected<sup>Q</sup></b> <b>Poorest Prognosis<sup>Q</sup></b>
<b>Lymphocyte predominance</b>	Lymphocytic & Histiocytic (popcorn cell) <sup>Q</sup>	<b>CD20+,</b> CD15-, C30-;	<b>EBV-</b>	Uncommon; young males with cervical or axillary L. nodes, <b>Best Prognosis<sup>Q</sup></b>

#### **Clinical Staging of Hodgkin's and Non-Hodgkin's Lymphomas (Ann Arbor Classification)**

Stage	Distribution of Disease
I	Involvement of a <b>single lymph node region</b> (I) or a single extra-lymphatic organ or site (IE).
II	Involvement of <b>two or more lymph node</b> regions on the <b>same side of diaphragm</b> alone (II) or localized involvement of an extra-lymphatic organ or site (IIE).
III	Involvement of lymph node regions on <b>both sides of the diaphragm</b> without (III) or with (IIIE) localized involvement of an extra-lymphatic organ or site.
IV	Diffuse involvement of <b>one or more extra-lymphatic organs</b> or sites with or without lymphatic involvement.

All stages are further divided on the basis of: **Absence (A)** or **Presence of (B)** symptoms:<sup>Q</sup> Unexplained fever, Drenching night sweats, and/or, Unexplained weight loss > 10%

#### **Treatment**

ABVD (Adriamycin, Bleomycine, Vineblastine & Dacarbazine) regimen is standard line of treatment<sup>Q</sup>






## MYELOID NEOPLASMS

Three broad categories of myeloid neoplasia exist:

- Acute myeloid leukemias (AML): discussed previously
- Myelodysplastic syndromes (MDS)
- Chronic Myeloproliferative Neoplasms (CMPN)

### Myelodysplastic Syndromes


- Definition:**
    - Group of **clonal stem cell disorders** characterized by **cytopenias**, **dysplasias** in either lineage, **ineffective erythropoiesis** and a **high risk of transformation to AML**.<sup>Q</sup>
  - Cytogenetics**
    - del 5q (MC best prognosis)**<sup>Q</sup>—Adults
    - Monosomy 5<sup>Q</sup>, **Monosomy 7**, **del 7q (MC treatment related MDS;**<sup>Q</sup> **both have poor prognosis)**<sup>Q</sup>
    - p53 mutation—aggressive disease
    - y, del 11q—very good prognosis
- WHO 2017**



Pseudo-Pelger Huet neutrophil

- Bone marrow Morphology:**
  - Cytopenias with features of dysplasia can be seen in either of the series like:

Erythroid series	Myeloid series	Megakaryocytic series
<ul style="list-style-type: none"> <li>Ring sideroblasts<sup>Q</sup></li> <li>Megaloblastic maturation</li> <li>Nuclear budding</li> <li>Nuclear bridging</li> </ul>	<ul style="list-style-type: none"> <li>Hypo or defective granulation</li> <li>Toxic granulations</li> <li><b>Döhle bodies</b></li> <li><b>Pseudo-Pelger-Huet</b><sup>Q</sup></li> <li><b>neutrophils</b></li> </ul>	<ul style="list-style-type: none"> <li>Micromegakaryocytes</li> <li>Single nuclear lobes</li> <li>Multiple separate nuclei</li> <li><b>(Pawn ball Megakaryocytes)</b><sup>Q</sup></li> </ul>



Pawn Ball megakaryocyte

### Chronic Myeloproliferative Neoplasms (CMPN)

- Definition:**
  - Presence of **mutated tyrosine kinases**<sup>Q</sup> or other **acquired aberrations** in signaling pathways that lead to growth factor independence leading to:
    - Increased proliferation of bone marrow
    - Extra-medullary hematopoiesis
    - Marrow fibrosis** and peripheral blood cytopenias
    - Transformation to acute leukemia**

We will now discuss the types of CMPN:


### Chronic Myelogenous Leukemia (CML)

- Characterized by:**
  - BCR-ABL gene**<sup>Q</sup> (210 kDa in size)
  - ABL gene on chr 9q translocates to BCR gene on chr 22q** which **activates tyrosine kinase**<sup>Q</sup>
- Cell of origin:**
  - Pluripotent hematopoietic stem cell**<sup>Q</sup>
- Etiology:**
  - Radiation exposure, No genetic predisposition
- Epidemiology:**
  - M> F; **Median age-5<sup>th</sup> -6<sup>th</sup> decade**
- Clinical features:**
  - Presents with **massive splenomegaly**, hepatomegaly and lymphadenopathy.<sup>Q</sup>
- Peripheral smear:**
  - Increased TLC (30,000/uL – 10,00,000/uL), **myeloid bulge**<sup>Q</sup> (myelocytes and metamyelocytes) and **basophilia**.<sup>Q</sup>
- LAP score:**
  - Low**<sup>Q</sup>



- **Bone marrow Morphology:**
  - Hypercellular marrow (Not required for diagnosis but **for staging**)<sup>Q</sup>
  - Increased small, **dysplastic forms of megakaryocytes**.
    - Scattered macrophages with abundant wrinkled, green-blue cytoplasm so-called **sea-blue histiocytes/ Pseudo-Gaucher cells**<sup>Q</sup>
- **Diagnosis of choice**
  - Cytogenetics -FISH/PCR

### The Philadelphia Chromosome



### WHO Diagnostic Criteria for Different Phases of CML (WHO 2017)

Accelerated Phase	Blast phase/ blast crisis
<ul style="list-style-type: none"> <li>• <b>Blasts 10–19%</b> in blood or marrow<sup>Q</sup></li> <li>• Peripheral blood <b>basophilia <math>\geq 20\%</math></b><sup>Q</sup></li> <li>• Newer Cytogenetic clonal evolution</li> <li>• Persistent thrombocytopenia (<math>&lt;100 \times 10^9/L</math>)</li> <li>• Persistent thrombocytosis (<math>&gt;1,000 \times 10^9/L</math>) unresponsive to therapy</li> <li>• Increasing splenomegaly &amp; WBC count unresponsive to therapy</li> <li>• Provisional response to TKI-Tyrosine kinase inhibitors</li> <li>• <b>Provisional” response-to-TKI (tyrosine kinase inhibitors) criteria is added- Occurrence of 2 or more mutations or resistance to therapy are added</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b><math>&gt;20\%</math> blasts<sup>Q</sup></b> (*Wintrobe’s latest 13<sup>th</sup> ed: <math>&gt;30\%</math>)</li> <li>• Clusters of blasts on BM biopsy</li> <li>• Extramedullary myeloid tumors (granulocytic sarcomas, <b>chloromas</b>)<sup>Q</sup></li> <li>• Lymphoblasts in any number should be reported as they signify poor prognosis</li> </ul>



### High Yield Facts

Conditions Associated with Abnormal Leukocyte Alkaline Phosphatase (LAP) Scores	
High LAP score ( $>130$ )	Low LAP Score ( $<15$ )
<ul style="list-style-type: none"> <li>• Infections (Leukemoid reaction)<sup>Q</sup></li> <li>• Growth factor therapy</li> <li>• Myeloproliferative disorders other than CML (ET, PCV, Myelofibrosis)<sup>Q</sup></li> <li>• AML<sup>Q</sup></li> <li>• Hodgkin’s disease<sup>Q</sup></li> <li>• Inflammatory disorders</li> <li>• Pregnancy,<sup>Q</sup> oral contraceptives</li> <li>• Stress</li> <li>• Drugs (lithium, corticosteroids, estrogen)</li> </ul>	<ul style="list-style-type: none"> <li>• CML<sup>Q</sup></li> <li>• Paroxysmal nocturnal hemoglobinuria<sup>Q</sup></li> <li>• Hereditary Hypophosphatemic Rickets</li> <li>• Myelodysplastic syndromes</li> <li>• Rare infections or toxic exposures</li> </ul>






## Polycythemia Vera

- **Epidemiology:**
  - Mean age = 60yrs; M:F=1-2:1
- **Genetic abnormality:**
  - Most frequent genetic abnormality in PCV is **JAK2 V617F**<sup>Q</sup>
- **Clinical Features:**
  - **CVS:** **Hypertension**<sup>Q</sup>, venous or arterial **thrombosis**, myocardial ischemia or stroke, pruritus after bath
  - **CNS:** Headache, dizziness, visual disturbances, **paraesthesias**
  - **Others:** Pruritus, **erythromelalgia**,<sup>Q</sup> **gout**<sup>Q</sup>

- **Diagnosis:** *Requires all 3 major criteria or first 2 major + minor criteria*

Major Criteria	Minor Criteria
<ul style="list-style-type: none"> <li>• Hemoglobin &gt;16.5 g/dl<sup>Q</sup> in men, &gt;16 g/dL in women or HCT &gt;49% in men or &gt;48% in woman. Increased red cell mass &gt;25% above mean normal predicted value.</li> <li>• Presence of <b>JAK2 mutation</b><sup>Q</sup></li> <li>• Hypercellular bone marrow biopsy with <b>panmyelosis</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Low serum erythropoietin level<sup>Q</sup></li> </ul>







## Essential Thrombocythemia/Primary Thrombocytosis

- **Epidemiology**
  - Mean age = 50-60 yrs; M=F
- **Clinical Feature**
  - Microvascular occlusion may lead to **transient ischemic attacks**<sup>Q</sup>, digital ischemia with **paraesthesia** & gangrene. **Bleeding**<sup>Q</sup> may also be seen, due to platelet function defects
- **Peripheral smear**
  - **Thrombocytosis** with **abnormalities in size, shape & granularity** of platelets
- **Diagnosis:**
  - All 4 or first 3 major + 1 minor criteria

Major criteria	Minor criteria
<ul style="list-style-type: none"> <li>• Sustained platelet count <math>\geq 4.5</math> Lakhs/<math>\mu\text{L}</math><sup>Q</sup></li> <li>• Bone marrow biopsy showing proliferation of megakaryocytes,</li> <li>• <b>Exclusion</b> of WHO criteria for PV, PMF, CML, MDS</li> <li>• <b>JAK2 mutation</b><sup>Q</sup>, CALR or MPL mutations</li> </ul>	<ul style="list-style-type: none"> <li>• Absence of evidence of reactive thrombocytosis</li> </ul>

## Chronic Idiopathic Myelofibrosis/Agnogenic Myeloid Metaplasia (AMM)

- **Epidemiology:**
  - Mean age: 6<sup>th</sup>-7<sup>th</sup> decade; M=F
- **Hallmark:**
  - Hallmark of primary myelofibrosis is: **obliterative marrow fibrosis**<sup>Q</sup>
- **Peripheral smear:**
  - Marrow distortion due to fibrosis leads to the premature release of nucleated erythroid and early granulocyte progenitors (**leukoerythroblastosis**)<sup>Q</sup>
  - Erythroids damaged in fibrotic marrow results in :Tear drop-shaped RBCs (**dacrocytes**)<sup>Q</sup>
- **Diagnosis requires: All 3 Major + at least 1 minor criteria**

Major Criteria	Minor Criteria
<ul style="list-style-type: none"> <li>• Atypical megakaryocytic hyperplasia, with <b>collagen fibrosis</b><sup>Q</sup></li> <li>• Exclusion of WHO criteria for PV, CML, MDS, or other MPDs</li> <li>• <b>JAK2V617F mutation</b><sup>Q</sup>, CALR, MPL</li> </ul>	<ul style="list-style-type: none"> <li>• Leukoerythroblastosis<sup>Q</sup></li> <li>• <math>\uparrow</math> LDH</li> <li>• Anemia</li> <li>• Palpable splenomegaly</li> <li>• Leucocytosis <math>&gt;11,000</math></li> </ul>

### High Yield Facts

- **Ph chr** discovered by Nowell & Hungerford in 1960
- **Ring sideroblasts** are erythroblasts with **iron-laden mitochondria**<sup>Q</sup> visible as **perinuclear granules** in **Iron/Prussian blue/Perl's**<sup>Q</sup> staining
- **Pseudo-Pelger-Huet neutrophils**<sup>Q</sup>: **bilobed hypogranular dysplastic neutrophils**<sup>Q</sup>

## MYELOYDYSPLASTIC/ MYELOPROLIFERATIVE (MDS/MPN) NEOPLASMS

- **Definition:**
  - Group of disorders with features of **both myeloproliferative and myelodysplastic syndromes**<sup>Q</sup>
- **Includes:**
  - Chronic myelomonocytic leukemia (CMML)
  - Atypical CML (a-CML)
  - Juvenile myelomonocytic leukemia (JMML)

### Other

Newer Myeloproliferative Neoplasms (MPNs)	
MPN	Mutation seen
• Systemic mastocytosis	• Constitutive <b>c-KIT</b> kinase activation
• Chronic eosinophilic leukemia	• Constitutive <b>PDGFR<math>\alpha</math>/<math>\beta</math></b> kinase activation
• Stem cell leukemia	• Constitutive <b>FGFR1</b> kinase activation

## JUVENILE MYELOMONOCYTIC LEUKEMIA (JMML)

- **Definition**
  - It is a **childhood mixed MDS/MPD** that includes childhood leukemias previously classified as CMML, juvenile CML, and infantile monosomy 7 syndrome.
- **Diagnostic Criteria:**

Genetic Criteria (Any 1 is sufficient)	Required Criteria (All 4 needed)	Other criteria
<ul style="list-style-type: none"> <li>• NFI mutation</li> <li>• CBL mutation (germ line)</li> <li>• Somatic mutation of KRAS/NRAS/PTPN11</li> </ul>	<ul style="list-style-type: none"> <li>• Peripheral blood monocytes <math>&gt; 1.0 \times 10^9/\text{L}</math></li> <li>• Blasts + promonocytes <math>&lt;20\%</math> in blood and marrow</li> <li>• Absence of Philadelphia chromosome or BCR/ABL fusion gene</li> <li>• Splenomegaly</li> </ul>	<ul style="list-style-type: none"> <li>• Increased hemoglobin F for age</li> <li>• Immature granulocytes in peripheral blood</li> <li>• Clonal chromosomal abnormality (i.e., includes monosomy 7)</li> <li>• GM-CSF hypersensitivity of myeloid progenitors in vitro</li> <li>• Hypophosphorylation of STAT 5 or Monosomy 7</li> </ul>

### High Yield Facts


- **JMML is the most common MDS/MPD of children**
- Increased hemoglobin F is seen in JMML
- JMML is associated with NF1





## LANGERHANS CELL HISTIOCYTOSIS (LCH) HISTIOCYTOSIS X<sup>Q</sup>

- **Current classification:**
  - **Unifocal** (i.e., single-system, single-site disease)
  - **Multifocal** (i.e., single-system, multiple-site disease)
  - **Disseminated** histiocytosis (i.e., multisystem disease)
- **Diagnosis: Light microscopy**
  - Basic histologic lesion: **granulomas**, containing histiocytes or Langerhans cells, mature eosinophils, lymphocytes, giant cells, neutrophils & plasma cells
  - The **Langerhans cell**<sup>Q</sup> (large mononuclear cells with few cytoplasmic vacuoles) is the '**sine qua non**' (**essential**) of the diagnostic lesion
- **Electron microscopy**
  - **Birbeck granules (tennis racket appearance)**<sup>Q</sup>
- **Immunohistochemistry**
  - **CD1a<sup>Q</sup>, S-100<sup>Q</sup> or Langerin (CD 207)<sup>Q</sup>** demonstration on the surface of LCH cells
- **Prognosis**
  - LCH may be a **self-limiting** disease, which may resolve spontaneously<sup>Q</sup>
  - For most patients with LCH, the prognosis is **excellent**
  - Patients with **multisystem** disease may experience fatal **organ failure**



Langerhans cells with convoluted nuclei with longitudinal grooves ("coffee-bean" shaped)



### High Yield Facts

- **Bone** is the **most commonly involved organ**<sup>Q</sup>
- **Skull** is the most commonly involved **site in both children and adults**<sup>Q</sup>
- Unifocal eosinophilic granuloma of bone is the most common form of the disease<sup>Q</sup>
- "**Punched out**" appearance of skeletal lesions is typically seen in LCH<sup>Q</sup>
- **Pulmonary LCH** occurs more commonly in **males** & is associated with **smoking**
- **Pneumothorax** is seen in **25–40%** cases of **pulmonary LCH**
- **Eosinophilic granuloma**:<sup>Q</sup> bone lesions, with no visceral involvement
- **Letterer-Siwe disease**:<sup>Q</sup> When granulomas involve multiple viscera
- **Hand-Schüller-Christian disease**<sup>Q</sup>: Triad of multiple bone lesions, exophthalmos and diabetes insipidus (DI)

### Thymoma

- **Definition:**
  - Tumors of **thymic epithelial cells**
- **Epidemiology:**
  - Usually seen in adults **older than 40 years** of age; rare in children; Males = females
- **Location:**
  - **Anterior superior mediastinum**, neck, thyroid
- **Gross Morphology:**
  - **Lobulated**, firm, gray-white masses of **up to 15 to 20 cm** in size.
  - Sometimes have areas of cystic **necrosis and calcification**.
  - Most are **encapsulated**, but 25% of the tumors penetrate the capsule & **infiltrate perithymic structures**
- **Histology:**
  - **Sheets of epithelial cells** giving **arborizing pattern** of reactivity along with interspersed **lymphoid cells**.
  - **IHC** –CK + CD45



### High Yield Facts

#### Post Transplant Lymphoproliferative Disorder (PTLD)

- Post-transplant lymphoma occurs due to **proliferation of B cells**
- 90% of **early** (<1 year post-transplant) PTLDs are **EBV positive**, when EBV-CTL immunity is lowest
- **Late** (>2 years post-transplant) PTLDs are **frequently EBV negative**, can be of T-cell origin, and may have a poorer prognosis.
- Majority of PTLDs are **CD20+**, but not all PTLDs are of B-cell phenotype and not all are EBV positive.
- **T-cell PTLD tends to occur late**, often more than 10 years after transplantation.



## 2018 Revision to the World Health Organization Classification of Leukemia & Lymphoma

Category	Latest modification	Category	Latest modification
<b>New Acute Myeloid Leukemia Subtypes 2016</b>	AML with RUNX1 mutation AML with BCR-ABL 1 mutation AML with biallelic CEBPA mutations Familial AML/MDS multiple types- EBPA, RUNX, GATA	<b>ALL</b>	Early precursor T-ALL –CD7, CD2, CD3 , Myeloid markers+
<b>Myeloid neoplasms</b>	Refer to CML in text	<b>Essential thrombocythemia and Primary myelofibrosis</b>	CALR and MPL mutation is needed in addition to JAK-2 mutation
<b>CML</b>		<b>Polycythemia Vera</b>	Hb cut off reduced to 16.5 gm% in males and 16 gm% in females or Hematocrit >49% (m) and 48% (f)
<b>CNL (chronic neutrophilic leukemia)</b>	CSF3R mutations added	<b>Systemic mastocytosis</b>	Removed from Chronic myeloproliferative neoplasms
<b>MDS</b>	Del9q is an MDS related entity only in the absence of NPM1 mutations SF3B1 mutation is strongly associated with ringed sideroblast		

## WHO 2018 Update

### The Cancer Genome Atlas (TCGA) project showing 9 classes of AML

**Class 1:** Transcription factor fusions  
e.g. t(8;21), inv(16), and t(15;17)

**Class 2:** Nucleophosmin 1  
NPM1 mutations

**Class 3:** Tumor suppressor genes  
e.g. TP53 and PHF6 mutations

**Class 4:** DNA methylation-related genes  
DNA hydroxymethylation e.g. TET2 , IDH1 and IDH2  
DNA methyltransferases


**Class 5:** Activated signalling genes  
e.g. FLT3, KIT RAS mutation..

**Class 6:** Chromatin-modifying, genes  
e.g. ASXL1 and EZH2 mutations., fusions, KMT2A-PTD

**Class 7:** Myeloid transcription factor genes  
e.g. CEBPA, RUNX1 mutations


**Class 8:** Cohesin complex genes  
e.g. STAG2, RAD21, SMC1, SMC2 mutations

**Class 9:** Spliceosome-complex genes  
e.g. SRSF2, U2AF1, ZRSR2 mutations




## Image-Based Questions

1. A 4-year-old male presents with fever, bleeding gums and fatigue for 4 days. CBC shows Hb of 8 gm%, TLC 86,000/UL, Platelet count of 25,000/ul. DLC shows Neutrophils 20%, Lymphocyte 40%, Eosinophils 10%, Basophils 0%, Monocytes 5%, Abnormal cells 25%. Bone marrow aspiration shows cells as shown in figure 60%. What is your diagnosis?




- Chronic myeloid leukemia
- Chronic lymphoid leukemia
- Acute leukemia
- Myelofibrosis

3. Peripheral smear showing the given figure is likely to be seen in all except:



- Hypergammaglobulinemia
- Severe anemia
- Multiple myeloma
- Hemolytic anemia

2. A 10-year-old boy presents to AIIMS OPD with mass in the abdomen. On imaging the paraaortic LN is enlarged. Biopsy from the lymph node suggests a pattern as shown in the figure. What is the underlying abnormality?




- p53 gene mutation
- RB gene mutation
- Translocation involving BCR-ABL genes
- Translocation involving MYC gene

4. For which procedure it is used?




- Bone marrow examination
- Liver biopsy
- Pleural biopsy
- Lumbar puncture






5. 40/F presented to Medicine OPD with fever and mucosal bleeding of 3 days. CBC shows Hb-9.8 gm%, TLC = 15,700/cumm, Platelet count = 15,000/cumm. Peripheral smear showed findings as shown in figure. His cytogenetics revealed t(8;21). What is your diagnosis?




- a. AML  
b. CML  
c. MDS  
d. ALL

7. 55/M presented with fatigue and dragging sensation in the abdomen to AIIMS OPD. He send the patients sample to a pathologist initially performed hemogram which revealed Hb = 8 gm%, TLC = 1500/cumm, platelet count = 79,000/cumm. He also reported some bizzare looking cells which are shown below. Which stain will the pathologist like to do to diagnose the condition?




- a. PAS  
b. NSE  
c. MPO  
d. TRAP

6. The most important investigation in the given case to diagnose if the condition is a neoplasm?




- a. JAK-2  
b. EPO level  
c. PaO<sub>2</sub>  
d. Bone marrow aspiration and biopsy

8. A 10-year-chid with bilateral cervical lymphadenopathy. Lymph node biopsy was performed, which showed cells as given in the figure. Which of the following is true regarding this condition?



- a. Hodgkin lymphoma: EBV and embryo cell  
b. Non-Hodgkin lymphoma HIV and Giant B cell  
c. TB, Mycobacteria and tiny granuloma  
d. Hodgkin lymphoma: EBV and Reed Sternberg cell





## Answers of Image-Based Questions

1. Ans. (c) **Acute leukemia**

- The image shows large cells with high N:C ratio, immature chromatin which are blasts. With >20% blasts, the diagnosis is acute leukemia.

2. Ans. (d) **Translocation involving MYC gene**

- The arrow marked shows cleared area (stars as macrophages) amidst hugely proliferating tumor cells (sky). This appearance of starry sky is seen in Burkitts lymphoma having MYC gene translocation.

3. Ans. (d) **Hemolytic anemia**

- The smear shows rouleux formation seen in multiple myeloma, severe anemia & cases of hypergammaglobinemia.

4. Ans. (a) **Bone marrow examination**

- This is Klima bone marrow aspiration needle for marrow aspiration and biopsy

5. Ans. (a) **AML**

- The blasts cells shows Auer rods inside them which are a hallmark of AML.

6. Ans. (a) **JAK-2**

- The figure shows increased platelets in smear and increased megakaryocytes in bone marrow. To diagnose this as essential thrombocythemia (neoplasm); JAK-2 mutation analysis should be done.

7. Ans. (d) **TRAP**

- The peripheral smear in the question shows hairy cells which can be diagnosed with TRAP stain.

8. Ans. (d) **Hodgkin lymphoma: EBV and Reed Sternberg cell**


- Figure shows Reed Sternberg cells in Hodgkins lymphoma which are EBV infected.



## Multiple Choice Questions

### NON-NEOPLASTIC WBC DISORDERS


1. A patient presented with intermittent fever, no wt loss, no anorexia, retroperitoneal mass. Peripheral smear finding were normal. Gross and microscopy of the mass is given. What is the most likely diagnosis?



- a. Non-Hodgkin's lymphoma (AIIMS Nov 18)  
b. Castleman disease  
c. Angiolymphoid hyperplasia

2. HHV-8 is related to all except: (AIIMS May 18)  
a. Primary effusion lymphoma  
b. Kaposi sarcoma  
c. Castleman disease  
d. Adult T cell lymphoma


3. Which of the following represents the marked area in the histology of lymph node? (AIIMS May 2017)



- a. Mantle zone  
b. Marginal zone  
c. Paracortical area  
d. Germinal centre


4. Dohle bodies with giant platelets are seen in: (JIPMER 2017)  
a. May heggalin anomaly  
b. Pelger huet anomaly  
c. Chediak higashi syndrome  
d. Bernard solier syndrome

5. Which of the following cells will increase in case of parasite infection? (AIIMS Nov 2016)



- a. A  
b. B  
c. C  
d. D

6. Identify the arrow marked cell in the given condition below? (AIIMS May 2016)



- a. Macrophage  
b. Lymphocyte  
c. Plasma cell  
d. Eosinophil

7. About the given instrument below all of the following statements are true except? (AIIMS May 2016)



- a. Done for diagnosis of infiltrative and granulomatous diseases  
b. No need of breath holding during the procedure  
c. Can be done in prone or lateral position  
d. Platelet count of <40000/ul is contraindication

8. A Warthin-Finkeldey cell is a type of giant multinucleate cell found in hyperplastic lymph nodes early in the course of: (Recent Question 2016-17)

- a. Measles  
b. Hodgkins  
c. Kala azar  
d. Syphilis

9. For which procedure is the following instrument used?



- a. Bone marrow examination (AIIMS Nov 2015)  
b. Liver biopsy  
c. Pleural biopsy  
d. Lumbar puncture

10. Marker of T-lymphocyte is: (Recent Question 2016)

- a. CD8  
b. CD20  
c. CD19  
d. CD45




11. **Host receptor for streptococcus pyogenes is?** (Recent Question 2015)
  - a. CD4
  - b. CD21
  - c. CD44
  - d. CD46
12. **Serious infections can occur when absolute neutrophil count decreases below?** (Recent Question 2016)
  - a. Less than 500/ul
  - b. Less than 800/ul
  - c. Less than 1000/u
  - d. less than 2000/ul
13. **Eosinophilia is found in?** (Recent Question 2016)
  - a. Cryptococcus
  - b. HPV
  - c. Strongyloides
  - d. Typhoid
14. **1st cell of RBC development** (Recent Question 2016)
  - a. Pro erythroblast
  - b. Intermediatenormoblast
  - c. Reticulocyte
  - d. Basophilic erythroblast
15. **Dilated endoplasmic reticulum is called as?** (Recent Question 2015)
  - a. Asteroid bodies
  - b. Bamboo bodies
  - c. Hirano bodies
  - d. Dohle bodies
16. **In infant, Bone Marrow biopsy is done from?** (Recent Question 2015)
  - a. Tibia
  - b. Sternum
  - c. Posterior superior Iliac Spine
  - d. Iliac crest
17. **In a case of anemia with thrombocytopenia and PMN showing inclusions. What is the most probable diagnosis?** (Recent Question 2015)
  - a. May Hegglin anomaly
  - b. Evan syndrome
  - c. Alder-Reilly anomaly
  - d. Pegler Huet Anomaly
18. **Which of the following is a B cell marker?** (PGI May 2014)
  - a. CD1
  - b. CD 10
  - c. CD1a
  - d. CD19
  - e. CD20
19. **Pan B cell marker:** (WB PG 2014)
  - a. CD 19p
  - b. CD 19q
  - c. CD 16
  - d. CD 21
20. **Infectious mononucleosis affects?** (JIPMER 2014)
  - a. B-cells
  - b. T-cells
  - c. NK cells
  - d. Macrophages
21. **The peripheral blood eosinophil count in Eosinophilia-myalgia syndrome is usually** (Bihar PG 2014)
  - a. Between 500 to 2000 cells/ microiliter
  - b. 2000 to 5000 cells/microliter
  - c. Less than 500 cells/microliter
  - d. More than 5000 cells/microliter
22. **In an ablated animal, myeloid series cells are injected. Which of following is seen after incubation period -** (AIIMS May 12)
  - a. RBC
  - b. Fibroblast
  - c. T lymphocytes
  - d. Hematopoietic stem cell
23. **All of the following stem cell populations are found within the bone marrow, except -** (AI 12)
  - a. Endothelial Progenitor cells
  - b. Myoblast Progenitor cells
  - c. Mesenchymal stem cells
  - d. Hematopoietic stem cells
24. **Leukocyte common antigen is:** (WB PG 2012)
  - a. CD 45
  - b. CD 20
  - c. CD 19
  - d. CD 41

25. **Generalized necrotising lymphadenopathy is -** (AI 11)
  - a. Kimura disease
  - b. Kikuchi disease
  - c. Non-Hodgkin's lymphoma
  - d. Castleman's disease
26. **Eosinophilic abscess in lymph node is characteristically seen in -** (DPG 11)
  - a. Kimura's disease
  - b. Hodgkin's lymphoma
  - c. Tuberculosis
  - d. Sarcoidosis
27. **Which of the following surface glycoproteins is most often expressed in human hematopoietic stem cell?** (DPG 10)
  - a. CD22
  - b. CD45
  - c. CD15
  - d. CD34


#### ACUTE LEUKEMIA

28. **Which of the following is not a provisional entity according to Revised WHO-2016 classification of Acute leukemia?** (AIIMS May 18)
  - a. AML with BCR-ABL
  - b. ALL with hyperploidy
  - c. AML with RUNX<sub>1</sub>
  - d. Early T cell precursor leukemia
29. **How will you differentiate mediastenal mass from thymoma differentiating it from ALL?** (AIIMS Nov 18)
  - a. Cytokeratin
  - b. CD1a
  - c. CD3
  - d. Tdt
30. **A patient presented with painless b/l proptosis. What is the next investigation to diagnose it as chloroma?** (AIIMS Nov 2017)
  - a. Blood haemoglobin
  - b. Peripheral smear
  - c. Platelets
  - d. Bone marrow (reticulin)
31. **All of the following are seen in the development of T lymphocyte at a point except?** (AIIMS May 2017)
  - a. Tdt
  - b. CD34
  - c. PAX5
  - d. Cd1a
32. **A 7 year old presents with fever, weight loss. On examination he was pale and had significant lymphadenopathy. Bone marrow histology is as given below. What is the most probable diagnosis?** (AIIMS May 2017)




- a. ALL
- b. AML
- c. Aplastic anaemia
- d. Juvenile myelomonocytic leukemia






- 33. Good prognosis for ALL is?** (JIPMER 2017)  
 a. Age > 10 yrs                      b. Hyperdiploidy  
 c. T cell variant                      d. TLC >1,00,000/uL
- 34. Best candidate for Autologous bone marrow transplant?** (JIPMER 2017)  
 a. Multiple myeloma  
 b. Leukemia  
 c. Thalassemia  
 d. Congenital Immunodeficiency
- 35. Most common translocation in acute promyelocytic leukemia?** (PGI May 2017)  
 a. t(8;14)                                  b. t(15;17)  
 c. t(9;22)                                  d. t(8;11)  
 e. t(11;14)
- 36. Which one is best prognostic factor for ALL?** (AIIMS Nov 2015)  
 a. Hyperploidy  
 b. Orgnomegaly  
 c. TLC more than 50,000/uL  
 d. Response to treatment
- 37. Auer rods are specific for** (Recent Question 2015)  
 a. Acute myeloid leukemia  
 b. Acute lymphocytic leukemia  
 c. Chronic lymphocytic leukemia  
 d. Hodgkin's lymphoma
- 38. True about ALL** (Recent Question 2015)  
 a. tdT positive                              b. Gamma globulins  
 c. t(8,14)                                      d. Insidious onset
- 39. Dohle bodies are seen in** (Recent Question 2015)  
 a. Neutrophils                              b. Macrophages  
 c. Plasma cells                              d. Histiocytes
- 40. All are true regarding hemophagocytic lymphohistiocytosis (HLH) except?** (Recent Question 2015)  
 a. Activation of macrophages and CD 8+ T cells  
 b. Cytopenias due to phagocytosis of progenitors in bone marrow  
 c. HTLV-1 is a cause in immunodeficient patients  
 d. Abnormal liver function tests
- 41. Stain used for diagnosis of granulocytic sarcoma** (Recent Question 2015)  
 a. Myeloperoxidase  
 b. Leukocyte alkaline phosphatase  
 c. Nonspecific esterase  
 d. Neuron specific enolase
- 42. Most common type of AML in Down's syndrome** (Recent Question 2015)  
 a. M2    b. M3  
 c. M6    d. M7
- 43. The following parameter in ALL indicates poor prognosis** (Recent Question 2015)  
 a. Age >10 years  
 b. WBC count <50000/mm<sup>3</sup> at diagnosis  
 c. Hyperdiploidy  
 d. Early pre-B phenotype
- 44. Drug that is not used in the treatment of ALL** (Recent Question 2015)  
 a. Rituximab                                  b. Methotrexate  
 c. Vincristine                                  d. Daunorubicin
- 45. Most common type of all in children** (Recent Question 2015)  
 a. Pre-B cell ALL                              b. Mature B cell ALL  
 c. Pre-T cell ALL                              d. Mature T cell ALL
- 46. Chloroma is a** (Recent Question 2015)  
 a. Lymphoma                                  b. Leukemia  
 c. Sarcoma                                      d. Carcinoma
- 47. A 35 year old male presented with complaints of bleeding gums. There is history of recurrent infections in the past 1 year. On examination, pallor present. Peripheral smear of the same patient is shown below. Identify the arrow marked structure?** (Recent Question 2015)  
 a. Dohle body                                  b. Normoblast  
 c. Auer Rod                                      d. Heinz bodies



- 48. In Acute lymphoblastic leukaemia favourable prognostic factors includes all except?** (MH PG 2014)  
 a. Age of 2 to 10 yrs                      b. Low white cell count  
 c. Presence of t (12; 21)                      d. Presence of t (9; 22)
- 49. Which one of the following is the likely diagnosis based on the smear given above:** (AP PGME 2015)  
 a. Acute myelogenous leukemia  
 b. Acute lymphoblastic leukemia  
 c. Hairy cell leukemia  
 d. Chronic lymphocytic leukemia



- 50. Good prognosis of AML are all except?** (Recent Question 2015)  
 a. t(15;17)                                      b. t(8;21)  
 c. inv 16    d. t(12;21)
- 51. Most common malignancy of blood is?** (Recent Question 2016)  
 a. ALL    b. CLL  
 c. AML    d. CML
- 52. AML causing Gum hypertrophy-** (Recent Question 2014)  
 a. M1    b. M2  
 c. M3    d. M4
- 53. DIC is common in which AML-** (Recent Question 2014)  
 a. Monocytic (M5)                              b. Promyelocytic (M3)  
 c. Erythrocytic (M6)                              d. Megakaryocytic (M7)






54. 40/F presented to AIIMS OPD with the following findings—Hb-9.8gm%, TLC= 15,700/cumm, Platelet count= 3 lac/cumm. Peripheral smear showed increased neutrophils with 14 % blasts, 15% myelocytes and metamyelocytes with some dysplasia. Cytogenetic study revealed t(8;21). What is your diagnosis? (AIIMS May 2014)
- AML
  - CML
  - MDS
  - ALL
55. Good prognostic factor of ALL are all except? (JIPMER 2013)
- Hyperdiploidy
  - Female
  - Pre B ALL
  - t(12;21)
56. Which of the following statements is true- (AIIMS May 12)
- Peak incidence of Chronic myeloid leukemia is in the fifth to sixth decades of life
  - Hairy cell leukemia in more than 50 years has a good prognosis
  - Acute lymphoid leukemia in less than 1 year has good prognosis
  - Chronic lymphocytic leukemia occurs in less than 50 years of age
57. A 10 year old child presents with pallor & history of blood transfusion 2 months back. On investigation, Hb -4.5gm/dl, total count 60,000/cu mm and blasts. platelet count- 2 lacs/cumm and CD 10(+)ve, CD 19 (+) ve, CD 117 (+) ve, MPO (+) ve& CD 33(-)ve. What is the most likely diagnosis? (AIIMS Nov 11)
- ALL
  - AML
  - Undifferentiated leukemia
  - Mixed phenotypic acute leukemia
58. Least likely to be Pre-leukemic condition is:
- Paroxysmal nocturnal hemoglobinuria(AIIMS Nov 11)
  - Paroxysmal cold hemoglobinuria
  - Aplastic anemia
  - Myelodysplastic syndrome
59. Most Common extranodal site of Lymphoma in HIV is? (DNB Dec 10)
- CNS
  - GIT
  - Retroperitoneum
  - Mediastinum
60. Poor prognostic indicator in ALL- (DNB June 10)
- Age < 2 year
  - TLC 4,000-10,000/mm3
  - Presence of testicular involvement at presentation
  - Presence of blasts in peripheral smear

#### HODGKIN'S LYMPHOMA

61. Lymphohistocytic variant of Reed Sternberg seen in which subtype of Hodgkin lymphoma? (Recent exam 2018)
- Nodular sclerosis
  - Lymphocyte rich
  - Lymphocyte predominant
  - Lymphocyte depleted
62. A 20 year old male presented with cervical lymphadenopathy. Histology of lymph node shows RS cell with vague nodule formation and background T reactive lymphocytes. The cells were positive for CD20, LCA, EMA and negative for CD 15 and CD30. Diagnosis is? (AIIMS May 2017)
- NLPHL
  - T cell rich B cell lymphoma
  - Nodular sclerosis Hodgkin
  - CLL
63. True about lymphoma? (PGI Nov 2016)
- Mantle cell origin is in germinal centre
  - DLBCL is most common in India
  - CD5/3/8 are markers of B cell lymphoma
  - Burkitts lymphoma arises from Germinal centre
64. True about CML is? (PGI Nov 2016)
- If Imatinib not working then Dasatinib can be used
  - BCR-ABL activates tyrosine kinase
  - BM biopsy is essential for diagnosis
  - Blast crisis has > 10% blasts
65. EBV is not associated with? (AIIMS May 2016)
- Lymphocyte predominant HD
  - Plasmablastic lymphoma
  - Nasopharyngeal Ca
  - Mixed cellularity HD
66. True about Hodgkin's lymphoma: (PGI May 2016)
- Often localized to single axial group of lymph node
  - Hepatomegaly is always present
  - Contiguous spread of lymph node
  - Can be cured by chemotherapy & radiotherapy
  - Commonly presents with painless lymphadenopathy
67. CD 30 is/are marker for: (Recent Question 2016-17)
- Anaplastic large cell lymphoma
  - Embryonal cell carcinoma
  - Squamous Cell Carcinoma
  - Seminoma
  - Hodgkin's lymphoma
68. Bimodal distribution is seen in? (Recent Question 2016-17)
- Hodgkins lymphoma
  - DLBCL
  - ALL
  - CML
69. RS cell having same immunophenotyping are present in which subtypes of Hodgkin's lymphoma: (Recent Question 2016-17)
- Nodular sclerosis
  - Lymphocyte predominant
  - Lymphocyte rich
  - Mixed cellularity
  - Lymphocyte depletion
70. In which following subtypes of Hodgkin lymphoma, the diagnostic R-S giant cells are usually negative for CD 15 and CD 30? (AP PGME 2013)
- Mixed cellularity
  - Lymphocyte rich
  - Lymphocyte depletion
  - Lymphocyte predominance



71. A 10 year child presented with bilateral cervical lymphadenopathy. Lymph node biopsy was performed, which showed cells as given in the figure. Which of the following is true regarding this condition? (AIIMS Nov 2015)
- Hodgkin lymphoma; EBV and embryo cell
  - Non Hodgkin lymphoma; HIV and Giant B cell
  - TB, Mycobacteria and tiny granuloma
  - Hodgkin lymphoma: EBV and Reed Sternberg cell



72. All are true regarding Reed Sternburg cell immunophenotype in classical Hodgkin lymphoma (Recent Question 2015)
- Positive for CD15 and CD30
  - Negative for other B-cell markers, T-cell markers, and CD45
  - Positive for PAX 5
  - Overexpression of BCL-6
73. Hodgkin lymphoma type that more commonly presents as fever of unknown origin (Recent Question 2015)
- Nodular sclerosis
  - Mixed cellularity
  - Lymphocyte predominance
  - Lymphocyte depletion
74. Hodgkins lymphoma type not associated with EBV (Recent Question 2015)
- Nodular sclerosis
  - Lymphocyte rich
  - Lymphocyte depleted
  - Mixed cellularity
75. Sea blue histiocytes are seen in (Recent Question 2015)
- Chronic lymphoblastic leukemia
  - Chronic myeloid leukemia
  - Langerhan cell histiocytosis
  - Burkitt lymphoma
76. Which of the following types of Hodgkin's lymphomas is not associated with Epstein Barr virus? (Recent Question 2015)
- Lymphocyte depletion
  - Mixed cellularity
  - Lymphocyte rich
  - Lymphocyte predominance
77. A 35 year old female presents with cervical and axillary lymphadenopathy. There is history of fever and drenching night sweats. She is diagnosed to have hodgkin's lymphoma. What is the stage of the disease? (Recent Question 2015)
- II-A
  - II-B
  - IIE-A
  - IIE-B
78. True about Hodgkin's lymphoma is/are? (PGI Nov 2015)
- Axial lymphadenopathy
  - hepatomegaly is common
  - Contiguous spread of lymph node
  - Can be cured by chemotherapy
  - An arbor classification is useful


79. Reticular variant of Reed Sternberg found in which subtype of hodgkin's? (Recent Question 2015)
- Lymphocyte rich Hodgkins lymphoma
  - Lymphocyte poor Hodgkins lymphoma
  - Lymphocyte predominant Hodgkins lymphoma
  - Nodular Sclerosis
80. Popcorn cell is seen with which Hodgkins lymphoma? (Recent Question 2015)
- Lymphocyte rich Hodgkins lymphoma
  - Lymphocyte poor Hodgkins lymphoma
  - Lymphocyte predominant Hodgkins lymphoma
  - Nodular Sclerosis
81. CD 15+ / CD30+ lymphoma among the following are? (PGI May 2014)
- Mixed cellularity Hodgkin lymphoma
  - Mantle cell lymphoma
  - Diffuse T- cell lymphoma
  - NLPHL
  - Acute lymphoblastic leukemia
82. Choose the FALSE statement regarding Hodgkin's lymphoma (APPGMEE 14)
- Affected lymph nodes become painful with alcohol ingestion
  - Ann Arbor Stage II is involvement of two or more lymph node groups on both sides of the diaphragm
  - 'B symptoms' are fever, night sweats and  $\geq 10\%$  weight loss in 6 months
  - ABVD regimen is standard line of treatment
83. A person is having painless lymphadenopathy. On biopsy, binucleated owl shaped nuclei with clear vacuolated area is seen. On IHC CD 15 and CD 30 were positive. What is the most probable diagnosis? (AIIMS Nov 2013)
- Nodular sclerosis
  - Large granular lymphocytic lymphoma
  - Lymphocyte depletion type
  - Lymphocyte predominant HD
84. Most common Non Hodgkins lymphoma is (AIIMS Nov 2013)
- Diffuse large B cell lymphoma
  - Follicular lymphoma
  - Anaplastic large cell lymphoma
  - Large T-cell leukemia/lymphoma
85. Flow cytometry is done on: (AIIMS May 2013)
- Polycythemia
  - Thrombocytosis
  - Basophil
  - Lymphocytes
86. Reed Sternberg like cell are seen in: (PGI May 2013)
- Adult T cell lymphoma
  - Extranodal NK-Cell Lymphoma
  - Marginal zone lymphoma
  - Diffuse large B cell lymphoma
  - Infectious mononucleosis
87. All are poor prognostic factors for Hodgkin's lymphoma except: (PGI Nov 2011)
- Young age
  - Involvement of stomach
  - Lymphocyte depletion
  - Extranodal metastasis
  - Large mediastinal mass



88. Which of the following malignancy is associated with underlying progression and spreads characteristically in a stepwise fashion and hence staging the disease is an important prognostic factor?  
 a. Hodgkin's lymphoma (MH 11, DNB Dec 08)  
 b. Multiple myeloma  
 c. Mature T cell NHL  
 d. Mature B cell NHL
89. True about nodular lymphocytic predominant Hodgkin's lymphoma- (PGI Nov 10)  
 a. Consists predominantly of classical RS cells  
 b. CD 15 & CD 30 positive  
 c. Made up of T lymphocytes  
 d. EBV positive  
 e. Has good prognosis
90. L & H variants of the Reed-Sternberg cells are positive for (MH 16)  
 a. CD 20 b. CD 15  
 c. CD 30 d. EBV

### NON-HODGKIN LYMPHOMA


91. In Alibert bazin syndrome, origin of lymphoma is from? (JIPMER 18)  
 a. Eosinophill b. B lymphocyte  
 c. Monocyte d. T lymphocyte
92. CLL/SLL arises from which cell? (AIIMS Nov 2017)  
 a. Mature B cell  
 b. Naive B cell  
 c. Centrocytes of germinal center  
 d. Progenitor B-cell
93. Which of the following Immunohisto chemistry marker is used in Cyclin D1 negative Mantle cell lymphoma?  
 a. SOX11 b. ITRA 1 (AIIMS May 2017)  
 c. MYD88 d. Annexin V
94. A 35 year old presented with fever. On examination he had enlarged and ulcerated tonsils. His peripheral blood smear showed lymphocytosis. Monospot test was negative. Tonsillectomy was done. The biopsy of the same showed large cells mixed with lymphocytes. The cells were positive for CD20, EBV LMP1, MUM1, CD 79a. Background cells were positive for CD3. The cells are negative for CD15. Your most probable diagnosis?



- a. Infectious mono-nucleosis (AIIMS May 2017)  
 b. Hodgkin lymphoma  
 c. EBV positive - DLBL  
 d. EBV positive mucocutaneous ulcer

95. PAX-5 is a marker for? (AIIMS May 2017)  
 a. Diffuse large B-cell lymphoma  
 b. Hodgkins lymphoma  
 c. Anaplastic lymphoma  
 d. Lymphoblastic lymphoma  
 e. AML
96. TRAP positivity is seen in? (JIPMER 2017)  
 a. Hairy cell leukemia b. ALL  
 c. Burkitts lymphoma d. T-cell leukemia
97. Primary extranodal neoplasms is/are? (PGI Nov 2017)  
 a. Burkitt lymphoma b. Waldeyer's ring lymphoma  
 c. Gastric lymphoma d. Thyroid lymphoma  
 e. Mycosis fungoides
98. BRAF mutation is seen in? (Recent Question 2016-17)  
 a. LCH b. Colon Ca  
 c. Hairy cell leukemia d. AML M7
99. A 10 yr old boy with mass in the abdomen. On imaging the paraaortic LN is enlarged. On biopsy starry sky appearance is seen. What is the underlying abnormality? (AIIMS May 2015)  
 a. p53 gene mutation  
 b. RB gene mutation  
 c. Translocation involving BCR-ABL genes  
 d. Translocation involving MYC gene
100. All are true about CLL except? (PGI Nov 2015)  
 a. Most common leukemia of adults in west  
 b. Diagnosed incidentally  
 c. Not treated in stage A  
 d. CD38 is poor prognostic marker  
 e. Most common age group is pediatric
101. CD5 is expressed in (Recent Question 2015)  
 a. Mantle cell lymphoma  
 b. Chronic myeloid lymphoma  
 c. Follicular lymphoma  
 d. Burkitt lymphoma
102. CD 30 positivity and t(2;5) is characteristic of (Recent Question 2015)  
 a. Langerhans cell histiocytosis  
 b. Lymphoplasmacytic lymphoma  
 c. Null cell lymphoma  
 d. Follicular lymphoma
103. Cyclin D1 is expressed in (Recent Question 2015)  
 a. Follicular lymphoma  
 b. Chronic lymphoid lymphoma  
 c. Mantle cell lymphoma  
 d. Splenic marginal zone lymphoma
104. CD20 is positive in all the following lymphomas except (Recent Question 2015)  
 a. Mantle cell lymphoma  
 b. Lymphocyte rich HL  
 c. Follicular lymphoma  
 d. Butkitt lymphoma
105. Find the false statement about diffuse large B cell lymphoma (Recent Question 2015)  
 a. Most common form of NHL  
 b. Waldeyer ring is involved commonly  
 c. Extranodal sites are also involved  
 d. Bone-marrow involvement in early phase





- 106. True about endemic Burkitt lymphoma:** (Recent Question 2015)
- All are associated with EBV infection
  - Abdominal mass involving ileocaecum and peritoneum and peritoneum
  - Most aggressive form
  - Bone marrow is commonly involved
- 107. Not true about anaplastic large T cell lymphoma** (Recent Question 2015)
- t(2;5) translocation
  - CD30 (ki-1) positive
  - Large anaplastic cells containing horseshoe-shaped nuclei and voluminous cytoplasm
  - ALK positive tumors carry worst prognosis
- 108. Neoplastic cells with multilobated nuclei (cloverleaf or flower cells) are seen in** (Recent Question 2015)
- Diffuse large B cell lymphoma
  - Adult T cell leukemia
  - Anaplastic large T cell lymphoma
  - Mycosis fungoides
- 109. Hallmark cells are seen in:** (Recent Question 2015)
- Anaplastic large cell lymphoma
  - Burkitt's lymphoma
  - Hairy cell leukaemia
  - Mantle cell lymphoma
- 110. Clover leaf cells are seen in:** (Recent Question 2015)
- Burkitt's lymphoma
  - Adult T cell leukemia-lymphoma
  - Hairy cell lymphoma
  - Mantle cell lymphoma
- 111. Gastric MALTomas may express all of the following except:** (Recent Question 2015)
- CD5
  - CD19
  - CD20
  - CD43
- 112. IgA lymphoma is seen in?** (Recent Question 2016)
- Spleen
  - Lymph nodes
  - Small Intestine
  - Large Intestine
- 113. Spleniculi means** (Recent Question 2015)
- Splenic calculi
  - Splenic atrophy
  - Splenic malignancy
  - Accessory spleen
- 114. A 60 year old male presents with generalized lymphadenopathy and hepatosplenomegaly. Immunophenotype: CD5 and CD19 are positive and CD10 negative. Diagnosis** (Recent Question 2015)
- Follicular lymphoma
  - Burkitt lymphoma
  - Hairy cell leukemia
  - CLL
- 115. Translocation t(8;14) of c-MYC gene is seen in** (Recent Question 2015)
- Follicular lymphoma
  - Burkitt lymphoma
  - Mantle cell lymphoma
  - Diffuse large B cell lymphoma
- 116. Characteristic translocation in mantle cell lymphoma** (Recent Question 2015)
- t(11;14)
  - t(15;17)
  - t(9;22)
  - t(8;14)
- 117. True about follicular lymphoma:** (PGI May 2015)
- Lymphadenopathy is the most common presentation
  - BCL-1 positive
  - CD5 positive
  - More common in males than females
- 118. True about Chronic Lymphocytic Leukemia:** (PGI May 2015)
- Most common leukemia in adult
  - Proliferation center is pathognomonic
  - Massive splenomegaly
  - Radiotherapy & chemotherapy are given in treatment
- 119. EBV is associated with?** (Recent Question 2015)
- Burkitts Lymphoma
  - Adamantinoma
  - Follicular lymphoma
  - CLL
- 120. CD 30 marker for**
- Anaplastic large cell lymphoma
  - Seminoma
  - Embryonal cell ca
  - Hodgkins lymphoma
- 121. An elderly male presents with anemia and fatigue. O/E splenomegaly-2 cm palpable below costal margin. Hemogram showed Pancytopenia. Which is the most common etiology?** (Recent Question 2014, DNB July 2014)
- Hairy cell leukemia
  - CML
  - Thalassemia
  - Follicular lymphoma
- 122. Mantle cell lymphomas are positive for all of the following, except -** (Recent Question 2014)
- CD23
  - CD20
  - CD5
  - CD45
- 123. Lethal midline granuloma is -** (Recent Question 2014)
- T cell lymphoma
  - B-cell lymphoma
  - NK/T cell lymphoma
  - LCH
- 124. Most common ocular lymphoma-** (Recent Question 2014)
- T-cell lymphoma
  - Hodgkin's lymphoma
  - B-cell NHL
  - Pre T-cell lymphoma
- 125. Cyclin-D & IGH fusion gene is associated with?** (Recent Question 2014)
- Mantle cell Lymphoma
  - Follicular carcinoma
  - Melanomas
  - Burkitt lymphoma
- 126. Commonest site for extranodal lymphoma is**
- Liver
  - Stomach (APPGMEE 14)
  - Small intestine
  - Large intestine
- 127. Following gene when mutated, protects tumor cells from Apoptosis** (APPGMEE 14)
- BCL - 2
  - BRCA
  - RB
  - TGF -  $\beta$
- 128. One of the following leukemia almost never develops after radiation** (APPGMEE 14)
- Acute myeloid leukemia
  - Chronic myeloid leukemia
  - Acute lymphoblastic leukemia
  - Chronic lymphocytic leukemia
- 129. Histological presence of "Hallmark Cells" with horse shoe-like or embryoid like nuclei and voluminous cytoplasm are seen in** (APPGMEE 14)
- Anaplastic large cell lymphoma (ALKpositive)
  - Familial Medullary Carcinoma
  - Familial Neuroblastoma
  - Lymphocytopredominant type Hodgkin's lymphoma
- 130. "Smudge cells" in the peripheral smear are characteristic of** (APPGMEE 14)
- Chronic myelogenous leukemia
  - Chronic lymphocytic leukemia
  - Acute myelogenous leukemia
  - Acute lymphoblastic leukemia






- 131. Burkitt's lymphoma is positive for?** (Recent Question 2013)  
 a. CD5  
 b. CD 15  
 c. CD20  
 d. CD25
- 132. A patient of 70 years, presented with generalized lymphadenopathy. WBC count was 20,000/mm<sup>3</sup> and blood film showed >70% mature looking lymphocytes. Next investigation that should be done:** (AIIMS May 2013)  
 a. LN biopsy  
 b. Peripheral blood Immunophenotyping  
 c. Bone marrow aspiration  
 d. Peripheral blood cytogenetics
- 133. Marker for Lymphoma is:** (AIIMS May 2013)  
 a. S-100  
 b. HMB-45  
 c. Leukocyte common antigen  
 d. Cytokeratin
- 134. All are true about Mantle cell lymphoma except:**  
 a. Associated with (11;14) translocation (PGI May 2013)  
 b. Overexpression of the BCL protein  
 c. CD 5 positive  
 d. CD 23 positive  
 e. Centroblasts frequently seen
- 135. Lymphoma associated with translocation of c-myc is?** (JIPMER 2013)  
 a. Follicular Lymphoma  
 b. Mantle cell Lymphoma  
 c. Burkitts Lymphoma  
 d. Anaplastic large cell lymphoma
- 136. Most common Non-Hodgkin's lymphoma of orbit:**  
 a. B cell  
 b. T cell (AIIMS May 2012)  
 c. NK cell  
 d. Plasma cell
- 137. International prognostic index for lymphomas includes the following prognostic factors, except:** (AIIMS May 11)  
 a. Stage of disease  
 b. Number of extralymphatic sites involved  
 c. LDH  
 d. Hemoglobin and albumin
- 138. True about abdominal lymphoma:** (PGI Nov 2011)  
 a. GIT lymphoma most commonly has polypoid appearance  
 b. Primary small intestinal lymphoma are most commonly located in ileum  
 c. Lymphoma is most common primary malignant neoplasm of spleen  
 d. Stomach is most common site for extranodal lymphoma  
 e. MALT lymphoma is associated with H. pylori infection
- 139. Features of hairy cell leukemia are all except:** (PGI May 2011)  
 a. Splenomegaly  
 b. Hepatomegaly  
 c. Vasculitic syndromes  
 d. Pancytopenia  
 e. Erythema multiforme
- 140. Massive splenomegaly is found in:** (PGI May 2011)  
 a. Hairy cell leukemia  
 b. CML  
 c. Typhoid  
 d. Sickle cell anemia  
 e. ITP


- 141. Compared to the other leukemias, hairy cell leukemia is associated with which of the following infections -** (MH 10)  
 a. Parvovirus D 19  
 b. Mycoplasma  
 c. Atypical mycobacteria  
 d. Salmonella

### PLASMA CELL NEOPLASMS

- 142. All of the following are feature of waldenstrom macroglobulinemia except:?** (PGI May 18)  
 a. Hyperviscosity  
 b. Polycythemia  
 c. Treated with alkylating agents  
 d. Low grade lymphoplasmacytoid lymphoma  
 e. IgG paraprotein
- 143. Myeloma associated with good prognosis? (JIPMER 18)**  
 a. t (11 : 14)  
 b. t (14 : 18)  
 c. del 17  
 d. t (14, 16)
- 144. Origin of lymphoplasmacytic lymphoma (Waldenstrom Macroglobulinemia) is from?** (JIPMER 18)  
 a. Germinal centre T cell  
 b. Germinal centre B cell  
 c. Post germinal centre B cell  
 d. Pre germinal centre B cell
- 145. Which of the following hematological condition is depicted below?** (Recent exam 2018)



- a. Multiple myeloma  
 b. Megaloblastic anemia  
 c. Chronic myeloid leukemia  
 d. Metastasis
- 146. Multiple myeloma causes?** (AIIMS May 2017)  
 a. Cast nephropathy  
 b. Amyloidosis  
 c. Cryoglobulinemia  
 d. Interstitial nephritis  
 e. Nephrocalcinosis
- 147. An elderly male presented with history of intractable diarrhea. His bone marrow and renal biopsy as shown below. Which of the following is the most appropriate diagnosis?** (AIIMS May 2017)



- a. Leishmaniasis  
 b. Multiple myeloma  
 c. Lymphoma  
 d. Urate nephropathy



- 148. MYD88 L265P mutation is seen in?** (JIPMER 2017)  
a. Hairy cell leukemia  
b. Waldenstrom Macroglobulinemia  
c. Multiple Myeloma  
d. AML
- 149. Large homogenous eosinophilic inclusions in plasma cells are called** (Recent Question 2015)  
a. Dutcher bodies  
b. Councilman bodies  
c. Russell bodies  
d. Mallory hyaline bodies
- 150. Russell bodies are seen in** (Recent Question 2015)  
a. Mast cells  
b. Plasma cells  
c. Histiocytes  
d. Langerhan cells
- 151. The single most important predictor of survival in multiple myeloma** (Recent Question 2015)  
a. IL-6 levels  
b. Bence jones proteinuria  
c. CD 138 positivity  
d. Serum  $\beta_2$ -microglobulin
- 152. In multiple myeloma treatment, the following drug is avoided during induction therapy for transplant candidates** (Recent Question 2015)  
a. Thalidomide  
b. Bortezomib  
c. Melphalan  
d. Dexamethasone
- 153. False statement about monoclonal gammopathy of unknown significance** (Recent Question 2015)  
a. <3g/dL of monoclonal protein  
b. No bence jones proteinuria  
c. Bone marrow plasma cells < 10%  
d. Does not progress to multiple myeloma
- 154. M spike in waldenstrom macroglobulinemia is due to** (Recent Question 2015)  
a. IgM  
b. IgG  
c. IgA  
d. IgD
- 155. False statement about MGUS** (Recent Question 2015)  
a. Few progress to multiple myeloma  
b. Asymptomatic  
c. Secrete M protein  
d. Bence jones proteinuria
- 156. POEM syndrome. E stands for:** (Recent Question 2016)  
a. Endocrinopathy  
b. Edema  
c. Eosinophilia  
d. Erythema
- 157. Life span of plasma cell** (Recent Question 2016)  
a. 12 hrs  
b. 24 hrs  
c. 48 hrs  
d. Days to weeks
- 158. Multiple myeloma is a tumor of?** (Recent Question 2015)  
a. B-lymphocyte  
b. T-lymphocyte  
c. Lymph nodes  
d. Plasma cell
- 159. A patient presents with bone pain. X-ray reveals destructive lesions. Lab investigations show hypercalcemia. Serum electrophoresis shows M spike, while Bone marrow shows 35% plasma cells. What is your diagnosis?** (JIPMER 2014)  
a. MGUS  
b. Smoldering myeloma  
c. Multiple myeloma  
d. Plasma cell leukemia
- 160. Beta-2 -microglobulin is a tumor marker for** (Bihar PG 2014)  
a. Multiple myeloma  
b. Lung cancer  
c. Colonic neoplasm  
d. Choriocarcinoma
- 161. Proliferation and survival of myeloma cells are dependent on which of the following cytokines?** (APPGMEE 14)  
a. IL-1  
b. IL-6  
c. IL-2  
d. IL-5
- 162. Multiple myeloma-all are true except?** (JIPMER 2013)  
a. Proteinuria  
b. Visual disturbance  
c. Bleeding  
d. Dystrophic calcification
- 163. Bence jones proteinuria is derived from?** (JIPMER 2013)  
a. Alpha globulins  
b. Light chain globulins  
c. Gamma globulins  
d. Delta globulins
- 164. Malignancy associated with Waldenstrommacroglobulinemia?** (JIPMER 2013)  
a. Mycosis fungoides  
b. Smoldering myeloma  
c. Primary effusion lymphoma  
d. Lymphoplasmacytic lymphoma
- 165. Which of the metabolic abnormality is seen in multiple myeloma?** (DNB 08/ DPG 11)  
a. Hyponatremia  
b. Hypokalemia  
c. Hypercalcemia  
d. Hyperphosphatemia
- 166. Multiple myeloma is diagnosed by -** (JIPMER 11)  
a. 24 hours urine protein  
b. Kidney biopsy  
c. > 10% plasmacytosis  
d. Rouleaux formation in blood
- 167. Lymphoplasmacytoid lymphomas may be associated with** (AIPGMEE 10)  
a. IgG  
b. IgM  
c. IgA  
d. IgE
- 168. Which of the following is not a minor diagnostic criteria for multiple myeloma?** (AIIMS Nov 10, 08)  
a. Lytic bone lesions  
b. Plasmacytosis greater than 20%  
c. Plasmacytoma on biopsy  
d. Monoclonal globulin spike on serum electrophoresis of > 2.5 g/dl for IgG, > 1.5 g/dl for IgA
- 169. Not a feature of multiple myeloma** (AIIMS May 05) (WB PG 2016)  
a. Hypercalcemia  
b. Anemia  
c. Hyperviscosity  
d. Elevated alkaline phosphatase

#### CHRONIC MYELOPROLIFERATIVE NEOPLASMS

- 170. True about BCR-ABL 'traits' are all except?** (JIPMER 18)  
a. P190 has an indolent course  
b. P190 is a bad prognostic factor  
c. P230 is positive in chronic neutrophilic leukemia  
d. P230 has an indolent course
- 171. A 40-year-old woman is on treatment for CLL. Over the past few months she noticed swellings in the neck and axilla which was rapidly increasing in size. She complains of feeling feverish and experiences weight loss. Which of the following is responsible?** (JIPMER 18)  
a. Richter transformation  
b. Progression of CLL  
c. EBV infection  
d. Immunodeficient hemolytic anemia  
e. MHC non-expression



172. A 60-year-old male living in hilly area has Hb of 16 gm% ,TC 21000/ul. DLC showed metamyelocytes and myelocytes 40%, N25% L40%, E5%. Platelet count 3.25 lakh/u. He presented with hypertension and on examination, spleen was just palpable below costal margin. What is the next step? (AIIMS Nov 2017)
  - a. Bone marrow with reticulin stain
  - b. JAK STAT mutation assessment
  - c. Philadelphia chromosome
  - d. Erythropoietin levels
173. 45/m presented with leuko-erythroblastic blood picture with dacrocytes. What is bone marrow finding? (JIPMER 2017)
  - a. Fatty degeneration with erythroid cell hyperplasia with megakaryocytes
  - b. Abundant fat cells
  - c. Focal cellular marrow with hypocellular areas and atypical megakaryocytes.
  - d. Hypercellular marrow with prominent blasts
174. Which of the following statements is true regarding juvenile chronic myeloid leukemia? (Recent Question 2016-17)
  - a. Philadelphia chromosome is negative.
  - b. Thrombocytopaenia is uncommon.
  - c. The prognosis is better than the adult form of chronic myeloid leukemia.
  - d. Single agent chemotherapy with busulfan or hydroxyurea can achieve remission.
175. Robertsonian translocation is seen in? (Recent Question 2016)
  - a. AML
  - b. CML
  - c. ALL
  - d. CLL
176. True about Robertsonian translocation is? (Recent Question 2016)
  - a. Acrocentric chromosome involved
  - b. Balanced translocation
  - c. Large part is lost
  - d. Poor prognosis
177. Which of the following is not a characteristic feature of Myelodysplastic syndrome? (Recent Question 2015)
  - a. Leucoerythroblastic blood picture
  - b. Pawn ball megakaryocytes
  - c. Pseudo pelger heut cells
  - d. Transformation to AML
178. The following is true regarding polycythemia vera ? (Recent Question 2015)
  - a. Raised ESR
  - b. Decrease LAP score
  - c. Thrombocytopenia
  - d. Leukocytosis
179. Best investigation for BCR-ABL (Recent Question 2015)
  - a. Flow cytometry
  - b. Fluorescent in situ hybridization
  - c. EISA
  - d. Polymerase chain reaction
180. Hyposegmented neutrophils are seen in (Recent Question 2015)
  - a. Megaloblastic anemia
  - b. Sideroblastic anemia
  - c. Accelerated phase of CML
  - d. Blast crisis phase of CML
181. Erythromelalgia is polycythemia vera is a complication of (Recent Question 2015)
  - a. Erythrocytosis
  - b. Thrombocytosis
  - c. Granulocytosis
  - d. Lymphocytosis
182. Treatment of choice for CML (Recent Question 2015)
  - a. Sorafenib
  - b. Imatinib mesylate
  - c. Sunitinib
  - d. Erlotinib
183. Not seen in polycythemia vera (Recent Question 2015)
  - a. Platelet function abnormalities
  - b. Normal red cell morphology
  - c. Abnormal oxygen saturation
  - d. Low ESR
184. Which of the following is NOT commonly seen in polycythemia vera ? (MH 16)
  - a. Thrombosis
  - b. Hyperuricemia
  - c. Prone for Acute Leukemia
  - d. Spontaneous severe infection
185. Not seen in essential thrombocythosis (Recent Question 2015)
  - a. Activating mutation in JAK2 gene
  - b. Abnormally large platelets
  - c. Erythromelalgia
  - d. Marrow fibrosis
186. Triad of leukoerythroblastosis, tear drop erythrocytes and large platelets is seen in (Recent Question 2015)
  - a. Essential thrombocythosis
  - b. Primary myelofibrosis
  - c. Myelodysplastic syndrome
  - d. Langerhan cell histiocytosis
187. False regarding myelodysplastic syndromes (Recent Question 2015)
  - a. Hypercellular bone marrow
  - b. Increased neutrophil alkaline phosphatase
  - c. Ringed sideroblasts
  - d. Pawn ball megakaryocytes
188. Mutation seen in systemic mastocytosis (Recent Question 2015)
  - a. FGFR1 fusion genes
  - b. BCR-ABL fusion gene
  - c. JAK 2 point mutation
  - d. c-kit point mutation
189. Leukocyte alkaline phosphatase score is decreased in (Recent Question 2015)
  - a. Pregnancy
  - b. Polycythemia vera
  - c. Infections
  - d. Myelodysplastic syndrome
190. Highest LAP score is seen in? (Recent Question 2015)
  - a. Acute Myeloid Leukemia
  - b. Polycythemia Vera
  - c. Chronic myeloid Leukemia
  - d. Paroxysmal Nocturnal Hemoglobinuria
191. Dwarf megakaryocytes with unilobed nucleus is characteristic of (Recent Question 2015)
  - a. Myelodysplastic syndrome
  - b. Essential thrombocytosis
  - c. Chronic myeloid leukemia
  - d. Polycythemia vera
192. 45/m presents with leukoerythroblastic blood picture in PBS with drytap, What is your diagnosis? (Recent Question 2015)
  - a. AML
  - b. CML
  - c. ALL
  - d. Myelofibrosis
193. BCR-ABL fusion gene is detected by? (Recent Question 2015)
  - a. Flow cytometry
  - b. FISH
  - c. Karyotyping
  - d. RT-PCR



**194. All are true about Polycythemia vera except-**  
(Recent Question 2014, AIIMS 01)

- a. Increased vit B<sub>12</sub>
- b. Decreased LAP score
- c. Leucocytosis
- d. Increased platelets

**195. Bone marrow finding in myelofibrosis-**  
(Recent Question 2014)

- a. Dry tap (hypocellular)
- b. Megaloblastic cells
- c. Microcytic cells
- d. Thrombocytosis

**196. Pseudo-Pelger-Huet cells are seen in-**  
(Recent Question 2014, 2013)

- a. Hairy cell leukemia
- b. Multiple myeloma
- c. Myelodysplastic syndrome
- d. Hodgkin's lymphoma

**197. Polycythemia is not caused by-** (Recent Question 2014)

- a. Renal carcinoma
- b. Liver carcinoma
- c. Cerebellar hemangioma
- d. Lung carcinoma

**198. Which of the following is not a myeloproliferative disease**  
(Recent Question 2014)

- a. Polycythemia vera
- b. Acute myeloid leukemia
- c. Chronic myeloid leukemia
- d. Essential thrombocytosis

**199. In patients with Chronic Myeloid Leukemia**  
(AP PGME 14)

- a. ABL gene on Chr. 22 is trans-located to BCR gene on Chr.9
- b. The fusion gene bcr-abl forms a protein with tyrosine kinase activity
- c. Splenomegaly is unusual
- d. Philadelphia chromosome positive patients respond poorly to Imatinib

**200. Which of the following is not a chronic myeloproliferative disorder?**  
(AP PGME 14)

- a. Polycythemia vera
- b. Myeloid metaplasia
- c. CML
- d. Essential thrombocytopenia

**201. About CML in children true is:** (AIIMS Nov 2013)

- a. Translocation between long arm of chr 9 and short arm of chr 22
- b. Protein tyrosine kinase inhibitor are the drug of choice
- c. Most commonly presents in blast crisis
- d. 2nd most common malignancy

**202. Myelofibrosis leading to a dry tap on bone marrow aspiration is seen with which of the following condition?**

- a. Burkitt's lymphoma
- b. Acute erythroleukemia
- c. Acute Megakaryocytic Leukemia
- d. Acute Myelomonocytic Leukemia

**203. In myelodysplastic syndrome, ring sideroblast is seen in:** (WB PG 2011)

- a. Mitochondria
- b. Golgi body
- c. Nuclear membrane
- d. ER

### HEAVY CHAIN DISEASES

**204. Palatal edema is significant for?** (IIPMER 18)

- a. Alpha heavy chain disease
- b. Gamma heavy chain disease
- c. Mu chain disease
- d. Beta heavy chain disease

### LCH

**205. Which of the following cell is seen in LCH?**  
(Recent Question 2016-17)

- a. Eosinophil
- b. Basophils
- c. Neutrophil
- d. Langerhans cell

**206. In Langerhans Cell Histiocytosis, the characteristic abnormality seen is**  
(AIIMS May 2015)

- a. Birbeck's granules
- b. Macrophages
- c. Plasma cell
- d. Giant cell

**207. Langerhans cell shows which of these?**  
(Recent Question 2015)

- a. Badminton racquet appearance
- b. CD 100a
- c. MPO +
- d. Birbeck's granules

**208. Localised langerhans cells histiocytosis affecting head & neck is -**  
(Recent Question 2014)

- a. Letterer-siwe disease
- b. Pulmonary langerhans cell histiocytosis
- c. Hand-schuller-christian disease
- d. Eosinophilic granuloma

**209. CD marker for Langerhans cell histiocytosis is?**  
(Recent Question 2013, DNB Aug 12)

- a. CD 17
- b. CD 23
- c. CD 1a
- d. CD 117

**210. About Pulmonary Langerhans cell histiocytosis, all are true EXCEPT:** (WB PG 2012)

- a. Associated with smoking
- b. Pneumothorax in 10% cases
- c. 20-40 years male predominance
- d. Corticosteroids have role in treatment





## Answers with Explanations

### 1. Ans. (b) Castleman disease

Based on clinical presentation, Castleman disease has been divided into a solitary and a multicentric form. The solitary form presents as a mass located most commonly in the mediastinum, neck, lung, axilla, mesentery, broad ligament, retroperitoneum, and several other sites. Grossly, it is round, well circumscribed, with a solid gray cut surface and can measure 15 cm or more in diameter. The follicles show marked vascular proliferation and hyalinization of their abnormal or atrophic germinal centers, surrounded by concentrically arranged small lymphocytes imparting an 'onion-skin' pattern.

Remember angiolymphoid hyperplasia shows thick walled blood vessels with prominent endothelial cells and inflammatory eosinophilia infiltrates.

### 2. Ans. (d) Adult T cell leukemia

HHV 8 is associated with

- Castleman disease
- Primary effusion lymphoma
- Kaposi sarcoma

### 3. Ans. (a) Mantle zone (Ref: R 9/p 583)

A normal lymph node has 2 areas: Cortex → Lymphoid follicles and paracortical areas, Medulla → predominantly blood vessels

#### Structures in a Follicle

- **Germinal center:** Round/oval zone containing pale staining cells, surrounded by darker cells
- **Mantle zone:** Small dark coloured unchallenged B cells surrounding pale staining germinal centers
- **Marginal zone:** Light zone surrounding follicles; contains postfollicular memory B cells derived after stimulation of recirculating cells from T cell dependent antigen; named "marginal cells" due to location

### 4. Ans. (a) May heggalin anomaly

May-Hegglin anomaly (MHA) is an autosomal dominant disorder characterized by thrombocytopenia, giant platelets containing few granules; and large, well-defined, basophilic, cytoplasmic inclusion bodies in granulocytes that resemble Döhle bodies

### 5. Ans. (c) C


(Ref: Wintrob's 13th ed. Pg. 303; Wintrob's Atlas)

**Key to the figure:**

- |                 |                 |
|-----------------|-----------------|
| • A: Lymphocyte | • C: Eosinophil |
| • B: Neutrophil | • D: Basophil   |

Eosinophils are increased in parasitic infection

### 6. Ans. (c) Plasma cell (Ref: Wintrob's 13th ed. Pg. 303; Wintrob's Atlas)



Plasma cells are spherical or ellipsoid and range from 5 to 30  $\mu\text{m}$  in size. The cytoplasm is abundant & basophilic (deep blue); with a well-defined perinuclear clear zone that contains Golgi apparatus.

### 7. Ans. (d) Platelet count of <40000/ul is contraindication

(Ref: Dacie Practical Hematology 10<sup>th</sup> ed/ pg 163)

The given image is of **Jamshedi Bone marrow biopsy needle**. Most are **14 to 18 gauge**;

In most of the situations where bone marrow aspiration is indicated (eg suspected Acute Leukemia, thrombocytopenia is usually present; So, platelet count of < 40,000/uL is not a contraindication to bone marrow aspiration

### 8. Ans. (a) Measles

### 9. Ans. (a) Bone marrow examination

(Ref: Dacie Practical Hematology, 10<sup>th</sup>ed/163; Complete review of Pathology 1st/761)

This is a Sahl's bone marrow aspiration needle. (Sahl's needle has a Screw on the side; S for S)



Klima's bone marrow aspiration needle (no side screw)

### 10. Ans. (a) CD8 (Ref: Robbins 9th/pg 590; 8th/pg 600)

### 11. Ans. (d) CD46 (Ref: Wintrob's 12th/pg 2523)

CD46 (Complement Membrane Cofactor Protein): It is a receptor to a number of pathogens, such as herpes virus 6, M protein of group A streptococci, Neisseria gonorrhoeae, and Escherichia coli.

### 12. Ans. (a) Less than 500/ul (Ref: Robbins 9th/pg 583)

Serious infections are most likely when the neutrophil count falls below 500 per mm<sup>3</sup>.

### 13. Ans. (c) Strongyloides

(Ref: Robbins 9th/pg 583 8th/pg 593)

**Parasitic** infestations: Ascariasis, Hookworm, Strongyloides, Filariasis, Trichinosis can cause eosinophilia



14. Ans. (a) **Pro erythroblast** (Ref: Robbins 9th/pg 580-581)

15. Ans. (d) **Dohle bodies**

(Ref: Robbins 9th/pg 583 9th/pg 593)

In sepsis or severe inflammatory disorders, there can be morphologic changes in the neutrophils:

- Cytoplasmic vacuoles
- Toxic granules
- Döhle bodies

16. Ans. (a) **Tibia** (Ref: Wintrobe's 12th/pg 10)

Iliac crest<sup>Q</sup> is the **most common** site for **bone marrow sampling overall in adults**, while in children it is Anterior medial **tibial**<sup>Q</sup> area, below tibial tuberosity

17. Ans. (a) **May Hegglin anomaly**

(Ref: Wintrobe's 12th/pg 1549)

- Anemia with thrombocytopenia & inclusions in neutrophils is suggestive of May Hegglin anomaly
- **Evan syndrome:** Autoimmune hemolytic Anemia with thrombocytopenia

18. Ans. (b, d, e); **b. CD 10; d. CD19; e. CD20**

(Ref: Robbins 9th/pg 590; 8th/pg 600)

19. Ans. (a) **CD 19p**

(Ref: Robbins 9th/pg 590; 8th/pg 600; Wintrobe's 12th/pg 2504)

CD19 is the pan B cell marker; Since it is present on Chr 16p, so the best suitable answer here is CD 19p

20. Ans. (a) **B-cells** (Ref: Wintrobe's 12th/pg 1589-1593)

**Pathogenesis of Infectious Mononucleosis:** Caused by Epstein Barr Virus (EBV)<sup>Q</sup> infection

- **Entry** of EBV in the **oral cavity**<sup>Q</sup>
- EBV initially **infects oral epithelial cells** à symptoms of **pharyngitis**.<sup>Q</sup>
- **Intracellular**<sup>Q</sup> **viral replication** and cell lysis with **release of new virions**
- Virus **spreads to contiguous structures** such as the **salivary glands**<sup>Q</sup>
- Eventual **viremia & infection of B lymphocytes**<sup>Q</sup> in the peripheral blood & entire lymphoreticular system, including **liver & spleen**.
- **DOWNY cells** are **atypical CD8+ T lymphocytes** that are characteristic of infectious mononucleosis<sup>Q</sup>
- DOWNY cells exhibit **both suppressor & cytotoxic**<sup>Q</sup> functions that develop **in response to the infected B lymphocytes**.

21. Ans. (b) **2000 to 5000 cells/microliter**

(Ref: Medscape (<http://emedicine.medscape.com/article/329614-overview>))

The peripheral blood eosinophil count in Eosinophilia-myalgia syndrome is usually **2000 to 5000 cells/microliter**

#### CDC definition of 'Eosinophilia-myalgia syndrome'

- Incapacitating myalgias,
- Blood **eosinophil count greater than 1000 cells/μl**, and
- **No evidence of infection** (eg, trichinosis) or **neoplastic** conditions that could account for these findings.

22. Ans. (a) **RBC** (Ref: Robbins 9th/pg 580-581)

- **Myeloid cells** are WBCs like **neutrophils, monocytes, Basophils, RBCs and Platelets**.
- **Lymphoid cells** include **B-Lymphocyte, T-lymphocyte and NK cells**.

So following injection of myeloid series cells, RBCs will be released into peripheral blood.

23. Ans. (b) **Myoblast Progenitor cells**

(Ref: Robbins 9th/pg 580-581)

- Endothelial Progenitor cells, Mesenchymal stem cells and Hematopoietic stem cells are found in the bone marrow.
- Mesenchymal stem cells are multipotent stromal cells constituting 0.001-0.01% of bone marrow cells.

24. Ans. (a) **CD 45**

(Ref: Robbins 9th/pg 590; 8th/pg 600; Wintrobe's 12th/pg 2522)

CD 45

- Found in all hematopoietic cells except erythrocytes.
- CD45 plays an essential role in lymphocyte activation.
- Peripheral blood naïve T cells are CD45RA, whereas memory (activated) T cells are CD45RO+

25. Ans. (b) **Kikuchi disease**

(Ref: Rheumatology: Diagnosis and Therapeutics; Edited by John J. Cush, Arthur Kavanaugh, Charles Michael Stein; Lippincott Williams & Wilkins, 2005; pg 228)

Kikuchi Disease

- **Kikuchi-Fujimoto disease or histiocytic necrotizing lymphadenitis** is a **benign**, rare disorder that affects young women (more so than men) with **recurrent necrotizing lymphadenitis**.
- MC involves **cervical region**, **Histology:** Zone of **necrosis** surrounded by blast-like plasmacytoid lymphocytes

26. Ans. (a) **Kimura's disease**

(Ref: Joachim's Lymph Node Pathology. Harry L. Joachim, L. Jeffrey Medeiros; Lippincott Williams & Wilkins, 2009. Pg 190)

**Kimura's Disease**

- **Eosinophilic abscess in lymph node is seen**
- A chronic inflammatory disorder involving subcutaneous tissue & lymph nodes predominantly in the head & neck region & is characterized by **angiolymphoid proliferation & eosinophilia**
- In **Kimura disease**, **salivary gland involvement, Glomerulonephritis, Nephrotic syndrome, Eosinophilia & increased IgE** are more common than in Kikuchi disease



- Angiolymphoid hyperplasia with eosinophilia (ALHE), owing to some histologic similarities, can be confused with or mistaken for an early stage of **Kimura disease**.

27. **Ans. (d) CD34** (Ref: Robbins 9th/pg 590; 8th/pg 600)

28. **Ans. (b) ALL with hyperploidy**

All with hyperploidy is known terminology  
Provisional entities in new WHO classification

Leukemia	Provisional entities
AML	AML with BCR-ABL-1 AML with mutated RUNX <sub>1</sub>
ALL	Early T cell precursor ALL

29. **Ans. (a) Cytokeratin**

A mediastinal mass has a differential of thymoma and T cell ALL. T -ALL will have the markers TDT (Lymphoblast), CD 1a & CD3. Thymoma has 2 components epithelial and lymphoid component so is positive for EMA, **cytokeratin 7 & 20**, CD57 CD5, bcl-2, calretinin, vimentin, **CD3**, CD1a, CD20, CD99 and Ki67 & **TdT**.

30. **Ans. (b) Peripheral smear**

Myeloid (granulocytic) sarcoma, or myeloblastoma are extramedullary blast proliferation. These tumors are called chloromas because some appear green or turn green in dilute acid secondary to expression of MPO. The tumors are usually localized; they often involve bone, periosteum, soft tissues, lymph nodes, or skin. Common sites are the orbit and the paranasal sinuses.

The diagnosis can be made if Auer rods are detected on blasts in peripheral smear or if myeloid origin is confirmed by cytochemical or immunohistochemical methods. The diagnosis should be suspected if eosinophilic myelocytes are present in hematoxylin and eosin-stained biopsy sections. Imprint preparations can be helpful.

31. **Ans. (c) PAX5** (Ref: R 9/ p 590)

PAX-5 is a B Cell marker.

32. **Ans. (a) ALL** (Ref: R 9/ 611)

- The picture is that of a lymphoblast. Lymphoblasts will be 3-4x larger than a mature RBC, High nuclear to cytoplasmic ratio, Round Nucleus with immature chromatin (not clumped), Prominent nucleoli, Cytoplasm is scant, light blue and lacks granules

33. **Ans. (b) Hyperdiploidy**

34. **Ans. (b) Leukemia**

(Ref: Wintrob's 13/p 734)

Autologous stem cell support after myeloablative therapy has been successful for treatment of acute myelogenous leukemia (AML), non-Hodgkin lymphoma, and Hodgkin disease

35. **Ans. (b) t(15:17)**

36. **Ans. (a, d) a. Hyperploidy; d. Response to treatment**

(Ref: Robbins 9th/ 590-592; Complete review of Pathology 1<sup>st</sup>/280)

Discussing options one by one:

• Hyperploidy	<b>Intermediate prognosis; can be considered good</b>
• Organomegaly	L. Node, liver, spleen enlargement, Testicular enlargement → poor prognosis
• TLC more than 50000/ul	Poor prognosis; TLC <10,000/ul has good prognosis
• Response to treatment	Early response to treatment is a good prognostic factor but non responsive is a poor prognostic factor

*The best answer suited here is Response to treatment > hyperploidy as remission status at 14 days is the best guide to prognosis; and so the best prognostic factor.*

37. **Ans. (a) Acute Myeloid Leukemia**

(Ref: Robbins 9th/pg 612 8th/622)

38. **Ans. (a) tdt positive** (Ref: Robbins 9th/pg 590)

39. **Ans. (a) Neutrophils** (Ref: Robbins 9th/pg 583/ 8th/593)

40. **Ans. (c) HTLV is a cause in immunodeficient patients**

(Ref: Robbins 9th/pg 239-242)

EBV rather than HTLV-1 is implicated in HLH

41. **Ans. (a) Myeloperoxidase** (Ref: Robbins 9th/pg 612; 8th/622)

CD117 is a Myeloid series marker

- **Granulocytic(Myeloid ) sarcoma, or Myeloblastoma, is an extramedullary tumor**
- Also called **chloromas** because some appear/turn green in dilute acid secondary to expression of MPO
- Usually localized; often involve bone, periosteum, soft tissues, lymph nodes, or skin.
- Common sites are the **orbit & paranasal sinuses;**
- Can involve GIT, genitourinary tract, breast, cervix, salivary glands, mediastinum, pleura, peritoneum & bile duct

42. **Ans. (d) M7** (Ref: Robbins 9th/pg 612; 8th/622)

43. **Ans. (a) Age >10 years** (Ref: Robbins 9th/pg 590-592)

44. **Ans. (a) Rituximab** (Ref: Robbins 9th/pg 590-592)

45. **Ans. (a) Pre-B cell ALL** (Ref: Robbins 9th/pg 590-592)

46. **Ans. (b) Leukemia** (Ref: Robbins 9th/pg 612; 8th/622 )

47. **Ans. (c) Auer Rod** (Ref: Robbins 9th/pg 590-592)



**48. Ans. (d) Presence of t (9;22)**

(Ref: Robbins 9th/pg 590-592)

**49. Ans. (a) Acute myelogenous leukemia**

(Ref: Robbins 9th/pg 590-592)

The peripheral smear shows myeloblasts having **delicate nuclear chromatin**, 2-4 nucleoli, and moderate cytoplasm. One of them shows Auer rods. So it is a case of AML.

**50. Ans. (d) t(12;21) (Ref: Robbins 9th/pg 612; 8th/pg 622)**

AML with t(8;21)<sup>q</sup> in V 16 and t(15, 17) have Favorable Prognosis

**51. Ans. (a) ALL**

(Ref: Robbins 9th/pg 590-592)

**52. Ans. (d) M4 (Ref: Robbins 9th/pg 612; 8th/pg 622)**

- AML causing **gum hypertrophy** are AML-M5, M4
- AML causing **extramedullary blast proliferations (Chloromas)** are AML M2, M4, M5
- AML causing blast infiltrations in skin (leukemia cutis) are AML M5, M4

**53. Ans. (b) Promyelocytic (M3)**

(Ref: Robbins 9th/pg 612; 8th/pg 622)

Acute Promyelocytic Leukemic (APML, M3) cells can induce **Disseminated intravascular coagulation (DIC)**

**54. Ans. (a) AML (Ref: Robbins 9th/pg 612; 8th/pg 622)**

This **40 yr old** female is presenting with **leukocytosis** and **increased blast counts** in the peripheral smear. The cytogenetic study done here shows **t(8;21)**. This finding is **suggestive of Acute myeloid leukemia** even if there are **<20% blast counts**.

- It cytogenetic abnormalities like **t(15;17)**, **t(8;21)**, **inv(16)** are encountered in a patient with symptomatic myeloid disease, **AML should be diagnosed despite the lower blast percent**.

**55. Ans. (c) Pre B ALL (Ref: Robbins 9th/pg 590-592)**

**56. Ans. (a) Peak incidence of Chronic myeloid leukemia is in the fifth to sixth decades of life**

(Ref: Robbins 9th/pg 590-592, 616-618)

Discussing the options one by one,

a.	True
b.	False; <b>Prognosis is worse for those &gt; 50 yrs age &amp; those with a Hb level &lt;10 g/dL &amp; white cell counts &lt;2 × 10<sup>9</sup>/L</b>
c.	False; ALL in children < 1 yr & >10 yrs has poor prognosis
d.	False; <b>Median age at diagnosis of CLL is 60 years</b>

**57. Ans. (d) Mixed phenotypic acute leukemia**

(Ref: WHO Classification of Hemato-Lymphoid Tumors, 4th edition, 2008, pg 150)

The child has severe Anemia & Leukocytosis with blasts on peripheral smear. So this is a case of leukemia.

Immunophenotyping suggests:

Immunophenotype	Lineage
CD 10(+ve),	B-Cell marker
CD 19 (+ve),	B-Cell marker (most specific)
CD 117 (+) ve,	Myeloid Cell marker
MPO (+) ve	Myeloid Cell marker (most specific)
CD 33(-) ve	Myeloid Cell marker

As both B-cell and myeloid lineages are positive, this is a case of **Mixed phenotypic acute leukemia (MPAL)**.

**58. Ans. (b) Paroxysmal cold hemoglobinuria**

(Ref: Wintrobe's 12th/pg chap 78)

- Paroxysmal nocturnal hemoglobinuria (PNH), Aplastic anemia & Myelodysplastic syndrome predispose to Leukemia

**59. Ans. (a) CNS (Ref: Harrison 18th/pg Chapter 189)**

**Lymphoma in HIV**

- **90% of lymphomas in HIV** are **B cell** in phenotype; more than half contain **EBV DNA**.
- **Immunoblastic lymphomas** account for **60% of the cases** of lymphoma in patients with AIDS.
- **Primary CNS lymphoma** accounts for **20%** of the cases of lymphoma in patients with HIV infection.
- **Most common extranodal site involved in Lymphoma in HIV is the CNS**, which is involved in one-third of all patients with lymphoma.

**60. Ans. (c) Presence of testicular involvement at presentation**

(Ref: Robbins 9th/pg 590-592; 8th/pg 600-603)

**61. Ans. (c) Lymphocyte predominant**

(Ref: Robbins 9th ed p 608)

Lymphohistiocytic variants (L&H cells) with polypoid nuclei, inconspicuous nucleoli, and moderately abundant cytoplasm are characteristic of the lymphocyte predominance subtype Hodgkin lymphoma.

**62. Ans. (a) NLPHL (Ref: R 9/p 606)**

This is classical description of **NLPHL**. (Nodular lymphocyte predominant Hodgkin lymphoma)





63. **Ans. (b, d) b. DLBCL is most common in India; d. Burkitts lymphoma arises from Germinal centre**

(Ref: Robbins 9th/pg 602-603)

**About other options,**

- a. Mantle cell lymphoma arises from mantle layer & not germinal centre  
c. CD 5, CD 3 & CD 8 are T cell markers

64. **Ans. (a) a. If Imatinib not working then Dasatinib can be used; b. BCR-ABL activates tyrosine kinase**

(Ref: Dacie Practical Hematology, 10th ed/163;  
Robbins 9th/pg 616-618)

Note: Bone marrow aspirate/biopsy is used for staging of CML & is not an essential criteria for diagnosis; Blast crisis has > 20% blasts in bone marrow/peripheral smear

65. **Ans. (a) a. Lymphocyte predominant HD**

(Ref: Robbins 9th/pg 325)

EBV is associated with Plasmablastic lymphoma, Nasopharyngeal carcinoma & Mixed cellularity Hodgkin disease

66. **Ans. (a, c, d, e) a. Often localized to single axial group of lymph node; c. Contiguous spread of lymph node d. Can be cured by chemotherapy & radiotherapy e. Commonly presents with painless lymphadenopathy**

(Ref: Harrison 19th/708-09; Robbins(SAE) 9th/607,610-11; Oxford Textbook of Haematology 2nd/211; CMDT 2016/ 530-31; Ref: Harrisons 19e/ pg 700)

In stage E of Hodgkin Lymphoma (Ann Arbor staging), localized, solitary involvement of extralymphatic tissue, excluding liver and bone marrow is seen; So hepatomegaly may not be always present in Hodgkin disease.

67. **Ans. (a, b, e) a. Anaplastic large cell lymphoma; b. Embryonal cell carcinoma; e. Hodgkin's lymphoma**

(Ref: Robbins (SAE) 9th/590,605)

68. **Ans. (a) Hodgkins lymphoma**

(Ref: Robbins 9th/pg 606-611)

69. **Ans. (a, c, d, e) a. Nodular sclerosis; c. Lymphocyte rich; d. Mixed cellularity; e. Lymphocyte depletion**

70. **Ans. (d) Lymphocyte predominance**

(Ref: Robbins 9th/pg 606-611/ 8th pg 616-621)

Lymphocyte predominance is CD20+, CD15-, CD30-;

71. **Ans. (d) Hodgkin Lymphoma: EBV and Rees Sternberg cell** (Ref: Robbins 9th/pg 606-611/ 8th pg 616-621)

Figure show binucleated R-S cell seen in Hodgkin lymphoma

72. **Ans. (d) Overexpression of BCL-6** (Ref: R 9th/pg 606-611)

73. **Ans. (b) Mixed cellularity** (Ref: Robbins 9th/pg 606-611)

74. **Ans. (a) Nodular sclerosis** (Ref: R 9th/pg 606-611)

75. **Ans. (b) Chronic myeloid leukemia**

(Ref: Robbins 9th/pg)

**Sea-blue-colored histiocytes:** Results from increased cell death and subsequent deposition of phospholipids in the macrophages in bone marrow.

Seen in: High rates of intramedullary cell death due to: *lipid storage diseases, myelodysplastic syndromes, lymphomas, chronic myelogenous leukemia, idiopathic thrombocytopenic purpura, autoimmune neutropenia, and  $\beta$ -thalassemia major.*

76. **Ans. (d) Lymphocyte predominance**

(Ref: Robbins 9th/pg 606-611)

77. **Ans. (b) II-B**

(Ref: Robbins 9th/pg 606-611/ 8th pg 616-621)

Involvement of **two or more lymph node** regions on the **same side of diaphragm** alone (II) or localized involvement of an extra-lymphatic organ or site (IIE).

**Presence of (B) symptoms:**<sup>9</sup>

- Unexplained fever,
- Drenching night sweats, and/or
- Unexplained weight loss > 10%

78. **Ans. (a, b, c, d, e); a. Axial lymphadenopathy; b. hepatomegaly is common; c. Contiguous spread of lymph node; d. Can be cured by chemotherapy; e. An arbor classification is useful**

(Ref: Robbins 9th/pg 606-611)

79. **Ans. (b) Lymphocyte poor Hodgkins lymphoma**

(Ref: Robbins 9th/pg 606-611; 8th/pg 616-621)

Reticular variant of Reed Sternberg cell is found in Lymphocyte depletion of Hodgkin's disease

80. **Ans. (c) Lymphocyte predominant Hodgkins lymphoma** (Ref: Robbins 9th/pg 606-611)

81. **Ans. (a) Mixed cellularity Hodgkin lymphoma**

(Ref: Robbins 9th/pg 606-611; 8th/pg 616-621)

In Mixed cellularity type, which is a **Classical Variety** of Hodgkin lymphoma, CD 15+ / CD30+

82. **Ans. (b) Ann Arbor Stage II is involvement of two or more lymph node groups on both sides of the diaphragm**

(Ref: Robbins 9th/pg 606-611; 8th/pg 616-621; Harrison 18th/ Chapter 110)

- Affected lymph nodes become painful with alcohol ingestion: TRUE



- Ann Arbor Stage II is involvement of two or more lymph node groups on both sides of the diaphragm: FALSE, as it is involvement of **two or more lymph node** regions on the **same side of diaphragm**.
- 'B symptoms' are fever, night sweats and  $\geq 10\%$  weight loss in 6 months: TRUE
- ABVD regimen is standard line of treatment for Hodgkin disease: True

**83. Ans. (a) Nodular sclerosis** (Ref: Robbins 9th/pg 606-611)

- Binucleated owl shaped nuclei with clear vacuolated area refers to Lacunar cells.
- **Lacunar cells** are seen in the **nodular sclerosis** subtype of Hodgkin's disease
- Lacunar cells have delicate, folded, or multilobate nuclei and abundant pale cytoplasm that is often disrupted during the cutting of sections, leaving the nucleus sitting in an empty space (lacuna)

**84. Ans. (a) Diffuse large B cell lymphoma**

(Ref: Robbins 9th/pg 595-596; 8th/pg 606-607)

Most common Non Hodgkins lymphoma is Diffuse large B cell lymphoma

*Diffuse large B cell lymphoma (DLBCL)*

**85. Ans. (d) Lymphocytes** (Ref: Robbins 9th/pg 592)

Flow cytometry is done to detect surface molecules like CD markers on:

- **Lymphocytes** for the diagnosis of **chronic lymphoproliferative disorders**
- **Blasts** for diagnosis of **Acute Leukemias**

**86. Ans. (a, d, e); a. Adult T cell lymphoma; d. Diffuse large B cell lymphoma; e. Infectious mononucleosis**

(Ref: Wintrobe's 12th/pg 2312)

Reed-Sternberg cells are **not absolutely specific** for HL

Reed Sternberg like cell are seen in

- Adult T cell lymphoma,
- Diffuse large B cell lymphoma &
- Infectious mononucleosis

**EBV-infected B cells resembling Reed Sternberg cells** are found in the lymph nodes of individuals with **infectious mononucleosis**, strongly suggesting that EBV-encoded proteins play a part in the remarkable metamorphosis of B cells into Reed-Sternberg cells.

**87. Ans. (a) Young age** (Ref: Wintrobe's 12th/pg 2319)

Involvement of stomach (stage IV disease), Lymphocyte depletion, Extranodal metastasis (stage IV disease), Large mediastinal mass & old age are some of the poor prognostic factors in HD

**Poor prognostic factors for Hodgkin's lymphoma**

- Albumin  $<4.0$  g/dl, Hemoglobin  $<10.5$  g/dl, Male sex, 45 years of age or more, Stage IV disease
- Leukocytosis at or above  $15,000/\text{mm}^3$ , Lymphocytopenia (lymphocytes  $\leq 600/\text{mm}^3$  and/or  $<8\%$  of TLC).

**88. Ans. (a) Hodgkin's lymphoma** (Ref: R 9th/pg 606-611)

Ann Arbor staging is used for Hodgkin's lymphoma staging;

**89. Ans. (e) Has good prognosis**

(Ref: Robbins 9th/pg 606-611; 8th/pg 616-621)

- a. False, popcorn cells are present
- b. False, it is CD 15 & CD 30 negative
- c. False, because both B & T lymphocytes, plasma cells & eosinophils are present;
- d. False, because it is EBV negative
- e. True;

**90. Ans. (a) CD 20** (Ref: Robbins 9th/pg 606-611)

**91. Ans. (d) T lymphocyte**

Mycosis fungoides, also known as **Alibert-Bazin syndrome** or granuloma fungoides, is the most common form of cutaneous T-cell lymphoma. It generally affects the skin, but may progress internally over time. Symptoms include rash, tumors, skin lesions, and itchy skin.

**92. Ans. (b) Naive B cell** (Ref: Robbins 9e pg 593)

**93. Ans. (a) SOX11**

WHO 2016 Hematolymphoid textbook.

**Mantle Zone Lymphoma**

- Positive stains - CD5, CD19 (strong), CD20 (strong), cyclin D1/BCL1 (variable nuclear staining since cells are at different stages of cell cycle) also CD22, CD43, CD79a, FMC7, surface IgM or IgD, kappa or lambda, BCL2
- SOX 11 positive in cyclin D1 negative mantle lymphoma.
- Negative stains - CD23, usually CD10; also BCL6, CD11c, TdT, T cell antigens

**94. Ans. (c) EBV positive - DLBL**

(Ref: Hematological malignancies, WHO. Page No. 223)

EBV positive DLBL	EBV positive muco-cutaneous ulcer (WHO 2016 entity)
Immunocompetent	Immuno-compromised
Patients > 50 years old	30-50 years old
The cells are positive for CD20, CD79a, IRF4/MUM1 positive, EBV LMP1 and EBNA negative, CD15 negative. Background are predominantly T lymphocytes	B-cell immunophenotype with uniform expression of CD30, MUM1, PAX5, and OCT-2, and variable CD20, CD45, CD15, CD79a, and BCL-6 expression

We can rule out infectious mononucleosis as monospot test is positive and Hodgkins lymphoma as CD 15 is negative.

**95. Ans. (a, b) a. Diffuse large B-cell lymphoma; b. Hodgkins lymphoma**

PAX-5 is a B-cell differentiation including lymphoblasts (used as a novel pan B-cell marker); also seen in R-S cells in NLPHL (+) vs. classic HL (weak).



96. Ans. (a) **Hairy cell leukemia**

97. Ans. (c, d, e) c. **Gastric lymphoma; d. Thyroid lymphoma; e. Mycosis fungoides**

At least one quarter of non-Hodgkin's lymphomas (NHL) arise from tissue other than lymph nodes and even from sites which normally contain no lymphoid tissue. These forms are referred to as primary extranodal lymphomas. Gastrointestinal localizations, CNS, skin (Mycosis fungoidosis), thyroid, testis. Primary nodal NHL have presentation in lymph node, Waldeyer's ring, spleen or bone marrow

98. Ans. (c) **Hairy cell leukemia**

99. Ans. (d) **Translocation involving MYC gene**

(Ref: Robbins 9th/pg 597; 8th/607)

- Macrophages with abundant clear cytoplasm showing characteristic **"starry sky" pattern** are seen in Burkitt lymphoma

100. Ans. (e) **Most common age group is pediatric**

(Ref: Robbins 9th/pg 593; 8th/pg 603)

Indications to treat in CLL are as follows:

- Rai stage 0–II disease in patients who are symptomatic, have progressive anemia/thrombocytopenia or lymphocytosis
- Rai stages III/IV
- Bulky or progressive lymphadenopathy or splenomegaly
- AIHA/ITP

101. Ans. (a) **Mantle cell lymphoma** (Ref: R 9th/pg 602-603)

102. Ans. (c) **Null cell lymphoma**

(Ref: Wintrobe's 12th/pg 2169)

Features of Anaplastic large cell lymphoma (ALCL):

- ALK rearrangements present
- CD30+
- t(2;5) seen but t(1;2) is most common.
- Because the variants are of **T- or null-cell origin** and occur in a similar age group as the t(2;5): they are also called ALKoma

103. Ans. (c) **Mantle cell lymphoma** (Ref: R 9th/pg 602-603)

104. Ans. (b) **Lymphocyte rich HL** (Ref: R 9th/pg 597)

105. Ans. (d) **Bone - marrow involvement in early phase**

(Ref: Robbins 9th/pg 608)

106. Ans. (a) **All are associated with EBV infection**

(Ref: Robbins 9th/pg 591; 8th/pg 691)

107. Ans. (d) **ALK positive tumors carry worst prognosis**

(Ref: Robbins 9th/pg 605; 8th/615)

Anaplastic Large-Cell Lymphoma (ALK Positive)

- **Defined by the presence of rearrangements in the ALK gene on chromosome 2p23** → break the *ALK* locus → formation of chimeric genes encoding ALK fusion proteins → Activate tyrosine kinases → trigger **JAK/STAT pathway**
- Typically composed of **large anaplastic cells**, with **horseshoe-shaped nuclei & voluminous cytoplasm (so-called hallmark cells)**
- Tumor cells cluster about venules & infiltrate lymphoid sinuses, **mimicing a metastatic carcinoma**.
- **Detection of ALK protein in tumor cells is a reliable indicator of an ALK gene rearrangement** as ALK is not expressed in normal lymphocytes or other lymphomas
- **Have good prognosis**

108. Ans. (b) **Adult T cell leukemia** (Ref: Robbins 9th/pg 605)

109. Ans. (a) **Anaplastic large cell lymphoma**

(Ref: Robbins 9th/pg 591)

110. Ans. (b) **Adult T cell leukemia lymphoma**

(Ref: Robbins 9th/pg 591; 8th/pg 691)

111. Ans. (a) **CD5** (Ref: Robbins 9th/pg 603; 8th/pg 613)

- **MALT lymphomas** express **B-cell antigens (CD19 and CD20) & monotypic surface Ig (IgM without IgD)**.
- **MALTomas** may be **CD43+** but lack other small B-cell lymphoma markers (CD5, CD10, CD23 & cyclin D1)

112. Ans. (c) **Small intestine**

(Ref: WHO Hemato-lymphoid Tumors 4th ed 2008 pg 197-199)

Heavy chain disease (HCD) comprises of 3 rare B-cell neoplasms that produces monoclonal heavy chains (IgG, IgA, IgM) and no light chains.

**Alpha HCD/IgA lymphoma**

- Most common HCD
- Variant of extranodal marginal zone lymphoma
- Also called immunoproliferative small intestine disease (IPSID); Caused by **C. jejuni** infection
- **Site:** GIT: **small intestine (MC)** and mesenteric lymph nodes, gastric and colonic mucosa

113. Ans. (d) **Accessory spleen** (Ref: Robbins 9th/pg)

Accessory spleens or spleniculi are seen in


- Gastrosplenic ligament
- Lienorenal ligament
- Gastrophrenic ligament
- Greater omentum
- Broad ligament of uterus
- Spermatic cord.

114. Ans. (d) **CLL** (Ref: Robbins 9th/pg 593; 8th/pg 603)

CD5 + tumors are CLL and Mantle zone lymphoma.

115. Ans. (b) **Burkitt lymphoma** (Ref: Robbins 9th/pg 597)

116. Ans. (a) **t(11;14)** (Ref: Robbins 9th/pg 597; 8th/pg 607)



117. Ans. (a) **Lymphadenopathy is the most common presentation** (Ref: Robbins 9th/pg 594; 8th/pg 604)

118. Ans. (a, b, c) **a. Most common leukemia in adult; b. Proliferation center is pathognomonic; c. massive splenomegaly** (Ref: Robbins 9th/pg 593)

119. Ans. (a) **Burkitts Lymphoma**

(Ref: 9th/pg 597; 8th/pg 607)

120. Ans. (a) **Anaplastic large cell lymphoma**

(Ref: Robbins 9th/pg 605; 8th/pg 615)

121. Ans. (a) **Hairy cell leukemia** (Ref: R 9th/pg 603-604)

In the given scenario an elderly male presents with Anemia, Splenomegaly & Pancytopenia.

Discussing the options one by one:

- **Hairy cell Leukemia:** Chronic B-cell leukemia characterized by **hairy cells, pancytopenia and splenomegaly**<sup>Q</sup>
- **CML:** Massive splenomegaly with Basophilic Leukocytosis seen
- **Thalassemia:** Jaundice, severe anemia (requiring transfusion), hepatosplenomegaly with leukoerythroblastosis, usually presents in childhood
- **Follicular lymphoma:** Lymphadenopathy ± Leukocytosis

122. Ans. (a) **CD23** (Ref: Robbins 9th/pg 602-603)

- CLL is positive for CD23;

123. Ans. (c) **NK/T cell lymphoma**

(Ref: Wintrobe's 12th/pg 2169)

Nasal NK/T lymphoma may present with facial swelling/destruction, so called **lethal midline granuloma** or **polymorphic reticulosis**. Occurs more commonly in males, with a median age of **50 to 55 years**.

124. Ans. (c) **B-cell NHL** (Ref: Wintrobe's 12th/pg 2154)

- Most common ocular lymphoma is B-cell NHL
- Most **orbital lymphomas** are of **B-cell origin** and are **low-grade**, particularly in the conjunctiva or eyelids, but can be a large B-cell lymphoma in the lacrimal gland or retrobulbar area.

125. Ans. (a) **Mantle cell lymphoma**

(Ref: Robbins 9th/pg 602-603; 8th/pg 612-613)

126. Ans. (b) **Stomach** (Ref: Wintrobe's 12th/pg 2177)

- Commonest site for **extranodal lymphoma** is **Stomach**
- **Gastrointestinal tract** is the **most common site for extranodal NHL (10 to 15% of all NHL)**.
- **Stomach** accounts for **50% of gastrointestinal lymphomas**

127. Ans. (a) **BCL - 2** (Ref: Robbins 9th/pg 594-595)

- **BCL2** antagonizes apoptosis and **promotes survival of follicular lymphoma cells**.
- **BCL** stands for '**B Cell Lymphoma**'

128. Ans. (d) **Chronic lymphocytic leukemia**

(Ref: Robbins 9th/pg 593; 8th/pg 603; Wintrobe's 12th/pg 2214)

- **Unlike other leukemias, there is no firm evidence linking an occupational exposure or radiation with an increased incidence of CLL**

129. Ans. (a) **Anaplastic large cell lymphoma (ALK positive)**

(Ref: Robbins 9th/pg 605; 8th/pg 615)

Histological presence of "**Hallmark Cells**" with horse shoe-like or embryoid like nuclei and voluminous cytoplasm are seen in Anaplastic large cell lymphoma (ALK positive)

130. Ans. (b) **Chronic lymphocytic leukemia**

(Ref: Robbins 9th/pg 593; 8th/pg 603)

**"Smudge cells"**

- Also called '**Basket cells**', '**shadow cells of Gumprecht**'
- Caused by **decrease in Vimentin**
- May predict **good prognosis**
- Seen mainly in **CLL**; Also have been reported in **AML, CML, ALL** & in normal peripheral smear, but rare;

131. Ans. (c) **CD20**

(Ref: Robbins 9th/pg 597; 8th/pg 607)

132. Ans. (b) **Peripheral blood Immunophenotyping**

(Ref: Robbins 9th/pg 593-604; 8th/pg 613-614)

Most likely this is a case of **Non-Hodgkin Lymphoma** with **spill over in the peripheral blood**.

Hence, next investigation that should be done is **Immunophenotyping of Peripheral blood Lymphocytes**

133. Ans. (c) **Leukocyte common antigen**

(Ref: R 9th/pg 590)

Since Lymphoma arises from mature leukocytes, so **Lymphoma cells are positive for Leukocyte common antigen (CD45)**.

134. Ans. (d, e) **d. CD 23 positive; e. Centroblasts frequently seen**

(Ref: Robbins 9th/pg 602-603; 8th/pg 612-613)

Discussing the options about Mantle cell lymphoma, one by one,


- Associated with t(11;14) translocation: TRUE, t(cyclin D1;Ig heavy chain)
- Overexpression of the BCL protein: TRUE, as Bcl2 is overexpressed
- CD 5 positive: TRUE
- **CD 23 positive: FALSE, It is CD 23 -ve**
- **Centroblasts frequently seen: FALSE, Centroblasts & centrocytes are seen in Follicular lymphoma**

135. Ans. (c) **Burkitts Lymphoma** (Ref: Robbins 9th/pg 597)

Lymphoma associated with **translocation of c-myc** is **Burkitt's Lymphoma**

136. Ans. (a) **B cell** (Ref: Wintrobe's 12th/pg 2154)





**137. Ans. (d) Hemoglobin and albumin**

(Ref: Wintrobe's 12th/pg 2154)

**International prognostic index for lymphomas** does not include Hemoglobin and albumin

Adverse Factor	Risk Group
Performance status $\geq 2$	Low
LDH > normal	Low-intermediate
Extra-nodal sites $\geq 2$	High-intermediate
Stage III / IV disease	High
Age > 60 years	High

**138. Ans. (b, c, d, e) b. Primary small intestinal lymphoma are most commonly located in ileum; c. Lymphoma is most common primary malignant neoplasm of spleen; d. Stomach is most common site for extranodal lymphoma; e. MALT lymphoma is associated with H. pylori infection**

(Ref: Wintrobe's 12th/pgs 12th ed -2152, Sternburg 1462-1464)

GIT involvement in NHL

- **Gastrointestinal tract** is the **most common extranodal site** of Lymphoma, at presentation
- Involved in **10 to 15% of adults with NHL**.
- **Stomach** is most frequently involved (option D), followed by the **small intestine, the colon, and esophagus**.
- **H. pylori** play a role in the development of most **MALT lymphomas (option E)**
- **Abdominal pain** is the most common presenting symptom.
- Dyspepsia, nausea & early satiety, suggest stomach involvement.
- Gross appearance of low-grade lymphoma resemble **peptic ulcers**; others have **enlarged mucosal folds; flat and either hyperemic or normal mucosa may also be seen**.
- Patients with rectal involvement usually present with hematochezia or a change in bowel habits
- **Obstruction, intussusception, or perforation** are associated with **aggressive small bowel lymphomas**, particularly Burkitt lymphoma and intestinal T-cell lymphoma.
- **Ileum is the most common site (60%-65%) (option B)** involving **small intestine lymphoma** followed by jejunum (20%-25%), duodenum (6%-8%)
- **Mantle cell lymphoma** presents with **GI symptoms in 20 to 30% of patients & multiple polyposis**
- Lymphomatous polyposis of GIT is not restricted to MCL & is also seen in follicular lymphoma (FL) & MALToma (But this is a rare presentation)
- Abnormal histology in the gastrointestinal tract is found in >80% of MCL patients

**For option C:**

- Most common benign tumors of spleen: **Cavernous hemangioma**
- Most common malignant tumor of spleen: **lymphoma**

**139. Ans. (c, e); c. Vasculitic syndromes; e. Erythema multiforme (Ref: Robbins 9th/pg 603-604; 8th/pg 613-614)**

Features of hairy cell leukemia include splenomegaly, Hepatomegaly & Pancytopenia.

**140. Ans. (a, b) a. Hairy cell leukemia; b. CML**

(Ref: Harrison 18th/Chapter 59)

**Diseases Associated with Massive Splenomegaly**

- |   |  |
|---|--|
| <ul style="list-style-type: none"> <li>• <b>Chronic myeloid leukemia</b></li> <li>• Lymphomas</li> <li>• <b>Hairy cell leukemia</b></li> <li>• Myelofibrosis with myeloid metaplasia</li> <li>• <b>Polycythemia vera</b></li> </ul> | <ul style="list-style-type: none"> <li>• Gaucher's disease</li> <li>• Chronic lymphocytic leukemia</li> <li>• Sarcoidosis</li> <li>• Autoimmune hemolytic anemia</li> <li>• Diffuse splenic hemangiomatosis</li> </ul> |
|---|--|

**141. Ans. (c) Atypical mycobacteria**

(Ref: Robbins 9th/pg 603-604; 8th/pg 613-614)

Hairy cell leukemia is associated with Atypical Mycobacteria.

**142. Ans. (b, e) b. Polycythemia; e. IgG paraprotein**

**143. Ans. (a) t (11 : 14)**

**144. Ans. (c) Post germinal centre B cell**

**145. Ans. (a) Multiple myeloma (Ref: Robbins 9th ed p 599)**

Normal plasma cell has characteristic eccentrically placed nucleus. Malignant plasma cells have a perinuclear clearing due to a prominent Golgi apparatus and an eccentrically placed nucleus. Relatively normal-appearing plasma cells, plasma-blasts with vesicular nuclear chromatin and a prominent single nucleolus, or bizarre, multinucleated cells may predominate. Other cytologic variants stem from the dysregulated synthesis and secretion of Ig, which often leads to intracellular accumulation of intact or partially degraded protein. Such variants include flame cells with fiery red cytoplasm

**146. Ans. (a, b, c, d, e) a. Cast nephropathy; b. amyloidosis; c. Cryoglobulinemia; d. Interstitial nephritis; e. nephrocalcinosis (Ref: Wintrobe 13/p 2377)**

Approximately 5% of myeloma gammaglobulins show reversible precipitation in the cold, so-called cryoglobulins, forming either a flocculent precipitate or a gel-like coagulum when the serum is cooled. The most common findings on autopsy include tubular atrophy and fibrosis (77%), tubular hyaline casts (62%), tubular epithelial giant cell reaction (48%), and nephrocalcinosis. Acute and chronic pyelonephritis can also be seen.

**147. Ans. (b) Multiple myeloma (Ref: R 9/ 595)**

Key to the image: 1 → Increase in plasma cells in marrow, 2 → Kidney biopsy with pink deposits in glomerulus, 3 → Fibrils in electron microscope. So the diagnosis is **Multiple myeloma, probably AL amyloidosis of kidney**



**148. Ans. (b) Waldenstrom Macroglobulinemia**

MYD88 (L265P) mutation is detectable in all patients with Waldenström's macroglobulinemia, therefore representing a hallmark of the disease.

**149. Ans. (c) Russell bodies** (Ref: Robbins 9th/pg 598-602)

The globular inclusions in the plasma cells are referred to as **Russell bodies** (if cytoplasmic) or **Dutcher bodies** (if nuclear).

**150. Ans. (b) Plasma Cells** (Ref: 9th/pg 598-602/8th 608-612)

**151. Ans. (d) Serum b2-microglobulin**

(Ref: Robbins 9th/pg 598-602/8th 608-612)

$\beta_2$ -M concentration is the strongest and most reliable prognostic factor for multiple myeloma. It depends not only on tumor burden but also on renal function. Elevated  $\beta_2$ -M values predict early death

**152. Ans. (d) Dexamethasone**

(Ref: Robbins 9th/pg 598-602)

**153. Ans. (d) Does not progress to multiple myeloma**

(Ref: Robbins 9th/pg 598-602/8th 608-612)

1% monoclonal gammopathy of unknown significance progress to multiple myeloma /year

**154. Ans. (a) IgM** (Ref: Robbins 9th/pg 598-602)

**155. Ans. (d) Bence jones proteinuria** (Ref: R 9th/pg 598-602)

**156. Ans. (a) Endocrinopathy** (Ref: Robbins 9th/pg 598-602)

**157. Ans. (d) Days to week** (Ref: Robbins 9th/pg 598-602)

Life span of plasma cells is almost same as that of Memory lymphocytes (days to months or even years)

**158. Ans. (d) Plasma cell** (Ref: Robbins 9th/pg 598-602)

Multiple myeloma is a **malignant proliferation of plasma cells** derived from a **single clone**.

**159. Ans. (c) Multiple myeloma** (Ref: Robbins 9th/pg 598-602)

In the given scenario, patient has **bone pain, destructive bony lesions** on X Ray, **hypercalcemia**, **M spike** and **35% plasma cells in Bone marrow** → Diagnostic of Multiple Myeloma

**160. Ans. (a) Multiple myeloma** (Ref: Robbins 9th/pg 598-602)

Serum  $\beta_2$  microglobulin level  $<3.5\text{mg/L}$  indicates **good prognosis**, in multiple myeloma.

**161. Ans. (b) IL-6** (Ref: Robbins 9th/pg 598-602)

IL-6 helps in **survival and proliferation** of myeloma cells

**162. Ans. (d) Dystrophic calcification** (Ref: R 9th/pg 598-602)

Calcification in multiple myeloma is due to hypercalcemia (metastatic calcification) and not dystrophic calcification.

**163. Ans. (b) Light chain globulins**

(Ref: R 9th/pg 598-602)

Excretion of **light chains** in the **urine** has been referred to as **Bence Jones proteinuria**.

Light chains includes  $\kappa$  and  $\lambda$  (kappa and lambda)

**164. Ans. (d) Lymphoplasmacytic lymphoma**

(Ref: Robbins 9th/pg 598-602; 8th/pg 608-612)

**Waldenstrom Macroglobulinemia**

**Indolent lymphoproliferative disorder** characterized by:

- Lymphoplasmacytic cell proliferation in marrow (**Lymphoplasmacytic lymphoma**) with secretion of **IgM**
- **Clinical Features** Anemia, lymphadenopathy, Hepatosplenomegaly and hyperviscosity
- CD138+, cyIgM+, CD19+

**165. Ans. (c) Hypercalcemia**

(Ref: Robbins 9th/pg 598-602; 8th/pg 608-612)

**166. Ans. (c) > 10% plasmacytosis**

(Ref: Robbins 9th/pg 598-602; 8th/pg 608-612)

Multiple myeloma is diagnosed by **> 10% plasmacytosis**

**167. Ans. (b) IgM**

(Ref: Robbins 9th/pg 598-602)

Lymphoplasmacytic cell proliferation in marrow (**Lymphoplasmacytic lymphoma**) with secretion of **IgM**

**168. Ans. (c) Plasmacytoma on biopsy**

(Ref: Wintrobe's 12th/pg 2377)

**Durie and Salmon criteria for diagnosis of Multiple myeloma.**

A minimum of 1 major and 1 minor criterion needed, although (1) + (a) is not sufficient, or 3 minor criteria that must include (a) and (b).

However, **International myeloma working group 2011** has **revised** the criteria as mentioned in pretexts.

**169. Ans. (d) Elevated alkaline phosphatase**

(Ref: Robbins 9th/pg 598-602; 8th/pg 608-612)

**Alkaline Phosphatase levels in multiple myeloma is normal** and not raised, as there is no bone formation and only bone lysis.

**170. Ans. (a) P190 has aggressive course**

CML can have 3 types of transcripts.

- P190–Poor prognosis Seen in ALL
- P210–Seen in CML
- P230–Seen in CNL good prognosis

**171. Ans. (a, d) a. Richter transformation; d. MHC non-expression is way to escape immunity but of to mark one a > d.**



**172. Ans. (d) Erythropoietin levels**

In this case the d/d of the clinical details given is polycythemia. Since he is from hilly area, it may be due to secondary causes. To rule out primary (low epo) vs secondary cause (high epo), erythropoietin must be the next logical step.

**173. Ans. (c) Focal cellular marrow with hypocellular areas and atypical megakaryocytes.**

(Ref: WHO Textbook of hematology 2016/ p 47)

**174. Ans. (a) Philadelphia chromosome is negative**

(Ref: Robbins 9th/pg 616-617)

**175. Ans. (b) CML**

(Ref: Robbins 9th/pg 616- 618; 8th 626-628)

**176. Ans. (a) Acrocentric chromosome involved**

(Ref: Robbins 9th/pg 160, Refer Genetics chapter)

**177. Ans. (a) Leukoerythroblastic blood picture**

(Ref: Robbins 9th/pg 614- 615; 8th 624-625)

Myelodysplastic syndrome usually results in pancytopenia rather than leukoerythroblastic blood picture.

**178. Ans. (d) Leukocytosis (Ref: Robbins 9th/pg 618- 619)**

Polycythemia vera results in Panmyelosis (increase in all myeloid components including leukocytes)

**179. Ans. (b) Florescent in situ hybridization**

(Ref: Robbins 9th/pg 170)

In PCV: Reddening, swelling, and pain in the digits (erythromelalgia) may occur and are typically associated with extreme platelet elevations.

**180. Ans. (d) Blast crisis phase of CML**

(Ref: Robbins 9th/pg 616- 618)

**181. Ans. (b) Thrombocytosis**

(Ref: Robbins 9th/pg 618)

**182. Ans. (b) Imatinib mesylate**

(Ref: Robbins 9th/pg 617)

**183. Ans. (c) Abnormal oxygen saturation**

(Ref: Robbins 9th/pg 618)

**184. Ans. (d) Spontaneous severe infection**

(Ref: Robbins 9th/pg 618)

**185. Ans. (d) Marrow fibrosis (Ref: Robbins 9th/pg 620/630)**

**186. Ans. (b) Primary myelofibrosis**

(Ref: Robbins 9th/pg 620/630)

The most characteristic peripheral blood finding in PMF is myelophthisis, defined by the presence of **leukoerythroblastosis** (presence of nucleated red blood cells, metamyelocytes, myelocytes, myeloblasts, and megakaryocytes) and dacryocytosis (**Tear drop RBCs**)

**187. Ans. (b) Increased neutrophil alkaline phosphatase**

(Ref: Robbins 9th/pg 617)

**188. Ans. (d) c-kit point mutation (Ref: R 9th/pg 616-620)**

Myeloproliferative disorders include:

Disease	Tyrosine kinase involvement
Chronic myeloid leukemia	ABL1
Myeloid and lymphoid neoplasms with eosinophilia	PDGFRA/Bor FGFR4
Polycythemia vera	JAK2 V617 F, JAK EXON 12
Primary myelofibrosis	JAK2 V 6174 , MPL W515 K/L
Essential thrombocythemia	JAK2 V617 F , MPL W 515 K/L

**189. Ans. (d) Myelodysplastic syndrome**

(Ref: R 9th/pg 614)

**190. Ans. (b) Polycythemia Vera**

(Ref: Wintrobe's 12th/pg 180-190)

**191. Ans. (a) Myelodysplastic Syndrome**

(Ref: Robbins 9th/pg 614- 615)

**192. Ans. (d) Myelofibrosis (Ref: Robbins 9th/pg 620)**

**193. Ans. (b) FISH (Ref: Robbins 9th/pg 616-618)**

- **Flow cytometry:** is used for **immunophenotyping** in Leukemias & Lymphomas
- **FISH-** used for detecting **submicroscopic chromosomal aberrations of BCR-ABL fusion**
- **Karyotyping:** Used for detecting **chromosomal defects of at least 5 mb size** (low sensitivity)
- **RT-PCR:** used for **quantitative DNA estimation**

**194. Ans. (b) Decrease LAP score (Ref: R 9th/pg 618-619)**

Increased vit B12- is seen in all myeloproliferative neoplasms including polycythemia vera due to increased levels of transcobalamin II

**195. Ans. (a) Dry tap (hypocellular) (Ref: Robbins 9th/pg 620)**

In Myelofibrosis bone marrow is replaced by fibrosis, giving a dry tap on bone marrow aspiration

**196. Ans. (c) Myelodysplastic syndrome**

(Ref: R 9th/pg 614-615)

Pseudo-Pelger-Huet neutrophils: bilobed hypogranular dysplastic neutrophils<sup>Q</sup> seen in Myelodysplastic syndromes.



**197. Ans. (d) Lung carcinoma**

(Ref: Robbins 9th/pg 618-619)

Polycythemia is not caused by Lung Carcinoma.

**198. Ans. (b) Acute myeloid leukemia**

(Ref: Robbins 9th/pg 616; 8th/pg 626)

AML is not a myeloproliferative disease

**199. Ans. (b) The fusion gene bcr-abl forms a protein with tyrosine kinase activity** (Ref: Robbins 9th/pg 616-618)

- FALSE, as **ABL gene is on chr 9q** while BCR is on chr 22q
- The **fusion gene bcr-abl forms a protein with tyrosine kinase activity**: TRUE
- FALSE; **Massive splenomegaly is a feature of CML**
- FALSE; as **response to Imatinib is good in patients with Philadelphia chromosome +ve**

**200. Ans. (d) Essential thrombocytopenia**

(Ref: Robbins 9th/pg 616-618; 8th/pg 626-628)

- Myeloid metaplasia is other name of myelofibrosis.
- Essential thrombocytosis is CMPD (not thrombocytopenia)

**201. Ans. (b) Protein tyrosine kinase inhibitor are the drug of choice** (Ref: Robbins 9th/pg 616-618)

- FALSE; as translocation **is between long arms of chr 9 & 22**
- TRUE; **Imatinib- a tyrosine kinase inhibitor is the drug of choice**
- FALSE; **Most commonly presents in the chronic phase**
- FALSE; **Commonest tumor in children is ALL, followed by CNS tumors.**

**202. Ans. (d) Acute Megakaryocytic Leukemia**

(Ref: Robbins 9th/pg 620; 8th/pg 630)

**Myelofibrosis leading to a dry tap** on bone marrow aspiration is seen with **Acute Megakaryocytic Leukemia**.

Conditions in which Myelofibrosis is seen are:

- Primary Myelofibrosis (PMF)
- Secondary Myelofibrosis:**
- Myeloproliferative Neoplasms- CML, PCV, ET
- Myeloid Neoplasms- AML M7, MDS
- Hairy cell Leukemia, Hodgkin's, Multiple Myeloma
- Metastasis
- Granulomatous diseases- TB, Sarcoidosis
- Autoimmune disease- SLE, Systemic sclerosis

- Others-Paget's disease, Renal osteodystrophy, Radiation, HyperPTH

**203. Ans. (a) Mitochondria**

(Ref: Robbins 9th/pg 614-615)

In **myelodysplastic syndrome**, ring sideroblast is seen in **Mitochondria**

**204. Ans. (b) Gamma heavy chain disease**

Franklin's disease (**gamma heavy chain disease**) It is a very rare B-cell lymphoplasma cell proliferative **disorder** which may be associated with autoimmune **diseases** and infection is a common characteristic of the **disease**. It is characterized by lymphadenopathy, fever, anemia, palatal edema, malaise, hepatosplenomegaly, and weakness.

**205. Ans. (d) Langerhans cell**

(Ref: Robbins 9th/pg 631-632)

**206. Ans. (a) Birbeck's granules**

(Ref: Robbins 9th/pg 531)

**Birbeck granules**

- Also known as Langerhans bodies or **X granules**
- Seen on Electron microscopy
- Rod-shaped organelles with a central striation and terminal vesicular dilation, giving them a 'tennis racket' appearance.

**207. Ans. (d) Birbeck's granules**

(Ref: Wintrobe's 12th/pg 1573)

**208. Ans. (d) Eosinophilic granuloma**

(Ref: 9th/pg 621-622)

Localised langerhans cells histiocytosis affecting head & neck is **Eosinophilic granuloma**

**209. Ans. (c) CD 1a**

(Ref: Robbins 9th/pg 621-622)

CD marker for Langerhans cell histiocytosis is CD 1a

**210. Ans. (b) Pneumothorax in 10% cases**

(Ref: Wintrobe's 12th/pg pg 1577)

- Pneumothorax is seen in 25-40% cases of pulmonary LCH**



[illegible]This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Red Blood Cells and its Disorders

## Key Points


- » **Hemoglobin** can be first visualized at **polychromatic (Intermediate) erythroblast<sup>o</sup>** stage
- » **Early recovery in acute blood loss is marked by thrombocytosis**
- » Decreased free serum haptoglobin, Hemoglobinuria, Methemoglobinuria, Hemosiderinuria are hallmark of intravascular hemolysis
- » **Most common** mutation in Hereditary spherocytosis are **AnkyrinQ > band 3 > Spectrin and band 4.2**
- » Both **intravascular and extravascular** episodic hemolysis is seen in<sup>o</sup> in G6PD-deficient individuals
- » **Sickle cell anemia results from point mutation in 6<sup>th</sup> codon of  $\beta$ -globin<sup>o</sup>** → replacement of a **glutamate** residue with a **valine** residue
- »  **$\alpha$  and  $\beta$  Thalassemia results from decreased production of  $\alpha$  and  $\beta$  chains respectively**
- » PNH results from acquired somatic clonal mutation of **PIG-A gene<sup>o</sup>** resulting in loss of **GPI**
- » **Divalent metal transporter type 1 (DMT-1) helps in absorption of Iron from Intestine but is not specific for Iron absorption**
- » **Hypersegmented neutrophils is the earliest abnormality to appear in macrocytic anemia**
- » **Aplastic anemia** refers to a **marrow hypoplasia** and **pancytopenia**

## Key Recent Updates

- » Eosin 5-maleimide (EMA) binding test is recent screening test for diagnosis of HS
- » FLAER is an Alexa488-labeled inactive variant of aerolysin to diagnose and monitor patients with PNH.



## DEVELOPMENT OF RBCs



- Reticulocytes takes **4–5 days (1–3 days in bone marrow and 1–2 days in peripheral circulation)**<sup>Q</sup> to **mature** into RBCs
- **Ferritin** molecules can be seen at **proerythroblast stage**<sup>Q</sup> by **electron microscopy**
- **Hemoglobin** can be first visualized at **polychromatic (Intermediate) erythroblast**<sup>Q</sup> stage



## REGULATION OF ERYTHROPOIESIS

Hypoxia → Kidney → Erythropoietin → Bone marrow → Manufactures and releases RBCs

## RBC INDICES

Mean Cell Volume (MCV) 80–100 m (fl)	Mean Cell Hemoglobin (MCH) 30 ± 3 (pg)	Mean Cell Hemoglobin Concentration (MCHC) 34 ± 2 (g/d)	Red Cell Distribution Width (RDW) 12–16
Average volume of a red cell	Average content (mass) of hemoglobin per red cell, $MCH = Hb/RBC \text{ count}^Q$	Average concentration of hemoglobin in a given volume of packed red cells, $MCHC = Hb/MCV^Q$	<b>Coefficient of variation</b> of red cell volume <sup>Q</sup>

- **Hematocrit or Packed cell volume:** ratio of the volume of RBCs to total volume of blood
- **Hct or PCV =  $MCV \times RBC \text{ concentration}$**



## ESR

- **Definition:**
  - Measurement of **sedimentation of RBCs** in diluted blood (EDTA or Citrate) after standing for **1 hr** in an open-ended glass tube of 30 cm length mounted vertically on a stand.



- **Characteristics:**
  - It is **slower to respond** to acute disease activity
  - It is **insensitive to small changes** in disease activity.
  - It is **less specific than CRP** because it is influenced by immunoglobulins & Anemia.
- **Stages of sedimentation:**
  - Formation of **rouleaux**
  - Period of **fast settling**
  - Period of **packing of the rouleaux** at the bottom of the tube
- **Normal value:**
  - Men: < 15 mm/hr
- **Very low ESR:** (0–1 mm)
  - Polycythemia, hypo- or afibrinogenemia, congestive cardiac failure & abnormalities of red cells such as poikilocytosis, spherocytosis, or sickle cells.
- **Very high ESR** (>100 mm)
  - Tuberculosis, Hodgkin's disease, multiple myeloma, chronic infective or inflammatory conditions
- **Factors influencing ESR**

Increased ESR	Lower ESR
<ul style="list-style-type: none"> <li>• Old age</li> <li>• Female</li> <li>• <b>Pregnancy</b></li> <li>• <b>Anemia</b></li> <li>• Paraprotein (Multiple Myeloma)</li> <li>• Hypergammaglobulinemia</li> <li>• Macrocytosis</li> <li>• Elevated fibrinogen (infection, inflammation malignancy)</li> <li>• Technical factors: dilution, high temperature</li> </ul>	<ul style="list-style-type: none"> <li>• Extreme leucocytosis</li> <li>• <b>Polycythemia</b></li> <li>• <b>Spherocytosis</b>, microcytosis</li> <li>• Hyperviscosity</li> <li>• Low Protein; fibrinogen, gammaglobulins</li> <li>• Technical factors; dilution, clotted sample</li> <li>• <b>Afibrinogenemia</b></li> </ul>
	No effect
	<ul style="list-style-type: none"> <li>• Obesity</li> <li>• Body temperature</li> <li>• Recent meal</li> <li>• Aspirin, NSAIDs</li> </ul>


**Erythrocyte Sedimentation Rate is zero in Afibrinogenemia**

## ANEMIA

**Definition:** Functionally defined as **an insufficient RBC mass** to adequately **deliver oxygen** to peripheral tissues

### WHO Criteria

Category	Hb (g/dl) <sup>o</sup> cut-off for Anemia	Mean Normal Hb (g/d)
<b>Adult male</b>	< 13	14.5
<b>Adult female</b>	< 12	13.5
<b>Pregnant female</b>	< 11	12.5
<b>Child &lt; 6 yr</b>	< 11	12



## Anemia Due to Blood Loss

### Acute Blood loss

- **Etiology:**
  - **External** (trauma, or obstetric hemorrhage)
  - **Internal** (e.g., from bleeding in the gastrointestinal tract, rupture of the spleen, rupture of an ectopic pregnancy, subarachnoid hemorrhage)
- **Pathophysiology:**
  - Loss of intravascular volume → Intravascular shift of water from the interstitial fluid compartment → **Hemodilution** → **Lowering of hematocrit**
- **Features:**
  - **Early recovery is marked by thrombocytosis<sup>o</sup>**
  - On day 5- **reticulocytosis starts<sup>o</sup>**
  - **Massive bleeding- granulocytosis** (leukocytosis)<sup>o</sup>
  - Day 7: reticulocytosis (10–15%)<sup>o</sup>
  - **MCV increases<sup>o</sup>** as newly produced RBC's are larger


### Chronic Blood Loss

Anemia is the only manifestation

## Hemolytic Anemias: Due to increased RBC Destruction

Characterized by:

- **Shortened RBC life span<sup>o</sup>** (normal = 120 days)<sup>o</sup>
- **Increase in erythropoiesis<sup>o</sup>**
- Accumulation of **hemoglobin degradation<sup>o</sup>** products







### Difference between Intra- and Extra-vascular Hemolysis


	Extravascular hemolysis	Intravascular hemolysis
<b>Site of hemolysis</b>	Mononuclear phagocytes <sup>a</sup> RE system (spleen <sup>a</sup> )	Within the circulation <sup>a</sup>
<b>Diseases causing hemolysis</b>	Thalassemia, <sup>a</sup> sickle cell anemia <sup>a</sup>	PNH, <sup>a</sup> G6PD deficiency <sup>a</sup> angioaphatic
<b>Serum haptoglobin</b>	Normal	Decreased <sup>a</sup>
<b>Hemoglobinuria</b>	Not seen	Positive <sup>a</sup>
<b>Methemoglobinuria</b>	Not seen	Positive <sup>a</sup>
<b>Hemosiderinuria</b>	Not seen	Positive <sup>a</sup>
<b>Serum unconjugated bilirubin</b>	Moderately elevated	Mildly elevated
<b>Splenomegaly</b>	Usual <sup>a</sup>	Uncommon

### Classification of Hemolytic Anemias

Hereditary	Acquired
<b>Membrane defects</b> <ul style="list-style-type: none"> <li>Hereditary spherocytosis</li> <li>Hereditary elliptocytosis</li> </ul> <b>Enzyme defects</b> <ul style="list-style-type: none"> <li>G6PD deficiency</li> <li>Pyruvate kinase deficiency</li> </ul> <b>Hemoglobin</b> <p>Genetic abnormalities (HbS, HbC, unstable)</p>	<b>Immune</b> <ul style="list-style-type: none"> <li><b>Autoimmune:</b> Warm or Cold antibody type</li> <li><b>Alloimmune</b> <ul style="list-style-type: none"> <li>Hemolytic transfusion reaction</li> <li>Hemolytic disease of the newborn</li> <li>Allografts, especially stem cell transplantation</li> </ul> </li> </ul> <ul style="list-style-type: none"> <li>Drug associated</li> <li><b>Red cell fragmentation syndromes</b></li> <li><b>March hemoglobinuria</b></li> <li><b>Infections:</b> Malaria, Clostridia</li> <li><b>Chemical and physical agents:</b> Drugs, industrial substances, burns</li> <li><b>Secondary:</b> Liver and renal disease</li> <li><b>Paroxysmal nocturnal hemoglobinuria</b></li> </ul>


### Hereditary Spherocytosis (HS)

- **Definition:**
  - **Inherited** intrinsic defect in the red cell membrane leading to **spherocytes**<sup>a</sup>
- **Inheritance pattern:**
  - **Autosomal Dominant**-75%<sup>a</sup>
  - **Compound heterozygous**-25%<sup>a</sup> (more severe, presentation at birth)



- **Spectrin** ( $\alpha/\beta$ )-major protein forming tetramers<sup>a</sup>
- **Actin**-binds spectrin tetramers<sup>a</sup>
- **Protein 4.1**- binds **spectrin** to inner surface of RBC membrane
- **Band 3**- RBC 's transmembrane ion transporter<sup>a</sup>
- **Ankyrin**-bridge between Spectrin and Band 3 helped by **Pallidin (band 4.2)**<sup>a</sup>

### Pathogenesis



### Clinical feature

- **Triad of** Jaundice + hemolytic anemia + splenomegaly (with a **family history of gall stones** )
- **Pigment type**<sup>a</sup> of gall stones are common



## ■ Diagnosis

- **PS:** Micro-spherocytosis of uniform size ( **variable size<sup>Q</sup>** in **Immune Hemolytic Anemia**).
- **Decreased MCV, Increased MCHC<sup>Q</sup>, Increased 'glycerol lysis test'<sup>Q</sup>**
- **Increased fragility** on Osmotic fragility testing (**OFT**): shift of curve to **right<sup>Q</sup>**
  - **24-hr incubated OFT** is the **most sensitive** test to diagnose HS<sup>Q</sup>
  - Diagnosis of choice -eosin-5-maleimide flow cytometric test



## High Yield Facts

- Life span of RBC in HS is **20 days<sup>Q</sup>** against a normal of 120 days
- **Most common** mutation in HS: **Ankyrin<sup>Q</sup> > band 3 > Spectrin and band 4.2<sup>Q</sup>**
- **α-spectrin** mutations (Autosomal recessive) leads to **most severe** phenotypes<sup>Q</sup>
- In **Spectrin** mutations, the most common RBC abnormality is **Hereditary Elliptocytosis > spherocytosis<sup>Q</sup>**



## Hemolytic Disease Due to Red Cell Enzyme Defects

### Four Types

- Glucose-6-Phosphate Dehydrogenase (**G6PD**) deficiency (**most common**)<sup>Q</sup>, Pyruvate Kinase deficiency, Pyrimidine-5' nucleotidase deficiency, Hereditary methemoglobinemia

### Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency

- **Inheritance:**
  - **X-linked recessive** trait (Chr Xq28)<sup>Q</sup>
- **Epidemiology:**
  - **Highest prevalence** in **Kurdish Jews** > African
- **Pathogenesis:**
  - **Mechanism:** **Hemolysis** occur due to **oxidative stress<sup>Q</sup>** due to **deficient G6PD** activity
  - **Defective pathway:** hexose monophosphate (**HMP**) shunt → leads to **deficient NADPH<sup>Q</sup>**
  - Normally NADPH reduces **oxidized glutathione (GSSG)** to **reduced glutathione (GSH)**, which **neutralizes H<sub>2</sub>O<sub>2</sub>**



## ■ Variants of G6PD

- Most common **normal variant**, **G6PD B+<sup>Q</sup>**
- **Most severe variants:** **G6PD A- and G6PD Mediterranean, Canton<sup>Q</sup>**

## ■ Inciting events

- Refer Ans

## ■ Lab Diagnosis

### Peripheral smear:

- **Heinz body on supravital staining<sup>Q</sup>**
- **Anisopoikilocytosis** with polychromasia<sup>Q</sup>
- **"Bite cells"**<sup>Q</sup>
- **Microspherocytes**
- **Increased Reticulocyte %<sup>Q</sup>**

### Screening assay:<sup>Q</sup>

- Methemoglobin reduction test
- Fluorescent spot test

### Diagnostic assay:<sup>Q</sup>

- Quantitative G-6PD enzyme assay by **electrophoresis**
- DNA analysis by PCR



## High Yield Facts

- **G6PD deficiency causes self-limited hemolysis**, which stops with release of younger G6PD abundant RBC's (**even in presence of offending drug**)<sup>Q</sup>
- Both **intravascular and extravascular** hemolysis<sup>Q</sup> in G6PD-deficient individuals.
- G6PD deficient individuals are **protected** against **malaria<sup>Q</sup>**
- Most frequently implicated food is the **fava bean**, which generates oxidants when metabolized called **"Favism"**<sup>Q</sup>
- WHO classification of G6PD is **Class I (most severe)** to **Class V (least severe)**<sup>Q</sup>



## DISORDERS OF HEMOGLOBIN

**Structure of Hemoglobin:** Composed of **two  $\alpha$ -** and **two non- $\alpha$ -globin chains**, each associated with a heme group

- Two sets of genes for the  $\alpha$  chains: **chr 16<sup>q</sup>**
- Two pairs of genes for  $\beta, \gamma, \delta$  chains: **chr 11<sup>q</sup>**

### Hemoglobins (embryonic)


- Gower 1 ( $\zeta_2\epsilon_2$ )
- Portland ( $\zeta_2\gamma_2$ )
- Gower 2 ( $\alpha_2\epsilon_2$ )

### Hemoglobins (at Birth)

- Hb F  $\alpha_2\gamma_2$  (75)%
- Hb A  $\alpha_2\beta_2$  (25)%

### Hemoglobins (adults)

- Hb A  $\alpha_2\gamma_2$  (96.5-97.5)%
- Hb A  $\alpha_2\delta_2$  (2.5-3.5)%
- Hb F  $\alpha_2\gamma_2$  (<1)%





## Sickle Cell Anemia

### Defect:

- Point mutation** in 6<sup>th</sup> codon of  $\beta$ -globin<sup>q</sup> → replacement of a **glutamate** residue with a **valine** residue.<sup>q</sup>
- Production of **HbS** with abnormal physiochemical properties that promotes the **polymerization** of deoxygenated hemoglobin.

### Pathogenesis:



### Factors affecting Sickling:

Factors INCREASING Sickling	Factors DECREASING Sickling
<ul style="list-style-type: none"> <li><b>Dehydration<sup>q</sup></b></li> <li>Increase in MCHC<sup>q</sup></li> <li>Decrease in pH<sup>q</sup>, Increase in ionic strength</li> <li>De-oxygenation</li> <li>Inflammation</li> </ul>	<ul style="list-style-type: none"> <li>Other Hemoglobins like: HbA, HbF*<sup>q</sup></li> <li><b>Concomitant alpha thalassemia<sup>q</sup></b></li> <li>Re-oxygenation</li> </ul>


### Clinical features

- Hand-foot syndrome or dactylitis<sup>q</sup>**
- Priapism and erectile dysfunction<sup>q</sup>**
- Stroke and retinopathy<sup>q</sup>**
- Cardiomegaly is a feature**
- Autosplenectomy<sup>q</sup>**: multiple infarcts in splenic artery causes spleen to be reduced to fibrous tissue (splenic atrophy → **spleen may become nonpalpable**)<sup>q</sup>
- Chronic hemolysis**: impaired growth and development.
- Renal involvement**: Papillary necrosis & **hyposthenuria<sup>q</sup>** (inability to concentrate urine)
- Infection
- Vaso-occlusive crises/Acute painful crises
- Acute chest syndrome<sup>q</sup>
- Sequestration crises
- Aplastic crises

### Diagnosis

- PS**: Irreversibly **sickle RBCs** and **target cells** (increased after autosplenectomy). **Howell-Jolly bodies** if asplenia.<sup>q</sup>
- Sickling test**: Mixing a blood sample with an oxygen consuming reagent, such as **metabisulfite** or **dithionite** induces sickling of RBCs, if HbS is present.<sup>q</sup>
- Spleen biopsy**: **Gamma gandy bodies<sup>q</sup>**
- X-ray**: "**Crew-cut**" appearance of skull, **fish-mouth vertebra<sup>q</sup>**
- Hemoglobin electrophoresis**
- Hb-HPLC** (High Performance Liquid Chromatography): **investigation of choice<sup>q</sup>**
- Prenatal diagnosis**: by analysis of fetal DNA obtained by amniocentesis or chorionic biopsy.





## High Yield Facts

### Hemoglobins with abnormal solubility

Abnormal Hb	Residue	Mutation	Molecular pathology
HbS	β6	Glu → Val	Polymenization
HbC	β6	Glu → Lys	Crystallization
HbD-Punjab	β121	Glu → Gln	Increases polymer is S/D heterozygote


Severity of the hemolysis<sup>Q</sup> correlates with the percentage of irreversibly sickled cells

- Sickle RBCs are **mechanically fragile**, leading to some **intravascular hemolysis**<sup>Q</sup>
- Sickle cell anemia, G6PD deficiency, Thalassemias, absence of Duffy antigen on RBCs protect against malaria<sup>Q</sup>
- HbSC disease is milder than sickle cell disease
- Sickle cell anemia usually **presents after first 6 months** of life
- **Splenomegaly** is first noted **after 6 months** of age
- **Gamma Gandy bodies** contains fibrous tissue, foci of fibrosis with Iron or Calcium salts
- **Gamma Gandy bodies** seen in **portal hypertension, Sickle cell anemia, Splenic congestion, CML** and some cases of leukemia and lymphoma


## Thalassemia

- **Definition**
  - A heterogeneous group of disorders caused by **inherited mutations** that **decrease the synthesis** of globin chains.
- **Distribution**
  - Worldwide: **Mediterranean**, Africa, South east Asia.
  - In India- **Sindh<sup>Q</sup>, Punjab, Gujrat and Bengal**
- **Classification (PS)**
  - **α Thalassemia: decreased production of α chains**
  - **β Thalassemia: decreased production of β chains**
  - Others: δβ-Thalassemias, γδβ-thalassemias, αβ-thalassemias
- **Diagnosis**
  - Moderate to severe **microcytic hypochromic anemia with aniso-poikilocytosis**
  - **Polychromasia, Target RBCs, Basophilic stippling, Howell jolly bodies<sup>Q</sup>**
  - Reticulocyte count is **increased but <5%<sup>Q</sup>** due to **ineffective erythropoiesis**
- **NESTROFT: Naked Eye Single Tube Red cell Osmotic Fragility Test<sup>Q</sup>**

- Assesses osmotic fragility of RBCs at a single concentration of buffered saline (**0.36% in single tube**)<sup>Q</sup>



Thalassemia: anisopoikilocytosis with target RBCs (A)



Basophilic stippling

### Screening test

**Mentzer Index<sup>Q</sup> = Mean corpuscular volume (MCV)/RBC count**  
This test helps in differentiating iron deficiency from thalassemia

In Thalassemia trait	In Iron deficiency Anemia
Mentzer's index < 13 <sup>Q</sup> (RBC count is normal with a low MCV)	Mentzer's index > 13 <sup>Q</sup> as (RBC count as well as MCV is low)

- **Diagnosis of choice:** HPLC; High performance liquid, and chromatography/Electrophoresis






## $\alpha$ -Thalassemia

- Normally 4  $\alpha$ -genes<sup>Q</sup> synthesize 2  $\alpha$  chains:  $\alpha$ ,  $\alpha/\alpha$ ,  $\alpha$
- Most common cause of  $\alpha$ -thalassemia is  $\alpha$ -gene deletion<sup>Q</sup> (frame-shift mutation)<sup>Q</sup>

No. of $\alpha$ genes deleted	Genotype
1 gene deletion trait	$-\alpha/\alpha$ , $\alpha^Q$
2 gene deletion trait	$-\alpha/-\alpha$ homozygous $\alpha$ thal trait or $-\alpha/\alpha$ , $\alpha$ heterozygous $\alpha$ thal trait
3 gene deletion (Hb-H) <sup>Q</sup>	$-/-$ , $\alpha$
4 gene deletion (Hb-BART) <sup>Q</sup>	$-/-/-/-$



Howell Jolly body

## $\beta$ -Thalassemias

### Types of Mutations in $\beta$ -Thalassemias

- Splicing mutations:** Most common cause of  $\beta(+)$  thalassemia.<sup>Q</sup>
- Promoter region mutations:**  $\beta(+)$  thalassemia<sup>Q</sup>
- Terminator mutations:** Most common cause of  $\beta(0)$  thalassemia<sup>Q</sup>



### High Yield Facts


Most common mutation in  $\beta$  thalassemia in India:

- IVS-1, position 5 (G  $\rightarrow$  C) (most common mutation)<sup>Q</sup>
- 619-bp deletion<sup>Q</sup>
- Codons 8/9, frameshift mutation
- Codons 41/42, frameshift mutation
- IVS-1, position 1 (G  $\rightarrow$  T)

### $\beta$ -Thalassemias:

Clinical classification	Manifestations	Globin genotype (normal= $\beta/\beta$ )
<b>Thalassemia trait (minor)</b>	<ul style="list-style-type: none"> <li>Asymptomatic</li> <li>May have mild anemia</li> </ul>	$\beta/\beta_0$ $\beta/\beta^+$
<b>Thalassemia intermedia</b>	<ul style="list-style-type: none"> <li>Mild to moderate anemia</li> <li>Variable need of blood transfusions</li> </ul>	$\beta^+/\beta^+^Q$
<b>Thalassemia major</b>	<ul style="list-style-type: none"> <li>Severe anemia &amp; Jaundice</li> <li>Presents in childhood</li> <li>Requires multiple transfusion</li> <li>Hemolytic facies</li> <li>Hepatosplenomegaly</li> </ul>	$\beta^+/\beta^+$ $\beta^+/\beta_0$ $\beta_0/\beta_0^Q$


## Pathophysiology of $\beta$ -Thalassemias






## Paroxysmal Nocturnal Hemoglobinuria (PNH)

- **Definition**
  - A triad of Intravascular hemolysis, Thrombosis & Pancytopenia
- **Underlying defect**
  - Acquired somatic clonal mutation of **PIG-A gene**<sup>Q</sup> resulting in **loss of GPI (Glycosyl phosphatidyl inositol)** linked proteins like CD55, CD59
- **Pathophysiology**
  - Deficiency of **Regulators of complement activity** → **Inappropriate complement activation** → **Intravascular hemolysis** & destruction of WBCs & platelets causing **pancytopenia**.<sup>Q</sup>
- **Diagnosis**
  - **Acidified serum hemolysis test (HAM's test)**:<sup>Q</sup>
  - Patient's RBC gets lysed in acidic pH in presence of complements
  - **Alkaline phosphatase: Low LAP** score<sup>Q</sup> in PNH patients
  - **Gold standard: Flow cytometry: absence of GPI-linked** proteins such as **CD59/CD55**<sup>Q</sup>
  - **FLAER (Fluorescent aerolysin)**- Most recently introduced test to diagnose PNH<sup>Q</sup> – **Diagnosis of choice**



PNH pathophysiology



PNH Card test

## High Yield Facts

- PIG-A gene is **X-linked** which **encodes Glycosyl phosphatidyl inositol (GPI)** anchor protein.
- 5% to 10% of PNH patients develop **AML or MDS**<sup>Q</sup>
- PNH has high association with **aplastic anemia**<sup>Q</sup>
- **Thrombosis is the leading cause of disease-related death in PNH**<sup>Q</sup>
- In PNH, hemolysis occurs at night due to slight decrease in blood pH during sleep, which increases the activity of complement

## Immuno-hemolytic Anemias

- **Definition**
  - Hemolytic anemias caused by antibodies that bind to red cells, leading to their premature destruction.
- **Classification**
  - Autoimmune (*see below*)
  - Alloimmune Eg: Hemolytic disease of New born, Rh Incompatibility
- **Etiology**

Warm Antibody Type (IgG Ab Active at 37°C) <sup>Q</sup>	Cold Antibody Type (IgM Ab Active Below 37°C) <sup>Q</sup>
<b>Primary (idiopathic)</b> <b>Secondary</b> <ul style="list-style-type: none"> <li>• Autoimmune (SLE, Evans syndrome)</li> <li>• Lymphoid neoplasms eg: CLL<sup>Q</sup></li> <li>• Drugs: <ul style="list-style-type: none"> <li>■ <b>Antigenic type</b><sup>Q</sup>: Penicillin, Cephalosporins</li> <li>■ <b>Innocent bystander type</b> (Immune complex): Penicillin</li> <li>■ <b>Tolerance-breaking</b>: Alpha methyl dopa<sup>Q</sup></li> </ul> </li> </ul>	<b>Primary (idiopathic)</b> <b>Secondary:</b> <ul style="list-style-type: none"> <li>• <b>Acute</b>:<sup>Q</sup> Mycoplasma pneumoniae, EBV, CMV, Influenza and HIV</li> <li>• <b>Chronic</b>: Lymphoid neoplasms</li> </ul>
	<b>Cold Hemolysin Type</b> <b>(Ab Active Below 37°C)</b> <ul style="list-style-type: none"> <li>• Viral infections.</li> </ul>

- **Diagnosis:**
  - **PS:** Anisopoikilocytosis, Polychromasia, **Spherocytes**<sup>Q</sup>
  - **Increased Reticulocyte %**
  - **Coomb's test is positive in Immune hemolytic anemia**<sup>Q</sup>
    - **Direct** Coomb's test- Detects **Antibody on RBC Surface**<sup>Q</sup>
    - **Indirect** Coomb's test- Detects **Antibody in Serum**<sup>Q</sup>



- **Evans Syndrome:** Auto Immune Hemolytic Anemia (AIHA) + thrombocytopenia
- **Paroxysmal cold hemoglobinuria** is seen in children following viral infections.
- In **Paroxysmal cold hemoglobinuria**, intravascular hemolysis and hemoglobinuria seen
- **Paroxysmal cold hemoglobinuria** is due to **Donath Landsteiner Antibodies (IgG)<sup>Q</sup>** that bind to **P blood group<sup>Q</sup>** antigen on RBC at 4°C → **Complement-mediated lysis of RBCs at 37°C**

## High Yield Facts

- In **Cold agglutinin disease**, chronic **intravascular hemolysis** occur on **exposure to cold (0-5 C)<sup>Q</sup>**
- **Cold agglutinin disease** is due to binding of **IgM** to "**i**" Ag on RBC's at 32°C<sup>Q</sup> → activates complements at 37°C
- Both intravascular & extravascular hemolysis is seen in **Cold agglutinin disease**

## Red Cell Fragmentation Syndromes

Macroangiopathic Hemolytic Anemia	Microangiopathic Hemolytic Anemia <sup>Q</sup>
<b>Cardiac hemolysis</b> <ul style="list-style-type: none"> <li>• Prosthetic heart valves</li> <li>• Perivalvular leaks</li> </ul> <b>Arteriovenous malformations</b>	<ul style="list-style-type: none"> <li>• <b>Hemolytic Uremic Syndrome (HUS)<sup>Q</sup></b></li> <li>• <b>Thrombotic Thrombocytopenic Purpura (TTP)<sup>Q</sup></b></li> <li>• <b>Disseminated intravascular coagulation (DIC)<sup>Q</sup></b></li> <li>• <b>Malignant hypertension<sup>Q</sup></b></li> <li>• Pre-eclampsia/HELLP</li> <li>• March hemoglobinuria</li> <li>• Vasculitis</li> <li>• Connective tissue disorders Eg. SLE,<sup>Q</sup> Scleroderma</li> <li>• Giant hemangioma (Kasabach-Merritt syndrome)<sup>Q</sup></li> </ul>




## Microangiopathic Hemolytic Anemia (MAHA) For detail Discussion, refer Bleeding Disorders

- **Characterized by:** Hemolytic anemia due to RBC fragmentation.
- **Pathophysiology:** Microvascular lesion → luminal narrowing, due to deposition of fibrin and platelets → traumatic damage → red cell fragmentation → Appearance of **schistocytes<sup>Q</sup>, burr cells, helmet cells, triangle cells** in blood smears.
- **Schistocytes:** Fragmented RBC's with 1-3 sharp spicules; (>3/5000)

## Anemias of Decreased Erythropoiesis

**NUTRITIONAL-DEFICIENCY ANEMIA:** Important examples are:

- Iron-deficiency anemia
- Folic acid and B<sub>12</sub> deficiency anemia
- Vit C, Copper and Zinc deficiency






## Iron Deficiency Anemia

### Iron Metabolism

**Site of Iron absorption:** Mainly occurs in **proximal small intestine (duodenum and jejunum)**<sup>Q</sup>



### High Yield Facts

- **Amount of iron** needed to replace RBC's lost through senescence amounts to **20 mg/day**
- Normally, an adult male absorbs **1 mg** of elemental iron daily to meet needs<sup>Q</sup>
- Iron-deficiency anemia is the **most common anemia prevalent in India**<sup>Q</sup>
- Iron-deficiency anemia has higher incidence in females, particularly **pregnant females**<sup>Q</sup>
- **Hepcidin** is the **principal iron regulatory hormone**; it is **negatively regulated by ferroportin**<sup>Q</sup>

\***Divalent metal transporter type 1 (DMT-1)**: also known as natural resistance macrophage-associated protein type 2 (Nramp 2) or DCT-1<sup>Q</sup>

\*\***Hephaestin** is similar to **ceruloplasmin**, the copper-carrying protein.<sup>Q</sup>

### Iron Deficiency Anemia

Etiology	Increased Demand for Iron	Increased Iron Loss		Decreased Iron Intake or Absorption
	<ul style="list-style-type: none"><li>• Rapid growth in infancy or adolescence</li><li>• Pregnancy<sup>Q</sup></li><li>• Erythropoietin therapy</li></ul>	<ul style="list-style-type: none"><li>• Acute or Chronic blood loss (Hookworm,<sup>Q</sup> Carcinoma colon<sup>Q</sup>)</li><li>• Menstruation</li><li>• Phlebotomy as treatment for polycythemia vera</li></ul>		Inadequate diet Malabsorption (Celiac ds, <sup>Q</sup> Crohn’s ds) Malabsorption due to surgery (postgastrectomy, <sup>Q</sup> Bilioth II <sup>Q</sup> ) Acute or chronic inflammation Chronic Renal Failure <sup>Q</sup>
Clinical features	<ul style="list-style-type: none"><li>• Depend on the severity &amp; chronicity of anemia</li><li>• <b>Fatigue, pallor, reduced exercise capacity, cheilosis<sup>Q</sup></b> (fissures at the corners of the mouth)</li></ul>		<ul style="list-style-type: none"><li>• <b>Koilonychia<sup>Q</sup></b>(spooning of the fingernails); <b>Platynychia<sup>Q</sup></b>: flattening of the fingernails</li><li>• Pharyngeal webs: <b>Plummer Vinson syndrome<sup>Q</sup></b></li></ul>	
Stages of Iron Deficiency	Features	Stage 1 (Prelatent)	Stage 2 (Latent)	Stage 3 (Anemia)
	Symptoms	None	Fatigue, malaise	Pallor ± pica
	Hemoglobin	Normal	<b>Normal<sup>Q</sup></b>	<b>Reduced<sup>Q</sup></b>
	MCV	Normal	Normal	Reduced
	Serum ferritin	<b>Reduced<sup>Q</sup></b>	<b>&lt;12 µg/L<sup>Q</sup></b>	<b>&lt;12 µg/L<sup>Q</sup></b>
	Transferrin saturation	Normal	<16% <sup>Q</sup>	<16%
	Free erythrocyte protoporphyrin	Normal	↑ <sup>Q</sup>	↑ <sup>Q</sup>
	Serum transferrin receptor	<b>Normal<sup>Q</sup></b>	↑	↑
	Bone marrow iron	Reduced	Absent	Absent
Lab diagnosis	Peripheral smear: <ul style="list-style-type: none"><li>• Anisocytosis, microcytic hypochromic RBCs</li><li>• <b>Pencil RBCs/cigar<sup>Q</sup></b> shaped RBCs are seen</li><li>• <b>Target RBCs</b> in severe cases</li><li>• <b>Reticulocyte %<sup>Q</sup>-not raised</b></li><li>• <b>Platelets-often increased<sup>Q</sup></b></li></ul>		RBC Indices: <ul style="list-style-type: none"><li>• <b>MCV &lt; 80fl</b></li><li>• <b>MCH &lt; 25pg</b></li><li>• <b>MCHC&lt; 27g/dl</b></li><li>• <b>Increased RDW (&gt;14)<sup>Q</sup></b></li><li>• <b>Serum Iron studies:</b> Refer to table below</li></ul>	
Response to therapy	<ul style="list-style-type: none"><li>• <b>Rapid subjective improvement</b>, with decrease in fatigue, <b>before any improvement in anemia</b></li><li>• <b>Earliest hematologic evidence: increase in reticulocyte % &amp; their hemoglobin content.</b></li><li>• <b>Reticulocytes attain maximal value (5 to 10%) on 5<sup>th</sup> to 10<sup>th</sup> day</b> of therapy</li></ul>			






## Serum Iron Studies: (to Distinguish the Differential Diagnosis)

Differential Diagnosis of Microcytic Anemia					
Tests	Normal Values	Iron Deficiency	Anemia of chronic disease	Thalassemia	Sideroblastic Anemia
<i>Peripheral Smear</i>	N/N	Micro/hypo	N/N or micro/hypo	Micro/hypo with target RBCs	Micro/hypo or Dimorphic
<i>S. Iron (ug/dl)</i>	50-150	<30	<50	Normal to high	Normal to high
<i>(TIBC) (ug/dl)</i>	30-300	>360	<300	Normal	Normal
<i>Transferrin saturation %</i>	33%	<10	10-20	30-80	30-80
<i>Ferritin (µg/L; ng/ml)</i>	50-300	<15	30-200	50-300	50-300
<i>Hb electrophoresis</i>	—	Normal	Normal	Abnormal	Normal

\*TIBC: Total Iron Binding Capacity



## ANEMIA OF CHRONIC DISEASE

Hypoferremia in presence of Adequate Reticuloendothelial Stores

### Etiology (for > 1-2 months)

- **Chronic Inflammation:** Rheumatoid Arthritis, Systemic Lupus Erythematosus
- **Chronic Infections:** TB, Chronic osteomyelitis, HIV
- **Malignancy:** Hodgkin's disease and Non-Hodgkin Lymphoma
- **Miscellaneous:** Alcoholic liver disease

## SIDEROBLASTIC ANEMIA

- **Definition**
  - A heterogeneous group of disorders characterized by amorphous iron deposits in erythroblast mitochondria

### Classification

Hereditary	X-linked (XLSA): (Pearson syndrome)
Acquired	Refractory anemia with ring sideroblasts (RARS)/Pure SA (PSA)
Reversible	Alcoholism Lead poisoning Drugs (isoniazid, pyrazinamide, chloramphenicol) Copper deficiency (zinc ingestion, copper chelation Hypothermia

- Sideroblasts can be seen in Iron overload



## MEGALOBlastic ANEMIA

- **Definition:**
  - Impairment of DNA synthesis that leads to ineffective hematopoiesis & distinctive morphologic changes,



including **abnormally large erythroid precursors** in bone marrow & RBCs in peripheral smear

#### ■ Etiology

- **Cobalamin Deficiency**
  - **Dietary deficiency**
  - **Malabsorption:** Ileal resection, Fish tapeworm infection
  - **Intrinsic factor Deficiency:** Pernicious anemia
  - **Increased requirements:** Pregnancy
- **Folic acid deficiency**
  - **Dietary deficiency:** Malnutrition, Alcoholics<sup>Q</sup>
  - **Impaired absorption:** Celiac Sprue<sup>Q</sup>, Small bowel resection/disease, Anticonvulsants/ OCPs
  - **Increased requirements:** Infancy, Pregnancy<sup>Q</sup>, Hemolytic anemias
- **Drug-induced suppression of DNA synthesis**
  - **Folate antagonists,** Alkylating agents
  - Metabolic inhibitors of synthesis of:
    - Purine: **hydroxyurea**,<sup>Q</sup> **6-MP**, **azathioprine**
    - Pyrimidine: **5-fluorouracil**; **cytosine arabinoside**
- **Inborn errors**
  - Defective folate metabolism
  - Defective vitamin B<sub>12</sub> metabolism
  - Hereditary orotic aciduria<sup>Q</sup>
  - Lesch-Nyhan syndrome<sup>Q</sup>

#### ■ Peripheral smear


- **Anisopoikilocytosis, macrocytic RBCs; Macro-ovalocytosis** are highly **characteristic**.<sup>Q</sup>
- Tear drop RBCs, Howel Jolly bodies, Cabot ring & Basophilic stippling may be seen
- **Macrocytes** appear “hyperchromic,” but the **MCHC is not elevated**.
- Even though there is **mild hemolysis**, **reticulocyte count is low**
- Neutrophils are also larger than normal (**macropolymorphonuclear**)<sup>Q</sup> and show **nuclear hypersegmentation**<sup>Q</sup> (**>=5 lobes in >5% neutrophils**) **Earliest abnormality to appear**<sup>Q</sup>

#### ■ Bone marrow

- Markedly **hypercellular** with **megaloblastic changes**<sup>Q</sup> at all stages of erythroid development
- Derangement in **DNA synthesis** → **apoptosis** of precursors of all 3 lineages in marrow (**ineffective hematopoiesis**) → **pancytopenia**.<sup>Q</sup>
- **Hallmark: nuclear cytoplasmic maturation asynchrony**<sup>Q</sup>
- **Megaloblast with sieve like chromatin**<sup>Q</sup>

#### ■ Other tests

- In vit B12 deficiency, serum **MMA (Methyl malonic acid) level is raised**.<sup>Q</sup>
- **Serum homocysteine is raised** in both early **cobalamin and folate deficiency**<sup>Q</sup>
- **Increased LDH levels**<sup>Q</sup>



#### Schilling test

##### Step 1: oral vitamin B<sub>12</sub> plus intramuscular vitamin B<sub>12</sub>

- Patient is given radiolabeled vit B<sub>12</sub> followed by i.m injection of unlabeled vit B<sub>12</sub> an hour later to temporarily saturate B<sub>12</sub> receptors in the liver.
- The patient's urine is then collected over the next 24 hours to assess the absorption.
- Normally, the ingested radiolabeled vitamin B<sub>12</sub> will be absorbed into the body and get excreted in the urine.
- Normal result: **> 10% of radiolabeled vit B<sub>12</sub> in the urine** over the first 24 hours.
- In patients with pernicious anemia or with deficiency due to impaired absorption, less than 10% of the radiolabeled vitamin B<sub>12</sub> is detected → we proceed to step 2

##### Step 2: Vitamin B<sup>12</sup> with oral intrinsic factor

- Normal excretion indicates lack of intrinsic factor production, or pernicious anemia.
- Low value on the second test implies **abnormal intestinal absorption** (malabsorption):






- Coeliac disease, biliary disease, Whipple's disease, small bowel bacterial overgrowth syndrome, fish tapeworm infestation (*Diphyllobothrium latum*), or liver disease.

**Step 3: Vitamin B<sub>12</sub> with antibiotics:** to identify patients with bacterial overgrowth syndrome.

**Step 4: Vitamin B<sub>12</sub> with pancreatic enzymes:** to identify patients with pancreatitis

## Aplastic Anemia



Aplastic anemia: Pancytopenia with hypocellular BM

## Pure Red Cell Aplasia (PRCA)

Pure red cell aplasia is a primary marrow disorder in which only erythroid progenitors are suppressed.

### Classification of Pure Red Cell Aplasia


**Self-limited:** Transient erythroblastopenia of childhood

**Hereditary:** Congenital pure red cell aplasia (**Diamond-Blackfan syndrome**)<sup>o</sup>

### Acquired

- Thymoma**
- Lymphomas:** CLL, Large granular lymphoma,<sup>o</sup> Hodgkin's,<sup>o</sup> Non Hodgkin's
- Paraneoplastic** to solid tumors
- Connective tissue disorders: **SLE, JRA, RA**
- Virus:** Parvovirus B19<sup>o</sup>, hepatitis, HTLV, EBV
- Pregnancy**<sup>o</sup>





- **Drugs:** Phenytoin, azathioprine, chloramphenicol, procainamide, isoniazid<sup>Q</sup>
- **Post ABO incompatible transplant<sup>Q</sup>**

## MYELOPHTHISIC ANEMIA

- **Definition**
  - A form of marrow failure in which space-occupying lesions replace normal marrow elements.
- **Etiology**
  - **Metastasis** from breast, lung, and prostate Carcinoma
  - **Infiltrative process** (e.g., granulomatous disease)
  - Spent phase of **myeloproliferative** disorders
  - Storage disorders: Eg. Gaucher's disease

### Peripheral Smear

- **Leukoerythroblastosis:** Abnormal release of nucleated RBC's and immature granulocytic forms into peripheral smears.
- Tear drop-shaped red cells (**Dacrocytes**)




### High Yield Facts


#### Diagnostic Criteria for Severe Aplastic Anemia<sup>Q</sup>

Bone marrow cellularity of <25% along with 2 of the following:


- Neutrophil count <500/ $\mu\text{l}^Q$
- Platelet count <20,000/ $\mu\text{l}^Q$
- Absolute reticulocyte count of <60,000/ $\mu\text{l}^Q$




Normal marrow



Hypercellular marrow



PRCA




Leukoerythroblastosis





## Image-Based Questions

1. The given figure shows a special stain on RBCs. Which of the following stains cannot be used for this?



- a. New Methylene blue      b. Crystal violet  
c. Brilliant cresyl blue      d. Methyl violet

3. In the above case, the following investigation was done. Identify the test ?




- a. APT test  
b. Kleihauer-Bethke acid elution for fetal hemoglobin  
c. Kleihauer-Bethke alkali elution for fetal hemoglobin  
d. Rosetting

2. The given figure shows urine sample from a patient who presented with anemia and Jaundice. (compare with normal sample on left). In which of the following condition this can be seen?




- a. Thalassemia      b. Hereditary Spherocytosis  
c. PNH      d. Sickle cell anemia


4. A male aged 20 years, native from Orissa had presented to AIIMS Medicine OPD with high grade fever with chills and rigor. His routine CBC revealed mild neutrophilia and increased TLC while Peripheral smear findings have been given below. He was given treatment for the same after he suffered with anemia and jaundice, and a repeat smear was done (as shown in next figure). What is your diagnosis?



- a. *P. vivax* and bite cells  
b. *P. falciparum* and bite cells  
c. Babesia and schistocytes  
d. *P. vivax* and scistocytes




5. A 5-year-old male child presented to AIIMS pediatrics OPD with severe transfusion requiring anemia and Jaundice. On examination Liver and spleen were palpable 5 cm below the costal margin. Peripheral smear analysis showed the following? What is your diagnosis?



- Nutritional anemia
- Aplastic anemia
- Autoimmune hemolytic anemia
- Thalassemia

6. A 45-year-old male presented to OPD with severe fatigue. On examination he was found to have severe anemia, and petechial spots. Given below is the picture of Peripheral smear and bone marrow biopsy. What is your diagnosis?



- Iron deficiency anemia
- Megaloblastic anemia
- Aplastic anemia
- Pure red cell aplasia



## Answers of Image-Based Questions


- Ans. (d) **Methyl violet**
  - This is staining of reticulocyte with supravital stains which includes New Methylene blue, Crystal violet, Brilliant cresyl blue
- Ans. (c) **PNH**
  - A case of anemia and Jaundice suggests hemolytic anemia. Red coloured urine suggests hemoglobinuria seen in a case of intravascular hemolysis like PNH.
- Ans. (b) **Kleihauer-Bethke acid elution for fetal hemoglobin**
  - In this figure the dark red colored cells are RBC which have resisted acid elution and have HbF, while ghost RBCs (light colored) lack HbF and so denatured with alkali and eluted with adding acid. **Kleihauer-Betke acid elution method to quantify the magnitude of fetomaternal bleed.**
- Ans. (a) ***P. vivax* and bite cells**
  - The given condition is malaria and shows trophozoite stage of *Plasmodium vivax*. On giving anti malarials like Primaquine, hemolysis may be precipitated showing bite cells in peripheral smear.
- Ans. (d) **Thalassemia**
  - Severe transfusion requiring anemia and Jaundice with smear showing Anisopoikilocytosis and targets on smear suggests Thalassemia.
- Ans. (c) **Aplastic anemia**
  - The given peripheral smear shows pancytopenia while bone marrow biopsy shows hypocellular marrow seen in a case of Aplastic anemia.




## Multiple Choice Questions

### RBCs AND ANEMIA


- Maltese cross can be seen in:** (AIIMS Nov 2019)
  - Pneumocystis carinii
  - Leishmaniasis
  - Cryptococcosis
  - Babesiosis
- Which of the following tubes contain Sodium fluoride as anticoagulant?** (AIIMS Nov 18)
 




a.



c.




b.




d.
- Formula for calculating reticulocyte production index:** (JIPMER Nov 2019)
  - Retic  $\times$  hct / normal hct
  - Corrected ret  $\times$  hct / 45 | maturation index
  - Reticulocyte percentage  $\times$  RBC count
  - Reticulocyte counted  $\times$  100 / number of red cells
- Ring sideroblast in myelodysplastic syndrome is associated with which gene mutation?** (JIPMER 18)
  - ASXL1
  - EZH2
  - TET2
  - SF3B1
- What should be the sequence of events during collection of blood sample?** (AIIMS Nov 2017)
  - Ask the patient his name\*verify from file Label the sample at bedside  $\rightarrow$  collect blood
  - Look at the file  $\rightarrow$  Collect sample  $\rightarrow$  Label the sample at bedside
  - Prelabel the sample vials  $\rightarrow$  Check the file patient details  $\rightarrow$  Collect sample
  - Collect sample  $\rightarrow$  Confirm name from file  $\rightarrow$  Label the sample vial
- Which vacutainer is used for electrolyte estimation?**
  - Na Citrate
  - EDTA (AIIMS Nov 2017)
  - Fluoride
  - Lithium heparin
- What does the red cell distribution width (RDW) deal with?** (AIIMS May 2017)
  - Hypochromia
  - Anisocytosis
  - Poikilocytosis
  - Anisochromia
- A 25 year old female came to OPD 1 year after postpartum. She was treated for iron deficiency anemia while pregnancy. Now she is pale and her Hb was 5% and reticulocyte count was 9%. Her corrected retic count is?** (AIIMS May 2017)
  - 6
  - 4.5
  - 3
  - 1
- Acetone free methanol in leishman stain does which of the following?** (AIIMS May 2017)
  - Fixes cells to slide
  - Colour of cell
  - Causes metabolic and enzymatic activity to stop
  - Washes the slide

- A 26-year-old patient having RBC count 2 lakhs/mm<sup>3</sup>, PCV 30% and haemoglobin 9.3, what is the type of anemia?** (AIIMS Nov 2016)
  - Iron deficiency anemia
  - Folic acid deficiency anemia
  - Thalassemia
  - Sideroblastic anemia
- Identify the organism in the given Image?**



- P. vivax
  - P. ovale (AIIMS Nov 2016)
  - P. falciparum
  - P. malariae
- The rate of production of red blood cells by the human bone marrow is?** (Recent Question 2016)
    - $1.5 \times 10^6$  cells/second
    - $1.5 \times 10^3$  cells/second
    - $7.5 \times 10^6$  cells/second
    - $7.5 \times 10^3$  cells/second
  - Identify the predominant abnormally shaped red cell type seen in the peripheral smear given below:** (Recent Question 2016)



- Macrocyte
  - Acanthocyte
  - Dacrocyte
  - Elliptocyte
- Erythropoietin in fetus is secreted by?** (Recent Question 2016)
    - Liver
    - Marrow
    - Spleen
    - Kidney
  - Most mature normoblast is:** (JIPMER Nov 2019)
    - Orthochromic normoblast
    - Pronormoblast
    - Polychromatic normoblast
    - Basophilic normoblast
  - 1st stage of RBC development is?** (Recent Question 2016)
    - Pro erythroblast
    - Intermediate normoblast
    - Reticulocyte
    - nRBC



- 17. Spur cells are seen in** (Recent Question 2015)  
a. Hereditary spherocytosis b. G6PD deficiency  
c. Liver disease d. Sideroblastic anemia
- 18. Spiculated RBCs with evenly spaced spikes are called** (Recent Question 2015)  
a. Acanthocytes b. Stomatocytes  
c. Echinocytes d. Dacryocytes
- 19. Most common cause of splenic rupture is:** (Recent Question 2015)  
a. Malaria b. ITP  
c. Thalassemia d. Cirrhosis
- 20. Supravital staining is used for?** (Recent Question 2015)  
a. Nucleated RBCs b. Reticulocytes  
c. Basophils d. Myeloblasts
- 21. When osmotic fragility is normal, RBCs begin to hemolyse when suspended in saline:** (Recent Question 2015, DNB 99)  
a. 0.33% b. 0.48%  
c. 0.9% d. 1.2%
- 22. True about reticulocyte is?** (Recent Question 2014)  
a. Stained by supravital staining  
b. Myeloid cell  
c. Romanowsky stains are used  
d. 5% is normal
- 23. Ratio of fat cells and blood cells in bone marrow is-** (Recent Question 2014, 2013)  
a. 1: 4 b. 1: 2  
c. 1: 1 d. 2: 1
- 24. Howel-Jolly bodies may be seen after-** (Recent Question 2014, AI 99)  
a. Hepatectomy b. Splenectomy  
c. Pancreatectomy d. Cholecystectomy
- 25. Cabot's ring is seen in?** (Recent Question 2014)  
a. Megaloblastic anemia  
b. Sickle cell disease  
c. Iron deficiency anemia  
d. Autoimmune anemia
- 26. Cyanosis is not seen in severe anemia because:** (AIIMS May 2012)  
a. Anemic hemoglobin has greater oxygen carrying capacity per unit gram of hemoglobin  
b. A critical concentration of reduced hemoglobin is required  
c. Increased RBC number counterbalances the oxygen shortage  
d. Blood flow through the skin is decreased in anemia
- 27. ESR is a very critical investigation in the diagnosis of TB. Which of the following is true about ESR in TB?** (AIIMS May 2012)  
a. No change in ESR  
b. Confirms recovery from TB  
c. ESR is raised because of increased RBC aggregate  
d. ESR is raised due to decreased RBC size
- 28. All are true about polycythemia vera except?** (DNB Aug. 12 Pattern)  
a. Increased ESR  
b. Decreased erythropoietin  
c. Increased LAP score  
d. Increased blood volume
- 29. Cabot's ring in RBC is seen in?** (DNB Aug 12, Dec 09)  
a. Acquired hemolytic anemia  
b. Hemochromatosis  
c. Thalassemia  
d. After splenectomy
- 30. If you are in PHC, which anticoagulant is used to sent the blood sample for blood glucose estimation'?** (AIIMS Nov 11)  
a. EDTA  
b. Heparin  
c. Potassium oxalate  
d. Potassium oxalate + sodium fluoride
- 31. True about RBC and anemia:** (PGI Nov 2011)  
a. In female of reproductive age 12.5% is cut of value for anemia  
b. In newborn Hb level is 15 gm%  
c. Microcytosis occurs in folic acid deficiency  
d. Macrocytosis may occur in Celiac disease  
e. Normocytic normochromic anemia is found in chronic diseases
- 32. Which of the following indicates hemolysis?** (JIPMER 11)  
a. Target cells b. Schistocytes  
c. Acanthocytes d. Basophilic stippling
- 33. Pappenheimer bodies are composed of ?** (DNB June 10)  
a. Copper b. Zinc  
c. Iron d. Lead

#### ESR

- 34. In which of the following condition(s) erythrocyte sedimentation rate is increased:** (PGI May 2019)  
a. Increased serum immunoglobulin level  
b. Spherocytosis  
c. Increased plasma viscosity  
d. Sickle cell anemia  
e. Increased level of C-reactive protein

- 35. ESR is decreased in all the following conditions except** (Recent Question 2015)  
a. Polycythemia  
b. Sickle cell anemia  
c. Congestive cardiac failure  
d. Multiple myeloma
- 36. Decreased ESR is seen in** (Recent Question 2016)  
a. Sickle cell Anemia  
b. Increased fibrinogen  
c. Anemia  
d. Hypoviscosity


#### HEMOLYTIC ANEMIAS

- 37. In hereditary spherocytosis, which of the following is/are increased?** (PGI May 18)  
a. Haptoglobin b. Spleen size  
c. MCHC d. MCV  
e. LDH
- 38. Secondary hemolytic anemia is seen in all except?** (JIPMER 18)  
a. CML b. CLL  
c. CRRT d. ECMO





39. **Eosin-5-Maleamide flow cytometry is used for diagnosis of?** (JIPMER 18)  
 a. G6PD  
 b. Hereditary spherocytosis  
 c. Sickle cell anemia  
 d. Alpha thalassemia
40. **Schistocytes seen in which of the following?** (PGI Nov 2018)  
 a. HUS  
 b. TTP  
 c. Prosthetic heart valves  
 d. Alpha thalassemia
41. **Acanthocytes are seen in?** (PGI MAY 18)  
 a. Hallervorden-Spatz disease  
 b. Abetalipoproteinemia  
 c. Severe liver disease  
 d. Ataxia telangiectasia  
 e. McLeod syndrome
42. **Blister cells seen in which condition?** (JIPMER 2017)  
 a. G6PD deficiency      b. Thalassemia  
 c. Sickle cell anemia      d. AIHA
43. **Cold agglutinins are directly against which of the following RBC antigens?** (JIPMER Nov 2019)  
 a. I  
 b. P  
 c. Le  
 d. Rh
44. **A 25-year-old female presented in December month with chronic fatigue and cyanosis with bluish lips and arthralgia. Peripheral blood film is shown below. What is the likely cause?** (AIIMS Nov 2016)



- a. Cold AIHA      b. Warm AIHA  
 c. Hemoglobinopathy      d. G6PD Deficiency
45. **All of the following statement is true regarding hemolytic anemia except?** (AIIMS Nov 2016)  
 a. Autosplenectomy      b. Increased LDH  
 c. Increased bilirubin      d. Decreased reticulocyte
46. **Which of the following is absent in hemolytic anemia?** (PGI Nov 2016)  
 a. Increased indirect bilirubin  
 b. Increased direct bilirubin  
 c. Increased reticulocyte count  
 d. Jaundice  
 e. Increased LDH
47. **True about sickle cell anemia is?** (JIPMER 2016)  
 a. More susceptible to infections especially by Pneumococcus  
 b. Sickling is irreversible  
 c. Bone pain is common in small bones  
 d. HbA2 is increased
48. **Direct coomb's test positive is seen in all except?** (AIIMS May 2016)  
 a. ABO incompatibility  
 b. Hemolytic d/s of newborn  
 c. Aplastic anemia  
 d. Autoimmune hemolysis
49. **Which one of the following is not associated with a high reticulocyte count?** (Recent Question 2016-17)  
 a. Acute bleed  
 b. Haemolytic anemia  
 c. Megaloblastic anemia  
 d. Response to treatment in 'nutrition - deficiency' anemia
50. **What happens when normal erythrocytes (blood - group matched) are transfused into a patient with anemia secondary to an intracorporeal defect?** (Recent Question 2016-17)  
 a. Donor cells are destroyed  
 b. Donor cells have normal survival  
 c. Depends on the severity of anemia  
 d. Depends on whether the donor cells are fresh or stored (older than a week)
51. **Adult hemoglobin consists of the following chains?** (Recent Question 2016-17)  
 a.  $2\alpha + 2\beta$       b.  $2\alpha + 2\delta$   
 c.  $2\beta + 2\gamma$       d.  $2\alpha + 2\gamma$
52. **Haemolysis in G6PD (glucose 6 phosphate dehydrogenase) enzyme deficiency may occur with all of the following drugs except:** (Recent Question 2016-17)  
 a. Primaquine      b. Phenacetin  
 c. Probenecid      d. Penicillin
53. **Warm antibody autoimmune hemolytic anemia, true is?** (Recent Question 2016-17)  
 a. Does not need complements  
 b. Antibody active at 2-8 C  
 c. IgM type Ab is involved  
 d. CLL is an important cause
54. **A 21-year-old male presents with complaints of fatigue and abdominal pain since birth. O/E jaundice and splenomegaly present. On USG gall stones are seen. What is your diagnosis?** (Recent Question 2015)  
 a. Hereditary spherocytosis      b. Sickle cell anemia  
 c. Cholangitis      d. Acute pancreatitis
55. **Fanconi anemia can lead to?** (Recent Question 2015)  
 a. B12 deficiency      b. Folate deficiency  
 c. Iron deficiency      d. Aplastic anemia
56. **In G6PD deficiency which cells are more prone for hemolysis** (Recent Question 2015)  
 a. Older red cells      b. Young red cells  
 c. Reticulocytes      d. All are susceptible
57. **Most common mutation in hereditary spherocytosis** (Recent Question 2015)  
 a. Spectrin      b. Ankyrin  
 c. Glycophorin A      d. Band 3
58. **Most common mutation in hereditary elliptocytosis** (Recent Question 2015)  
 a. Spectrin      b. Ankyrin  
 c. Glycophorin A      d. Band 4.2
59. **Drug that is safe in G6PD deficiency** (Recent Question 2015)  
 a. Primaquine      b. Acetanilid  
 c. Quinidine      d. Dapsone



- 60. Duffy antigen is associated with:** (Recent Question 2016)  
a. Plasmodium vivax      b. Falciparum  
c. Ovale      d. Malariae
- 61. In hereditary spherocytosis, which gene is altered?** (Recent Question 2016)  
a. Spectrin      b. Laminin  
c. Desmin      d. Vimentin
- 62. Warm Antibodies are?** (Recent Question 2016)  
a. Complete Antibody      b. Incomplete Antibody  
c. Heterophilic antibody      d. IgM
- 63. Best treatment of atypical HUS is?** (Recent Question 2016)  
a. Plasmapheresis      b. Antibiotics  
c. IVIG      d. Dialysis
- 64. Direct coombs test is positive in hemolytic anemia due to** (Recent Question 2015)  
a. Paroxysmal cold hemoglobinuria  
b. Paroxysmal nocturnal hemoglobinuria  
c. Idiopathic thrombocytopenic purpura  
d. Hemolytic uremic syndrome
- 65. Not used in the treatment of PNH** (Recent Question 2015)  
a. Cyclosporine  
b. Eculizumab  
c. Bone marrow transplantation  
d. Leucocyte depleted blood transfusion
- 66. Hemolytic crisis in hereditary spherocytosis is precipitated by** (Recent Question 2015)  
a. Parvovirus B19 infection  
b. Infectious mononucleosis  
c. Human T-cell leukemia virus  
d. Cytomegalovirus
- 67. A 25 year old patient presents with the history of dyspnea on exertion for 3 weeks. Investigations revealed Hb-7g/dl, reticulocyte count 18% and positive coombs test. Diagnosis** (Recent Question 2015)  
a. Autoimmune hemolytic anemia  
b. Paroxysmal nocturnal hemoglobinuria  
c. Sickle cell anemia  
d. Hereditary spherocytosis
- 68. Kleihauer Bethke test is done for?** (APPGMEE 2015)  
a. Cephalopelvic disproportion  
b. Fetomaternal haemorrhage  
c. Determining karyotype of normal fetus  
d. Diagnosing fetal infections
- 69. Features of hemolytic anemia are all except?** (Recent Question 2015)  
a. Hemoglobinemia      b. Bilirubinemia  
c. Reticulocytosis      d. Haptoglobin increased
- 70. Direct globulin test is positive in?** (Recent Question 2014)  
a. PNH  
b. Sickle cell anemia  
c. Thalassemia  
d. Paroxysmal cold hemoglobinuria
- 71. Intracorporeal hemolytic anemia is seen in-** (Recent Question 2014)  
a. Autoimmune hemolytic anemia  
b. TTP  
c. Thalassemia      d. Infection
- 72. Not true about hereditary spherocytosis-** (Recent Question 2014)  
a. Defect in ankyrin      b. Decreased MCV  
c. Decreased MCHC      d. Reticulocytosis
- 73. In sickle cell anemia all are true except:** (Recent Question 2014)  
a. Sickle cells      b. Target cells  
c. Howell jolly bodies      d. Ringed sideroblast
- 74. Person having heterozygous sickle cell trait is protected from infection of -** (Recent Question 2014)  
a. Plasmodium falciparum      b. P. vivax  
c. Pneumococcus      d. Salmonella
- 75. All are features of haemolytic uremic syndrome, except-** (Recent Question 2014)  
a. Hyperkalemia  
b. Anemia  
c. Renal microthrombi  
d. Neuro psychiatric disturbances
- 76. Thalassemia gives protection against-** (Recent Question 2014)  
a. Filariasis      b. Kala-azar  
c. Malaria      d. Leptospirosis
- 77. Donath Landsteiner antibody is seen in-** (Recent Question 2014)  
a. PNH  
b. Waldenström's macroglobulinemia  
c. Paroxysmal cold hemoglobinuria  
d. Malaria
- 78. HUS is differentiated from TTP by?** (JIPMER 2014)  
a. Presence of Microangiopathic hemolytic anemia  
b. Renal failure  
c. Neurological symptoms  
d. Absence of fever
- 79. Maternal blood from fetal blood can be differentiated by?** (JIPMER 2014)  
a. Osmotic fragility test  
b. Kleihauer-Bethke test  
c. APT test  
d. Bubble test
- 80. Reagent used in APT test?** (JIPMER 2014)  
a. Sodium hydroxide      b. Sodium chloride  
c. Sodium acetate      d. Sodium bicarbonate
- 81. Severe Hereditary spherocytosis can be seen due to defect of the following protein?** (JIPMER 2014)  
a.  $\alpha$ -Spectrin      b. Ankyrin  
c. Band 3      d. Band 4.2
- 82. Kleihauer-Betke test is used to detect** (APPGMEE 14)  
a. Fernald pattern in follicular phase  
b. Cephalopelvic disproportion  
c. Fetomaternal blood leak  
d. Sperm-cervical mucus interaction
- 83. A 16-year-old Afro-American boy presenting with non-healing ulcer of foot with recurrent pneumonia, chronic hemolytic anemia. The peripheral blood erythrocytes showed some RBCs with peculiar appearance. Most likely cause is:** (AIIMS Nov 14/ Nov 2013)  
a. Single amino acid base substitution  
b. Trinucleotide repeat  
c. Antibody to RBC membrane  
d. Genomic imprinting





- 84. Person having heterozygous sickle cell trait is protected from infection of:** (AIIMS May 2013/Nov 12)  
 a. *P. falciparum* b. *P. vivax*  
 c. *Pneumococcus* d. *Salmonella*
- 85. True about  $\beta$ -thalassemia:** (PGI May 2013)  
 a. Common in India b. Change in  $\beta$ -globin gene  
 c. Microcytosis d. Increased HbF  
 e. Secondary hemochromatosis may occur
- 86. Autosplenectomy is seen in?** (DNB Aug 12 Pattern, DNB June 10)  
 a. Hereditary spherocytosis  
 b. G6 PD deficiency  
 c. Sickle cell anemia  
 d. Thalassemia major
- 87. False about Sickle cell anemia is?** (JIPMER 2012)  
 a. Fetal Hb persists at high conc in adult life as it is protective  
 b. Co existent alpha thal is milder disease  
 c. Aplastic crisis is related to spleen  
 d. Sequestration crisis is related to spleen
- 88. Which is false about hemolytic anemia?** (JIPMER 2012)  
 a. Decreased LDH  
 b. Decreased Haptoglobin  
 c. Decreased RBC survival  
 d. Increased Unconjugated Bilirubin
- 89. Which of the following is not true about HS?** (JIPMER 2012)  
 a. Extravascular hemolysis  
 b. Mutation in protein of RBC membrane  
 c. Young RBCs are normal in shape  
 d. After splenectomy, Spherocytes disappear
- 90. Which is false about G6PD deficiency?** (JIPMER 2012)  
 a. X linked recessive  
 b. Young RBCs are more prone to hemolysis  
 c. Episodic hemolysis  
 d. Both intravascular & extravascular hemolysis is seen
- 91. A 23-year-old female presented with jaundice and pallor for 2 months. Her peripheral blood smear shows the presence of spherocytes. The most relevant investigation to arrive at a diagnosis is** (AIIMS May 11)  
 a. Reticulocyte count b. Osmotic fragility test  
 c. Coombs test d. Tests for PNH
- 92. Reticulocyte index  $\geq 2.5$  is/are seen in:** (PGI Nov 2011)  
 a. Hemolysis b. Blood loss  
 c. Hemoglobinopathy d. Iron deficiency anemia  
 e. Macrocytic anemia
- 93. Drugs that do not carry risk of hemolysis in persons with G-6-PD deficiency:** (PGI May 2011)  
 a. Primaquine b. Paracetamol  
 c. Ceftriaxone d. Nitrofurantoin  
 e. Vitamin K analogues
- 94. Schistocytes are found in:** (PGI May 2011)  
 a. Microangiopathic hemolytic anemia  
 b. HUS  
 c. Hereditary schistocytosis  
 d. Thrombotic thrombocytopenic purpura  
 e. DIC

- 95. Most common cause of hereditary spherocytosis?** (DNB Dec 11)  
 a. Spectrin b. Glycophorin  
 c. Ankyrin d. Band 4
- 96. Heinz bodies are seen in:** (MAHA 11)  
 a. Thalassemia  
 b. G6PD deficiency  
 c. Hereditary spherocytosis  
 d. Paroxysmal nocturnal hemoglobinuria
- 97. Unconjugated hyperbilirubinemia with increased urobilinogen is seen in:** (AI 10)  
 a. Hemolytic anemia b. Liver cirrhosis  
 c. Bile duct obstruction d. Sclerosing cholangitis
- 98. In hereditary spherocytosis mutation not seen is?** (DNB Dec 10)  
 a. Ankyrin b. Spectrin  
 c. Band-3 d. Na<sup>+</sup> Cl<sup>-</sup> channel protein
- 99. A Newborn with ABO incompatibility will characteristically show the presence of following on peripheral smear:** (AIIMS Nov 10)  
 a. Schistocytes b. Elliptocytes  
 c. Microspherocytes d. Polychromasia
- 100. A 5 year old male child presents with episodic jaundice and anemia since birth. Which of the following is least likely diagnosis?**  
 a. Hereditary spherocytosis  
 b. Sickle cell disease  
 c. PNH  
 d. G6PD deficiency

#### ABNORMALITIES OF HEMOGLOBIN

- 101. A 6-year-old patient with anemia, on electrophoresis shows HbF 90%, Hb A<sub>2</sub>/=3%. Which of the following will be seen on peripheral smear?** (AIIMS Nov 2017)




- a. A, B  
 c. ABC  
 b. A, C  
 d. B, C
- 102. CBC done on a patient who had come for blood transfusion revealed MCV 56fl, Hb 13 gm%, MCHC 32 gm/dl & RDW 14, RBC count 6million/ul**  
 a. Sideroblastic anaemia b.  $\beta$ -Thalassemia disease  
 c.  $\beta$ -Thalassemia trait d. Iron deficiency anemia






- 103. In a patient suffering with sickle cell anemia, electrophoretic mobility of HbS in relation to HbA will be?**  
(AIIMS Nov 2015)  
a. Retarded  
b. Accelerated  
c. Same  
d. Will depend upon concentration of HbS
- 104. Screening test for thalassemia** (Recent Question 2015)  
a. Alkali denaturation test  
b. Kleihauer test  
c. Hb electrophoresis  
d. NESTROFT
- 105. Bart hemoglobin is tetramer of** (Recent Question 2015)  
a.  $\alpha$  chain  
b.  $\beta$  chain  
c.  $\gamma$  chain  
d.  $\delta$  chain
- 106. Hemoglobin H disease is caused by deletion**  
(Recent Question 2015)  
a. Single  $\alpha$  globin chain  
b. Two  $\alpha$  globin chains  
c. Three  $\alpha$  globin chains  
d. All  $\alpha$  globin chain
- 107. All the following features precipitate sickling of HbS except**  
(Recent Question 2015)  
a. Hypoxia  
b. Dehydration  
c. Infections  
d. Alkalosis
- 108. Autosplenectomy is seen in**  
(WB PG 2016) (Recent Question 2015)  
a. Hereditary spherocytosis  
b. Sickle cell anemia  
c. Thalassemia  
d. G6PD deficiency
- 109. Sickle cell disease is less severe in**  
(Recent Question 2015)  
a. HbSC  
b. Females  
c. Thalassemia  
d. All the above
- 110. A young adult presents with history of bone pains, recurrent chest infections and dyspnea. Peripheral blood smear shows. (refer Q149) what is your diagnosis?**  
(Recent Question 2015)  
a. Megaloblastic anemia  
b. Hereditary spherocytosis  
c. Sickle cell anemia  
d. Thalassemia
- 111. In HbM the position of point mutation**  
(Recent Question 2015)  
a.  $\beta$  chain, 87th codon, Histidine  $\rightarrow$  Tyrosine  
b.  $\alpha$  chain, 87th codon, Histidine  $\rightarrow$  Tyrosine  
c.  $\beta$  chain, 6th codon, Glutamine  $\rightarrow$  Valine  
d.  $\beta$  chain, 6th codon, Glutamine  $\rightarrow$  Lysine
- 112. Mutation in sickle cells disease is an example of**  
(Recent Question 2015)  
a. Silence mutation  
b. Missense mutation  
c. Nonsense mutation  
d. Frameshift mutation
- 113. In  $\alpha$ -thalassemia, HbBarts is said when number of gene loci affected is:**  
(PGI May 2015)  
a. 1  
b. 2  
c. 3  
d. 4  
e. None

- 114. 25/f came to OPD with anemia, jaundice and joint pain. Which of the following statements is not true about the condition?**  
(APPGMEE 2015)



- a. She can have pulmonary syndrome  
b. She can have retinopathy  
c. Hydroxyurea can help her  
d. HbF and HbA2 will be undetectable
- 115. Boy born to mother who is O-ve has blood group B+ve. He developed jaundice on day 1, Peripheral blood smear is given, which cell is absent?**  
(Recent Question 2015)  
a. Sickle cell  
b. Anisocytosis  
c. Target cell  
d. Schistocytes



- 116. All are seen in sickle cell anemia EXCEPT?**  
a. Target cells  
b. Jaundice (DNB June 11)  
c. Reticulocytosis  
d. High hematocrit
- 117. Heterozygous sickle cell anemia gives protection against:**  
(AI 10)  
a. G6PD  
b. Malaria  
c. Thalassemia  
d. Dengue fever
- 118. HbH is associated with:**  
(AI 11)  
a. Deletion of 3 alpha genes  
b. Deletion of 4 alpha genes  
c. Deletion of 2 beta genes  
d. Deletion of 1 beta genes
- 119. NESTROFT test is used in screening of -**  
(AIIMS Nov 11)  
a. Thalassemia  
b. Autoimmune hemolytic anemia  
c. Spherocytosis  
d. G6PD deficiency





## OTHER HEMOLYTIC ANEMIAS

- 120. Which of the following tests is gold standard to diagnose a case of PNH?** (Recent exam 2018)
- HAMS test
  - Sucrose lysine test
  - Flow cytometry
  - Bone marrow
- 121. CD59 deficiency leads to:**
- PNH
  - Chédiak-Higashi disease
  - Hairy cell leukemia
  - Hemolytic uremic syndrome
- 122. An abnormal Ham test is most likely associated with which of the following?** (AIIMS Nov 11)
- Defect in spectrin
  - Defective GPI anchor
  - Defect in complement
  - Mannose-binding residue defect
- 123. PNH is associated with a deficiency of:** (AI 10)
- DAF
  - MIRL
  - GPI anchored protein
  - All of the above
- 124. Mutation of which of the following gene is most important in paroxysmal nocturnal hemoglobinuria** (AI 10)
- Decay accelerating factor (DAF)
  - Membrane inhibitor of reactive lysis (MIRL)
  - Glycosylphosphatidyl inositol (GPI)
  - CD8 binding protein
- 125. Schistocyte is/are found in:** (PGI May 10)
- TTP
  - DIC
  - Severe iron deficiency
  - March hemoglobinuria
  - Iron deficiency anemia
- 126. Warm antibody hemolytic anemia is seen in?** (DNB June 10)
- Methyldopa
  - EBV infection
  - Quinine
  - Mycoplasma infection

## NUTRITIONAL ANEMIAS

- 127. Which of the following is the best indicator to assess iron deficiency anemia:** (AIIMS Nov 2019)
- TIBC increased, Ferritin increased, Transferrin saturation increased, Serum transferrin receptors increased
  - TIBC reduced, Ferritin reduced, Transferrin saturation reduced, Serum transferrin receptors decreased
  - TIBC increased, Ferritin reduced, Transferrin saturation reduced, Serum transferrin receptors decreased
  - TIBC increased, Ferritin reduced, Transferrin saturation reduced, Serum transferrin receptors increased
- 128. Hepcidin inhibits the function of which of the following?** (AIIMS Nov 2019)
- DMT-1
  - Hephaestin
  - Ceruloplasmin
  - Ferroportin
- 129. Iron enters enterocyte by:** (AIIMS Nov 2019)
- DMT-1
  - Ferroportin
  - Ferritin
  - Hephaestin

- 130. Hypoproliferative anemia(s) is/are:** (PGI May 2019)
- Anemia of chronic inflammation
  - Anemia due to chronic renal disease
  - Iron deficiency anemia
  - Myelodysplastic syndrome
  - Sickle cell anemia
- 131. Which of the following is/are lower in anaemia of chronic diseases as compared to iron deficiency anaemia?** (PGI May 18)
- TIBC (Total iron binding capacity)
  - Serum ferritin
  - Serum iron
  - Transferrin saturation
  - Iron stores
- 132. Microcytosis on peripheral blood smear is/are seen in?** (PGI May 18)
- Iron deficiency anemia
  - Thalassaemia
  - Sideroblastic anemia
  - Vit. B12 deficiency
  - Folate deficiency
- 133. Differential diagnosis for microcytic anemia is?** (PGI Nov 2018)
- Lead poisoning
  - Sideroblastic anemia
  - Occult blood loss
  - Atransferrinemia
- 134. All decrease in iron deficiency anemia except?** (AIIMS Nov 2017)
- Ferritin
  - TIBC
  - Iron
  - Transferrin
- 135. Which of these is not involved in iron metabolism?** (AIIMS Nov 2017)
- Hepcidin
  - Ferroportin
  - Transthyretin
  - Ceruloplasmin
- 136. Which of the following would be the findings of sideroblastic anemia?** (AIIMS May 2017)
- Coarse basophilic stippling in lead poisoning
  - erythroid hypoplasia in marrow
  - Dimorphic anaemia
  - Increase transferrin saturation
  - Increase MCHC
- 137. Which of the following is accurate regarding the internal iron homeostasis in iron deficiency anemia** (JIPMER 2018)
- Transferrin receptor 1 - iron responding elements increases transferrin receptor mRNA concentration and synthesis
  - Ferritin mRNA concentration - iron response element increases and ferritin synthesis decreases
  - Ferritin mRNA concentration - iron response element decreases and Ferritin synthesis increases
  - Transferrin receptor 1 - iron responding elements decreases transferrin receptor mRNA concentration and increases synthesis
- 138. Which of the following are true regarding a child with iron deficiency anaemia** (JIPMER 2017)
- Raised MCV
  - Raised transferrin saturation
  - Increased TIBC
  - Increased ferritin



- 139. Following are causes of megaloblastic anemia except?**  
 a. Defect in DNA synthesis (PGI Nov 2016)  
 b. Folic acid deficiency  
 c. Lead toxicity d. Vit B12 deficiency

- 140. Match List-I with List-II and select the correct answer using the code given below the Lists?**

(Recent Question 2016-17)

List-I (Blood picture)	List-II (Type of Anemia)
A. Microcytic, hypochromic red cells	1. Vitamin B12 deficiency anemia
B. Macrocytic, hypochromic red cells	2. Thalassemia major
C. Large number of early, intermediate and late erythroblasts	3. Aplastic anemia
D. Low reticulocyte count	4. Iron-deficiency anemia

**Code:**

	A	B	C	D
a.	1	4	2	3
b.	2	3	4	1
c.	4	1	2	3
d.	3	1	2	4

- 141. Which of the following conditions does not cause pancytopenia?** (Recent Question 2016)

- a. Hypersplenism  
 b. Aplastic anemia  
 c. Cancer infiltrating the bone-marrow  
 d. Hemolysis from G6PD enzyme deficiency

- 142. Which of the following is least likely?** (Recent Question 2016-17)

- a. Celiac disease  
 b. Thalassemia major  
 c. Nutritional anemia  
 d. Paroxysmal nocturnal haemoglobinuria

- 143. Iron metabolism and regulation are important for RBC precursor cell. Which of the following helps in regulation of iron metabolism but is not specific for iron?**

- a. Hepsidin b. DMT-1 (AIIMS Nov 2015)  
 c. Ferroportin d. Ferritin

- 144. Transfer of Iron from enterocyte to plasma is inhibited by?** (Recent Question 2016-17)

- a. Hepsidin b. DMT-1  
 c. Ferroportin d. Hepestin

- 145. Iron stores are best indicated by?** (Recent Question 2016)

- a. S. ferritin b. S. Iron  
 c. S. transferrin d. TIBC

- 146. Release ferroportin store is controlled by?** (Recent Question 2016)

- a. Hepsidin b. Transferrin  
 c. Ferritin d. Hepoxin

- 147. Which of the following correctly describes principle of Prussian blue stain?** (Recent Question 2016)

- a. Ferrocyanide to ferricyanide  
 b. Ferrocyanide to ferroferric cyanide  
 c. Ferroferric cyanide to ferrocyanide  
 d. Ferrocyanide to ferricferro cyanide

- 148. True about Hereditary Hemochromatosis a/e:** (Recent Question 2016)

- a. Mutations in the HFE gene  
 b. Autosomal recessive disorder  
 c. Excess iron affects organ function  
 d. Diagnosed by serum ferritin

- 149. Pappenheimer body is seen in:** (Recent Question 2016)

- a. Sideroblast b. Siderocyte  
 c. Reticulocyte d. Bite cells

- 150. Calculate iron deficit for a 50 kg person, with Hb-5g/dL. Add 1000 mg for stores.** (Recent Question 2015)

- a. 2150 mg b. 1650 mg  
 c. 1150 mg d. 1575 mg

- 151. A 50 year old male has Hb 9.2g/dL and MCV112fL. Next step:** (Recent Question 2015)

- a. Iron supplement b. Folate supplement  
 c. Transfusion d. Check B12 and folate level

- 152. A 40 year old female presents with signs of heart failure. Hb-6 g/dL, MCV-112. Next step** (Recent Question 2015)

- a. Check B12 and folate levels  
 b. Blood transfusion  
 c. Start iron tablets  
 d. Start folate supplementation

- 153. MCV is increased in the following anemia** (Recent Question 2015)

- a. Iron deficiency anemia  
 b. Sideroblastic anemia  
 c. Anemia of chronic disease  
 d. Folate deficiency anemia

- 154. Find the false statement regarding megaloblastic anemia** (Recent Question 2015)

- a. Hypersegmented neutrophils are the earliest manifestation  
 b. Reticulocyte count decreased  
 c. Hypercellular bone marrow  
 d. MCHC is increased

- 155. 20-year-old female present with features of anemia. Blood tests: Hb-5g/dL, MCV - 52 fL, MCHC-20 g/dL, PCV - 32%. Diagnosis** (Recent Question 2015)

- a. Phenytoin toxicity  
 b. Fish tape worm infection  
 c. Hook worm infection  
 d. Blind loop syndrome

- 156. Most sensitive indicator of iron deficiency anemia** (Recent Question 2015)

- a. Packed cell volume b. Hemoglobin  
 c. Serum ferritin d. Serum iron

- 157. Macrocytic anemia with MCV> 110 fL is seen in** (Recent Question 2015)


- a. Thiamine deficiency b. B12 deficiency  
 c. Hypothyroidism d. Phenytoin toxicity

- 158. Iron deficiency anemia and anemia of chronic disease can be differentiated by the following parameter** (Recent Question 2015)

- a. Microcytic, hypochromic anemia  
 b. Serum iron  
 c. TIBC  
 d. Transferrin saturation



159. In a patient with thalassemia major, who had received multiple blood transfusions, the serum iron overload can be detected by (Recent Question 2015)
- Serum ferritin level
  - Blood iron level
  - Total iron binding capacity
  - Blood hemoglobin level
160. Iron absorption is increased by (Recent Question 2015)
- Phytates
  - Tannates
  - Plant food
  - Ascorbic acid
161. Microcytic anemia is not seen in the following condition (Recent Question 2015)
- Osteomyelitis
  - Leukopenia
  - Papillary necrosis
  - Stroke
162. True statement about anemia of chronic disease (Recent Question 2015)
- Increased serum ferritin
  - Increased serum iron
  - Increased TIBC
  - Increased transferrin saturation
163. Average serum ferritin value in males (Recent Question 2015)
- 50 mg/dL
  - 100 mg/dL
  - 200 mg/dL
  - 500 mg/dL
164. Most important inflammatory mediator, involved in anemia of chronic disease? (Recent Question 2015)
- IL-1
  - IL-6
  - TNF $\alpha$
  - IFN- $\gamma$
165. True about iron deficiency anemia (JIPMER 2015)
- Parenteral iron is indicated when anemia response slowly to oral iron
  - Ferrous sulphate 200mg has less elemental iron than the same dose of ferrous gluconate
  - Sustained release iron is a useful way of giving larger doses
  - Absorption of iron is increased by ascorbic acid
166. This patient came with anemia. This peripheral smear shows. (APPGMEE 2015)



- Microcytic hypochromic anemia due to iron deficiency
  - Macrocytic anemia
  - Target cells of thalassemia
  - Spherocytes of autoimmune hemolytic anemia
167. Which of the following is not associated with microcytic hypochromic anemia? (APPGMEE 2015)
- Chronic Lead poisoning
  - Thalassemia
  - Hereditary spherocytosis
  - Iron deficiency
168. Red cells containing granules of non-heme iron, which gives positive Prussian blue reaction with Perl's stain as well as stain with Romanowsky dyes (referred to as pappenheimer bodies) are known as: (AP 2012)
- Schistocytes
  - Spherocytes
  - Sideroblasts
  - Siderocytes
169. A 5-year-old child presents with pallor and constipation with Hb-6 gm%. Which is the most appropriate therapy? (Recent Question 2015)
- Packed cell RBC's
  - Oral iron therapy
  - Parenteral iron
  - Hematologist referral
170. Most sensitive marker in iron deficiency anemia: (Recent Question 2014)
- TIBC
  - Serum ferritin
  - Serum Iron
  - Serum transferrin saturation
171. Normal transferrin saturated with iron is ? (Recent Question 2014)
- 20%
  - 35%
  - 50%
  - 70%
172. Response to iron in iron deficiency anemia is denoted by? (Recent Question 2014, DNB 09)
- Restoration of enzymes
  - Reticulocytosis
  - Increase in iron binding capacity
  - Increase in hemoglobin
173. Iron deficiency causes (Recent Question 2014)
- Megaloblastic anemia
  - Microcytic hypochromic anemia
  - Macrocytic hypochromic anemia
  - Microcytic hypochromic anemia
174. Bone marrow in lead poisoning contains? (JIPMER 2014)
- Ringed siderocytes
  - Giant metamyelocytes
  - Dwarf megakaryocytes
  - Fibrotic changes
175. All of the following statements about iron deficiency anemia are true except? (AIIMS Nov 2013)
- Latent iron deficiency is most common presentation in India
  - Transferrin saturation is less than 16%
  - Serum ferritin is the earliest marker
  - It can present without detectable abnormalities
176. Leptocyte in blood smears seen in? (Recent Question 2013)
- Sickle cell anemia
  - Thalassemia
  - Post splenectomy
  - Uremia
177. First sign of improvement in oral iron therapy is? (DNB Aug. 12 Pattern) (WBPB 2015)
- Reticulocytosis
  - Raise of hemoglobin
  - Raise in RBC count
  - Increase in ESR
178. Parameter increased in IDA? (JIPMER 2012)
- RBC protoporphyrin
  - Serum Iron level
  - Serum ferritin
  - Transferrin saturation



- 179. A 35-year-old lady on treatment for rheumatoid arthritis has following lab findings: Hb-9 gm/dl, MCV- 55 fl, serum iron-30 ug/dl, ferritin- 200 ng/ml, TIBC- 298 ug/dl. What is the most probable diagnosis? (AIIMS Nov 11)**  
 a. Thalassemia minor      b. Thalassemia major  
 c. Anemia of chronic disease  
 d. Iron deficiency anemia
- 180. 33-years-old alcoholic on ATT presents with increased serum iron and increased transferrin saturation. Diagnosis? (JIPMER 11)**  
 a. Iron deficiency anemia      b. Sideroblastic anemia  
 c. Megaloblastic anemia      d. Anemia of chronic disease
- 181. A patient with microcytic hypochromic anemia. Hb- 9 g%, serum iron-20 microg/dl, ferritin level-800 ng/ml, transferrin percentage saturation is 64%. What is possible diagnosis? (AIIMS Nov 10)**  
 a. Atransferrinemia  
 b. Iron deficiency anemia  
 c. DMT 1 mutation  
 d. Anemia of chronic disorder
- 182. A 60-years-old female presents with history of 8 blood transfusions in 2 yrs. Her Hb-6.0gm/dl, TLC-5800/cumm, platelet- 3.4 lakhs/cumm, MCV-60 fl, RBC-2.1 lakhs/mm<sup>3</sup>. She is having hypochromic microcytic anemia. Which investigation is not needed? (AIIMS Nov 10)**  
 a. Evaluation for pulmonary hemosiderosis  
 b. Urinary hemosiderin  
 c. Bone marrow examination  
 d. GI endoscopy
- 183. A man presents with fatigue. Hemogram analysis done suggested low Hb, high MCV. The next investigation is? (Recent Question 2015)**  
 a. Vit B<sub>12</sub>/folate levels      b. Bone Marrow  
 c. S. Iron studies      d. Reticulocyte count
- 184. Which of the following does not indicate Megaloblastic anemia? (AIIMS May 2013/ Nov 2012)**  
 a. Increased reticulocyte count  
 b. Raised Bilirubin  
 c. Mild splenomegaly  
 d. Nucleated RBC
- 185. Symptoms of folate deficiency occur in association with which of the following? (PGI May 2011)**  
 a. Nitrous oxide      b. Pyrimethamine  
 c. Sodium nitroprusside      d. Sodium valproate  
 e. 5-FU
- 186. Macrocytic anemia is seen in all EXCEPT. (DNB June 10)**  
 a. Vitamin B<sub>12</sub> deficiency  
 b. Hemolytic anemia  
 c. Post hemorrhagic anemia  
 d. Anemia of chronic disease
- 187. Criteria for severe aplastic anemia are all except? (JIPMER 2017)**  
 a. BM cellularity <25%  
 b. Reticulocyte <60,000/ul  
 c. Platelet <20,000/ul  
 d. Absolute neutrophil count <1500/ul
- 188. Bone marrow failure with neutropenia and exocrine pancreatic deficiency is a feature of? (JIPMER 2017)**  
 a. Fanconi anemia  
 b. Dyskeratosis congenita  
 c. Diamond Blackfan syndrome  
 d. Diamond Schwachman syndrome
- 189. Parvovirus preferentially involves in which of the following cells:**  
 a. Erythroid progenitors      b. Myeloid precursors  
 c. Megakaryocytes      d. Both a & c
- 190. All of the following are causes of increased reticulocyte count except?**  
 a. Treatment of megaloblastic anaemia with Vit B12  
 b. Congenital dyserythropoetic anaemia  
 c. Hereditary spherocytosis  
 d. Aplastic anaemia
- 191. The following drug is not associated with pure red cell aplasia: (Recent Question 2015)**  
 a. Phenytoin      b. Isoniazid  
 c. Erythropoietin      d. None of the above
- 192. Pancytopenia with hypocellular bone marrow seen in: (Recent Question 2015)**  
 a. Fanconi's anemia  
 b. Paroxysmal nocturnal hemoglobinuria  
 c. Hairy cell leukemia  
 d. Myelophthisis
- 193. 2-year-old child presents with short stature and café-au-lait spots. Bone marrow aspiration yields a little material and mostly containing fat. What is your diagnosis: (Recent Question 2015)**  
 a. Fanconi anemia  
 b. Dyskeratosis congenita  
 c. Tuberous sclerosis  
 d. Osteogenesis imperfecta
- 194. Dry tap is a feature: (Recent Question 2015)**  
 a. Anemia of chronic disease  
 b. Megaloblastic anemia  
 c. Aplastic anemia  
 d. Sickle cell anemia
- 195. MC tumor associated with pure red cell aplasia: (Recent Question 2015)**  
 a. Hepatoma  
 b. Hodgkins lymphoma  
 c. Thymoma  
 d. Bronchogenic carcinoma
- 196. 2/F presented with maculopapular rash 24hrs after onset of mild fever. There was prominent erythema over the cheek. The causative organism also causes?**  
 a. CMV      b. ALL (JIPMER 2014)  
 c. DIC      d. PRCA
- 197. Pure red cell aplasia is associated with all except? (AIIMS Nov 2013)**  
 a. ABO incompatibility after renal transplant  
 b. 5q- syndrome  
 c. Drugs  
 d. Large granular lymphocytic leukemia
- 198. Pancytopenia with cellular bone marrow is seen in all except: (WBPG 2015)**  
 a. Megaloblastic Anemia      b. MDS  
 c. PNH      d. G6PD deficiency

#### HYPOPROLIFERATIVE ANEMIAS





199. **Prominent reticulocytosis is a feature of:** (WBPG 2015)
- Aplastic Anemia
  - Hemolytic Anemia
  - Nutritional Anemia
  - Anemia of chronic disease
200. **Aplastic anemia can progress to all except:** (DPG10)
- AML
  - Myelodysplastic anemia
  - Pure red cell aplasia
  - Paroxysmal nocturnal hemoglobinuria

201. **True about Dyskeratosis congenita:** (PGI May 2011)
- Pancytopenia
  - Nail dystrophy
  - Hyperkeratosis
  - X-linked
  - Leukoplakia



## Answers with Explanations

1. **Ans. (d) Babesiosis**



Babesia microti in a thin blood smear. Note the classic "Maltese Cross" tetrad-form in the infected RBC in the lower part of the image.

2. **Ans. (c) Gray**

Hemoguard stopper	Tube content	Determination
	Serum separator tube (SST)	All biochemistry not mentioned elsewhere (1 tube), microbiology (1 tube)
	Heparin	Chromosome studies, lead, amino acids, troponin
	Fluoride/oxalate	Glucose
	EDTA	Full blood count (FBC) and ESR, C3/C4, hemoglobin A1c, homocysteine, ACTH
	Plain (No additive)	LDH, Ca, drugs (Phenytoin, theophylline, lithium), endocrine testing (except thyroid)
	Sodium citrate	Coagulation testing, PT, INR, APTT, D-Dimer, etc...
	ESR	Westergren Sedimentation Rate; requires full draw

3. **Ans. (b) Corrected ret  $\times$  hct / 45 | maturation index**

(Ref: Harrison 19th ed/pg 396)

4. **Ans. (d) SF3B1**

5. **Ans. (a) Ask the patient his name\*verify from file Label the sample at bedside  $\rightarrow$  collect blood**

6. **Ans. (d) Lithium heparin**

(Ref: Henry Clinical chemistry)

Heparin, a mucopolysulfuric acid, is an effective anticoagulant available as lithium heparin (LiHep) and sodium heparin (NaHep) in green-top tubes. Heparin accelerates the action of antithrombin III, neutralizing thrombin and preventing the formation of fibrin. Heparin has an advantage over EDTA as an anticoagulant, as it does not affect levels of ions such as calcium. However, heparin can interfere with some immunoassays. Heparin should not be used for coagulation or hematology testing.

7. **Ans. (b) Anisocytosis**

### Red Cell Distribution width

- The RDW is an index of the variation in cell size and volume within the red cell population.

### Normal Value

- RDW-SD 39-46 fL, RDW-CV 11.6-14.6% in adult

### Condition with Increased RDW:

- Iron deficiency anaemia
- Folate/vit B12 deficiency anaemia

8. **Ans. (c) 3** (Ref: Harrison 19th Edition. Page No. 396)

### Reticulocyte Count has two Corrections

- Correction for the degree of anaemia (Absolute reticulocyte count) and Correction for the prolonged duration of stay of reticulocyte in peripheral smear.

Corrected reticulocyte count  $\rightarrow$  Reticulocyte count  $\times$  Hb or



Hct of patient/Normal Hb or Hct

$$= 9 \times 5/15$$

$$= 9/3 = 3$$

Corrected reticulocyte count = 3 %

9. Ans. (a) **Fixes cells to slide**

(Ref: Bancroft's Histochemical Techniques, Page 90-98)

### Acetone Free Methanol is the Fixative in Leishmann Stain

- If acetone is there it will destroy the cell membrane

10. Ans. (b) **Folic acid deficiency anemia**

(Ref: Robbins 9th/pg 631)

- **Hematocrit** or **Packed cell volume**: ratio of the volume of RBCs to total volume of blood
- Hct or PCV = MCV  $\times$  RBC concentration

Now coming back to the question:

- $MCV = PCV / RBC \text{ count (in millions)} = 30 / 0.2 = 150$  (Increased)
- $MCH = Hb / RBC \text{ count} = 9.3 / 0.2 = 46$  (Increased)
- $MCHC = Hb / MCV = 31$  (Normal)

The answer here is clearly macrocytic anemia (increased MCV as macrocytosis) so the correct option is b. Folic acid deficiency anemia

11. Ans. (c) **P. falciparum**

(Ref: Practical hematology, Dacie pg 110)

12. Ans. (a)  **$1.5 \times 10^6$  cells/second**

(Ref: Wintrobe's 12th ed/pg 107-108)

- Bone marrow produces approximately **2.4 million new erythrocytes per second** ( $2.4 \times 10^6/s$ ) in human adults.
- They circulate for about 100–120 days
- The closest to this data is option A (hence the answer)

13. Ans. (b) **Acanthocyte** (Ref: Dacie 11th/pg 85)

14. Ans. (a) **Liver** (Ref: Robbins 9th/pg 618; 8th/pg 628)

Sources of erythropoietin:

**Fetus:** Liver

**Adult:**

- Interstitial cells in peritubular capillary bed of kidneys (85%)
- Perivenous hepatocytes in the liver (15%)
- Brain (protective effect against excitotoxic damage triggered by hypoxia)
- Uterus and oviducts (mediate estrogen-dependent angiogenesis).

15. Ans. (a) **Orthochromic normoblast** (Ref: R 9th/pg 580)

16. Ans. (a) **Pro erythroblast** (Ref: Robbins 9th/pg 580)

17. Ans. (c) **Liver disease** (Ref: Dacie 11th/pg 85)

18. Ans. (c) **Echinocytes** (Ref: Dacie 11th/pg 85)

19. Ans. (a) **Malaria** (Ref: Robbins 9th/pg 625)

- Most common cause of splenic rupture is Trauma
- Most common cause of spontaneous splenic rupture in the world is Infectious Mononucleosis
- Most common cause of spontaneous splenic rupture in India is Malaria

20. Ans. (b) **Reticulocytes** (Ref: Dacie 11th/pg 34)

21. Ans. (b) **0.48 %** (Ref: Dacie 11th/pg 246)

When osmotic fragility is normal, RBC's begin to hemolyse when suspended in 0.48 % saline

### Osmotic Fragility Test

- Gives an indication of the **surface area/volume ratio** of erythrocytes.
- Normally RBCs start hemolysing at 0.5% NaCl and is completely hemolysed at 0.3% NaCl
- It is useful in the diagnosis of **hereditary spherocytosis** & screening for **thalassaemia**.
- Red cells that are **spherocytic** have **increased osmotic fragility** i.e. take up less water in a hypotonic solution before rupturing than do normal red cells.

Conditions associated with:

Increased osmotic fragility (OF)	Decreased (OF)
<ul style="list-style-type: none"> <li>• Hereditary spherocytosis</li> <li>• Hereditary elliptocytosis (HE)</li> <li>• Hereditary stomatocytosis</li> <li>• Autoimmune hemolytic anemia</li> </ul>	<ul style="list-style-type: none"> <li>• Thalassaemia</li> <li>• Enzyme abnormalities</li> <li>• Hereditary xerocytosis</li> <li>• Iron deficiency</li> </ul>

22. Ans (a) **Stained by supravital staining**

(Ref: Dacie 11th/pg 34; Refer Ans 2 Above)

- Reticulocytes stained on Romanowsky stain (used for usual hematological staining) are called polychromatophils;
- However it is not a specific stain;

23. Ans. (c) **1:1** (Ref: Robbins 9th/pg 582; 8th/pg 592)

Ratio of fat cells and blood cells in bone marrow is 1:1

24. Ans. (b) **Splenectomy**

(Ref: Dacie 11th/pg 85; Robbins 9th/pg 632; 8th/pg 643)

### Howell-Jolly Bodies

- Howell-Jolly bodies are **nuclear remnants**.
- They are **small, round cytoplasmic inclusions** that **stain purple** on a Romanowsky stain.
- Seen after **splenectomy**, in **splenic atrophy**, **pernicious anemia**, **coeliac disease**

25. Ans. (a) **Megaloblastic anemia**

(Ref: Wintrobe's Atlas of Clinical Hematology, 1st Edition, 2007, chapter 1)



26. Ans. (b) **A critical concentration of reduced hemoglobin is required** (Ref: Harrison 18th/pg Chapter 35)

- Cyanosis becomes apparent when the concentration of **reduced hemoglobin in capillary blood** > 4 g/dL.
- It is the **absolute**, rather than the **relative**, quantity of **reduced Hb** that is **important in producing cyanosis**.
- In a patient with severe anemia, the **relative** quantity of reduced Hb in the venous blood may be very large, but the **absolute** quantity of reduced Hb is still small. Therefore, patients with **severe anemia & even marked arterial desaturation** may not display cyanosis.

27. Ans. (c) **ESR is raised because of increased RBC aggregate**

(Ref: Dacie 11th/pg102)

In **TB**, **ESR is raised** because of increased RBC aggregates. ESR is increased by any cause or focus of inflammation.

28. Ans. (a) **Increased ESR** (Ref: Dacie 11th/pg105)

In **polycythemia vera**, **ESR is decreased** due to decrease in rouleaux formation or increase the RBC surface area to volume ratio.

29. Ans. (a) **Acquired hemolytic anemia**

(Ref: Wintrobe's 12th; Atlas of Clinical Hematology, 1st Edition, 2007 chapter 1)

**Cabot's ring** in RBC is seen in **Acquired hemolytic anemia**. **Pernicious Anemia** > **Hemolytic Anemias** > **Post Splenectomy**; **Lead poisoning**

30. Ans. (d) **Potassium oxalate + sodium fluoride**

(Ref: Teitz clinical chemistry)

**Anticoagulant** used to send blood sample for **glucose** estimation is **Potassium oxalate + sodium fluoride**

31. Ans. (b, d, e); **b. In newborn Hb level is 15 gm%; d. Macrocytosis may occur in Celiac disease and e. Normocytic normochromic anemia is found in chronic diseases**

(Ref: Robbins 9th/pg 645-654; Wintrobe's 12th/pg 1221)

32. Ans. (b) **Schistocytes**

(Ref: Hematology: Basic principles and practice, 6th edition, 2013 chapter 32; Robbins 9th/pg 640; 8th/pg 650)

Discussing the options one by one

- Target cells**: Seen in **Thalassemia**, **Liver disease**, **Abetalipoproteinemia**, **severe Iron deficiency anemia**
- Schistocytes**: Seen in **Microangiopathic hemolytic anemia**
- Acanthocytes**: See table below
- Basophilic stippling**: See table below

33. Ans. (c) **Iron** (Ref: Wintrobe's Atlas of Clinical Hematology, 1st Edition, 2007 chapter 1)

34. Ans. (a) **Increased serum immunoglobulin level**; (e) **Increased level of C-reactive protein** (Ref: Dacie 11th/pg 105)

**Erythrocyte Sedimentation Rate is zero in Afibrinogenemia**

Increased ESR	Lower ESR
<ul style="list-style-type: none"> <li>• Old age</li> <li>• Female</li> <li>• <b>Pregnancy</b></li> <li>• <b>Anemia</b></li> <li>• Paraprotein (Multiple Myeloma)</li> <li>• <b>Hypergammaglobulinemia</b></li> <li>• Macrocytosis</li> <li>• Elevated fibrinogen (infection, inflammation malignancy)</li> <li>• <b>Technical factors</b>: Dilution, high temperature</li> </ul>	<ul style="list-style-type: none"> <li>• Extreme leukocytosis</li> <li>• <b>Polycythemia</b></li> <li>• Spherocytosis, microcytosis</li> <li>• Hyperviscosity</li> <li>• Low protein, fibrinogen, gammaglobulins</li> <li>• Technical factors; dilution, clotted sample</li> <li>• Afibrinogenemia</li> </ul>
	<b>No effect</b> <ul style="list-style-type: none"> <li>• Obesity</li> <li>• Body temperature</li> <li>• Recent meal</li> <li>• Aspirin, NSAIDs</li> </ul>

35. Ans. (d) **Multiple myeloma** (Ref: Dacie 11th/pg 105)

36. Ans. (a) **Sickle cell Anemia** (Ref: Dacie 11th/pg 105)

37. Ans. (b, c, e) **b. Spleen size; c. MCHC; e. LDH**

Decreased MCV, Increased MCHC, and ↑ LDH. It also causes splenomegaly.

38. Ans. (a) **CML**

CLL IS USUALLY ASSOCIATED WITH AIHA.

**Extracorporeal membrane oxygenation** (ECMO) is a treatment that uses a pump to circulate blood through an artificial lung back into the bloodstream of very ill baby. This system provides heart-lung bypass support outside the baby's body. This can be associated with hemolysis. Continuous renal replacement therapy (CRRT) can also have secondary hemolysis.

39. Ans. (b) **Hereditary spherocytosis**

**A flow cytometry-based test using eosin-5-maleimide (EMA) dye is used for diagnosis of hereditary spherocytosis (HS)**. Eosin-5-maleimide (EMA) dye, which reacts covalently with lysine-430 on the first extracellular loop of band-3 protein helps detecting HS, as vertical bonds are usually broken in HS.

40. Ans. (a, b, c) **a. HUS; b. TTP; c. Prosthetic heart valves**

41. Ans. (a, b, c, e) **a. Hallervorden-Spatz disease; b. Abetalipoproteinemia; c. Severe liver disease; e. McLeod syndrome**

42. Ans. (a) **G6PD deficiency** (Ref: Wintrobe's 13/p 2089)

The peripheral blood smear IN **G6PD def** contains spherocytes and eccentrocytes or "blister" cells.



43. Ans. (a) **I** (Ref: Robbins 9th/pg 643)

44. Ans. (a) **Cold AIHA**

The given figure is of RBC agglutination. The history is suggestive of agglutination occurring in the month of December that is cold induced. Cold Agglutinin Type. This form of immunohemolytic anemia is caused by IgM antibodies that bind red cells avidly at low temperatures (0°–4°C).

45. Ans. (d) **Decreased reticulocyte**

46. Ans. (b) **Increased direct bilirubin**

Increased indirect bilirubin and not direct bilirubin are the findings in hemolytic anemia

	MCV	% S	% A	% A <sub>2</sub>	% F
AS	N	35–38	62–65	< 3.5	<1
SS	N	88–93	0	<3.5	5–10
S/β° thalassaemia	L	88–93	0	>3.5	5–10

47. Ans. (d) **HbA2 is increased**

(Ref: Practical hematology by Dacie 11th ed/ pg 312)

In sickle cell anemia, HbA2 is decreased and not increased.

48. Ans. (c) **Aplastic anemia**

(Ref: Wintrobe 13th ed / Page 576)

**Coomb's test is positive in Immune hemolytic anemia**

- They can be of 2 types:
    - **Direct Coomb's test**- Detects Antibody on RBC Surface<sup>Q</sup>
    - **Indirect Coomb's test**- Detects Antibody in Serum
- Out of the given options: Aplastic anemia is not a hemolytic anemia and so is automatically the answer here.

49. Ans. (c) **Megaloblastic anemia**

Megaloblastic anemia has a low normal Retic % as it is a nutritional anemia.

50. Ans. (b) **Donor cells have normal survival.**

(Ref: Robbins 9th ed. Pg. 631-632)

This Question is based on the knowledge that in intracorporeal defect of RBCs the defect lies in the RBCs itself and not in spleen or vessels and so any erythrocytes (blood – group matched) are transfused into a patient will not be destroyed. Hence the answer is **donor cells have normal survival.**

51. Ans. (a) **2 α + 2 β** (Ref: Robbins 9th/pg 638-642)

52. Ans. (d) **Penicillin** (Ref: Robbins 9th/pg 644-645)

Penicillin causes hemolysis by mechanism of autoimmune hemolytic anemia and not G6PD deficiency mediated.

53. Ans. (d) **CLL is an important cause**

54. Ans. (a) **Hereditary spherocytosis**

This is a case of anemia (fatigue), splenomegaly (abdominal pain) with jaundice and gall stones since birth. These findings suggest a inherited causes of extravascular hemolysis. The best option is Hereditary spherocytosis.

55. Ans. (d) **Aplastic anemia** (Ref: Robbins 9th/pg 653)

56. Ans. (a) **Older red cells** (Ref: Robbins 9th/pg 634; 8th/pg 644)

Because mature red cells do not synthesize new proteins, G6PD– or G6PD Mediterranean enzyme activities fall quickly to levels inadequate to protect against oxidant stress as red cells age. Thus, older red cells are much more prone to hemolysis than younger ones.

57. Ans. (b) **Ankyrin** (Ref: Robbins 9th/pg 632; 8th/pg 642)

Most common mutation:

- Hereditary spherocytosis is Ankyrin>Band-3>spectrin
- Hereditary elliptocytosis is Spectrin

58. Ans. (a) **Spectrin** (Ref: Robbins 9th/pg 632; 8th/pg 642)

59. Ans. (c) **Quinidine** (Ref: Robbins 9th/pg 634-635)

Quinidine do not carry risk of hemolysis in persons with G-6-PD deficiency

60. Ans. (a) **Plasmodium vivax** (Ref: Harrison 18th/pg 1014)

**Duffy antigen/chemokine receptor (DARC)**, also known as **Fy glycoprotein (FY)** or **CD234 (Cluster of Differentiation 234)**

- Duffy antigen is located on the surface of RBCs, and is named after the patient in which it was discovered.
- glycosylated membrane protein and a non-specific receptor for several chemokines.
- The protein is also the receptor for the human malarial parasites Plasmodium vivax and Plasmodium knowlesi.
- Polymorphisms in this gene are the basis of the Duffy blood group system

61. Ans. (a) **Spectrin** (Ref: Robbins 9th/pg 632; 8th/pg 642)

62. Ans. (b) **Incomplete Antibody**

(Ref: Wintrobe 12th/ed pg 959)

IgM-coated RBCs may spontaneously agglutinate because the pentameric antibody can cross-link RBCs. The capability for **IgM antibodies** to agglutinate saline-suspended RBCs without additional reagents has led to the traditional terminology of “**complete**” antibodies. In contrast, **IgG antibodies** typically require antihuman globulin (AHG) to agglutinate saline-suspended RBCs and are thus termed “**incomplete**” antibodies. Now as warm Antibodies are IgG, so they are “Incomplete antibodies”





**63. Ans. (a) Plasmapheresis**

(Ref: Robbins 9th/pg 941; 8th/pg 952; Harrison 19th/pg 740-745)

**Treatment of Atypical HUS**

- Fresh frozen plasma can induce remission.
- Plasma exchange may benefit patients with deficiency of **Complement Factor H or Factor I** by replenishing the missing protein or removing the antibodies.

**64. Ans. (a) Paroxysmal cold hemoglobinuria**

(Ref: Robbins 9th/pg 643; 8th/pg 653)

**65. Ans. (a) Cyclosporine** (Ref: Wintrobe's pg 1011)

**66. Ans. (a) Parvovirus B19 infection**

(Ref: Harrison 18th/pg 1478)

- Aplastic/hypoplastic crises resulting from parvovirus B19 infection may occur in Hereditary spherocytosis just as is seen in individuals with other chronic hemolytic disorders.
- Parvovirus B<sub>19</sub> selectively infects erythroid precursors and inhibits their growth.
- Erythropoietic arrest leads to a sudden decrease in hemoglobin concentration and reticulocytopenia.

**67. Ans. (a) Autoimmune hemolytic anemia** (Ref: 9th/pg 643)

**68. Ans. (b) Fetomaternal haemorrhage**

(Ref: Oski's Pediatrics: Principles & Practice, chapter 66; Dacie 11th/pg 307)

**APT test for differentiation of fetal & adult (maternal) hemoglobin in stool or vomitus**

- Mix 1 volume of sample with 5 volumes of water.
- Centrifuge mixture and remove clear-pink supernatant solution containing hemoglobin.
- Mix 1 ml of 1% sodium hydroxide with 4 ml supernatant and observe color change after 2 minutes.
  - Remains **pink: Hemoglobin F**
  - Turns **yellow-brown: Hemoglobin A** (maternal blood)

A common approach to evaluating FMH is the **rosette test to screen for the presence of fetal cells** followed by the **Kleihauer-Betke acid elution method to quantify the magnitude of feto-maternal bleed.**

The Kleihauer-Betke acid elution method:

- **Quantitative estimate of the volume of FMH** based on different solubility properties of HbF&HbA.
- On a peripheral smear **treated with an acidic solution** & counterstained, red cells containing acid-soluble **HbA appear as pale ghosts** compared to **deeply red stained red cells containing acid-resistant HbF**
- Can also be used to detect HbF containing cells in  $\beta$ -thalassemia, hereditary persistence of hemoglobin F, sickle cell disease & **myelodysplastic syndrome**.

- Increased HbF is also found in congenital red cell aplasia (Blackfan-Diamond syndrome) & congenital aplastic anemia (Fanconianemia), JMML, presence of erythropoietic stress (haemolysis, bleeding, recovery from acute bone marrow failure) and in pregnancy.

**69. Ans. (d) Haptoglobin increased**

(Ref: Robbins 9th/pg 631-632)

Haptoglobin

- A glycoprotein that is synthesized in the liver.
- It consists of two pairs of  $\alpha$  chains and two pairs of  $\beta$  chains.
- With hemolysis, free Hb readily dissociates into dimers of  $\alpha$  &  $\beta$  chains; the  $\alpha$  chains bind avidly with the  $\beta$  chains of haptoglobin in plasma or serum to form a complex

**70. Ans. (d) Paroxysmal cold hemoglobinuria**

(Ref: Robbins 9th/pg 642; 8th/pg 654)

**Direct globulin test is positive in Paroxysmal cold hemoglobinuria, as anti P antibody is formed.**

**71. Ans. (c) Thalassemia** (Ref: 9th/pg 631-632)

Intra-corporcular hemolytic anemia is seen in thalassemia

**72. Ans. (c) Decreased MCHC** (Ref: 9th/pg 632-633)

In hereditary spherocytosis, **increased** rather than decreased MCHC is seen

**73. Ans. (d) Ringed sideroblast**

(Ref: Robbins 9th/pg 635-636)

**Ring sideroblasts are not seen in sickle cell anemia**, but in Sideroblastic Anemia

**Peripheral smear finding in sickle cell anemia:**

- Anisopoikilocytosis, polychromasia, Increased Retic %
- Irreversibly **sickle RBCs** and **target cells** (increased after autosplenectomy).
- **Howell-Jolly bodies** due to asplenia.<sup>o</sup>

**74. Ans. (a) Plasmodium falciparum**

(Ref: Robbins 9th/pg 635-636; 8th/pg 645-646; Wintrobe's 12th/pg 1038)

Person having heterozygous sickle cell trait is protected from infection of Plasmodium falciparum

**Mechanism by which patients with sickle cell trait are protected against Malaria:**

- **Preferential sickling of parasitized cells** has been seen in the blood of children with sickle cell trait & malaria.
- **Selective removal of sickled cells** from the circulation probably reduces the degree of parasitemia and substantially limits the infectious process.

**75. Ans. (d) Neuro psychiatric disturbances**

(Ref: Robbins 9th/pg 941; 8th/pg 952-953)

Neuro-psychiatric disturbances are not a feature of hemolytic uremic syndrome, but may be seen in TTP



**76. Ans. (c) Malaria**

(Ref: Wintrobe's 12th/pg 1084)

RBCs with common Hemoglobinopathies (i.e.  $\alpha$ - &  $\beta$ -thalassemias, HbS, HbC, HbE) & enzyme (**glucose-6-phosphate dehydrogenase [G6PD]**) defects have shown a **reduced parasite invasion/growth** and an **increased susceptibility to phagocytosis** of the infected RBC as a **malaria-protective effect**.

**77. Ans. (c) Paroxysmal cold hemoglobinuria**

(Ref: Robbins 9th/pg 642; 8th/pg 654)

**Paroxysmal cold hemoglobinuria:**

- Rare **intravascular hemolysis** and **hemoglobinuria**.
- **IgGs Abs<sup>Q</sup>** that bind to the **P blood group<sup>Q</sup>** antigen on RBC (direct antibody testing positive) are called **Donath Landsteiner Antibodies<sup>Q</sup>**
- Binds to RBC's at a low temperature (optimally at 4°C), but when the temperature is shifted to 37°C, lysis of red cells takes place in the presence of complement
- Seen in children following **viral infections<sup>Q</sup>**.

**78. Ans. (d) Absence of fever** (Ref: Robbins 9th/pg 941)

- In both HUS & TTP, there is Microangiopathic hemolytic anemia, Renal failure & Neurological symptoms.
- Fever & neurologic findings are seen in TTP, not in HUS

**79. Ans. (c) APT test**

(Ref: Oski's Pediatrics: Principles & Practice, chapter 66; Dacie 11th/pg 307)

**80. Ans. (a) Sodium hydroxide**

(Ref: Oski's Pediatrics: Principles & Practice, chapter 66; Dacie 11th/pg 307)

**81. Ans. (a)  $\alpha$ -Spectrin**

(Ref: 9th/pg 632-633; 8th/pg 642-643)

**82. Ans. (c) Fetomaternal blood leak**

(Ref: Oski's Pediatrics: Principles & Practice, chapter 66; Dacie 11th/pg 307; Refer to Ans 81)

**83. Ans. (a) Single amino acid base substitution**

(Ref: Robbins 9th/pg 635-636; 8th/pg 645-647)

This is a case of 16 year old boy presenting with foot ulcer, recurrent pneumonia and chronic hemolytic anemia. This history of **non-healing ulcer** and **repeated infection** along with given peripheral smear finding is suggestive of **Sickle cell anemia**. **Point mutation in 6th codon of  $\beta$ -globin** that leads to **replacement of glutamate by valine** residue

**84. Ans. (a) P. falciparum** (Ref: Robbins 9th/pg 635-636)

- HbA/S heterozygotes (**Sickle cell trait**) have a **6-fold reduction** in risk of dying from **severe falciparum** malaria.

- Due to **impaired parasite growth at low oxygen tensions** and **reduced parasitized red cell cyto-adherence**.
- **Parasite multiplication** in HbA/E heterozygotes is **reduced** at high parasite densities.

**85. Ans. (a, b, c, d, e); a. Common in India; b. Change in  $\beta$ -globin gene; c. Microcytosis; d. Increased HbF; e. Secondary hemochromatosis may occur**

(Ref: Robbins 9th/pg 638-642)

a.	<b>True</b>	<b>Distribution:</b> <sup>Q</sup> of $\beta$ thalassemia: <ul style="list-style-type: none"> <li>• Mediterranean, Africa, middle-east, Pakistan, India and South east Asia.</li> <li>• In India- <b>Sindh<sup>Q</sup>, Punjab, Gujrat and Bengal</b> have higher prevalence</li> </ul>
b.	<b>True</b>	<b>Decreased synthesis</b> of $\beta$ -globin chains due to mutation of beta globin gene
c.	<b>True</b>	Marked <b>aniso-poikilocytosis, microcytic hypochromic</b> RBC's
d.	<b>True</b>	Selected survival of HbF cells in thalassemia causes increased level of HbF in blood
e.	<b>True</b>	<b>Repeated transfusion in thalassemia causes Iron overload</b> leading to Secondary hemochromatosis

**86. Ans. (c) Sickle cell anemia** (Ref: Robbins 9th/pg 635-636)

**Patients of sickle cell anemia may undergo auto-splenectomy<sup>Q</sup>** due multiple infarcts in splenic artery causes spleen to be reduced to fibrous tissue (splenic atrophy → **spleen may become non-palpable**)<sup>Q</sup>

**87. Ans. (c) Aplastic crisis is related to spleen**

(Ref: Robbins 9th/pg 635-636)

- Fetal Hb persists at high conc in adult life as it is protective: TRUE; HbF has increased O<sub>2</sub> affinity, so decreases sickling;
- Co-existent alpha thalassemia is milder disease: TRUE
- Aplastic crisis is related to spleen: FALSE, as aplastic crisis is related to Parvovirus B19 infection
- Sequestration crisis is related to spleen: TRUE, as it is due to hypersplenism

**88. Ans. (a) Decreased LDH** (Ref: 9th/pg 631-632)

In hemolytic anemia, there is decreased haptoglobin, decreased RBC survival, increased unconjugated bilirubin & Increased LDH.

**89. Ans. (d) After splenectomy, Spherocytes disappear**

(Ref: Robbins 9th/pg 632-633; 8th/pg 642-643)

Discussing options about HS one by one,

- TRUE, as hemolysis occurs in spleen
- TRUE, as Spectrin/Ankyrin/Band-3 defect is seen in HS
- TRUE, as RBCs get damaged & deformed in circulation
- FALSE; **After splenectomy, Spherocytes are still present, but hemolysis of these spherocytes in spleen is prevented**



**90. Ans. (b) Young RBCs are more prone to hemolysis**

(Ref: Robbins 9th/pg 634-635; 8th/pg 644-645)

Discussing the options about G6PD deficiency one by one,  
a. TRUE; It has X linked recessive: inheritance

b. Young RBCs are more prone to hemolysis: FALSE, rather **younger RBCs are more resistant to hemolysis** - Wintrobe's 12<sup>th</sup>/pg 935-937

- As red cells age, the activity of G6PD declines.
- The normal enzyme (G6PD B) has an in vivo half-life of 62 days
- In contrast, the **G6PD variants associated with hemolysis are unstable** and have **much shorter half-lives**.
- The **anemia is self-limited** because the old susceptible population of erythrocytes is replaced by **younger RBC with sufficient G6PD activity to withstand an oxidative assault**.

c. Episodic hemolysis: TRUE, as it occurs only when exposed to inciting factors like infection & certain drugs;

d. Both intravascular & extravascular hemolysis is seen: TRUE

**91. Ans. (c) Coombs test**

(Ref: 9th/pg 643-644; 8th/pg 653-654)

A 23-year-old female presented with jaundice and pallor for 2 months. Her peripheral blood smear shows the presence of spherocytes. This scenario is suggestive of Autoimmune hemolytic Anemia

The most relevant investigation to arrive at a diagnosis is Coombs test

**92. Ans. (a, b, c); a. Hemolysis; b. Blood loss; c. Hemoglobinopathy** (Ref: Robbins 9th/pg 631-632)

**93. Ans. (b, c); b. Paracetamol; c. Ceftriaxone**

(Ref: Robbins 9th/pg 634-635; 8th/pg 644-645)

Paracetamol & Ceftriaxone do not carry risk of hemolysis in persons with G-6-PD deficiency

**94. Ans. (a, b, c, d, e); a. Microangiopathic hemolytic anemia; b. HUS; c. Hereditary schistocytosis; d. Thrombotic thrombocytopenic purpura; e. DIC**

(Ref: Robbins 9th/pg 941)

Schistocytes are found in Microangiopathic hemolytic anemia, HUS, Hereditary schistocytosis, TTP, DIC

**95. Ans. (c) Ankyrin**

(Ref: Robbins 9th/pg 632-633)

**96. Ans. (b) G6PD deficiency** (Ref: Robbins 9th/pg 634-635)

Heinz bodies are seen in G6PD deficiency

**97. Ans. (a) Hemolytic anemia**

(Ref: Robbins 9th/pg 631-632)

Unconjugated hyperbilirubinemia with increased urobilinogen is seen in Hemolytic anemia

**98. Ans. (d) Na<sup>+</sup> Cl<sup>-</sup> channel protein** (Ref: R 9th/pg 632-633)

**99. Ans. (c) Microspherocytes** (Ref: Robbins 9th/pg 631-632)

**ABO incompatibility** causes immune hemolytic anemia which is characterized by Spherocytes on peripheral smear

**100. Ans. (c) PNH** (Ref: Robbins 9th/pg 631-632)

- The given scenario is suggestive of episodes of hemolysis since birth, so it is probably due to a hereditary cause.
- Out of the given options, **hereditary spherocytosis, Sickle cell disease & G6PD deficiency** are **inherited causes** of hemolytic anemia, so they usually present in early childhood.
- **PNH is due to an acquired somatic mutation in PIG A gene, which usually presents in adults**. So, PNH is the least likely diagnosis in this cause.
- Remember that, **Sickle cell disease usually presents beyond 6 months age, due to high HbF concentration before that**, which prevents sickling

**101. Ans. (c) ABC**

Looking at the history and HPLC finding, it looks Thalassemia major which will have severe anisopoikilocytosis, (slide C) nRBC (slide B) and Howell jolly body (slide A)

**102. Ans. (c) b-Thalassemia trait**

This patient has a normal Hb but low MCV. So if you calculate the Mentzer index= MCV/RBC count, it comes to 9.3, which is less than 13 which is indicative of Thalassemia trait.

**103. Ans. (a) Retarded**

(Ref: Complete review of Pathology 1st /343)

Pathogenesis of Sickle cell anemia:

Sequence of codon 6 of the Beta-chain changed from GAG in the normal gene to GTG in the sickle cell gene, resulting in substitution of **valine for glutamic acid**

**Concept:**

*So if you see, the charge of amino acid is changing from charged (glutamic acid, more negative) to neutral amino acid (valine), so when we do an electrophoresis, the mobility of HbS will be lesser than HbA towards Anode (i.e retarded) (which is positively charged).*

At alkaline pH;  $\text{HbA}_2 \rightarrow \text{HbC} \rightarrow \text{HbS} \rightarrow \text{HbF} \rightarrow \text{HbA}$   
Cathode(-)  $\xrightarrow{\hspace{1.5cm}}$  Anode (+)

**104. Ans. (d) NESTROFT**

(Ref: Robbins 9th/pg 638; 8th/pg 648)

Screening tests for Thalassemia trait: NESTROFT (Naked Eye Single Tube Red cell Osmotic Fragility Test)<sup>Q</sup> assesses osmotic fragility of red cells at a single concentration of buffered saline (0.36% in single tube)<sup>Q</sup>



105. Ans. (c)  **$\gamma$  chain** (Ref: Robbins 9th/pg 638; 8th/pg 648)

106. Ans. (c) **Three  $\alpha$  globin chains** (Ref: Robbins 9th/pg 638)

Classification of  $\alpha$ -thalassemia:

- Normally 4  $\alpha$ -genes synthesize: **2  $\alpha$  chains**  $\rightarrow \alpha, \alpha / \alpha, \alpha$
- Most common cause of  $\alpha$ -thalassemia is  **$\alpha$ -gene deletion (frame-shift mutation)<sup>o</sup>**

107. Ans. (d) **Alkalosis** (Ref: Robbins 9th/pg 635; 8th/pg 645)

108. Ans. (b) **Sickle cell anemia** (Ref: Robbins 9th/pg 635)

Multiple infarcts in splenic artery causes spleen to be reduced to fibrous tissue referred to autosplenectomy (splenic atrophy  $\rightarrow$  spleen may become non-palpable)

109. Ans. (d) **All the above**

(Ref: Robbins 9th/pg 635; 8th/pg 645)

110. Ans. (c) **Sickle cell anemia** (Ref: Robbins 9th/pg 635)

111. Ans. (b)  **$\alpha$  chain, 87th codon, Histidine  $\rightarrow$  Tyrosine**

(Ref: Wintrob's 12th ed 1060-1070)

- M (met) hemoglobins are characterized by heme-iron oxidation and result in cyanosis.
- HbM Iwate (HBA 87th position his  $\rightarrow$  tyr)

112. Ans. (b) **Missense mutation** (Ref: Robbins 9th/pg 635)

113. Ans. (d) **4** (Ref: Robbins 9th/pg 638; 8th/pg 648)

114. Ans. (d) **HbF and HbA2 will be undetectable**

(Ref: Wintrob's 12th ed p 1062)

The given peripheral blood picture shows sickle cells. Pulmonary syndrome & retinopathy are complications of Vaso-occlusive crises. Hydroxyurea increases HbF levels and so is used for treatment. HPLC diagnosis suggests **increase in HbS, HbA2, and HbF** while Hb A is decreased.

115. Ans. (a) **Sickle cell** (Ref: Robbins 9th/pg 635; 8th/pg 645)

The given scenario is of Rh incompatibility resulting in Hemolytic disease of newborn. The RBC morphology shows anisopoikilocytosis, polychromasia, target RBCs and Schistocytes.

116. Ans. (d) **High hematocrit** (Ref: Robbins 9th/pg 635-636)

In sickle cell anemia: Target cells, Jaundice & Reticulocytosis are seen  
But hematocrit is low (not high), as it is an anemia

117. Ans. (b) **Malaria** (Ref: 9th/pg 635-636; 8th/pg 645-647)

118. Ans. (a) **Deletion of 3 alpha genes** (Ref: 9th/pg 638-642)

119. Ans. (a) **Thalassemia**

(Ref: Robbins 9th/pg 638-642; 8th/pg 648-652)

120. Ans. (c) **Flow cytometry**

(Ref: Robbins 9th/pg 643)

PNH is diagnosed by flow cytometry, which provides a sensitive means for detecting red cells that are deficient in GPI-linked proteins such as CD59 & CD55

121. Ans. (a) **PNH**

(Ref: Robbins 9th/pg 642)

122. Ans. (b) **Defective GPI anchor**

(Ref: R 9th/pg 642-643)

An abnormal Ham test is seen in PNH, which is due to defective GPI anchor

123. Ans. (d) **All of the above**

(Ref: Robbins 9th/pg 642-643)

PNH is associated with a deficiency of DAF (CD55), MRL (CD59) & GPI anchored protein

124. Ans. (c) **Glycosyl phosphatidyl inositol**

(Ref: Robbins 9th/pg 642-643; 8th/pg 652-653)

125. Ans. (a, b, d); a. **TTP**; b. **DIC**; d. **March Hemoglobinuria**  
(Ref: Robbins 9th/pg 631-632)

126. Ans. (a) **Methyldopa** (Ref: Robbins 9th/pg 643-644)

Drugs causing warm Ab type Autoimmune Hemolytic Anemia:

- **Antigenic type<sup>o</sup>**: penicillin and cephalosporins
- **Innocent bystander type** (Immune complex type): Penicillin
- **Tolerance-breaking**: Alpha methyl dopa<sup>o</sup>

127. Ans. (d) **TIBC increased, Ferritin reduced, Transferrin saturation reduced, Serum transferrin receptors increased** (Ref: Robbins 9th/pg 659)

128. Ans. (d) **Ferroportin** (Ref: Robbins 9th/pg 659)

**Hepcidin** is a regulator of iron metabolism. **Hepcidin** inhibits iron transport by binding to the iron export channel ferroportin which is located on the basolateral surface of gut enterocytes and the plasma membrane of reticuloendothelial cells (macrophages).

129. Ans. (a) **DMT-1** (Ref: Robbins 9th/pg 659)

The DMT1 protein, also known as Nramp2, SLC11A2, and DCT1, conducts iron transport at two distinct compartments of the cell: (1) It facilitates iron uptake at the apical cell membrane in for instance duodenal enterocytes (2) It transports iron across erythroid precursors in the bone marrow





130. Ans. (a) **Anemia of chronic inflammation**; (b) **Anemia due to chronic renal disease**; (c) **Iron deficiency anemia**; (d) **Myelodysplastic syndrome** (Ref: Robbins 9th/pg 659)

**Anemias of Decreased Erythropoiesis: Important Examples are:**

- Iron deficiency anemia, Folic acid and B<sub>12</sub> deficiency anemia, vitamin C, Copper and Zinc deficiency. ACD, MDS, Aplastic and pre red cell aplasia

131. Ans. (a) **TIBC (Total iron binding capacity)**

132. Ans. (a, b, c) **a. Iron deficiency anemia; b. Thalassemia; c. Sideroblastic anemia**

133. Ans. (a) **Lead poisoning** (b) **Sideroblastic anemia** (c) **Occult blood loss** (d) **Atransferrinemia**

All 4 will cause microcytic anemia hypochromic

134. Ans. (b) **TIBC**

Transferrin is quantified in terms of the amount of iron it will bind, a measure called the total iron-binding capacity (TIBC; In the average subject, the plasma iron concentration is ~100 µg/dl), and the TIBC is ~300 µg/dl).

Serum iron concentration is reduced in iron deficiency, and the TIBC is often increased

135. Ans. (c) **Transthyretin** (Ref: Wintrobe's 14th/p 813)

Ceruloplasmin acts as an oxidase for a variety of substrates, one of which is ferrous iron. There is evidence that ceruloplasmin is required for the optimal mobilization of iron from cells to plasma.

136. Ans. (a, c, d) **a. Coarse basophilic stippling in lead poisoning; c. Dimorphic anaemia; d. Increase transferrin saturation**

Findings in sideroblastic anemia: Serum Iron: High; Increased ferritin levels; Normal total iron-binding capacity; High transferrin saturation, MCV is usually normal or low. lead poisoning will show coarse basophilic stippling of red blood cells on peripheral blood smear. Specific test: Prussian blue stain of RBC Erythroid hyperplasia is found on bone marrow examination

137. Ans. (a) **Transferrin receptor 1 - iron responding elements increases transferrin receptor mRNA concentration and synthesis**

138. Ans. (c) **Increased TIBC**

139. Ans. (c) **Lead toxicity**

In megaloblastic anemia there is defect in DNA synthesis lead toxicity causes iron deficiency anemia

140. Ans. (c) **4 1 2 3** (Ref: Robbins 9th/pg 643)

141. Ans. (d) **Haemolysis from G6PD enzyme deficiency**

(Ref: Robbins 9th/pg 643)

**G6PD deficiency causes anemia and not pancytopenia**

142. Ans. (b) **Thalassemia major** (Ref: Robbins 9th/pg 631)

Though all the options mentioned can cause Iron deficiency resulting in microcytic hypochromic RBCs. Thalassemia will result in raised ferritin (due to increased GIT absorption & multiple transfusions) and not low ferritin.

143. Ans. (b) **DMT-1**

(Ref: Wintrobe 13ed /pg 811; Robbins 9th/649; Complete review of Pathology 1<sup>st</sup>/314)

### About DMT-1:

- Mucosal uptake of **non-heme iron** begins at **brush border** of mucosal cell, mediated by divalent metal ion transporter 1 (**DMT1, also known as Nramp2, SLC11A2**).
- Levels of DMT1 are markedly increased in iron-deficient animals.
- **In addition to iron, DMT1 has been shown to transport a variety of divalent metal ions, including Mn<sup>2+</sup>, Co<sup>2+</sup>, Cu<sup>2+</sup>, Zn<sup>2+</sup>, Cd<sup>2+</sup>, and Pb<sup>2+</sup>. (Hence non-specific)**
- It acts as a proton symporter; protons accompany metal ions into the cell.

**Hepcidin: principal iron regulatory hormone; and is negatively regulates ferroportin on basolateral surface of enterocyte<sup>Q</sup>**

144. Ans. (a) **Hepcidin** (Ref: Robbins 9th/pg 649; 8th/pg 659)

145. Ans. (a) **S.ferritin** (Ref: Robbins 9th/pg 649; 8th/pg 659)

146. Ans. (a) **Hepcidin** (Ref: Robbins 9th/pg 649; 8th/pg 659)

Ferroportin which regulates the Iron release from tissue store house is regulated by hepcidin hormone released by liver.

147. Ans. (d) **Ferrocyanide to ferricferrocyanide**

(Ref: T. Singh 3rd ed/pg 21)

### Prussian Blue Stain:

**Purpose:** To demonstrate ferric iron in tissue sections.

**Principle:** The reaction occurs with the treatment of sections in acid solutions of ferrocyanides. Any **ferric ion (+3)** in the tissue combines with the **ferrocyanide** and results in the formation of a bright blue pigment called 'Prussian blue' or **ferric ferrocyanide**.

148. Ans. (d) **Diagnosed by serum ferritin**

(Ref: Robbins 9th/pg 649; 8th/pg 659)

Hereditary hemochromatosis (Autosomal recessive) is caused due to mutations of HFE gene on chr6p21.3.



149. Ans. (b) **Siderocyte** (Ref: Robbins 9th/pg 649)

**Pappenheimer bodies:** Siderotic mitochondria of the developing cell may be retained in some circulating erythrocytes (Pathognomonic siderocytes)

150. Ans. (a) **2150 mg** (Ref: Wintrob's 12ed/814)

Treatment of anemia due to iron deficiency.  
The total dose is calculated from the amount of iron needed to restore the hemoglobin deficit plus an additional amount to replenish stores.

**Total iron deficit [mg] =** body weight [kg] × (target Hb (15)-actual Hb) [g/dl] × 2.4 + **Iron for iron stores** (depot iron) [mg]

In this question:  $50 \times (15-5) \times 2.4 + 1000 = 2200$  mg  
Nearest correct option is A i.e 2150 mg

151. Ans. (b) **Folate supplement** (Ref: Robbins 9th/pg 645)

152. Ans. (d) **Start folate supplementation**

(Ref: Robbins 9th/pg 645)

153. Ans. (d) **Folate deficiency anemia** (Ref: R 9th/pg 645)

154. Ans. (d) **MCHC is increased** (Ref: Robbins 9th/pg 645)

MCHC: Mean Corpuscular Hemoglobin Concentration

- Low in iron deficiency anemia and thalassemia.
- Increased in spherocytosis

155. Ans. (c) **Hook worm infection** (Ref: Robbins 9th/pg 649)

The findings in the question are suggestive are Iron deficiency anemia. Hookworm infection can cause this among the given options.

156. Ans. (c) **Serum ferritin** (Ref: Robbins 9th/pg 649-652)

Among the options provided, **most sensitive marker in iron deficiency anemia is Serum ferritin.**  
Serum ferritin reflects the **storage of Iron** which is **decreased** even in the **pre-latent stage** of Iron deficiency Anemia and is the **most sensitive** marker.

157. Ans. (b) **B12 deficiency** (Ref: Robbins 9th/pg 645)

158. Ans. (c) **TIBC** (Ref: Robbins 9th/pg 649)

**Total Iron Binding Capacity (TIBC) is increased in IDA** while it is decreased in ACD.

159. Ans. (a) **Serum ferritin level** (Ref: Robbins 9th/pg 649)

Serum ferritin level is the best marker of Iron store levels.

160. Ans. (d) **Ascorbic acid** (Ref: Robbins 9th/pg 649-652)

#### Iron Absorption

- Site of iron absorption: duodenum and upper jejunum (proximal small intestine)<sup>a</sup>.
- Ferrous form (Fe<sup>2+</sup>) of Iron is absorbed<sup>a</sup>.
- Transport of Fe<sup>2+</sup> into enterocyte occurs via DMT1<sup>a</sup>.

Increased by	Decreased by
<ul style="list-style-type: none"> <li>• Acids<sup>a</sup></li> <li>• Ascorbic acid (Vitamin C)<sup>a</sup></li> <li>• Amino acid containing SH-group</li> <li>• Meat (these reduce Fe<sup>3+</sup> to Fe<sup>2+</sup>)</li> </ul>	<ul style="list-style-type: none"> <li>• Alkalies</li> <li>• Phosphates<sup>a</sup></li> <li>• Phytates<sup>a</sup></li> <li>• Tetracycline</li> <li>• Presence of other food in stomach</li> </ul>

161. Ans. (b) **Leukopenia** (Ref: Robbins 9th/pg 649)

162. Ans. (a) **Increased serum ferritin** (Ref: R 9th/pg 649)

163. Ans. (b) **100 µg/dL** (Ref: Robbins 9th/pg 649)

164. Ans. (b) **IL-6** (Ref: Wintrob's 13th ed/pg 1223)

Cytokines most often implicated in the pathogenesis of ACD are TNF, IL-1, IL-6, and the interferons. However, **IL-6 is the potent inducer of hepcidin**, proposed as a **major mediator of the iron abnormalities of ACD.**

165. Ans. (d) **Absorption of iron is increased by ascorbic acid**

(Ref: Robbins 9th/pg 649; 8th/pg 659; Wintrob's 12th ed/827)

- False Parenteral Iron is given in severe iron deficiency anemia or if patient is unable to tolerate oral iron.
- False as ferrous sulphate (65mg/325mg) has more Iron than gluconate (37.5mg/325mg)
- False as parenteral iron is best
- True, refer to table above

166. Ans. (b) **Macrocytic anemia** (Ref: Robbins 9th/pg 645)

The peripheral smear here shows hypersegmental neutrophil and macro-ovalocytes, suggestive of macrocytic anemia.

167. Ans. (c) **Hereditary spherocytosis** (Ref: R 9th/pg 632)

168. Ans. (d) **Siderocytes** (Ref: Wintrob's 12th ed/ 848)

169. Ans. (b) **Oral iron therapy** (Ref: Robbins 9th/pg 649-652)

The history provided is suggestive of Iron deficiency (nutritional) Anemia. Oral Iron therapy is the treatment of choice.

170. Ans. (b) **Serum ferritin** (Ref: Robbins 9th/pg 649-652)

171. Ans. (b) **35%** (Ref: Robbins 9th/pg 649-652)

172. Ans. (b) **Reticulocytosis**

(Ref: Robbins 9th/pg 649-652; 8th/pg 659-662; Wintrob's 12th/pg 829)

**Response to iron therapy:**

- **Rapid subjective improvement**, with disappearance or marked diminution of fatigue, lassitude, and other

Contd...



nonspecific symptoms, **before any improvement in anemia** is observed.

- **Earliest hematologic evidence** of response to treatment is an **increase in reticulocytes % & Hb content**.
- **Maximal reticulocytes % is on 5<sup>th</sup> to 10<sup>th</sup> day** after institution of therapy & thereafter returns to normal.
- Maximum value usually ranges from 5 to 10% and is inversely related to the level of hemoglobin.
- **Hemoglobin level** is the **most accurate** measure of the **degree of anemia** in iron deficiency

---

**173. Ans. (b) Microcytic hypochromic anemia**

---

(Ref: Robbins 9th/pg 649-652; 8th/pg 659-662)

Iron deficiency causes Microcytic hypochromic anemia

Differential diagnosis of microcytic hypochromic RBC's<sup>Q</sup>:

- Iron Deficiency Anemia
- Anemia of Inflammation/chronic disease
- Thalassemia
- Sideroblastic Anemia

---

**174. Ans. (a) Ringed siderocytes**

---

(Ref: Wintrobe's 12th/pg 848)

Bone marrow in lead poisoning contains Ringed siderocytes.

Hematologic findings in **Lead poisoning** include:

- Anisocytosis with microcytic Hypochromic Anemia
- Basophilic stippling
- Ringed sideroblasts in bone marrow may be seen

---

**175. Ans. (a) Latent iron deficiency is most common presentation in India**

---

(Ref: Robbins 9th/pg 649-652; 8th/pg 659-662)

**a. False- Manifest (Anemia), not Latent (pre-anemia) iron deficiency is most common presentation in India**

b. True- Saturation of transferrin is always reduced to <16%. At this level of saturation, iron delivery to erythroid precursors is limited.

c. True-Most sensitive marker in iron deficiency anemia is Serum ferritin

d. True-The onset of iron deficiency anemia is usually insidious & progression of symptoms is gradual. As a result, patients accommodate remarkably well to advancing anemia & may not present with clinical features

---

**176. Ans. (b) Thalassemia** (Ref: Robbins 9th/pg 649-652)

---

**Leptocytes** are **flattened red cells** in which the **volume-to-surface area ratio is decreased**. Seen in **iron deficiency anemia and thalassemia** in which the red cells with a low mean cell hemoglobin (MCH) and mean cell volume (MCV) are unusually resistant to osmotic lysis.

---

**177. Ans. (a) Reticulocytosis** (Ref: Robbins 9th/pg 649-652)

---

---

**178. Ans. (a) RBC protoporphyrin** (Ref: R 9th/pg 649-652)

---

Heme in Hb consists of Iron & Porphyrin.

In **Iron deficiency anemia**, **free erythrocyte protoporphyrin increases**, which is due to excess of protoporphyrin over

iron in heme synthesis. Each RBC contains less hemoglobin, resulting in microcytosis & hypochromia.

Stages in the Development of Iron Deficiency; Refer to pretexts of this chapter

---

**179. Ans. (c) Anemia of chronic disease**

---

(Ref: Robbins 9th/pg 649-652; 8th/pg 659-662)

This is a case of 35 year, female presenting with microcytic anemia. (Hb-9gm/dL, MCV- 55 fL).

Serum Iron studies reveal:

- Serum iron-30 ug/dL → Low
- Ferritin- 200 ng/mL and TIBC- 298 ug/dl levels are suggestive of Anemia of chronic disease.

---

**180. Ans. (b) Sideroblastic anemia**

---

(Ref: Wintrobe's 12th/pg 847-848)

This is an Alcoholic patient on Anti TB drugs (isoniazid, pyrazinamide) who presents with features of iron overload on iron studies (increased serum iron & increased transferrin saturation). These features are suggestive of sideroblastic anemia.

---

**181. Ans. (a) Atransferrinemia** (Ref: Wintrobe's 12th/pg 814)

---

This patient with **microcytic hypochromic** anemia, has **high ferritin level** (800 ng/ml) & **increased transferrin percentage saturation** (64%). The most possible diagnosis is Atransferrinemia.

### Atransferrinemia

- It is a rare **congenital autosomal recessive** condition with low transferrin level in the body.
- So whatever transferrin is left, is saturated.
- It presents with hypochromic microcytic anemia, decreased serum levels of iron, TIBC, and increased serum level of ferritin (as iron absorption is markedly increased but iron transfer to erythropoietic tissues are reduced)

**High transferrin saturation (64% in this case) & high ferritin level, excludes Iron deficiency Anemia (option B excluded)**

**In DMT 1 mutation, transferrin saturation is low (hence option C excluded)**

**High transferrin saturation excludes Anemia of chronic disease (option D)**

---

**182. Ans. (c) Bone marrow examination**

---

This is a 60 yr old female presents with history of 8 blood transfusions in 2 yrs presenting with severe degree of hypochromic microcytic anemia. The most probable diagnosis is **Iron deficiency anemia, for which one of the underlying causes can be chronic blood loss**.

The investigations required for assessing the patients conditions are:

- **Evaluation for pulmonary hemosiderosis:** to exclude bleeding from alveolus;
- **Urinary hemosiderin:** to assess iron overload (due to repeated blood transfusions)
- **G1 endoscopy:** to assess GI bleeds which may cause such severe iron deficiency over a period of time.

**Idiopathic pulmonary hemosiderosis:**

- **Recurrent alveolar bleeding** may eventually produce **pulmonary hemosiderosis and fibrosis**.
- Presents with **hemoptysis, dyspnea, alveolar opacities** on chest Xray & **anemia**.
- Iron in the shed blood is converted to hemosiderin by pulmonary macrophages, but it cannot be used for Hb synthesis.
- Thus, **repeated hemorrhages can lead to iron deficiency** despite a normal total amount of body iron. Bone marrow examination is not required to investigate for Iron deficiency anemia, hence the answer

**183. Ans. (d) Reticulocyte count**

(Ref: Robbins 9th/pg 645-648)

- This is a tricky question as the initial investigation looks like B<sub>12</sub>/ folate levels, as the patient has macrocytosis.
- But remember, Macrocytosis can also be seen in hemolytic anemia, as newer RBC precursors released are larger in size.
- So the initial investigation should be reticulocyte count
- If reticulocyte % is low it can be macrocytic anemia & if there is reticulocytosis, then it can be a case of hemolytic anemia.

**184. Ans. (a) Increased reticulocyte count**

(Ref: Robbins 9th/pg 645-648; 8th/pg 654-658)

Reticulocyte count does not increase in Megaloblastic anemia

**185. Ans. (e) 5-FU (Ref: Robbins 9th/pg 645-648)****Drug induced causes of macrocytic anemia: by drugs that suppress DNA synthesis**

- Folate antagonists
- Alkylating agents
- Metabolic inhibitors of:
  - Purine synthesis: Hydroxyurea, 6-mercaptopurine, azathioprine
  - Pyrimidine synthesis: 5-fluorouracil; cytosine arabinoside

**186. Ans. (d) Anemia of chronic disease (Ref: 9th/pg 652-653)**

Anemia of chronic disease leads to microcytic anemia and not macrocytic anemia

**187. Ans. (d) Absolute neutrophil count <1500/ul**

(Ref: Wintrob's 13/p 1189)

A bone marrow cellularity of <25% and markedly decreased values of at least two of three hematopoietic lineages (neutrophil count <500/ $\mu$ l, platelet count <20,000/ $\mu$ l, and absolute reticulocyte count of <60,000/ $\mu$ l) define SAA

**188. Ans. (d) Diamond Schwachman syndrome**

(Ref: Wintrob's 13/p 1523-1533)

Shwachman-Diamond syndrome (SDS) is an autosomal recessive disorder characterized by exocrine pancreatic insufficiency and persistent or intermittent neutropenia

and an increased susceptibility to infections. Hematologic problems also can include anemia, thrombocytopenia, and pancytopenia

**189. Ans. (a) Erythroid progenitors**

(Ref: Robbins 9th ed p 655)

A special form of red cell aplasia occurs in individuals infected with parvovirus B19, which preferentially infects and destroys red cell progenitors. Normal individuals clear parvovirus infections within 1 to 2 weeks; as a result, the aplasia is transient and clinically unimportant. However, in persons with moderate to severe hemolytic anemias, even a brief cessation of erythropoiesis results in rapid worsening of the anemia, producing an aplastic crisis.

**190. Ans. (b, d) b. Congenital dyserythropoetic anaemia; d. Aplastic anaemia****191. Ans. (d) None of the above**

(Ref: Robbins 9th/pg 653-655)

- **Drugs:** associated with PRCA are Phenytoin, azathioprine, chloramphenicol, procainamide, isoniazid<sup>Q</sup>

**192. Ans. (a) Fanconi's anemia**

(Ref: Harrison 18th/Chapter 107)

**193. Ans. (a) Fanconi anemia (Ref: Robbins 9th/pg 653)****194. Ans. (c) Aplastic anemia (Ref: Robbins 9th/pg 653)****195. Ans. (c) Thymoma (Ref: Robbins 9th/pg 653)****196. Ans. (d) PRCA (Ref: Robbins 9th/pg 635-636)**

In the case provided, the child develops maculopapular rash 24hrs after onset of mild fever. There was prominent erythema over the cheek. It suggests **Erythema infectiosum caused by parvovirus B19 infection**.

**Parvovirus also causes aplastic crisis in any hemolytic anemia** due to suppression of early erythroblasts by acting on P receptors.

**197. Ans. (b) 5q- syndrome (Ref: 9th/pg 653-655)**

Pure red cell aplasia is not associated with 5q- syndrome. Pure red cell aplasia is a primary marrow disorder in which only erythroid progenitors are suppressed.

**198. Ans. (d) G6PD deficiency**

(Ref: Harrison 18th/pg Chapter 107)

Pancytopenia with cellular bone marrow is not seen in G6PD deficiency, which is a type of hemolytic anemia

**199. Ans. (b) Hemolytic Anemia**

(Ref: Robbins 9th/pg 631-632)

Prominent reticulocytosis is a feature of Hemolytic Anemia, while all others in the options have normal/ low reticulocyte count





**200. Ans. (c) Pure red cell aplasia** (Ref: Robbins 9th/pg 653-655)

Aplastic anemia can progress to

- AML
- Myelodysplastic anemia
- Paroxysmal nocturnal hemoglobinuria

Aplastic anemia cannot progress to Pure red cell aplasia

**201. Ans. (a, b, c); a. Pancytopenia; b. Nail dystrophy; c. Hyperkeratosis**

(Ref: Robbins 9th/pg 653-655; 8th/pg 662-664)

In Dyskeratosis congenita, there is Pancytopenia, Nail dystrophy & Hyperkeratosis.

# Bleeding and Coagulation Disorders

## Key Points

- » **Hemoglobin** can be first visualized at **polychromatic (Intermediate) erythroblast<sup>o</sup>** stage
- » **Platelet adhesion** refers to attachment of vWF on endothelium with **GpIb/IX** on platelets
- » **Platelet aggregation** refers to attachment of **GpIIb/IIIa** on platelets that bridges adjacent platelets
- » **Primary hemostatic plug** consists of: **Platelets, Fibrinogen, Entrapped RBCs & WBCs**
- » **All clotting factors** are synthesized by liver, except factor VIII, from endothelium
- » Vitamin K-dependent factors are F- **II, VII, IX, and X<sup>o</sup>** and anticoagulant **proteins C, S and Z**
- » **Thrombomodulin** synthesized from liver **inhibits** activation of **factor V and VIII**
- » **ITP** results from platelet destruction by platelet-specific (**GpIb/IX; GpIIb/IIIa**) **autoantibodies**
- » **Von Willebrand disease** can be diagnosed by **decreased Ristocetin cofactor assay<sup>o</sup>/Ristocetin induced platelet aggregation**
- » **D-dimer test** is **more specific<sup>o</sup>** for detection of **FDPs** in diagnosis of **DIC**
- » **Factor V Leiden** is the **most common inherited thrombophilia**


## Key Recent Updates

Thromboelastography is used to assess viscoelastic changes in clotting used to assess hypo- and hypercoagulable states and to guide hemostatic therapies with fresh-frozen plasma, with platelet concentrates as well as with coagulation factor concentrates.



## HEMOSTASIS

Process of formation of a **blood clot**<sup>o</sup>, which **prevents or limits** bleeding.



- High Yield Facts**
- 1<sup>st</sup> response to endothelial injury is **vasoconstriction**
  - Endothelial injury **activates extrinsic pathway** by releasing **tissue factor**

### Formation of Primary Hemostatic Plug

#### Platelet Plug Formation: 2 Processes

- Platelet adhesion:**
  - vWF on **endothelium**<sup>o</sup> with Gp Ib/IX on **platelets**<sup>o</sup>
- Platelet aggregation:**
  - Gp IIb/IIIa on **platelets**<sup>o</sup> that **bridges adjacent platelets**<sup>o</sup>

**PLATELET:** Disc shaped anucleate<sup>a</sup> cell fragments shed from megakaryocytes

**α-Granules** contains:

- P-selectin (adhesion molecule)
- Fibrinogen<sup>a</sup>
- Coagulation factors: V<sup>a</sup>, VIII<sup>a</sup>, vWF<sup>a</sup>
- Platelet factor 4<sup>a</sup> (a heparin binding chemokine)
- Platelet derived growth factor (PDGF)
- Transforming growth factorβ (TGF-β)


**Dense (or δ) granules** contain: ("DENSE")

- aDENosine diphosphate (ADP)<sup>a</sup>
- Serotonin<sup>a</sup>
- Epinephrine<sup>a</sup>
- Ionized calcium<sup>a</sup>



### High Yield Facts

- Platelets life span is 7-10 d
- Amyloidosis causes pinch purpura
- Solar purpura is senile
- Gloves & stock syndrome is caused by Parvo B<sub>19</sub>, CMV
- F VIII & IX are x-linked while other are AR



### Mnemonic

"B comes before G & Ib comes before IIb"

Disease	Deficient Gp
Bernard Soulier syndrome	Ib / IX
Glanzmann thrombasthenia	IIb / IIIa


### Formation of Secondary Hemostatic Plug

#### Coagulation Cascade

Series of **amplifying**<sup>o</sup> enzymatic reactions that leads to the deposition of an **insoluble fibrin clot**.<sup>o</sup>

Each Reaction Step Involves

- Enzyme** (an **activated coagulation factor**)
- Substrate** (an **inactive proenzyme** form of a coagulation factor)
- Cofactor** (a reaction accelerator)





## Role of Coagulation Factors

- Clotting factors normally circulate in plasma in their **inactive forms**<sup>Q</sup>
- Components are assembled on a **negatively charged phospholipid surface**<sup>Q</sup>, provided by **activated platelets**<sup>Q</sup>
- Based on laboratory assays: Coagulation cascade can be of **extrinsic and intrinsic pathways**<sup>Q</sup>




## High Yield Facts

- All clotting factors are synthesized by **LIVER**,<sup>Q</sup> except **factor VIII**, from **endothelium**<sup>Q</sup>
- Factor **VIIIa/tissue factor** complex is the most important activator of **factor X**<sup>Q</sup>
- Factor **IXa/ VIIIa** complex is the most important activator of **factor X**<sup>Q</sup>
- Factor **Xa** converts **prothrombin to thrombin** which requires factor **Va**<sup>Q</sup> as cofactor
- Factor **Va** is the '**fundamental protease of the coagulation system**'<sup>Q</sup>
- Thrombin** is called '**Master regulator of clotting pathway**'<sup>Q</sup>
- F VIII dose** = VIII levels (Target – bleeding) × 0.5 IU/kg

## Overview of Coagulation Pathway


### Three Pathways that make the Classical Blood Coagulation Pathway



## Functions of Thrombin

- It converts soluble plasma **fibrinogen to insoluble fibrin**<sup>Q</sup>
- Activates factor **XIII (fibrin-stabilizing factor)** to factor **XIIIa**, which covalently cross-links & thereby **stabilizes the fibrin clot**.<sup>Q</sup>
- Thrombin is the most potent activator of platelets**<sup>Q</sup>
- Pro-inflammatory** effects-contribute to tissue **repair and angiogenesis**<sup>Q</sup>
- Some **anticoagulant** effects

## Factors that Limit Coagulation







## High Yield Facts


### Vitamin K-Dependent Coagulation Factors

- **Vitamin K**, a **fat-soluble vitamin** is a **cofactor for carboxylation** of the  $\gamma$ -carbon of the glutamic acid residues of vitamin K-dependent factors
- Vitamin K-dependent factors are **F- II, VII, IX, and X<sup>a</sup>** and anticoagulant **proteins C, S and Z<sup>a</sup>**
- A critical step for **calcium and phospholipid binding<sup>a</sup>** of these proteins
- The enzymes  $\gamma$  glutamyl carboxylase and **epoxide reductase** are critical for the **regeneration of vitamin K<sup>a</sup>**
- **Earliest** coagulation factor affected due to **Vit K def.** is **factor VII<sup>a</sup>** (**shortest  $t_{1/2}$** )
- **Warfarin blocks** the action of **epoxide reductase** and **competitively inhibits** the effects of **vitamin K<sup>a</sup>**
- **Abnormal plasma proteins** produced in **Vit K def.** are called **PIVKA**
- **Abnormal prothrombin** produced in **Vit K def.** are called **des  $\gamma$ -carboxy prothrombin (DCP)**
- **Mutations** in the genes encoding the **gamma-carboxylase (GGCX)<sup>a</sup>** or **vitamin K epoxide reductase complex 1 (VKORC1)<sup>a</sup>** result in **defective enzymes (1 to 30% of normal activity)<sup>a</sup>**
- Vit K deficiency bleeding can be treated by **high doses of vitamin K<sup>a</sup>**, replacement therapy with **FFP** or **Plasma concentrate<sup>a</sup>**

### Some Important Regulators of Coagulation System

- **Thrombomodulin**
  - Has a transmembrane **proteoglycan-binding site for thrombin<sup>a</sup>** (See figure below for mechanism).
- **Tissue Factor Pathway Inhibitor (TFPI)**
  - Plasma protease inhibitor
  - Regulates the TF-induced **extrinsic pathway<sup>a</sup>** of coagulation.
  - TFPI **inhibits** the **TF/FVIIa/FXa complex<sup>a</sup>**

## FIBRINOLYTIC SYSTEM



### Physiologic Regulation of Fibrinolysis


Occurs primarily at 3 levels

- **Plasminogen activator inhibitors (PAIs)** **inhibit plasminogen activators** (tPA and uPA)
- **Plasminogen Activators**
  - Cleave **plasminogen** to generate the active enzyme **plasmin**
  - Tissue type plasminogen activator (**tPA<sup>a</sup>**)
  - Urokinase type plasminogen activator (**uPA<sup>a</sup>**)
- **Plasmin** cleaves **fibrin to fibrin fragments** (fibrinolysis) releasing Fibrin Degradation products (FDP) and D-dimers
- Thrombin-activatable fibrinolysis inhibitor (**TAFI**) **limits fibrinolysis<sup>a</sup>**
- $\beta_2$ -**antiplasmin** **inhibits plasmin.<sup>a</sup>**



## High Yield Facts

- **Protein S** is the **cofactor for Protein C<sup>a</sup>**
- **Thrombomodulin, Protein C and Protein S** are synthesized from **liver<sup>a</sup>**
- **Thrombomodulin** **inhibits** activation of **factor V** and **factor VIII<sup>a</sup>**
- **D-dimer** is used to diagnose **deep-venous thrombosis (DVT)** and **pulmonary embolism.<sup>a</sup>**



### POINTS TO REMEMBER



- TAFI is Thrombin activated fibrinolysis inhibition
- Thrombin bound to thrombomodulin activates TAFI
- Removes lysine residues from fibrin & prevents fibrinolysis

### CLINICAL DIFFERENCES BETWEEN DISORDERS OF PLATELETS/VESSELS AND COAGULATION


Characteristics	Disorders of Platelets/Vessels	Disorders of Coagulation
<b>Petechiae<sup>a</sup></b>	Common	Rare
<b>Hematomas<sup>a</sup></b>	Rare	Common
<b>Ecchymoses<sup>a</sup></b>	Characteristic: small and multiple	Common: large and solitary
<b>Hemarthrosis<sup>a</sup></b>	Rare	Characteristic
<b>Sex of patient</b>	F>>M	M>>F
<b>Family history</b>	Rare (except vWD)	Common

### THROMBOCYTOPENIA


Definition < 1.5 lac/cu mm<sup>o</sup>



Normal platelets



Thrombocytopenia



Thrombocytosis



## IMMUNE THROMBOCYTOPENIC PURPURA (ITP)

Previously called: **Idiopathic thrombocytopenic purpura**<sup>Q</sup>  
Severe **thrombocytopenia** caused by **immune destruction**<sup>Q</sup> of platelets.


### Clinical forms:

- **Acute ITP:** Seen in children; usually **self-resolving**. 50% of cases are associated with a **history of viral infection 2-3 weeks before onset**.<sup>Q</sup>
- **Chronic ITP:** Seen in adults; usually **long-standing disorder (>6 months)**<sup>Q</sup>, characterized by **multiple relapses and remissions**<sup>Q</sup>
- ITP may be **secondary to:** SLE<sup>Q</sup>, Infections: **Viral infections**<sup>Q</sup>, HIV<sup>Q</sup> and hepatitis C<sup>Q</sup>, *Helicobacter pylori*<sup>Q</sup>

- **Pathophysiology:** Platelet-specific (**GpIb/IX; Gp IIb/IIIa**) **autoantibodies** that bind to platelets, which are then rapidly **cleared from circulation** by the **mononuclear phagocyte system**<sup>Q</sup> via macrophage **Fcγ receptors** predominantly in the **spleen and liver**.<sup>Q</sup>
- **Clinical feature: Mucocutaneous bleeding:** Oral mucosa, gastrointestinal, or heavy menstrual bleeding, petechiae & ecchymoses
- No splenomegaly

### Laboratory Testing in ITP

- **Peripheral smear:** Low platelet counts, **Large sized platelets**,<sup>Q</sup>
- **BM shows (not required otherwise):** Increased young form of **megakaryocytes**,<sup>Q</sup> **Clot retraction**, which depends on platelets, is **defective**.<sup>Q</sup>



## VON WILLEBRAND DISEASE

Also known as **Angiohemophilia**<sup>Q</sup>, **pseudohemophilia**<sup>Q</sup>

- vWD is the **most common inherited bleeding disorder**.<sup>Q</sup>
- Most **common type** of vWD is **type 1 disease**<sup>Q</sup>
- Most **severe type** of vWD is **type 3 disease**<sup>Q</sup>
- All vWD are **Autosomal dominant** except **type 3**<sup>Q</sup>


3 types	Types	Type 1	Type 2	Type 3
	<b>Defect</b>	Partial deficiency <sup>Q</sup> of vWF	Qualitative vWF defects	Severe deficiency of vWF <sup>Q</sup>
	<b>Inheritance</b>	Autosomal Dominant <sup>Q</sup>	Autosomal Dominant <sup>Q</sup>	Autosomal Recessive <sup>Q</sup>
<b>C/F</b>	• "Platelet-like" superficial bleeding Except in severe VWD-type 3			
<b>Investigations</b>	• Coagulation tests: <b>Prolonged aPTT</b> <sup>Q</sup> , Normal PT and TT; <b>Ristocetin induced platelet aggregation</b> <sup>Q</sup> defective • Serum electrophoresis: <b>Decreased vWF multimers in type 1 and 3</b> <sup>Q</sup>			
<b>Treatment</b>	• <b>DDAVP or Desmopressin</b> , causes a <b>transient rise in FVIII &amp; VWF; FFP</b>			

**Functions of vWF:** Adhesion of platelets to sub-endothelium<sup>Q</sup>, Binding protein for FVIII (Increases FVIII half-life in circulation).<sup>Q</sup>

### High Yield Facts

- **Primary hemostatic plug** consists of: **Platelets, Fibrinogen, Entrapped RBCs & WBCs**<sup>Q</sup>
- **Primary plug is reversible**<sup>Q</sup>
- **Platelet aggregation** is followed by **platelet contraction**<sup>Q</sup>
- **Large platelets**<sup>Q</sup> are seen in **Bernard Soulier syndrome**
- Bernard Soulier syndrome & Glanzmann thrombasthenia are **Autosomal recessive**<sup>Q</sup>
- Bleeding in **Glanzmann thrombasthenia** is **more severe**<sup>Q</sup> than Bernard Soulier syndrome
- **P-selectin** causes **adherence of WBCs**<sup>Q</sup> with **activated platelets**
- **Primary hemostatic plug is loose** → Stronger secondary hemostatic plug required
- **Weibel Palade bodies** on vWF is synthesized by endothelial cells and **α granules** on platelets






## Platelet Aggregation Studies

Disease	Platelet Aggregation studies			Other Features
	Collagen	ADP	Ristocetin	
<i>Bernard–Soulier disease</i>	N	N	Absent	Giant platelets <sup>q</sup>
<i>Glanzmann's thrombasthenia</i>	Absent <sup>q</sup>	Absent <sup>q</sup>	N/Reduced <sup>q</sup>	Absent clot retraction
<i>Von Willebrand's disease</i>	N	N	Absent <sup>q</sup>	Corrected by factor VIII:vWF
<i>Storage pool disease</i>	Absent	Absent	N	Absent dense bodies <sup>q</sup>

## HEPARIN INDUCED THROMBOCYTOPENIA AND THROMBOSIS (HITT)



### Mnemonic

#### 4 T's in the diagnostic of HITT

- **Thrombocytopenia** (decrease of  $\geq 50\%$  platelet count)
- **Timing** of : HIT develops after exposure to heparin for **5–14 days**<sup>q</sup>
- **Thrombosis** in **50%**<sup>q</sup>
- **Other** causes of thrombocytopenia not evident.

## THROMBOTIC MICROANGIOPATHIES

- Hemolytic-uremic syndrome (HUS)
- Thrombotic thrombocytopenic purpura (TTP)

### Hemolytic-Uremic Syndrome (HUS)

Characterized by Triad of:

- Acute onset of **microangiopathic hemolytic anemia**<sup>q</sup> (schistocytes<sup>q</sup>, burr cells<sup>q</sup>, or helmet cells<sup>q</sup> on peripheral blood smear) (**Hallmark of HUS**)<sup>q</sup>
- **Renal Failure**<sup>q</sup>
- **Thrombocytopenia**<sup>q</sup>


### Classification

Points	Typical HUS	Atypical HUS
<i>Also Called</i>	Epidemic <sup>q</sup> , diarrhea-positive, D+ HUS <sup>q</sup>	Non-epidemic <sup>q</sup> , diarrhea-negative, D- HUS <sup>q</sup>
<i>Etiology</i>	<ul style="list-style-type: none"> <li>• <i>E. coli</i><sup>q</sup> (O157:H7)-Shiga-like toxins<sup>q</sup></li> <li>• <i>Shigella dysenteriae</i><sup>q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Inherited <b>mutations</b> of complement-regulatory proteins: <b>Factor I, H and CD46</b><sup>q</sup> causing <b>hyperactivation of complement</b></li> <li>• <b>APLA</b>: (lupus anticoagulant)<sup>q</sup>, Complications of <b>pregnancy</b><sup>q</sup> or the postpartum period.</li> <li>• <b>Systemic sclerosis</b><sup>q</sup> &amp; <b>malignant hypertension</b>.<sup>q</sup></li> <li>• Drugs: e.g., cyclosporine<sup>q</sup>, FK-506<sup>q</sup>, OKT3, mitomycin C,<sup>q</sup> Ganciclovir, OCP'</li> </ul>





## Pathogenesis



## Thrombotic Thrombocytopenic Purpura (TTP)

### Pentad of findings:

- Microangiopathic hemolytic anemia
- **Thrombocytopenia<sup>Q</sup>**
- **Neurologic findings<sup>Q</sup>**
- **Renal failure<sup>Q</sup>**
- **Fever<sup>Q</sup>**

### Classification

Inherited	Acquired
<b>Upshaw-Schulman syndrome<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• <b>Deficiencies of ADAMTS13<sup>Q</sup></b>, a plasma metalloprotease that cleaves vWF multimers into smaller sizes.</li> <li>• ↑ in <b>HIV<sup>Q</sup></b> and in <b>pregnant women</b></li> <li>• Drugs (<b>ticlopidine</b> and <b>clopidogrel</b>)<sup>Q</sup></li> </ul>

## TESTS USED TO EVALUATE BLEEDING AND COAGULATION DISORDERS

Test	Description and Significance	Abnormal values in: platelet Abnormalities
<b>Bleeding time (BT)</b> (normal: 3–8 min) <sup>Q</sup>	<ul style="list-style-type: none"> <li>• Time taken for a standardized skin puncture to stop bleeding.</li> <li>• Tests the ability of vessels to vasoconstrict &amp; platelets to form a hemostatic plug.</li> </ul>	
<b>Tourniquet test</b> (Hess test) <sup>Q</sup>	<ul style="list-style-type: none"> <li>• BP cuff inflated to above diastolic pressure for 5 minutes</li> <li>• Appearance of ≥ <b>20 petechiae<sup>Q</sup></b> in 1 sq inch area of forearm indicates <b>capillary fragility, thrombocytopenia, or platelet abnormalities</b>.</li> </ul>	
<b>Prothrombin time (PT)<sup>Q</sup></b> (12-15 sec)	Time taken for clotting to occur when <b>tissue thromboplastin<sup>Q</sup></b> (brain extract) and <b>calcium</b> are added to the patient's plasma to <b>activate extrinsic pathway<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• <b>Deficiencies or inhibitors of factors VII, X, and V; II and I;<sup>Q</sup></b></li> <li>• Lupus inhibitors; heparin; <b>warfarin<sup>Q</sup></b></li> <li>• Liver disease</li> </ul>
<b>Activated Partial thrombo-plastin time (aPTT)<sup>Q</sup></b> (26-37 sec)	Time taken for clotting when <b>Kaolin/Silica</b> is added to <b>activate intrinsic pathway<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• Deficiencies or inhibitors of:</li> <li>• Prekallikrein, high-molecular-weight kininogen</li> <li>• Factors XII, XI, IX, VIII, X<sup>Q</sup>, V<sup>Q</sup>, II<sup>Q</sup> and I;</li> <li>• <b>Lupus inhibitors; heparin<sup>Q</sup>; warfarin therapy<sup>Q</sup></b></li> </ul>
<b>Thrombin time (TT)</b> (14-19 sec)	Time taken for clotting when <b>thrombin</b> is added to the patient's plasma to test for <b>conversion of fibrinogen to fibrin (depends on fibrinogen levels)<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>• <b>Afibrinogenemia<sup>Q</sup></b></li> <li>• <b>Dysfibrinogenemia<sup>Q</sup></b></li> <li>• <b>Hypofibrinogenemia,<sup>Q</sup></b></li> </ul>



## High Yield Facts

- **Clot retraction time & Prothrombin consumption index** are used to assess platelet function
- Clot retraction failure of clot retraction in 1–4 hours indicates **thrombocytopenia** or **abnormal platelet function**<sup>Q</sup>.
- **Ratio** of anticoagulant (**Trisodium citrate**): Blood = **1:9**<sup>Q</sup>
- Sample required: **Platelet poor plasma (PPP)**<sup>Q</sup>
- Storage: **Room temperature**<sup>Q</sup>
- Test should ideally be performed **within 2 hours**<sup>Q</sup> of sample collection
- **Prothrombin time (PT)** assay screens **extrinsic + common pathway**<sup>Q</sup> (factors VII, X, V, II, and fibrinogen).
- **Partial thromboplastin time (PTT)** assay screens **intrinsic pathway + common pathway**<sup>Q</sup> (factors XII, XI, IX, VIII, X, V, II, and fibrinogen).
- Clotting time (CT) (normal: 5–10 min)<sup>Q</sup> time taken for the patient's blood to **clot in a test tube**, very **insensitive test**; Not used now a days.

## DISSEMINATED INTRAVASCULAR COAGULATION (DIC)

DIC is an **acute, subacute, or chronic thrombo-hemorrhagic** disorder characterized by the **excessive activation of coagulation**<sup>Q</sup> and the **formation of thrombi** in the **microvasculature of the body**<sup>Q</sup>.

### Etiology of DIC

<b>Sepsis:</b> <ul style="list-style-type: none"> <li>• <b>Bacterial:</b> Staphylococci<sup>Q</sup>, meningococci<sup>Q</sup>, gram-negative bacilli, Anaerobic<sup>Q</sup></li> <li>• <b>Mycotic:</b> Histoplasmosis &amp; Aspergillosis, Parasitic, Viral</li> </ul>	<b>Immunologic disorders:</b> <ul style="list-style-type: none"> <li>• Acute hemolytic transfusion reaction</li> <li>• Transplant rejection</li> <li>• GVHD<sup>Q</sup></li> </ul>
<b>Trauma and tissue injury:</b> Brain injury (gunshot), Rhabdomyolysis <sup>Q</sup>	<b>Drugs:</b> Warfarin (especially in neonates with protein C deficiency) <sup>Q</sup>
<b>Vascular disorders:</b> Giant hemangiomas ( <b>Kasabach-Merritt syndrome</b> ) <sup>Q</sup>	<b>Envenomation:</b> Snake, Insect bites
<b>Obstetrical complications:</b> <b>Abruptio placentae</b> <sup>Q</sup> , Amniotic-fluid embolism, Dead fetus syndrome, <b>Septic abortion</b> <sup>Q</sup>	<b>Liver disease:</b> <b>Fulminant hepatic failure</b> <sup>Q</sup> Cirrhosis, <b>Fatty liver of pregnancy</b> <sup>Q</sup>
<b>Cancer (“SLAP”):</b> Stomach Ca, Lung Ca, Acute promyelocytic leukemia (APML) <sup>Q</sup> Pancreas (most common) & Prostate Adenocarcinoma <sup>Q</sup>	<b>Miscellaneous:</b> <b>Shock</b> <sup>Q</sup> , ARDS, <b>Massive transfusion</b> <sup>Q</sup>



## High Yield Facts


- **Most sensitive test** for DIC is elevated **Fibrin Degraded Products (FDPs) level**<sup>Q</sup>
- DIC is **unlikely diagnosis** in the **presence of normal levels of FDP** (High negative predictive value)<sup>Q</sup>
- **D-dimer test** is **more specific**<sup>Q</sup> for detection of FDPs; Indicates that the **cross-linked fibrin**<sup>Q</sup> has been **digested by plasmin**<sup>Q</sup>.
- **Decreased Fibrinogen level** is seen in DIC<sup>Q</sup>
- **High-grade DIC:** Increased levels of **antithrombin III** or **plasminogen activity <60% of normal**<sup>Q</sup>
- The reference concentration of D-dimer is less than 0.5 µg/mL fibrinogen-equivalent units (FEU)
- Sample collected in - plasma (with sodium citrate anticoagulant, 3.2%)

### Two Major Mechanisms Trigger DIC

- Release of **tissue factor**<sup>Q</sup> & other **pro-coagulants**, into the circulation, and
- **Widespread injury** to the **endothelial cells**<sup>Q</sup>




## Lab Diagnosis



- **Hemogram:**
  - Reduced platelet counts  $< 100,000/\mu\text{L}$
- **Peripheral smear:**
  - Presence of **schistocytes (fragmented red cells)**<sup>Q</sup> in peripheral blood smear
- **Coagulation tests:**
  - Prolongation of PT, aPTT and/or TT<sup>Q</sup>
- **Other investigations:**
  - Most sensitive test for DIC is elevated **FDP (Fibrin Degraded Products)** level<sup>Q</sup>
  - DIC is unlikely diagnosis in the presence of normal levels of FDP (High negative predictive value)<sup>Q</sup>
  - D-dimer test is more specific<sup>Q</sup> for detection of fibrin degradation products; indicates that the cross-linked fibrin<sup>Q</sup> has been digested by plasmin.<sup>Q</sup>
  - Decreased Fibrinogen level is also seen in DIC<sup>Q</sup>
  - High-grade DIC: Increased levels of antithrombin III or plasminogen activity  $< 60\%$  of normal.<sup>Q</sup>

## THROMBOSIS





## Endothelial Injury

- **Most important factor<sup>Q</sup>** causing thrombosis in **arterial and cardiac** circulation
- Endothelial injury → platelet activation → thrombus formation in **high stress circulation like arteries<sup>Q</sup>**
- **Endothelial dysfunction**- shifts pattern of gene expression in endothelium to “**prothrombotic**.” state


## Stasis or Turbulent Blood Flow

- Turbulence causes arterial,<sup>Q</sup> whereas stasis causes venous thrombosis.<sup>Q</sup>

- Aortic and arterial dilations called **aneurysms<sup>Q</sup>** result in local stasis<sup>Q</sup>
- A dilated atrium in Rheumatic mitral valve stenosis is a site of profound stasis<sup>Q</sup> and a prime location for thrombosis<sup>Q</sup>
- Hyperviscosity syndromes like polycythemia<sup>Q</sup> and with deformed red cells as in sickle cell anemia.<sup>Q</sup>

## Blood Hypercoagulability

- It can either be primary (inherited) or secondary (acquired) hypercoagulable state.



## Investigations for Thrombophilia

- **Kaolin Clotting Time (KCT)<sup>Q</sup>**
- **Dilute Russel Viper Venom Test (DRVVT)<sup>Q</sup>**
- **Platelet neutralisation test<sup>Q</sup>**




## High Yield Facts

- Glutamine → arginine substitution at amino acid residue 506 that renders **factor V resistant** to cleavage and inactivation by protein C (also called **Activated Protein C resistance**)
- Factor V Leiden<sup>Q</sup> is the most common inherited cause of hypercoagulability
- **Antithrombin III, Protein C or Protein S deficiency** causes **venous thrombosis only<sup>Q</sup>**
- **Factor V Leiden** causes most commonly **venous thrombosis<sup>Q</sup>** but also **arterial thrombosis**
- Hyperhomocystinemia, APLA, DIC, HIT, PNH, Polycythemia Vera and dysfibrinogenemia cause both-arterial and venous thrombus<sup>Q</sup>
- Thrombomodulin is produced by all endothelial cells except those of cerebral microcirculation<sup>Q</sup>
- Thrombocytopenia (decrease of  $\geq 50\%$  platelet count), Timing of : HIT develops after exposure to **heparin for 5–14 days<sup>Q</sup>**, Thrombosis in **50%<sup>Q</sup>**, Other causes of thrombocytopenia not evident.

## ANTI-PHOSPHOLIPID ANTIBODY (APLA) SYNDROME

**Autoantibody<sup>Q</sup>**-mediated **acquired<sup>Q</sup> thrombophilia** characterized by **recurrent<sup>Q</sup> arterial or venous thrombosis<sup>Q</sup>** and/or **pregnancy morbidity** in the presence of **autoantibodies against phospholipid (PL)-binding plasma proteins ( $\beta$ -2 GPI)<sup>Q</sup>**






## Pathophysiology


### Diagnostic Criteria

One clinical event and at least one laboratory abnormality.



## Image-Based Question

1. Identify the defect in RBC morphology.




- |                 |                 |
|-----------------|-----------------|
| a. Sickle cell  | b. Bilte cells  |
| c. Schistocytes | d. Elliptocytes |



## Answer of Image-Based Question

1. Ans. (c) Schistocytes

- The given smear shows helmet cells with spiculated ends in a case of microangiopathic hemolytic anemia.



## Multiple Choice Questions

### DISORDERS OF PRIMARY HEMOSTASIS


- Glanzmann thrombasthenia is due to defect in**  
(Recent Question 2019)  
a. Gp IIb/IIIa                      b. Gp Ib-IX  
c. CD68                              d. Von Willebrand factor
- Bleeding time increased in which of the following conditions?**  
(Recent Question 2018)  
a. Von Willebrand disease      b. Hemophilia A  
c. DIC                                d. Both a & c
- Which of the following is true regarding Bernard-Soulier syndrome?**  
(Recent Question 2018)  
a. It is due to defect in platelet adhesion  
b. It is due to defect in platelet aggregation  
c. It is due to defect in platelet receptor GpIb-IX  
d. Both a and c
- Which of the following is/are released from dense granules of platelets?**  
(PGI Nov 2017)  
a. Serotonin                        b. Histamine  
c. PDGF                              d. ATP  
e. Lysosome
- Platelet adhesion to vessel wall is due to?**  
(AIIMS May 2015)  
a. Factor IX                        b. Fibrinogen  
c. vWF                                d. Fibronectin
- Which of the following is not a component of the dense granules of platelets?**  
(Recent Question 2015)  
a. ADP                                b. Calcium  
c. Epinephrine                      d. Platelet factor 4
- Earliest event of vascular trauma is?**  
(Recent Question 2016)  
a. Vasoconstriction                b. Platelet adhesion  
c. Platelet aggregation            d. Vasodilatation
- Which of the following is a qualitative defect of platelet?**  
(Recent Question 2016)  
a. VWD  
b. Hemophillia A  
c. Hemophillia C  
d. Glanzman thrombasthenia
- PDGF is present in which granules of platelets?**  
(Recent Question 2016)  
a. Alpha                                b. Beta  
c. Delta                                d. None
- In von Willebrand disease, what is true ?**  
(Recent Question 2016)  
a. Ristocetin aggregation test is decreased  
b. Ristocetin aggregation is Increased  
c. No effect  
d. May be increased or decreased
- An adolescent female presents with palpable purpura. Her hemogram suggested only anemia. What is your diagnosis?**  
(Recent Question 2015)  
a. ITP                                    b. TTP  
c. HUS                                d. HSP

- Bernard Soulier syndrome is a defect in?**  
(Recent Question 2015)  
a. Platelet Aggregation  
b. Platelet Adhesion  
c. Platelet activation  
d. Platelet agglutination
- A 7 years old boy presented with sudden onset petechiae and purpura. There was a history of URTI 2 weeks back. On examination, there was no hepatosplenomegaly. He is most probably suffering from:**  
(Recent Question 2015)  
a. ALL                                b. Acute viral infection  
c. ITP                                 d. Aplastic Anemia
- Platelet aggregation is caused by?**  
(Recent Question 2014)  
a. Nitrous oxide                      b. Thromboxane A<sub>2</sub>  
c. Aspirin                              d. PGE<sub>2</sub>
- Glanzmanns disease is-**  
(Recent Question 2014)  
a. Congenital defect of platelets  
b. Congenital defect of RBCs  
c. Defect of neutrophils  
d. Clotting factor deficiency
- Vasoconstricting mediator is-**  
(Recent Question 2014)  
a. Prostacyclin                        b. Thromboxane-A<sub>2</sub>  
c. PGG<sub>2</sub>                                d. Lipoxins
- All the following conditions cause thrombocytopenia except**  
(Recent Question 2014)  
a. Giant hemangioma  
b. Infectious mononucleosis  
c. HIV infection  
d. Iron deficiency anemia
- vWF is useful in:**  
(Recent Question 2014)  
a. Platelet adhesion                b. Platelet aggregation  
c. Clot formation                    d. Fibrinolysis
- Fever, fluctuating neurological symptoms, renal failure and severe thrombocytopenia are characteristic of**  
(Recent Question 2015)  
a. Hemolytic uremic syndrome  
b. Thrombotic thrombocytopenic purpura  
c. Idiopathic thrombocytopenic purpura  
d. Disseminated intravascular coagulation
- All the following are associated with HUS except**  
(Recent Question 2015)  
a. E coli O157:H7  
b. Shigelladysenteriae type 1  
c. Staphylococcus aureus  
d. Streptococcus pneumoniae
- A 4-year-old boy with sudden onset of petechial rashes. History of viral illness 2 weeks ago present. Investigations reveal thrombocytopenia and anti-platelets antibodies. What is your diagnosis?**  
(Recent Question 2015)  
a. Immune thrombocytopenic purpura  
b. Henochschonleinpurpura  
c. Thrombotic thrombocytopenic purpura  
d. Haemolytic uremic syndrome



22. After tonsillectomy, a 9 year old child is having continuous bleeding. Bleeding time and PTT are prolonged. Platelet count and PT are normal. What is your diagnosis (Recent Question 2015)
- Von willebrand disease
  - Vitamin K deficiency
  - Immune thrombocytopenic purpura
  - Hemophilia A
23. Find the false statement about TTP (Recent Question 2015)
- Renal failure
  - Negative direct anti-globulin test
  - Antibodies to ADAMTS 13
  - PT and aPTT are prolonged
24. True regarding heparin induced thrombocytopenia (Recent Question 2015)
- Platelet counts usually <10,000/uL
  - Increased risk of thrombosis
  - Associated with severe bleeding
  - HIT antibodies disappear in 5-14 days
25. Anticoagulant used for complete blood count? (Recent Question 2015)
- Trisodium citrate
  - Heparin
  - EDTA
  - Potassium oxalate
26. Increased PT indicates (Recent Question 2015)
- Platelet function defect
  - Intrinsic pathway defect
  - Extrinsic pathway defect
  - Common pathway defect
27. The following feature differentiates TTP and DIC (Recent Question 2015)
- Elevated D-dimers
  - Schistocytes in peripheral smear
  - Thrombocytopenia
  - Increased thrombin time/Decreased fibrinogen
28. Glanzmann thrombasthenia is due to: (Recent Question 2015)
- Abnormal platelet granule formation
  - Deficiency of von willebrand factor
  - Dysfunction of GpIIb-IIIa
  - Dysfunction of GpIb-IX
29. Spontaneous bleeding usually occurs when the platelet counts fall below (Recent Question 2015)
- 20000/uL
  - 50000/uL
  - 100000/uL
  - 120000/uL
30. Disorder of platelet aggregation: (Recent Question 2015)
- Bernard soulier syndrome
  - Glanzmann's thrombasthenia
  - Idiopathic thrombocytopenic purpura
  - Gray platelet syndrome
31. All are true about chronic ITP except: (Recent Question 2015)
- Common in females
  - Anti platelet antibodies
  - Splenomegaly
  - I.V. immunoglobulin used in treatment
32. Evans syndrome is: (Recent Question 2015)
- Autoimmune hemolytic anemia + ITP
  - Autoimmune hemolytic anemia + TTP
  - Autoimmune hemolytic anemia + HUS
  - Autoimmune hemolytic anemia + vWD

33. Five years old child presents with oliguria. There is history of bloody diarrhea 2 weeks ago. Coagulation tests are normal. Peripheral smear is given. What is your diagnosis: (Recent Question 2015)



- Thrombotic thrombocytopenic purpura
  - Idiopathic thrombocytopenic purpura
  - G6PD deficiency
  - Hemolytic uremic syndrome
34. The following is not a platelet function test (Recent Question 2015)
- Bleeding time
  - Prothrombin time
  - Clot retraction time
  - Prothrombin consumption index
35. Giant platelets, thrombocytopenia and cytoplasmic inclusions in the neutrophils is characteristic of (Recent Question 2015)
- May hegglin anomaly
  - Pelgerheut anomaly
  - Wiskott Aldrich syndrome
  - Gray platelet syndrome
36. Which plasma protein is necessary for adhesion of platelets to subendothelialfibres? (APPGMEE 14)
- Glycoprotein IIb
  - Von Willebrand factor
  - Platelet factor 3
  - Factor X
37. Select the FALSE statement among the following: (APPGMEE 14)
- Anti-D is used in Immune Thrombocytopenic purpura
  - DDAVP is used in von Willebrand disease type 3
  - DDAVP is used in severe form of Hemophilia A
  - EACA is used in Factor XI deficiency for minor bleeds
38. HUS is differentiated from TTP by? (JIPMER 2014)
- Presence of MAHA
  - Renal failure
  - Neurological symptoms
  - Absence of fever
39. Platelet is attached to collagen in endothelium via: (AIIMS Nov 2013)
- Factor 8
  - Factor 9
  - vWF
  - Fibronectin
40. Not true about von Willebrand disease: (PGI May 2013)
- aPTT is normal
  - Bleeding time is normal
  - Most common pattern of inheritance is autosomal recessive
  - Bleeding from mucosa in oral cavity may present
  - Normal platelet count



- 41. A newborn baby presented with profuse bleeding from the umbilical stump after birth. Rest of the examination and PT, APTT are within normal limits. Most probable diagnosis is-** (AIIMS May 12)
- Factor X deficiency
  - Glanzmann thrombasthenia
  - Von Willebrand disease
  - Bernard Soulier disease
- 42. What abnormalities will be present in a patient having deficiency of Von Willebrand factor?** (DNB Aug 12 Pattern, DNB Dec 09)
- Increased aPTT, Increased PT
  - Decreased PT, Increased aPTT
  - Normal PT, Normal aPTT
  - Normal PT, Increased aPTT
- 43. Which is not true regarding Bernard Soulier syndrome** (AI II)
- Ristocetin aggregation is normal
  - Aggregation with collagen and ADP is normal
  - Large platelets
  - Thrombocytopenia
- 44. Which of the following statements about platelet function defects is true?** (AI II)
- Normal platelet count with prolonged bleeding time
  - Thrombocytopenia with prolonged bleeding time
  - Thrombocytosis with prolonged bleeding time
  - Normal platelet count with normal bleeding time
- 45. Increased bleeding time is seen in:** (PGI May 2011)
- Coumarin derivative administration
  - Thrombocytopenia
  - Congenital afibrinogenemia
  - DIC
  - Hemophilia A
- 46. Normal platelet count is/are seen in -** (PGI Nov 10)
- DIC
  - Shaken baby syndrome
  - Microangiopathic hemolytic anemia
  - Splenomegaly
  - Kaasabach-Merritt syndrome
- 47. A 25-year-old asymptomatic female underwent pre-op coagulation testing. Her BT was 3 minutes, PT was 15/14 seconds, aPTT 45/35 seconds. Platelet counts were 2.5 lac/uL, factor VIII levels 60 IU/dL. Most likely diagnosis is:** (AIIMS Nov 11)
- Factor IX deficiency
  - vWD type 3
  - Factor VIII inhibitor
  - Lupus anticoagulant
- 48. Which test is NOT used to assess the Platelet function?** (Recent Question 2013)
- Clot retraction time
  - Bleeding time
  - Ristocetin induced assay
  - Clot lysis time
- 50. True about Hemophilia B are?** (PGI Nov 2018)
- Autosomal recessive
  - Cryoprecipitate for treatment
  - Haemophilia B had factor 8 <5%
  - X linked
  - PT is raised
- 51. Increase PT is seen with?** (PGI May 18)
- Vitamin K deficiency
  - Factor V deficiency
  - Factor VIII deficiency
  - Factor IX deficiency
  - Warfarin administration
- 52. Which of the following is true regarding Von Willebrand disease?** (Recent exam 2018)
- Type 1 & 3 are associated with quantitative defects in vWF
  - Normal platelet count
  - Desmopressin stimulates release of VWF
  - All of the above
- 53. Which of the following is the most common manifestation of hemophilia?** (Recent exam 2018)
- Hemoptysis
  - Hemarthrosis
  - Hematemesis
  - Mucosal bleeding
- 54. In PT test, the addition of  $\text{Ca}^{2+}$  & tissue thromboplastin activates which pathway?** (AIIMS Nov 2017)
- Extrinsic
  - Intrinsic
  - Fibrinolytic
  - Common
- 55. Which coagulation factor is not in circulating form in blood?** (JIPMER 2017)
- F XI
  - F X
  - F III
  - F XIII
- 56. Which of the following is not associated with prolonged prothrombin time?** (PGI Nov 2017)
- Haemophilia A
  - Von-Willebrand disease
  - Factor VII deficiency
  - Disseminated intravascular coagulation (DIC)
- 57. Both PT and aPTT will be increased in deficiency of?** (PGI May 2017)
- Factor 2
  - Factor 5
  - Factor 8
  - Factor 10
  - Factor 12
- 58. Feature(s) of XIII factor deficiency is/are:** (PGI May 2016)
- Delayed wound closure
  - Clot solubility tests are abnormal
  - $\uparrow$ aPTT
  - $\uparrow$ PT
  - $\uparrow$ BT
- 59. Which of the following statement(s) is/are correct except:** (PGI May 2016)
- Increased PT in extrinsic pathways
  - Increased aPTT in intrinsic pathways
  - If platelet count is > 1.5 lac/microL, then normal homeostasis present
  - BT is decreased in platelet abnormality
- 60. Endotoxin first damages ?** (Recent Question 2016-17)
- Endothelium
  - RBC
  - Platelets
  - WBC

#### DISORDERS OF SECONDARY HEMOSTASIS

- 49. Which of the following is required for the function of heparin ?** (AIIMS May 18)
- Protein C
  - Protein S
  - Antithrombin III
  - Thrombomodulin





- 328



- 86. Hemophilia C is deficiency of** (WBPG 2015)  
 a. Factor VIII                      b. Factor IX  
 c. Factor X                         d. Factor XI
- 87. LakiLorand factor is** (Recent Question 2014)  
 a. Factor X                         b. Factor XI  
 c. Factor XII                      d. Factor XIII
- 88. Levels of all coagulation factors are increased in pregnancy EXCEPT** (APPGMEE 14)  
 a. Factor VIII                      b. Factor IX  
 c. Factor X                         d. Factor XI
- 89. Factor Xa is necessary for conversion of prothrombin to thrombin** (APPGMEE 14)  
 a. Only in the extrinsic pathway  
 b. Only in the intrinsic pathway  
 c. As part of both extrinsic and intrinsic pathways  
 d. Only if the normal blood clotting cascade is inhibited
- 90. Isolated rise in aPTT is seen in?** (Recent Question 2013)  
 a. Von Willebrand's disease  
 b. Factor 7 deficiency  
 c. Vitamin K deficiency  
 d. Anti phospholipid antibodies
- 91. Increased Prothrombin time is/are seen in:** (PGI May 12)  
 a. Pt on oral anticoagulant  
 b. Pt with liver disease  
 c. Factor X deficiency  
 d. Factor VII deficiency      e. Vit. K deficiency
- 92. Prolonged apTT, but asymptomatic is seen in deficiency of factor** (JIPMER 2012)  
 a. V                                      b. VII  
 c. X                                      d. XII
- 93. All of the following are true about blood coagulation, except:** (AI 11)  
 a. Factor X is part or both intrinsic and extrinsic pathways  
 b. Extrinsic pathway is activated by contact with negatively charged surfaces  
 c. Intrinsic pathway can be activated in vitro  
 d. Calcium is required in several steps of coagulation
- 94. True regarding prothrombin time measurement?**  
 a. Platelet rich plasma is required (AIIMS Nov 11)  
 b. Activate with kaolin  
 c. Should be measured within 2 hours  
 d. Immediate refrigeration to preserve coagulation factor viability
- 95. Increased PT and Normal PTT are found in?** (DNB Dec 11)  
 a. Von Willebrand disease      b. Factor 7 deficiency  
 c. Factor 8 deficiency          d. Thrombin deficiency
- 96. The best screening test for hemophilia -** (DNB Dec 11)  
 a. PT                                      b. CT  
 c. PTT                                    d. BT
- 97. Prolonged PT and Normal PTT may be seen in:**  
 a. Thrombocytopenia          b. DIC (DNB June 11)  
 c. Vit. K deficiency              d. Aspirin toxicity
- 98. A 9 years old boy with elevation in both PT and aPTT. What is the diagnosis?** (AIIMS Nov 10)  
 a. Defect in extrinsic pathway  
 b. Defect in intrinsic pathway  
 c. Platelet function defect  
 d. Defect in common pathway

## DIC

- 99. Which among the following laboratory investigation is best to reveal about bleeding in Disseminated Intravascular Coagulation (DIC)?** (AIIMS May 18)  
 a. Increased PT  
 b. Increased aPTT  
 c. Decreased fibrinogen  
 d. Increased FDPs
- 100. Most common cause of DIC** (Recent Question 2015)  
 a. Obstetric complications      b. Cyanotic heart disease  
 c. Malignancies                      d. Extensive burns
- 101. The most sensitive test for DIC is** (Recent Question 2015)  
 a. Bleeding time                      b. Clotting time  
 c. Prothrombin time                d. FDP level
- 102. The following is not a contraindication for heparin in DIC** (Recent Question 2015)  
 a. If the platelet count cannot be maintained at  $\geq 50,000/\text{mL}$   
 b. In cases of central nervous system/gastrointestinal bleeding  
 c. Placental abruption  
 d. DIC associated with malignancy
- 103. What is not associated with DIC** (Recent Question 2015)  
 a. Thrombocytopenia              b. Increased PT  
 c. Hyperfibrinogenemia          d. Increased FDP
- 104. DIC causes all except?** (Recent Question 2014)  
 a. Increased fibrinogen level  
 b. Increased d-dimer  
 c. Decreased platelet counts  
 d. Decreased clotting factors
- 105. In DIC, following are seen except** (Recent Question 2014)  
 a. Fibrinogen decreased          b. Thrombocytopenia  
 c. Normal APTT                      d. PT elevation

## THROMBOPHILIAS

- 106. Which of the following is true regarding Factor V Leiden mutation?** (Recent exam 2018)  
 a. Increased risk of deep vein thrombosis  
 b. Factor V becomes resistant to cleavage by protein C  
 c. Glutamine to Arginine substitution at 506  
 d. All of the above
- 107. True about hematological disorder?** (PGI May 2017)  
 a. In Haemophilia B factor 8 is useful.  
 b. Both PT and aPTT are increased in DIC  
 c. IV Ig is useful in ITP  
 d. Cryoprecipitate is useful in hemophilia A
- 108. Most common inherited hyper coagulation defect?** (Recent Question 2016-17)  
 a. Factor 5 mutation  
 b. Prothrombin mutation  
 c. Hyperhomocytinemia  
 d. Protein C deficiency
- 109. Tumors causing thrombosis are all except?** (Recent Question 2016-17)  
 a. Stomach                              b. Lung  
 c. Pancreas                            d. Breast
- 110. Antithrombin is activated by?** (Recent Question 2016)  
 a. Heparin                              b. Factor I  
 c. Fctor V                              d. Factor VIII



**111. Which of the following is not a usual cause of DVT?**

(Recent question 2016)

- a. Injury
- b. Hypothermia
- c. Stasis
- d. Hypercoagulability

**112. Treatment of antiphospholipid antibody syndrome**

(Recent Question 2015)

- a. Warfarin for life
- b. Warfarin for 1 year
- c. IV immunoglobulin
- d. Glucocorticoids

**113. Not a feature of antiphospholipid antibody syndrome**

(Recent Question 2015)

- a. Budd-Chiari syndrome
- b. Coomb's positive hemolytic anemia
- c. Libman sacks endocarditis
- d. Priapism

**114. Leiden mutation is related to?** (Recent Question 2015)

- a. Factor V
- b. Factor VIII
- c. Antithrombin III
- d. Severe inherited bleeding

**115. All antibodies are seen in APLA except?**

(Recent Question 2015)

- a. Anti- $\beta$ 2 glycoprotein
- b. Anti-Prothrombin
- c. Anti-Phospholipid
- d. Anti-Cardiolipin

**116. All are true about Virchow's triad except-**

(Recent Question 2014)

- a. Concerns the risk of intravascular thrombus
- b. Depends on endothelial injury
- c. Depends on platelet activation
- d. Depends on stasis

**117. Leiden factor is -**

(Recent Question 2014)

- a. Factor VI
- b. Factor VIII
- c. Factor IV
- d. Factor V

**118. All endothelial cells produce thrombomodulin except those found in -**  
(Recent Question 2013, AI 05)

- a. Hepatic circulation
- b. Cutaneous circulation
- c. Cerebral microcirculation
- d. Renal circulation

**119. Heparin treatment is monitored by:**

(Recent Question 2013)

- a. PT INR
- b. aPTT
- c. Vit K levels
- d. PT

**120. Presentation of antiphospholipid syndrome includes:**

(PGI May 2013)

- a. Recurrent abortion
- b. Fetal death
- c. Both arterial and venous thrombosis
- d. Prolonged aPTT
- e. Prolonged PT

**121. Which of the following antibodies is most frequently seen in Antiphospholipid Syndrome?** (AI 11)

- a. Beta 2 microglobulin antibody
- b. Anti-nuclear antibody
- c. Anti-centromere antibody
- d. Anti- beta 2 glycoprotein antibody

**122. Both arterial and venous thrombosis occur in:**

(PGI Nov 2011)

- a. Antiphospholipid antibodies
- b. Antithrombin III deficiency
- c. Hyperhomocysteinemia
- d. Protein C deficiency
- e. Mutation in factor V gene

**123. Hypercoagulability due to defective factor V gene is called**  
(AIIMS 10)

- a. Lisbon mutation
- b. Leiden mutation
- c. Aruiphospholipid syndrome
- d. Inducible thrombocytopenia syndrome

**124. Warfarin skin necrosis is caused by-**

(Recent Question 2014)

- a. Protein C / Protein S deficiency
- b. APLA
- c. Vitamin K deficiency
- d. Fibrinogen deficiency

**125. Tissue Factor Pathway Inhibitor (TFPI) inhibits?**

(Recent Question 2013)

- a. Factor V
- b. Factor VII
- c. Factor X
- d. Factor IV

**126. Screening test for patients suspected of having a hypercoagulable State includes?** (Recent Question 2013)

- a. Russell viper venom time
- b. Prothrombin time
- c. Heparin cofactor II
- d. Clotting Factor levels

**127. International Normalised Ratio (INR) is defined as?**

- a. (PT patient/ PT control)<sup>ISI</sup>
- b. (PT control/ PT patient)<sup>ISI</sup>
- c. (PT patient/ PT control)<sup>1/ISI</sup>
- d. (PT control/ PT patient)<sup>1/ISI</sup>

**128. Increased clotting time and decreased platelet count is associated with?** (WBPG 10, 12)

- a. Vascular injury
- b. Volume contraction
- c. Decreased platelet clumping
- d. Decreased platelet phospholipid

**129. Warfarin therapy is monitored by:**

(Recent Question 2013)

- a. PT INR
- b. apTT
- c. Vit K levels
- d. PT



## Answers with Explanations

1. Ans. (a) **Gp IIb/IIIa**

2. Ans. (d) **Both a & c**

BT	↑
Deficiency of platelets	Functional defect of platelets
VWD is adhesion defect as platelets cannot bind with vessel wall. So primary hemostatic plug is not formed.	Dic is hyper coagulable to state so many clots are formed which consume all platelets.

3. Ans. (d) **Both a and c**

(Ref: Robbins 9th/pg 660)

Bernard-Soulier syndrome illustrates the consequences of defective adhesion of platelets to subendothelial matrix. Bernard-Soulier syndrome is caused by an inherited deficiency of the platelet membrane glyco protein complex Ib-IX. This glycoprotein is a receptor for vWF and is essential for normal platelet adhesion to the subendothelial extracellular matrix.

4. Ans. (a, b, d) **a. Serotonin; b. Histamine; d. ATP**

The dense granules of platelets contain adenosine diphosphate (ADP), adenosine triphosphate (ATP), ionized calcium, serotonin and histamine. PDGF is a constituent of alpha granules.

5. Ans. (c) **vWF** (Ref: Robbins 9th/pg 660; 8th/pg 670)

Platelet adhesion to vessel wall is due to vWF on endothelium and GpIb/IX receptors on the surface of platelets.

6. Ans. (d) **Platelet factor 4** (Ref: Robbins 9th/pg 660)

**Dense (or δ) granules** of platelets contain: ("DENSE")

- aDENosine diphosphate (ADP)<sup>Q</sup>, Serotonin<sup>Q</sup>, Epinephrine<sup>Q</sup>, Ca

7. Ans. (a) **Vasoconstriction** (Ref: Robbins 9th/pg 661)

Endothelial Injury causes vasoconstriction<sup>Q</sup> (due to endothelin, serotonin) immediately and markedly reduces blood flow to the injured area.

8. Ans. (d) **Glanzman thrombosthenia** (Ref: R 9th/pg 660)

9. Ans. (a) **Alpha** (Ref: Robbins 9th/pg 660; 8th/pg 670)

α-Granules of platelets contains:

- Pselectin, Fibrinogen<sup>Q</sup>, V<sup>Q</sup>, VIII<sup>Q</sup>, and vWF<sup>Q</sup>, Platelet factor 4<sup>Q</sup>, (PDGF), TGF-β)

10. Ans. (a) **Ristocetin aggregation test is decreased**

(Ref: Wintrobe's 12th ed/1392)

**Ristocetin cofactor assay<sup>Q</sup>/Ristocetin induced platelet aggregation is decreased or absent in vWD**

11. Ans. (d) **HSP** (Ref: Harrison 18th/Chapter 53)

12. Ans. (b) **Platelet Adhesion**

(Ref: Robbins 9th/pg 660; 8th/pg 670)

13. Ans. (c) **ITP** (Ref: 9th/pg 657-658; 8th/pg 667-668)

A 7 year old boy presented with sudden onset petechiae and purpura with a history of URTI 2 weeks back and there is no hepatosplenomegaly. He is most probably suffering from ITP. For complete discussion on **ITP**; Refer to pretext of this chapter.

14. Ans. (b). **Thromboxane A2**

(Ref: Robbins 9th/pg 116-117)

### Thromboxane A2 (TXA2)

- It is **produced by activated platelets** and has prothrombotic properties
- It is an **Arachidonic acid metabolite** of COX pathway
- Functions: It is a potent **platelet aggregator & vasoconstrictor**
- Its effect is **neutralized by prostacyclin (PGI2)**

15. Ans. (a) **Congenital defect of platelets**

(Ref: Robbins 9th/pg 660; 8th/pg 670)

Glanzmanns disease is a congenital defect of platelets; Refer Ans 4 above;

16. Ans. (b) **Thromboxane-A2** (Ref: Robbins 9th/pg 116-117)

17. Ans. (d) **Iron deficiency anemia**

(Ref: Robbins 9th/pg 657-658; 8th/pg 667-668; Wintrobe's 12th/pg 1595)

Thrombocytopenia refers to platelet count < **1.5 lac/cu mm<sup>Q</sup>**

Refer to pretext of this chapter for important causes of Thrombocytopenia

- **Thrombocytopenia** is seen in **50%** of patients with **Infectious Mononucleosis**
- **Thrombocytosis** is seen in **Iron deficiency Anemia**, platelet counts rise to twice the normal value and **return to normal with Iron therapy**. Exact cause unknown, but probably **due to ↑ level of Erythropoietin**

18. Ans. (a) **Platelet adhesion** (Ref: Robbins 9th/pg 661)

2 major **functions of vWF** are: Platelet adhesion via GpIb/IX and Stabilizes factor VIII in circulation



**19. Ans. (b) Thrombotic thrombocytopenic purpura**

(Ref: Robbins 9th/pg 659; 8th/pg 669)

- Features of hemolytic uremic syndrome include Hyperkalemia (due to deranged renal function), Anemia (due to hemolysis) & renal microthrombi
- But Neuro psychiatric disturbances are seen in TTP & not in HUS

**20. Ans. (c) Staphylococcus aureus**

(Ref: Robbins 9th/pg 659)

**21. Ans. (c) Thrombotic thrombocytopenic purpura**

(Ref: Robbins 9th/pg 659; 8th/pg 669)

Thrombocytopenia and anti-platelets antibodies in a child presenting with purpura after 2 weeks of viral illness is characteristic of ITP.

**22. Ans. (a) Von Willebrand disease**

(Ref: Robbins 9th/pg 661)

- a. Von willebrand disease: True: Bleeding time and aPTT prolonged, PT and platelet count normal
- b. Vitamin K deficiency: False: PT prolonged
- c. Immune thrombocytopenic purpura: False aPTT is normal
- d. Hemophilia A: False: Bleeding time normal

**23. Ans. (d) PT and aPTT are prolonged** (Ref: R 9th/pg 662)

**PT and aPTT are normal and not prolonged.**

**24. Ans. (b) Increased risk of thrombosis** (Ref: R 9th/pg 657)

**Options:**

- a. Platelet counts usually  $<10,000/\mu\text{L}$ : **False:** Thrombocytopenia (decrease of  $>50\%$ )
- b. **Increased risk of thrombosis: True: seen in 50%**
- c. Associated with severe bleeding: False: Bleeding is uncommon in HIT
- d. HIT antibodies disappear in 5-14 days: False: they appear after 5-14 days.

**25. Ans. (c) EDTA** (Ref: Robbins 9th/pg 656)

K2 (Dipotassium)-EDTA is the common anticoagulant for hemogram analysis.

**26. Ans. (c) Extrinsic pathway defect**

(Ref: Robbins 9th/pg 661)

**27. Ans. (d) Increased thrombin time/Decreased fibrinogen**

(Ref: Robbins 9th/pg 663; 8th/pg 673)

**Increased thrombin time/Decreased fibrinogen is seen in DIC only not in TTP.**

**28. Ans. (c) Dysfunction of GpIIb-IIIa**

(Ref: Wintrobe's 13th ed/pg 689)

**29. Ans. (a) 20000/uL** (Ref: Robbins 9th/pg 662; 8th/pg 692)

In general, the risk of significant spontaneous hemorrhage increases gradually as the platelet count drops to  $<50 \times 10^9/\text{L}$  and is high at counts  $<5 \times 10^9/\text{L}$ . So among the options the best answer is 10,000/uL.

**30. Ans. (b) Glanzmann's thrombasthenia**

(Ref: Robbins 9th/pg 662)

**31. Ans. (c) Splenomegaly**

(Ref: Robbins 9th/pg 657; 8th/pg 667)

Splenomegaly is very rare in ITP

**32. Ans. (a) Autoimmune hemolytic anemia + ITP**

(Ref: Wintrobe's 13th ed/pg 968)

**Evans syndrome:** Antiplatelet antibodies in patients with ITP do not usually cross-react with RBCs. However, RBC fragmentation because of weak complement activation, may occur. Such patients may also have a positive Coombs test and autoimmune hemolytic anemia. This is referred as Evans syndrome.

**33. Ans. (d) Hemolytic uremic syndrome**

(Ref: R 9th/pg 659)

Features of oliguria following an episode of diarrhea with peripheral smear finding of Schistocytes (Helmet cells) is suggestive of Hemolytic uremic syndrome.

**34. Ans. (b) Prothrombin time**

(Ref: Robbins 9th/pg 662)

- a. **Bleeding time** assess ability of platelets to form platelet plug
- b. **Prothrombin time** assesses extrinsic pathway of coagulation
- c. **Clot retraction study** measures the time taken for a platelet plug to undergo last step of retraction, which indicates overall platelet function.
- d. An abnormal **Prothrombin consumption index** indicates impaired availability of platelet membrane phospholipid for coagulation.

**35. Ans. (a) May hegglin anomaly**

(Ref: Wintrobe's 12th ed/pg 1549)

May-Hegglin anomaly is a rare, dominantly inherited disorder characterized by large (2 to 5  $\mu\text{m}$ ), well-defined, basophilic and pyroninophilic inclusions in granulocytes (neutrophils, eosinophils, basophils, monocytes) and accompanied by thrombocytopenia and giant platelets containing few granules.

**36. Ans. (b) Von Willebrand factor**

(Ref: Robbins 9th/pg 661)

**Von Willebrand factor** is necessary for adhesion of platelets to subendothelial fibres.



**37. Ans. (b) DDAVP is used in von Willebrand disease type 3**

(Ref: Wintrobe's 12th/pg 1392, 1410)

**DDAVP is used in von Willebrand disease type 1 & 2a, but not in type 3**

**38. Ans. (c) Neurological symptoms**

(Ref: Robbins 9th/pg 659)

HUS is differentiated from TTP by **Absence of neurological symptoms**

- **Neurologic findings (due to microthrombi formed in vessels of CNS) are present in TTP in addition to the triad seen in HUS (Microangiopathic hemolytic anemia, Thrombocytopenia & Renal failure)**

**39. Ans. (c) vWF**

(Ref: Robbins 9th/pg 661; 8th/pg 671)

**Platelet is attached to collagen in endothelium via vWF**

**40. Ans. (a, b, c); a. aPTT is normal; b. Bleeding time is normal; c. Most common pattern of inheritance is autosomal recessive**

(Ref: Robbins 9th/pg 661; 8th/pg 671)

Discussing the options about von Willebrand disease, one by one-

A.	False	<b>vWF stabilizes factor VIII, which takes part in Intrinsic pathway. So in vWD, when vWF is deficient, aPTT is prolonged</b>
B.	False	Because there is <b>defective platelet adhesion in vWD</b> → ↑ BT
C.	False	Most common pattern of inheritance is <b>Autosomal dominant for vWD type I</b>
D.	True	Mucosal bleeding is seen in vWD.
E.	True	In vWD platelet count is normal

**41. Ans. (b) Glanzmann thrombasthenia**

(Ref: Robbins 9th/pg 660; 8th/pg 670)

A newborn baby presented with profuse bleeding from the umbilical stump after birth.

This is a case of severe inherited bleeding disorder. As **PT & aPTT are normal, Glanzmann thrombasthenia** is the most probable cause, which can present with **profuse bleeding soon after birth;**

It can be **diagnosed by platelet aggregation studies**, which will be defective.

The other differential diagnosis in this scenario is **Factor XIII deficiency** (not given in the option)

**42. Ans. (d) Normal PT, Increased a PTT**

(Ref: R 9th/pg 662)

In vWD, **Normal PT, Increased a PTT**

**vWF stabilizes factor VIII**, which takes part in Intrinsic pathway. So in vWD, when vWF is deficient, **Intrinsic**

**pathway of coagulation is affected, giving rise to prolonged aPTT**; but as extrinsic pathway is not affected, PT remains normal

**43. Ans. (a) Ristocetin aggregation is normal**

(Ref: Robbins 9th/pg 660; 8th/pg 670)

**44. Ans. (a) Normal platelet count with prolonged bleeding time** (Ref: Robbins 9th/pg 660; 8th/pg 670)

In **platelet function defects**, there is **normal platelet count with prolonged bleeding time**

**45. Ans. (b, c, d); b. Thrombocytopenia; c. Congenital afibrinogenemia; d. DIC** (Ref: Robbins 9th/pg 656)

↑ **bleeding time (BT)** is seen in:

A.	<b>Coumarin derivative administration</b>	No	<b>PT (initially) followed by both PT &amp; aPTT will be abnormal</b>
B.	Thrombocytopenia	Yes	Because of defective formation of primary hemostatic plug
C.	<b>Congenital afibrinogenemia</b>	No	<b>PT, TT &amp; aPTT will be abnormal, as formation of stable clot is only affected</b>
D.	DIC	Yes	Due to thrombocytopenia in DIC
E.	<b>Hemophilia A</b>	No	<b>aPTT abnormal, PT normal</b>

**46. Ans. (b) Shaken baby syndrome** (Ref: R 9th/pg 657-658)

**Normal platelet count is/are seen in Shaken baby syndrome.**

**47. Ans. (c) Factor VIII inhibitor**

(Ref: Wintrobe's 12th/pg 1452)

This is the case of an asymptomatic female with normal BT, PT & platelet counts; aPTT is ↑, factor VIII levels (N=50-150 IU/dl) are near normal.

This is a **controversial** MCQ, as both factor VIII inhibitor & lupus anticoagulant are possible answers

Discussing the options one by one:

A.	Factor IX deficiency	Also called Hemophilia B, excluded because patient is asymptomatic; Factor IX deficiency leads to clinical bleeding
B.	vWD type 3	Excluded because patient is asymptomatic & BT is normal
C.	<b>Factor VIII inhibitor</b>	<b>Best possible answer as patient is asymptomatic with prolonged aPTT due to mild deficiency of factor VIII</b>
D.	Lupus anticoagulant	Second best answer, as isolated ↑aPTT in an asymptomatic female can be due to Lupus anticoagulant. However, <b>usually the patient presents with thrombocytopenia &amp; prolonged PT also.</b>



48. Ans. (d) **Clot lysis time**

(Ref: Wintrobe's 12th/pg 499)

Discussing the options one by one,

A. Clot retraction time	Used to assess the <b>global contractile function</b> of platelets; hence it is a test for platelet function
B. Bleeding time	First screening test to assay <b>platelet function defects</b>
C. Ristocetin induced assay	Ristocetin induced <b>platelet adhesion</b> and <b>aggregation study</b> is used to assess <b>vWF and gplb/IX deficiency</b>
D. Clot lysis time	Used for assessing FXIII deficiency which is responsible for strength of clot

49. Ans. (c) **Antithrombin III**

50. Ans. (b) **Cryoprecipitate for treatment (c) Haemophilia B had factor 8 <5%**

Hemophilia B is deficiency of factor 9.

51. Ans. (a, b, e) **a. Vitamin K deficiency; b. Factor V deficiency; e. Warfarin administration**

52. Ans. (d) **All of the above**

53. Ans. (b) **Hemarthrosis**

(Ref: Robbins 9th/pg 663)

In all symptomatic haemophilia cases there is a tendency toward easy bruising and massive hemorrhage after trauma or operative procedures. In addition, "spontaneous" hemorrhages frequently occur in regions of the body that are susceptible to trauma, particularly the joints, where they are known as hemarthroses. Recurrent bleeding into the joints leads to progressive deformities that can be crippling. Petechiae are characteristically absent.

54. Ans. (a) **Extrinsic**

To PT we add of  $\text{Ca}^{2+}$  & tissue thromboplastin which activates extrinsic pathway, while to test for aPTT we add kaolin, silica which activates intrinsic pathway.

55. Ans. (c) **F III**

Tissue factor (factor III) is also called as platelet tissue factor. It is found on the outside of blood vessels and is not exposed to the bloodstream. It initiates the extrinsic pathway at the site of injury.

56. Ans. (a, b) **a. Haemophilia A; b. Von-Willebrand disease**

Haemophilia A is a deficiency of FVIII & Von-Willebrand disease is a defect of vWF so both of them are associated with aPTT rather than PT.

57. Ans. (a, b, d) **a. Factor 2; b. Factor 5; d. Factor 10**

Both PT and aPTT will be increased in deficiency of factors of common pathway.

58. Ans. (a, b) **a. Delayed wound closure, b. Clot solubility tests are abnormal**

(Ref: Harrison 19th/ 733, 736; CMDT 2016/ 556-57)

59. Ans. (d) **BT is decreased in platelet abnormality**

(Ref: Robbins 9th/118-119)

60. Ans. (a) **Endothelium** (Ref: Robbins 9th/pg 131-133)

61. Ans. (b) **aPTT** (Ref: Robbins 9th/pg 656-661)

62. Ans. (d) **PT-INR** (Ref: Wintrobe's 12th/pg 1489)

63. Ans. (c) **Defect in common pathway**

(Ref: Robbins 9th/pg 118-119)

64. Ans. (c) **Partial thromboplastin time**

(Ref: Robbins 9th/pg 656)

65. Ans. (c) **Factor XIII** (Ref: Robbins 9th/pg 662; 8th/pg 672)

66. Ans. (c) **VIII** (Ref: Wintrobe's 13th/pg 1389)

vWF is required for normal platelet adhesion, and also acts as a carrier of factor VIII in the plasma

67. Ans. (d) **VII** (Ref: Robbins 9th/pg 662; 8th/pg 672)

68. Ans. (d) **Factor XIII def** (Ref: Robbins 9th/pg 662)

- Abnormal bleeding manifests shortly after birth, when bleeding from the healthy umbilical cord remnant occurs.
- Umbilical stump bleeding is an uncommon presentation for other congenital bleeding disorders. Rebleeding at circumcision is also common.
- The most life-threatening complication of factor XIII deficiency is spontaneous intracranial hemorrhage

69. Ans. (d) **IIa** (Ref: Robbins 9th/pg 662; 8th/pg 672)

70. Ans. (b) **vWD-type 2**

(Ref: Robbins 9th/pg 663; 8th/pg 673)

71. Ans. (a) **Low platelet count** (Ref: Robbins 9th/pg 657)

- This test measures the amount of time it takes for a blood clot to pull away from the walls of a test tube (Shrinking).
- The edges of the blood vessel wall at the point of injury are slowly brought together again to repair the damage.
- It is used to evaluate and manage blood platelet disorders, including Glanzmann's thrombasthenia
- So Clot retraction depends primarily on the number and activity of the blood platelets



72. **Ans. (b) Hepatocytes**  
(Ref: Robbins 9th/pg 662; 8th/pg 672)
73. **Ans. (a) Von willebrand disease**  
(Ref: Robbins 9th/pg 660)
74. **Ans. (c) Conversion of fibrinogen to fibrin impaired**  
(Ref: Robbins 9th/pg 660; 8th/pg 670)
75. **Ans. (c) VIII** (Ref: Robbins 9th/pg 660; 8th/pg 670)
76. **Ans. (a) VII** (Ref: Robbins 9th/pg 660; 8th/pg 670)  
**Factor VII is involved in extrinsic pathway.**
77. **Ans. (c) Desmopressin**  
Endothelial stores of vWF can be released therapeutically with administration of desmopressin, So its administration can be helpful for treatment of mild vWD deficiency.
78. **Ans. (a) ↑ aPTT** (Ref: Robbins 9th/pg 662; 8th/pg 672)
79. **Ans. (d) Disseminated intravascular coagulation**  
(Ref: Wintrobe's 12th/pg 1437)
- Levels of fibrin degradation products, including cross-linked fibrin degradation products (D-dimer), are usually increased in the presence of acute venous thromboembolism.
  - Absence of an elevated level of D-dimer in patients undergoing an evaluation for acute DVT or PE has an excellent negative predictive value for thrombosis.
80. **Ans. (a) Normal PT, Elevated APTT**  
(Ref: Robbins 9th/pg 662; 8th/pg 672)
81. **Ans. (b, c, d, e); b. Factor 9 deficiency; c. X-linked disorder; d. Clinically indistinguishable from hemophilia A; e. Fresh frozen plasma given for treatment**  
(Ref: Robbins 9th/pg 662; 8th/pg 672)
- About option E. The ideal treatment for hemophilia B is Factor IX concentrate or Prothrombin Complex. But FFP is an alternative when these are not available, hence we will take option E as true.
82. **Ans. (b) Factor VIII deficiency** (Ref: R 9th/pg 662-663)
- Hemophilia A is Factor VIII deficiency
  - Hemophilia B is Factor IX deficiency
  - Hemophilia C is Factor XI deficiency
  - Pseudo-Hemophilia is vWD
  - Parahemophilia is factor V deficiency
83. **Ans. (a) Hemophilia A** (Ref: Robbins 9th/pg 662-663)  
**X-linked recessive disease in male with clotting defect is Hemophilia A**

84. **Ans. (a) Early solubiliztion of clot**  
(Ref: Wintrobe's 12th/pg)  
**Early solubiliztion of clot** is a feature of **Factor XIII deficiency**
85. **Ans. (b) Factor IX**  
(Ref: Wintrobe's 12th/pg 499)  
Platelet associated coagulation factors are: Fibrinogen, vWF, Factor V, XI, XIII & HMWK
86. **Ans. (d) Factor XI**  
(Ref: Robbins 9th/pg 662-663)  
**Hemophilia C is Factor XI deficiency**
87. **Ans. (d) Factor XIII** (Ref: Wintrobe's 12th/pg 1452)  
**LakiLorand factor** is the other name for Factor XIII
88. **Ans. (d) Factor XI** (Ref: Wintrobe's 12th/pg 1403)  
In pregnancy,
- Factor II & V remain unchanged
  - Factor XI & XIII are decreased
  - Other coagulation factors ↑ making pregnancy a hypercoagulable state
89. **Ans. (c) As part of both extrinsic and intrinsic pathways**  
(Ref: Robbins 9th/pg 118-119; 8th/pg 119)  
**Activated factor IX along with factor VIIIa (Intrinsic pathway) & factor VIIa (extrinsic pathway), converge to activate Factor X to Xa is necessary for conversion of prothrombin to thrombin (common pathway)**
90. **Ans. (a, d); a. Von Willibrand's disease; d. Anti phospholipid antibodies**  
(Ref: Robbins 9th/pg 662)  
Isolated rise in aPTT is seen in Von Willibrand's disease, as vWF deficiency causes factor VIII destabilization; Even Anti phospholipid antibodies (option D). seen in APLA can cause ↑aPTT, but prolongation of PT & TT can also be seen; So the best answer is A.
91. **Ans. (a, b, c, d, e); a. Pt on oral anticoagulant; b. Pt with liver disease; c. Factor X deficiency; d. Factor VII deficiency; e. Vit. K deficiency**  
(Refer to Pretext of this chapter)
92. **Ans. (d) XII**  
(Ref: Wintrobe's 12th/pg 1403)  
**Factor XII deficiency usually is not associated with hemorrhagic manifestations.** On the other hand, **myocardial infarction and thrombophlebitis** have been observed in patients with severe factor XII deficiency. However please note that the above theory is currently debated as other procoagulant factors have also been found to have a role.





**93. Ans. (b) Extrinsic pathway is activated by contact with negatively charged surfaces**

(Ref: Robbins 9th/pg 118-119; 8th/pg 119)

**Discussing options about blood coagulation one by one-**

- A. Factor X is part of both intrinsic and extrinsic pathways: True, refer Ans 50 above
- B. Extrinsic pathway is activated by contact with negatively charged surfaces: False, as negatively charged surfaces activate intrinsic pathway
- C. Intrinsic pathway can be activated in vitro: True
- D. Calcium is required in several steps of coagulation: True, Calcium acts as a cofactor for activation of coagulation factors

**94. Ans. (c) Should be measured within 2 hours**

(Ref: Dacie 11th/pg)

- **Ratio** of anticoagulant (**Trisodium citrate**): Blood = **1:9**<sup>Q</sup>
- Sample required: **Platelet poor plasma** (PPP)<sup>Q</sup>
- Storage: **Room temperature**<sup>Q</sup>
- Test should ideally be performed **within 2 hours**<sup>Q</sup> of sample collection
- **Prothrombin time (PT)** assay screens **extrinsic + common pathway**<sup>Q</sup> (factors VII, X, V, II, and fibrinogen).
- **Partial thromboplastin time (PTT)** assay screens **intrinsic pathway + common pathway**<sup>Q</sup> (factors XII, XI, IX, VIII, X, V, II, and fibrinogen).

**95. Ans. (b) Factor 7 deficiency** (Ref: Robbins 9th/pg 656)

Disease	PT	apTT
A. Von Willebrand disease	N	↑
B. Factor 7 deficiency	↑	N
C. Factor 8 deficiency	N	↑
D. Thrombin deficiency	↑	↑

**96. Ans. (c) PTT**

(Ref: Robbins 9th/pg 662-663)

**Best screening test for hemophilia (both A & B) is apTT, which is ↑.**

**97. Ans. (c) Vit. K deficiency** (Ref: Robbins 9th/pg 656)

Disease/ Condition	BT	PT	apTT
A. Thrombocytopenia	↑	N	N
B. DIC	↑	↑	↑
C. Vit. K deficiency	N	↑	↑ (late)
D. Aspirin toxicity	↑	N	N

**98. Ans. (d) Defect in common pathway**

(Ref: Robbins 9th/pg 656; 8th/pg 666)

Defect in common pathway leads to elevation of both PT & apTT.

**99. Ans. (c) Decreased fibrinogen**

(Ref: Wintrobe pg 684)

Though ↑ FDP is the most sensitive test for DIC bleeding occurs due to defective aggregation of platelets and decreased fibrinogen. (Remember aggregation also requires fibrinogen).

**100. Ans. (a) Obstetric complications**

(Ref: Harrison 18th ed/pg 1390)

The most common causes are bacterial sepsis, malignant disorders such as solid tumors or acute promyelocytic leukemia, and obstetric causes. DIC is diagnosed in almost one-half of pregnant women with abruptio placentae, or with amniotic fluid embolism

**101. Ans. (d) FDP level** (Ref: Robbins 9th/pg 663-664)

**102. Ans. (d) DIC associated with malignancy**

(Ref: Robbins 9th/pg 662; 8th/pg 672)

About use of Heparin in DIC:

- Low doses of continuous infusion heparin (5-10 U/kg per h) may be effective in patients with low-grade DIC associated with **solid tumor, acute promyelocytic leukemia, or in a setting with recognized thrombosis.**
- Heparin is also indicated for the **treatment of purpura fulminans** during the **surgical resection of giant hemangiomas** and **during removal of a dead fetus.**
- In **acute DIC**, the use of heparin is likely to **aggravate bleeding.**
- To date, the use of heparin in patients with severe DIC has no proven survival benefit.

**103. Ans. (c) Hyperfibrinogenemia**

(Ref: R 9th/pg 663-664)

**104. Ans. (a) ↑ fibrinogen level**

(Ref: Robbins 9th/pg 663-664)

**105. Ans. (c) Normal APTT**

(Ref: Robbins 9th/pg 663-664)

**106. Ans. (d) All of the above**

(Ref: Robbins 9th/pg 123)

Factor V Leiden mutation results in glutamine to arginine substitution at amino acid residue 506 that renders factor V resistant to cleavage and inactivation by protein C. As a result, an important antithrombotic counterregulatory pathway is lost and chances of thrombosis are highly raised.

**107. Ans. (b, c, d) b. Both PT and aPTT are increased in DIC; c. IV Ig is useful in ITP; d. Cryoprecipitate is useful in hemophilia A**



**108. Ans. (a) Factor 5 mutation**

Factor V Leiden is MC

**109. Ans. (d) Breast** (Ref: Robbins 9th/pg 663-664)

Thrombosis in breast carcinoma is least common among given options

**110. Ans. (a) Heparin** (Ref: Harrison 18th/pg 1403)

- Antithrombin, a serine protease inhibitor, regulates coagulation by inactivating thrombin and other procoagulant enzymes, including factors Xa, IXa, XIa, and XIIa.
- Once bound to Heparin, the natural anticoagulant effect of Antithrombin is potentiated, resulting in the accelerated binding and inactivation of serine proteases, and factor Xa and thrombin in particular.
- The inhibition of these factors affects the common pathway of coagulation, resulting in decreased formation of thrombin and fibrin.

**111. Ans. (b) Hypothermia**

(Ref: Robbins 9th/pg 663; 8th/pg 673)

**112. Ans. (a) Warfarin for life** (Ref: Harrison 19th/pg 2135)

- After the first thrombotic event, APS patients should be placed on **warfarin for life**, aiming to achieve an international normalized ratio (INR) ranging from **2.5 to 3.5**, alone or in combination with **80 mg of aspirin daily**.
- Pregnancy morbidity is prevented by a combination of **heparin with aspirin 80 mg daily**. **IV immunoglobulin (IVIg)** 400 mg/kg every day for 5 days may also prevent abortions, whereas glucocorticoids are ineffective

**113. Ans. (d) Priapism** (Ref: Robbins 9th/pg 123-124)

**Hypercoagulability due to defective factor V gene is called Leiden mutation**

- **Factor V Leiden**: glutamine → arginine substitution at amino acid residue 506 that renders **factor V resistant** to cleavage and inactivation by protein C.

**114. Ans. (a) Factor V** (Ref: Robbins 9th/pg 123; 8th/pg 123)

**115. Ans. (b) Anti-Prothrombin** (Ref: Robbins 9th/pg 124)

**116. Ans. (c) Depends on platelet activation**

(Ref: Robbins 9th/pg 122; 8th/pg 122)

**117. Ans. (d) Factor V** (Ref: Robbins 9th/pg 123-124)

**Leiden factor is factor V.**

**118. Ans. (c) Cerebral microcirculation**

(Ref: Robbins 9th/pg 663)

**Thrombomodulin is produced by all endothelial cells except those of cerebral microcirculation**

**119. Ans. (b) aPTT** (Ref: Dacie 11th/pg 472)

Therapeutic treatment with UFH is given by continuous intravenous infusion and is usually monitored using the APTT, which is repeated 6 h after every dose change.

**120. Ans. (a, b, c, d); a. Recurrent abortion; b. Fetal death; c. Both arterial and venous thrombosis; d. Prolonged aPTT**

(Ref: Robbins 9th/pg 124; 8th/pg 123)

**Autoantibody<sup>o</sup>-mediated acquired<sup>o</sup> thrombophilia** characterized by **recurrent<sup>o</sup> arterial or venous thrombosis<sup>o</sup>** and/or **pregnancy morbidity** in the presence of **autoantibodies against phospholipid (PL)-binding plasma proteins (β-2 GPI)<sup>o</sup>**

**APLA can result in ↑ phospholipid based assays like PT, aPTT, TT.**

**But isolated Increase in aPTT is most commonly seen**

**121. Ans. (d) Anti- beta 2 glycoprotein antibody**

(Ref: Robbins 9th/pg 124; 8th/pg 123)

**Anti- beta 2 glycoprotein antibody** is most frequently seen in Antiphospholipid Syndrome

**122. Ans. (a, c, e); a. Antiphospholipid antibodies; c. Hyperhomocysteinemia; e. Mutation in factor V gene**

(Ref: Wintrobe's 12th/pg 1465)

Both arterial and venous thrombosis occur in:

- |  |   |
|--|---|
| <ul style="list-style-type: none"> <li>• <b>APLA syndrome</b></li> <li>• <b>Hyperhomocystinemia</b></li> <li>• <b>Factor V Leiden</b></li> <li>• <b>DIC</b></li> </ul> | <ul style="list-style-type: none"> <li>• <b>Heparin induced Thrombocytopenia (HIT)</b></li> <li>• <b>(PNH)</b></li> <li>• <b>Polycythemia Vera (PCV)</b></li> <li>• <b>Dysfibrinogenemia</b></li> </ul> |
|--|---|

**123. Ans. (b) Leiden mutation**

(Ref: Robbins 9th/pg 123-124)

**124. Ans. (a) Protein C/Protein S deficiency**

(Ref: Wintrobe's 12th/pg 1490)

Warfarin skin necrosis

- Due to **warfarin-induced rapid reduction in protein C levels** in patients with a **pre-existing inherited protein C deficiency** that results in a **hypercoagulable state and thrombosis**.
- Not all heterozygous protein C-deficient patients receiving warfarin experience this complication, and not all patients with this complication have protein C deficiency.

**125. Ans. (c) Factor X** (Ref: Robbins 9th/pg 121; 8th/pg 116)

TFPI is a major anticoagulant, present in plasma and associated with vascular endothelium. TFPI binds to and neutralizes factor Xa to inhibit coagulation.

**126. Ans. (a) Russell viper venom time**

(Ref: R 9th/pg 123-124)

**Screening Laboratory Evaluation for Patients Suspected of Having a Hypercoagulable State**

- **Activated protein C resistance** (diluting patient plasma with factor V-deficient plasma)
- **Prothrombin G20210A mutation** testing by polymerase chain reaction
- Activity assays for **antithrombin, protein C, and protein S**
- Activated partial thromboplastin time, mixing studies, and **dilute Russell viper venom time**
- Fasting total plasma **homocysteine level**
- **Anticardiolipin &  $\beta_2$ -glycoprotein 1 antibody testing** by enzyme-linked immunosorbent assays
- **Factor VIII activity**

**127. Ans. (a) (PT patient/ PT control)ISI**

(Ref: Wintrobe's 12th/pg 1489)

**128. Ans. (a) Vascular injury**

(Ref: Robbins 9th/pg 663-664)

- DIC shows  $\uparrow$  CT and low Platelet counts
- DIC begins with factors that causes endothelial (vascular) injury (refer to pretext of this chapter)

**129. Ans. (a) PT INR**

(Ref: Wintrobe's 12th ed/pg 1489)

Patients with recurrent thrombosis should receive **long-term warfarin therapy** at a dosage to maintain an international normalized ratio (INR) value of **2.0 to 3.0**.

# Cardiovascular System and its Disorders

## Key Points

- » **Fatty streak is seen in Aortas of infants**, virtually all adolescents, even those without known risk factors
- » **Most common<sup>o</sup> vessel involved & Most common site<sup>o</sup> of atherosclerotic aneurysm is Lower abdominal aorta**
- » **Hyperplastic arteriolosclerosis is seen** in Malignant hypertension
- » **Hyaline arteriolosclerosis is seen in** Benign hypertension and Diabetes mellitus (DM)
- » Most important cause of true **aortic aneurysms are atherosclerosis**
- » **Most common cause of dissection is hypertension**
- » **Anti-endothelial cell antibodies are seen in Kawasaki's disease**
- » **Strawberry tongue is seen in Kawasaki disease**
- » **Microscopic hallmark** of HSP is the deposition of **IgA in the walls** of involved blood vessels.
- » **Most common vascular tumor: Capillary hemangioma**
- » **Time for reversible injury in heart is 30 mins**
- » **Intravenous drug abusers: Right side** of heart is affected
- » Dilated cardiomyopathy is **most common**
- » **HOCM is the leading cause of** unexplained left ventricular hypertrophy
- » **Native cardiac valve endocarditis-Staphylococcus aureus**
- » The most common cardiac tumor is the **secondaries or metastasis**
- » The most common primary cardiac tumor in the adults is **the myxoma**
- » The most common cardiac tumor in the children is the **rhabdomyoma**.

## Key Recent Updates

- » Most common vascular tumor is hemangioma
- » Aschoff's nodule is hallmark of Rheumatic heart disease.





## BLOOD VESSELS

### BLOOD VESSELS: OVERVIEW

The basic histological layers of the blood vessels (particularly arteries) are:



- Tunica intima (Innermost layer)-single layer of **endothelial cells**<sup>Q</sup>
- Internal elastic lamina
- Tunica media (Middle layer)- constituents depends on the type of artery as explained below
- External elastic lamina

- Tunica adventitia (Outermost layer)-loose connective tissue containing nerve fibers and the **vasa vasorum**<sup>Q</sup>





#### High Yield Facts

- All vessels **except capillaries**<sup>Q</sup> have three-layered architecture consisting of an **intima, media, and adventitia**.
- There are 3 types of capillaries-**continuous, fenestrated & sinusoidal**



#### Arteries are divided into three types

	Large or Elastic Arteries	Medium or Muscular Arteries	Small Arteries (0.2 mm) Arterioles (20-100 μm)
Type of vessel	Aorta & major branches <b>Pulmonary arteries</b>	Smaller branches of aorta (coronary & renal)	• Within tissues and organs.
Characteristic feature	Tunica media is <b>rich in elastin fibers</b> <sup>Q</sup>	Tunica media is rich in <b>smooth muscle cells</b> <sup>Q</sup>	<b>Resistance vessels</b> is other name of arterioles




### High Yield Facts

#### Fibromuscular dysplasia


- Cause of **renovascular hypertension** in **young females<sup>Q</sup>**
- No association with oral contraceptives or increased estrogen expression.<sup>Q</sup>**
- On angiography, the vessels have a “**string of beads<sup>Q</sup>**” appearance (due to **markedly attenuated media<sup>Q</sup>**)

## VESSEL SCLEROSIS

Denotes **arterial wall thickening** and **loss of elasticity**







### High Yield Facts

**Mönckeberg's arteriosclerosis, or Mönckeberg's sclerosis, also called medial calcific sclerosis**

- Example of **dystrophic calcification**<sup>Q</sup>
- Prevalence **increases with age**<sup>Q</sup>
- More frequent in **diabetes mellitus<sup>Q</sup>, chronic kidney disease, and systemic lupus erythematosus**
- **Non obstructive lesion**<sup>Q</sup>

**Hyperplastic arteriosclerosis is pathological change seen in malignant hypertension**

- Arterioles of all organs of body are affected but **avored sites** are:  
**Kidney<sup>Q</sup>, Small intestine<sup>Q</sup>, Gall bladder<sup>Q</sup>, Peri pancreatic fat<sup>Q</sup>, Periadrenal fat<sup>Q</sup>**

Atherosclerosis plaque is **present within the intima<sup>Q</sup>**, has a **core of lipid (cholesterol & its esters) & a covering of fibrous cap**.

### Histopathology of Atherosclerotic Plaque

- **Fibrous cap<sup>Q</sup>** – Consists of **smooth muscle cells, collagen**.
- **'Shoulder'<sup>Q</sup>** – Cellular area around cap having **macrophages, smooth muscle cells and T lymphocytes**.
- **Necrotic core**–Debris of **dead cells, foam cells, lipid (cholesterol, cholesterol clefts.<sup>Q</sup>) fibrin and variably organized thrombus**.

### Major Risk Factors for Atherosclerosis

Modifiable	Nonmodifiable (Constitutional)
• Hyperlipidemia	• Genetic abnormalities
• Hypertension	• Family history
• Cigarette smoking	• Increasing age

Contd...

- Diabetes
- Inflammation
- Physical inactivity

- Male gender
- Stress (Type A personality)


### Additional Risk Factors for Atherosclerosis

<b>a. Inflammation:</b> <ul style="list-style-type: none"> <li>■ Present during <b>all stages of atherosclerosis<sup>Q</sup></b>, linked with <b>atherosclerotic plaque formation and rupture<sup>Q</sup></b></li> <li>■ Marker-CRP</li> </ul>	<b>d. Metabolic syndrome:</b> <ul style="list-style-type: none"> <li>■ Characterized by <b>insulin resistance, hypertension, dyslipidemia (↑ LDL &amp; ↓ HDL), hypercoagulability, proinflammatory state<sup>Q</sup></b></li> </ul>
<b>b. Hyperhomocysteinemia</b>	<b>e. Factors Affecting Hemostasis:</b> <ul style="list-style-type: none"> <li>■ Elevated plasminogen activator inhibitor 1 and thrombinase</li> </ul>
<b>c. Lipoprotein a [Lp(a)]:</b> <ul style="list-style-type: none"> <li>■ Altered form of LDL that contains the <b>apolipoprotein B-100</b> portion of LDL linked to apolipoprotein A (apo A)</li> <li>■ Structural similarity to plasminogen.<sup>Q</sup></li> </ul>	<b>f. Infection:</b> <ul style="list-style-type: none"> <li>■ <b>Herpesvirus, Cytomegalovirus &amp; Chlamydomphila pneumoni</b></li> </ul>



### High Yield Facts

- **C-reactive protein (CRP): Most sensitive marker of inflammation** that correlate with **ischemic heart disease risk<sup>Q</sup>**
- **Lipoprotein a [Lp(a)]** is associated with **coronary & cerebrovascular disease risk, independent of total cholesterol or LDL levels<sup>Q</sup>**



## High Yield Facts

- **Endothelial injury and dysfunction** is the cornerstone of the **response-to-injury hypothesis**<sup>Q</sup>
- **Non-denuding endothelial dysfunction underlies most human atherosclerosis**<sup>Q</sup>
- Most important causes of endothelial dysfunction are hemodynamic disturbances<sup>Q</sup> & hypercholesterolemia<sup>Q</sup>
- Foam cells are **lipid laden smooth muscles cells/tissue macrophages or Blood monocytes**<sup>Q</sup>
- In foam cells **oxidized LDL** is ingested by the **scavenger receptors**<sup>Q</sup> present on macrophages & smooth muscle cells<sup>Q</sup>

## Morphology of Atherosclerosis

- **Fatty streak**
  - **Earliest lesion** of atherosclerosis is composed of **lipid filled foam cells**.<sup>Q</sup>
  - Begin as yellow flat spots **less than 1 mm**<sup>Q</sup>
  - **Donot**<sup>Q</sup> cause flow disturbances.
  - **Seen in Aortas of infants, virtually all adolescents, even those without known risk factors**.<sup>Q</sup>
- **Atherosclerotic Plaque**
  - Characterized by **intimal thickening and lipid accumulation**<sup>Q</sup>, which together form plaques
  - White-yellow and **encroach on the lumen**<sup>Q</sup> of the artery

In descending order<sup>Q</sup>, the most extensively involved vessels in Atherosclerosis are:

<b>Lower abdominal aorta</b>	<b>Most common</b> <sup>Q</sup> vessel involved & Most common <b>site</b> <sup>Q</sup> of atherosclerotic aneurysm
<b>Coronary arteries</b>	<b>Left Anterior Descending</b> is Most common <sup>Q</sup> coronary artery involved → causes MI

Contd...

<b>Popliteal arteries</b>	<b>Most common peripheral vessel</b> <sup>Q</sup> with aneurysm formation → ischemic gangrene of lower limbs
<b>Internal carotid arteries</b>	Cause stroke
<b>Vessels of circle of Willis</b>	Cause stroke

- **Vessels spared in Atherosclerosis** are upper extremities vessels,<sup>Q</sup> mesenteric & renal arteries except at their ostia<sup>Q</sup>

## Type of Plaques

Characteristics	Stable Plaques	Vulnerable/ Unstable Plaques
<b>Fibrous cap</b>	Dense	Thin
<b>Lipid cores</b>	Minimal	Large
<b>Inflammation</b>	Minimal	Marked
<b>Clinical Manifestations</b>	Due to <b>chronic ischemia</b>	<b>Fatal ischemic complications</b>

**Acute plaque change** is described as:

- Rupture, ulceration, or erosion, Hemorrhage into a plaque, Atheroembolism, Aneurysm formation


## ANEURYSM

- **Definition:**
  - Localized abnormal dilation of a blood vessel or the wall of the heart
- **Types:**
  - **True aneurysm**
  - **False/Pseudoaneurysm**
- **Characteristics:**
  - Involves **attenuated intact arterial wall** or **thinned ventricular wall** of the heart
  - Vascular wall defect leading to **extravascular hematoma that freely** communicates with intravascular space<sup>Q</sup>
- **Associated Conditions:**
  - Atherosclerosis<sup>Q</sup>
  - Syphilis<sup>Q</sup>
  - Congenital vascular aneurysms<sup>Q</sup>
  - Post MI ventricular aneurysms.<sup>Q</sup>
  - Post MI rupture contained by pericardial adhesion<sup>Q</sup>
  - Leakage at the suture junction of vascular anastomosis<sup>Q</sup>
- **Etiology:**
  - Most important cause of true **aortic aneurysms** are **atherosclerosis**<sup>Q</sup> > **hypertension**

The inherited causes of aneurysm are: M-L-E

<b>Marfan syndrome</b>	Defective synthesis of the protein <b>fibrillin</b> <sup>Q</sup>
<b>Loeys Dietz syndrome</b>	Defect in <b>elastin &amp; collagen type I &amp; III</b> due to mutation in <b>TGF-B receptor</b> <sup>Q</sup>
<b>Ehlers-Danlos syndrome</b>	Defect in <b>collagen type III</b> <sup>Q</sup>






**Latest Update**


Inflammatory aneurysms	Immunoglobulin G4 (IgG4)-related disease
<p><b>Younger patients<sup>Q</sup></b>; Present with back pain</p> <ul style="list-style-type: none"> <li>Elevated inflammatory markers (e.g., elevation of <b>C-reactive protein<sup>Q</sup></b>)</li> <li><b>Abundant lymphoplasmacytic inflammation<sup>Q</sup></b> with many macrophages &amp; sometimes giant cells</li> <li>Associated with <b>dense periaortic scarring that can extend into the anterior retroperitoneum<sup>Q</sup></b></li> <li>Cause: <b>Localized<sup>Q</sup></b> immune response to the abdominal aortic wall</li> </ul>	<ul style="list-style-type: none"> <li>High plasma levels of IgG4</li> <li><b>Tissue fibrosis associated with frequent infiltrating IgG4-expressing plasma cells.<sup>Q</sup></b></li> <li>Affects <b>pancreas, biliary system, and salivary gland.<sup>Q</sup></b></li> <li><b>Cause aortitis and periaortitis → aneurysms.<sup>Q</sup></b></li> <li>Responds well to <b>steroid therapy.</b></li> </ul>

## AORTIC DISSECTION

- **Definiton:**
  - Blood separates the laminar planes of the **media** to form a blood-filled channel within the aortic wall
- **Epidemiology:**
  - 40-60 years; at younger age in Marfan's syndrome
- **Etiology:**
  - **Most common cause of dissection is hypertension<sup>Q</sup>**
- **Types depending on level of aorta affected:**



- **Histology:**
  - **The most frequent preexisting histologically detectable lesion is cystic medial degeneration<sup>Q</sup>**
  - **Inflammation is characteristically absent.**



Aortic dissection: Dissecting aorta with two lumens: true lumen in the right, while false lumen in the left\*



## High Yield Facts



- **Double-barreled aorta**-Formed when the dissecting hematoma re-enters the lumen of the aorta through a **second distal intimal tear<sup>Q</sup>**, creating a new false vascular channel
- **Chronic dissection**: When false channels are **endothelialized<sup>Q</sup>**
- **Cystic medial necrosis (CMN)** involves large arteries, in particular the aorta, <sup>Q</sup>
- In CMN, there is accumulation of **basophilic ground substance in the media<sup>Q</sup>** with cyst-like lesions


### Diseases causing CMN

- Marfan's syndrome<sup>Q</sup>, Chronic aortic dissection<sup>Q</sup>, Bicuspid aortic valve<sup>Q</sup>, Scurvy<sup>Q</sup>, Aortic aneurysm<sup>Q</sup>, Atherosclerotic disease<sup>Q</sup>, Hypertension<sup>Q</sup>, Ehler danlos syndrome (type IV)<sup>Q</sup>


## VASCULITIS

Vasculitis is the **inflammation of vessel wall**.

### Classification



### Antibodies in Vasculitis






## Large Vessel Vasculitis

### Giant Cell (Temporal) Arteritis/Cranial Arteritis

- **Epidemiology:**
  - Most common type of vasculitis in adults (usually >50 years)
- **Pathophysiology:**
  - T-cell mediated immune response<sup>Q</sup> against vessel wall antigens<sup>Q</sup> that releases proinflammatory cytokines (particularly TNF<sup>Q</sup>)
- **Clinical features:**
  - Fever, anemia (normocytic normochromic)<sup>Q</sup>, high ESR, and headaches (most common)
  - Most specific symptom- jaw claudication<sup>Q</sup>
  - Sudden blindness (due to involvement of ophthalmic arteries)
  - Associated with polymyalgia rheumatica<sup>Q</sup>
- **Arteries affected:**
  - Most commonly: superficial temporal arteries<sup>Q</sup>, vertebral and ophthalmic arteries.
  - Lesions also occur in other arteries, including the aorta (giant cell aortitis<sup>Q</sup>)
- **Diagnosis:**
  - Biopsy and histological confirmation of temporal artery is the investigation of choice.
  - Granulomatous inflammation with giant cells & fragmentation of internal elastic lamina.<sup>Q</sup>
- **Treatment:**
  - Corticosteroids & anti-TNF therapies are the drug of choice




Granulomatous inflammation with giant cells

- **Clinical features:**
  - Ocular disturbances & marked **weakening of the pulses in the upper extremities** (hence the name **pulseless disease<sup>Q</sup>**)
- **Vessels involved:**
  - Most common vessel involved - subclavian artery<sup>Q</sup>
  - Least common vessel involved- coronary vessels<sup>Q</sup>
  - Aortic arch & its branches involved, so also called **Aortoarteritis or Aortic Arch syndrome<sup>Q</sup>**
  - Pulmonary artery involvement in half the cases can cause pulmonary hypertension<sup>Q</sup>

## Medium Vessel Vasculitis

### Polyarteritis Nodosa (PAN)

- Systemic vasculitis of small- or medium-sized muscular arteries<sup>Q</sup>
- In descending order-kidney, heart, liver and GIT vessels are involved. Typically **pulmonary circulation is spared.<sup>Q</sup>**
- Association with **Hepatitis B antigen<sup>Q</sup>** in serum (30% patients)-Deposits contain HBsAg-HBsAb complex (immune complex)<sup>Q</sup>
- Characterized by **segmental transmural necrotizing inflammation** accompanied by **fibrinoid necrosis<sup>Q</sup>**
- Characteristically, **all stages of activity<sup>Q</sup>** coexist in different vessels or within the same vessel
- **Glomerulonephritis<sup>Q</sup>** is NOT seen.
- Most common cause of mortality: **Renal involvement<sup>Q</sup>**



Fibrinoid necrosis: accumulation of amorphous, proteinaceous material in the tissue matrix

### Takayasu's Arteritis

- **Epidemiology:**
  - Seen in **adult female<sup>Q</sup>** <50 years of age.
- **Characteristic:**
  - Granulomatous vasculitis of medium and larger arteries<sup>Q</sup>

### Kawasaki's Disease (Infantile Polyarteritis, Mucocutaneous Lymph Node Syndrome)

- **Acute febrile<sup>Q</sup>**, usually self-limited illness of **infancy and childhood<sup>Q</sup>** (80% patients are ≤ 4 years)
- It is associated with an **arteritis** affecting **large to medium-sized, and even small vessels**





- **Diagnostic criteria:**
  - **Anti-endothelial** cell antibodies<sup>Q</sup>.
  - Leading cause of **acquired heart disease**<sup>Q</sup> in children.
  - **Coronary artery** involvement<sup>Q</sup> → aneurysms that rupture or thrombose → **Acute Myocardial Infarction**<sup>Q</sup>

Fever for > 5 days  
+  
4 from conjunctivitis, adenopathy, rash, erythema of hands/feet & mucosa involvement. (CARE – M)



Kawasaki's disease: Shows typical strawberry tongue

## Mnemonic

### Kawasaki's disease (CREAM)


- **C**-Conjunctivitis (non-exudative); non purulent conjunctivitis
- **R**-Rash (Polymorphous non-vesicular)
- **E**-Edema (or erythema of hands or feet)
- **A**-Adenopathy (cervical, often unilateral and non suppurative)
- **M**-Mucosal involvement (erythema or fissures or crusting at times referred as **strawberry tongue**<sup>Q</sup>)
- **F**-Fever (most important constitutional symptom<sup>Q</sup>)

## Small Vessel Vasculitis

### Microscopic Polyangiitis/Leukocytoclastic Vasculitis/Hypersensitivity Vasculitis

- **Necrotizing vasculitis**<sup>Q</sup> that generally affects capillaries, as well as small arterioles and venules
- All lesions are of **same age**<sup>Q</sup> in any given patient and are **distributed more widely**<sup>Q</sup>.
- **Necrotizing glomerulonephritis** (90% of patients) and **pulmonary capillaritis** are **common**.
- Typically **spare medium-sized and larger arteries**<sup>Q</sup>; consequently, **infarcts are uncommon**<sup>Q</sup>
- Granulomatous inflammation is **absent**<sup>Q</sup>
- **P-ANCA**<sup>Q</sup> is present in majority of the patients.
- **Leukocytoclastic vasculitis**<sup>Q</sup> -vessel wall infiltrated with intact and apoptotic neutrophils<sup>Q</sup>

- **Clinical features**-palpable cutaneous purpura<sup>Q</sup>, hemoptysis, hematuria and proteinuria



## High Yield Facts

### Hypersensitivity Vasculitis

- Seen most commonly in **Post capillary venules**<sup>Q</sup>
- Mic- **leukocytoclastic vasculitis**<sup>Q</sup>

## Churg-Strauss Syndrome

### (Allergic Granulomatosis and Angiitis)

- Small-vessel necrotizing vasculitis classically associated with **asthma, allergic rhinitis**<sup>Q</sup>
- **Lung infiltrates, peripheral hypereosinophilia, and extra-vascular necrotizing granulomata**<sup>Q</sup>
- **P-ANCA**<sup>Q</sup> present in less than 50% of patients
- Multisystem diseases with cutaneous involvement (**palpable purpura**<sup>Q</sup>), gastrointestinal tract bleeding, and renal disease (primarily as **focal and segmental glomerulosclerosis**)<sup>Q</sup>
- The heart is involved in 60% of patients and accounts for almost **half of the deaths** in the syndrome<sup>Q</sup>
- New name suggested is **EGPA**-Eosinophilic granulomatosis with polyangiitis

## Granulomatosis with Polyangiitis

- Previously called as **Wegener's granulomas**

**Necrotizing vasculitis** which is characterized by **triad of**

- Acute necrotizing granulomas of either **upper (more commonly) or lower respiratory tract**<sup>Q</sup> or both.
- **Necrotizing or granulomatous vasculitis**<sup>Q</sup> affecting small to medium-sized vessels (e.g., capillaries, venules, arterioles, and arteries), most prominent in the lungs and upper airways.
- **Focal necrotizing, often crescentic, glomerulonephritis**<sup>Q</sup>





Wegener's granulomatosis: Strawberry gums

**High Yield Facts**

- Limited Wegener's granulomatosis<sup>Q</sup> is characterized by only respiratory tract involvement<sup>Q</sup>
- Granulomas are seen in Wegener's in respiratory tract but not in renal parenchyma<sup>Q</sup>
- Patients with classical PAN<sup>Q</sup> are ANCA negative
- PAN is the commonest cause of mononeuritis multiplex.<sup>Q</sup>


**Henoch-Schonlein Purpura (HSP)/Anaphylactoid Purpura/Purpura Rheumatica**

- **Definition:**
  - Systemic vasculitis syndrome involving small size vessels.<sup>Q</sup>
- **Epidemiology:**
  - Most common small vessel vasculitis in children
  - Usually follows an upper respiratory tract infection<sup>Q</sup> (in 2/3<sup>rd</sup> cases)
- **Pathophysiology:**
  - Vasculitis is caused by immune complex deposition
  - Most common antibody seen in these immune complexes is IgA (elevated)
- **Clinical features (tetrad)**
  - Palpable purpura (Most common sites<sup>Q</sup>-buttocks and exterior surface of legs and arms),
  - Arthritis or arthralgia
  - Renal involvement (proteinuria,<sup>Q</sup> Microscopic hematuria with RBC casts in urine),
  - Colicky Abdominal pain and/or upper GI bleed
- **Diagnosis:**
  - Platelet count is normal or elevated (as Purpura does not occur due to a low platelet count but due to vasculitis)
  - Serum complement levels are normal
  - Microscopic hallmark of HSP is the deposition of IgA in the walls<sup>Q</sup> of involved blood vessels.
  - Morphology of vasculitis-leukocytoclastic vasculitis<sup>Q</sup>

- **Treatment:**
  - Administration of glucocorticoids (Prednisolone)
- **Prognosis:**
  - Excellent and disease is self limiting.
  - Incidence of renal involvement in adults is 45-85%, while it is 20-50% in children



Palpable Purpura



Leukocytoclastic vasculitis

**Behçet's Disease**

- Small-to-medium vessel neutrophilic vasculitis<sup>Q</sup>
- Clinical triad of recurrent oral aphthous ulcers, genital ulcers, and uveitis<sup>Q</sup>
- Sin qua non for diagnosis- Oral ulceration (hallmark)<sup>Q</sup>
- Positive pathergy test<sup>Q</sup>- if a papule or pustule >2 mm, 24-48 hours after needle prick, depth of 5 mm
- HLA-B51<sup>Q</sup> associated
- Microscopic findings are nonspecific- neutrophils are seen infiltrating vessels walls
- Immunosuppression with steroids or TNF-antagonist therapies is generally effective.



Image showing pathergy test


### Buerger's Disease (Thromboangitis Obliterans)

- **Segmental, thrombosing**, acute and chronic inflammation<sup>Q</sup> of **medium-sized and small<sup>Q</sup> arteries**,
- **Tibial and radial arteries<sup>Q</sup>**, with secondary extension into the **veins and nerves** of the extremities.
- Seen in **heavy cigarette smokers<sup>Q</sup>**; Onset is **before 35 years of age<sup>Q</sup>**
- Associated with **HLA B5 and HLA A9<sup>Q</sup>**.
- Associated with **hypersensitivity to intradermal injections of tobacco extracts<sup>Q</sup>**.
- Mic-segmental thrombosing vasculitis extending into **contiguous veins & nerves, encasing all in fibrous tissue<sup>Q</sup>**.
- Thrombus contains **microabscess** with **granulomatous inflammation<sup>Q</sup>**.

#### Clinically

- |   |  |   |
|---|--|---|
| • <b>Intermittent claudication<sup>Q</sup></b> -leg pain induced by exercise that is relieved on rest | • <b>Instep claudication<sup>Q</sup></b> -Instep foot pain induced by exercise | • Superficial nodular phlebitis and cold induced Raynaud's phenomenon |
|---|--|---|

- Treatment includes smoking cessation.



Thrombus containing abscess

Buerger's disease



### High Yield Facts

#### Henoch-Schonlein Purpura (HSP):


- HSP may present in a clinically identical fashion to **hypersensitivity vasculitis<sup>Q</sup>**
- **Biopsy with direct immunofluorescence<sup>Q</sup>** (IgA in HSP) is needed to distinguish the two.

#### Rheumatoid vasculitis:

- Occurs in the setting of long-standing, severe rheumatoid arthritis
- Affects small- and medium-sized arteries → Causes visceral infarction
- May also cause a clinically significant aortitis

### Infectious Vasculitis

Caused by vascular infections that can weaken arterial walls and culminate in mycotic aneurysms




### Mnemonic

#### Granulomatous vasculitis

WTC has TB  
W – **W**egener's  
T – **T**akayasu

C – **C**hurg- Strauss  
T – **T**emporal arteritis  
B – **B**uerger's disease.



Granulomatous vasculitis: we can see epithelioid cells and giant cells s/o granulomatous vasculitis

### VEINS AND LYMPHATICS

Varicose veins and phlebothrombosis/thrombophlebitis together account for at least 90% of clinical venous disease.






## Thrombophlebitis and Phlebothrombosis

- Involvement of **deep leg veins**<sup>Q</sup> accounts for more than 90% of cases.

- **Prolonged immobilization**<sup>Q</sup>-most important risk factor for deep venous thrombosis (DVT) in the lower extremities.
- **Systemic hypercoagulability**<sup>Q</sup>: often also plays a role in potentiating thrombophlebitis.


## VASCULAR TUMORS

Classification of Vascular Tumors and Tumor-Like Conditions



## Benign Tumors


### Hemangioma



A proliferation of vascular channels filled with red blood cells, consistent with a hemangioma



## High Yield Facts



### Port-wine stain or nevus flammeus

- It is the **most common form of Vascular ectasias**; it persists throughout life<sup>Q</sup>
- Caused by a somatic activating c.548G → A mutation in the **GNAQ**<sup>Q</sup> gene
- Part of Sturge-Weber syndrome<sup>Q</sup> or Klippel-Trénaunay-Weber syndrome<sup>Q</sup>

### Capillary Hemangioma


**Most common** vascular tumor, occurs in skin, mucus membrane & viscera.



Strawberry <sup>Q</sup> or Juvenile Hemangioma	Pyogenic Granulomas
<ul style="list-style-type: none"> <li>• Very common</li> <li>• Occurs in <b>new borns<sup>Q</sup></b></li> <li>• <b>Completely regress<sup>Q</sup></b> by age 7</li> </ul>	<p>Red pedunculated lesions on skin, gingival, or oral mucosa (resembling <b>granulation tissue<sup>Q</sup></b>)</p> <p><b>Pregnancy tumor</b> (granuloma gravidarum)</p> <ul style="list-style-type: none"> <li>• Occurs in &lt; 1% of patients; <b>Usually regresses<sup>Q</sup></b> after delivery</li> <li>• Pyogenic granuloma in the gingiva of pregnant women<sup>Q</sup></li> </ul>



Strawberry hemangioma or capillary hemangioma




Capillary lobular hemangioma

### Cavernous Hemangiomas


- Are more infiltrative, frequently involve deep structures
- **Do not<sup>Q</sup>** spontaneously regress.
- Intravascular thrombosis and associated dystrophic calcification are **common<sup>Q</sup>**.
- Component of **von Hippel-Lindau disease<sup>Q</sup>**, with vascular lesions in cerebellum, brain stem, retina, pancreas, and liver.

### Lymphangioma



### Glomus Tumor (Glomangioma)

- **Benign tumor<sup>Q</sup>** arising from the smooth muscle cells of the glomus body
- Most commonly present in the **distal portion of the digits (under fingernails).<sup>Q</sup>**
- **Histologically:** branching vascular channels & stroma containing nests /aggregates of **glomus cells** around vessels.
- **Excision is curative<sup>Q</sup>**



### Bacillary Angiomatosis

- **Vascular proliferation in immunocompromised hosts<sup>Q</sup>**
- Caused by opportunistic **Gram-negative bacilli of the Bartonella family**.
- Lesions can involve the **skin, bone, brain, and other organs**
- **Microscopy:** capillary proliferation with **epithelioid endothelial cells<sup>Q</sup>** exhibiting nuclear atypia and mitoses
- Stromal neutrophils, nuclear dust, and the causal bacteria can be identified

### Intermediate/Borderline Tumors

#### Kaposi Sarcoma (KS)

- It is caused by KS Herpes virus or **Human herpes virus 8 (HHV8)<sup>Q</sup>**





- Characterized by proliferation of **spindle cells**<sup>Q</sup> which are of **vascular origin**<sup>Q</sup> and has the following 4 forms:

Type of Kaposi Sarcoma	Age	Association with HIV	Organs Affected
<b>Classic KS</b>	Elderly	Absent	Skin
<b>African/Endemic KS</b>	<40 yrs	Absent	<b>Lymph nodes ± Viscera</b>
<b>Transplant associated/ Immunosuppression-associated KS</b>	Any	Absent	Lymph nodes, mucosa & visceral organs
<b>Epidemic/AIDS associated KS</b>	Any	Present; AIDS-defining illness	<b>Lymph nodes &amp; viscera involved</b>



### High Yield Facts

- Most common** kaposi tumor in central Africa- KS (Endemic African KS in combination with AIDS-associated KS)
- Most common immunoblastic lymphoma (DLBCL) followed by AIDS associated (epidemic) KS**<sup>Q</sup>
- Most common cause of death** in KS- **opportunistic infections**<sup>Q</sup> rather than KS

### Epithelioid Hemangioendothelioma

- Vascular tumor of adults occurring around medium and large-sized veins
- 20 % of these** can metastasize; cured by excision



**Malignant Tumors**

**Angiosarcoma**

- Malignant endothelial cell neoplasm **most commonly**<sup>Q</sup> seen in **skin**,<sup>Q</sup> **soft tissue**,<sup>Q</sup> **breast**<sup>Q</sup> and **liver**<sup>Q</sup>
- Hepatic angiosarcoma is associated with **arsenic**, **thorotrast** (a radioactive contrast) & **polyvinyl chloride (PVC)**<sup>Q</sup>
- Other important risk factor is **radiation**<sup>Q</sup>
- Endothelial cell origin is demonstrated by staining for CD31, CD34 or VWF. (Highly specific) FLI/ERG
- Locally invasive; readily metastasize

**Hemangiopericytoma**

- Tumor derived from **pericytes**<sup>Q</sup>- **perivascular cells** that wrap around blood capillaries
- These tumors most commonly arise from **pelvic retroperitoneum**<sup>Q</sup> or the **limbs**<sup>Q</sup> (particularly thighs).
- Capillaries are arranged in '**fish-hook pattern**';<sup>Q</sup> **seen best with silver stains**<sup>Q</sup>

Angiosarcoma: Irregularly shaped vascular lumina lined by highly atypical endothelial cells S/o angiosarcoma

Hemangiopericytoma showing staghorn (fishhook pattern)



### High Yield Facts

- Lymphangiosarcoma** arises from dilated lymphatic vessels;<sup>Q</sup> most common site: post mastectomy arm<sup>Q</sup>
- Rouget cells** —other name of **pericytes**<sup>Q</sup>
- Pericytic (perivascular) tumors**-Glomus tumor, Hemangiopericytoma, Myopericytoma & PEComa(Perivascular epithelioid cell tumor)



## CARDIOVASCULAR SYSTEM

### HEART

#### Valves

- The four cardiac valves—tricuspid, pulmonary, mitral, and aortic—maintain unidirectional blood flow.
- Cardiac valves:** Lined by endothelium and have similar, trilayered architecture:
  - A dense collagenous core (fibrosa) at the outflow surface
  - A central core of loose connective tissue (spongiosa)
  - A layer rich in elastin on the inflow surface
- Pathologic changes of valves are largely of three types:
  - Damage to collagen that weakens the leaflets, exemplified by mitral valve prolapse
  - Nodular calcification beginning in interstitial cells, as in calcific aortic stenosis
  - Fibrotic thickening, the key feature in rheumatic heart disease

R<sup>9th</sup>

#### Latest Update

- Cardiac stem cells are present **within the myocardium**<sup>Q</sup>, they are the **greatest in neonates**, and constitute up to **5% to 10% of normal atrial cellularity**<sup>Q</sup> and **1 in 100,000 cells**<sup>Q</sup> in a normal ventricle<sup>Q</sup>.

#### Effects of Aging on Heart


- Epicardial fat increases
- Basophilic degeneration** within cardiac myocytes
- Intracellular lipofuscin deposits**

- Sigmoid septum**<sup>Q</sup> bulging of the basal ventricular septum into the left ventricular outflow tract
- Lambl's excrescences**<sup>Q</sup> small filiform processes on closure lines<sup>Q</sup> of aortic mitral valves: due to **organization of small thrombi**<sup>Q</sup>
- Aorta becomes progressively **stiffer**-due to accumulation of **atherosclerotic plaque**
- Myocytes decrease in number, and deposition of **extracellular amyloid** (most commonly due to poorly catabolized transthyretin<sup>Q</sup>)

#### Cardiac Hypertrophy

Cardiac myocytes which cannot undergo hyperplasia in response to stress, so the only adaptation seen in a cardiac muscle can be hypertrophy<sup>Q</sup> which can be of the following types:


Types	Concentric hypertrophy	Eccentric hypertrophy
Pathophysiology	Pressure overload	Volume overload
Definition	Deposition of the sarcomeres in parallel <sup>Q</sup> to the long axis of the cells	Dilatation with increased ventricular diameter <sup>Q</sup>
Cause	<ul style="list-style-type: none"> <li>Aortic stenosis<sup>Q</sup></li> <li>Hypertension<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Valvular regurgitation<sup>Q</sup> (mitral/aortic)</li> <li>Thyrotoxicosis<sup>Q</sup></li> <li>Severe anemia<sup>Q</sup></li> </ul>







## Morphologic Findings in Heart failure



## ISCHEMIC HEART DISEASE

- Imbalance between **perfusion & demand** of heart for oxygenated blood.
- IHD can present as one or more of the following clinical syndromes:
  - Angina pectoris**-where ischemia is **not severe** to cause infarction, but the **symptoms show infarction risk**<sup>Q</sup>
  - Myocardial infarction (MI)** where ischemia causes frank cardiac necrosis
  - Chronic IHD with heart failure**
  - Sudden cardiac death (SCD)**



### High Yield Facts

- No hypertrophy or dilation of left ventricle**-In cases of failure caused by **mitral valve stenosis or unusual restrictive cardiomyopathies**<sup>Q</sup>
- Most common**<sup>Q</sup> underlying etiology of LVF-**Hypertension**<sup>Q</sup>
- Flash pulmonary edema**,<sup>Q</sup> **Rapid onset pulmonary edema**<sup>Q</sup> because the left ventricle cannot expand normally
- Most common cause of ischemic heart disease**<sup>Q</sup>- Atherosclerotic narrowing resulting in coronary arterial obstruction
- The level of cardiac markers remain unchanged in stable angina**<sup>Q</sup>

## Myocardial Infarction (MI)

Result of acute plaque change that induces an abrupt thrombotic occlusion, resulting in myocardial necrosis.

### Myocardial Infarction

Transmural Infarction	Subendocardial Infarction (Nontransmural)	Multifocal Microinfarction	
<ul style="list-style-type: none"> <li><b>Epicardial vessel occluded</b></li> <li><b>Full thickness</b><sup>Q</sup> of the ventricular wall involved</li> <li>Associated with a combination of chronic coronary atherosclerosis, acute plaque change and superimposed thrombosis</li> </ul>	<ul style="list-style-type: none"> <li><b>Inner 1/3<sup>rd</sup> of ventricular wall</b><sup>Q</sup></li> <li>Subendocardium-least <b>perfused</b><sup>Q</sup> region of myocardium: <b>most</b><sup>Q</sup> <b>vulnerable</b> to any reduction in coronary flow (<b>hypotension / shock</b>)<sup>Q</sup></li> </ul>	<p>Occur in the setting of <b>microembolization, vasculitis</b><sup>Q</sup>, or <b>vascular spasm</b><sup>Q</sup></p> <p><b>Outcome:</b></p> <ul style="list-style-type: none"> <li><b>Sudden cardiac death</b><sup>Q</sup> (usually caused by a fatal arrhythmia)</li> <li><b>Ischemic dilated cardiomyopathy</b><sup>Q</sup></li> </ul>	<p>Transmural infarction: Full thickness of the ventricular wall involved</p>



## Myocardial Response<sup>Q</sup>

Feature	Time
• Onset of <b>ATP depletion</b>	Seconds <sup>Q</sup>
• <b>Loss of contractility</b> <sup>Q</sup>	<2 min <sup>Q</sup>
• ATP reduced to <b>50%</b> of normal <sup>Q</sup>	10 min <sup>Q</sup>
• ATP reduced to <b>10%</b> of normal <sup>Q</sup>	40 min <sup>Q</sup>
• <b>Microvascular injury</b> <sup>Q</sup>	>1 hr <sup>Q</sup>


### Latest Update

**TAKOTSUBO** cardiomyopathy <sup>Q</sup>: (also called “**broken heart syndrome**” because of the association with emotional distress) - **ischemic dilated cardiomyopathy**<sup>Q</sup>

### What are the symptoms of takotsubo cardiomyopathy?

The name "takotsubo syndrome" comes from the Japanese word "takotsubo"—octopus trap, because the left ventricle takes on a shape resembling an octopus trap.

- Difficulty in breathing
- Nausea
- Fainting
- Abnormal heart rhythms
- Anxiety
- Chest pain



## Evolution of Morphological Changes in MI<sup>Q</sup>

Time	Gross	Light Microscopy	Electron microscopy
<b>Reversible injury</b> 0-30 min.	None	None	<ul style="list-style-type: none"> <li>• Relaxation of myofibrils<sup>Q</sup></li> <li>• Glycogen loss</li> <li>• Mitochondrial swelling<sup>Q</sup></li> </ul>
<b>Irreversible injury</b>			
30 min-4 hr	None	Waviness of fibers at border <sup>Q</sup> (earliest microscopic change)	<ul style="list-style-type: none"> <li>• Sarcolemmal disruption<sup>Q</sup></li> <li>• Mitochondrial amorphous densities<sup>Q</sup></li> </ul>
4-12 hr	Dark mottling appears <sup>Q</sup>	Beginning of coagulative necrosis <sup>Q</sup>	
12-24 hr	Dark mottling	<ul style="list-style-type: none"> <li>• Neutrophilic infiltration <sup>Q</sup></li> <li>• Myocyte hypereosinophilia<sup>Q</sup></li> <li>• Marginal contraction band necrosis</li> </ul>	
1-3 days	Yellow tan infarct center <sup>Q</sup>	Brisk interstitial infiltrate of neutrophils	
3-7 days	Hyperemic borders <sup>Q</sup> , central yellow tan softening	Macrophages <sup>Q</sup> at infarct border	
7-10 days	Maximum <sup>Q</sup> yellow tan	Early formation of fibrovascular granulation tissue <sup>Q</sup> at margins	
10-14 days	Red gray infarct borders	Well established granulation tissue <sup>Q</sup> Collagen deposition <sup>Q</sup>	
2-8 weeks	Gray-white scar <sup>Q</sup>	Collagen deposition <sup>Q</sup>	
>2 months	Scarring complete	Dense collagenous scar <sup>Q</sup>	






Image showing Waviness of fibers at border





Image showing Brisk interstitial infiltrate of neutrophils


## High Yield Facts

- Time for reversible injury in heart is 30 mins<sup>a</sup>
- Myocardial infarct **less than 12 hours<sup>a</sup> old- not apparent on gross examination<sup>a</sup>.**
- **Necrotic area** can be **visualized after 2–3 hours<sup>a</sup>** by immersion in **triphenyltetrazolium chloride (TTC)<sup>a</sup>**
- **Infarcted area** is revealed as **unstained pale zone<sup>a</sup>**
- TTC imparts brick red magenta color to **non-infarcted myocardium<sup>a</sup>** where **dehydrogenase enzymes<sup>a</sup>** are preserved.



Triphenyltetrazolium chloride, an enzyme substrate that colors viable myocardium magenta. Failure to stain is due to enzyme loss after cell death

- **Most common cause of SCD<sup>a</sup>-malignant ventricular arrhythmias<sup>a</sup>-myocardial ischemia-induced irritability.**



### Complication of MI (Mnemonics – ACT RAPID)

- Arrhythmia:
  - **Most common** within one hour<sup>Q</sup>-**Ventricular fibrillation**<sup>Q</sup>
  - **Most common** after one hour<sup>Q</sup>-**Supraventricular tachycardia**<sup>Q</sup>
- Contractile dysfunction
  - Leads to Cardiogenic shock
- Mural Thrombus
- Myocardial Rupture
  - **Most common** - Rupture of **ventricular free wall**<sup>Q</sup>
  - **Most common** site for **free wall rupture**- anterolateral wall at the midventricular level<sup>Q</sup>
  - **Most frequent** 3 to 7 days after MI<sup>Q</sup>
  - **Least common**-rupture of papillary muscles<sup>Q</sup>
- Ventricular Aneurysm
  - **False aneurysm**<sup>Q</sup> - **localized hematoma**<sup>Q</sup> communicating with the ventricular cavity
  - **Wall of a false aneurysm** consists only of **epicardium and adherent parietal pericardium**<sup>Q</sup>
- Pericarditis
  - **Post MI pericarditis-Dressler syndrome or post MI syndrome**<sup>Q</sup>
    - **Autoimmune reaction**<sup>Q</sup>
    - **Fibrinous or fibrinohemorrhagic pericarditis**<sup>Q</sup>
    - Develops **second or third day**<sup>Q</sup> following a transmural infarct
- Infarct expansion
  - **Seen with** anteroseptal infarcts
  - Papillary muscle **Dysfunction**: Leads to Post infarct mitral regurgitation.
  - **Progressive late heart failure (chronic IHD)**

R<sup>9th</sup>

### Latest Update

#### Infarct Modification by Reperfusion.

- **Reperfusion Injury**:<sup>Q</sup>Damage that occurs **after restoration of flow** to “vulnerable” myocardium<sup>Q</sup> that is ischemic but not irreversibly damaged
- **Stunned Myocardium**:<sup>Q</sup> State of prolonged **cardiac failure**<sup>Q</sup> induced by short-term ischemia that usually recovers after several days
- **Hibernation**:<sup>Q</sup> Myocardium that is subjected to **chronic, sublethal ischemia** may also enter into a **state of lowered metabolism**<sup>Q</sup>
- **Reperfused infarcts are usually hemorrhagic**<sup>Q</sup>
- Irreversibly injured myocytes exhibit<sup>Q</sup>: **Contraction bands (intensely eosinophilic intracellular stripes), composed of closely packed sarcomeres**<sup>Q</sup> due to exaggerated contraction of sarcomeres when perfusion is re-established because of high concentration of **calcium ions**<sup>Q</sup> from the plasma.

### Chronic Ischemic Heart Disease

- Also called **ischemic cardiomyopathy**
- **Progressive congestive heart failure**<sup>Q</sup> as a consequence of **accumulated ischemic myocardial damage**<sup>Q</sup>
- Appears post infarction due to the **functional decompensation of hypertrophied noninfarcted myocardium**<sup>Q</sup>
- Chronic IHD patients account for **50% of cardiac transplant recipients**<sup>Q</sup>
- Microscopic findings include **myocardial hypertrophy**<sup>Q</sup>, **diffuse subendocardial vacuolization**<sup>Q</sup> & **fibrosis**<sup>Q</sup>.



### High Yield Facts

- **Most commonly** affected valve in RHD is **mitral valve**<sup>Q</sup>
- **Least commonly** affected in RHD is **pulmonary valve**<sup>Q</sup>
- In acute rheumatic heart disease, the **most common valvular lesion** is **mitral regurgitation**<sup>Q</sup>
- In chronic rheumatic heart disease, **most common valvular lesion** is **mitral stenosis**<sup>Q</sup>
- **Mac-callum patches**- map-like areas of **thickened part of the endocardium in the posterior wall of left atrium**<sup>Q</sup> - caused by **regurgitant jets of blood flow**<sup>Q</sup>, due to incompetence of the mitral valve.

### RHEUMATIC FEVER AND RHEUMATIC HEART DISEASE (RHD)

- An acute **immunologically mediated multisystem inflammatory disease**<sup>Q</sup> that occurs **few weeks** after an attack of **group Aβ- hemolytic streptococcal pharyngitis**<sup>Q</sup>
- **Most common age group** affected is children between of **5-15 years**<sup>Q</sup>.
- It is **not an infective disease**<sup>Q</sup>.
- **Type II hypersensitivity reaction**
- Antibodies against ‘M’ protein of some streptococcal strains **cross-react with the glycoprotein antigens in the heart, joints and other tissues (molecular mimicry)**.


### Salient Features of the Major Criteria

#### Carditis

**Pancarditis**: Involves pericardium, myocardium and endocardium

- **Endocardium**:
  - **Fibrinoid necrosis**<sup>Q</sup> within the cusps or tendinous cords.
  - **Verrucae**- small (1 to 2 mm) vegetations **along the lines of closure**<sup>Q</sup>
- **Myocardium**- has **Aschoff's bodies**<sup>Q</sup> in the **perivascular location**
- **Pericarditis** is associated with **fibrinous/serofibrinous**<sup>Q</sup> exudates-‘**bread and butter**’ pericarditis.<sup>Q</sup>





**Aschoff Bodies:** Here u can see collection of foci of swollen eosinophilic collagen surrounded by Lymphocytes, giant cells<sup>o</sup> (marked with green arrow) and Antischkow cells (cells with caterpillar like chromatin shown in the inset)

### Migratory Polyarthritits

- Most commonly seen manifestation; More commonly seen in the adults<sup>o</sup> as compared to children
- Involvement of the large joints of the body; Arthritis involves one joint after the other<sup>o</sup> (migratory)
- Subsides spontaneously without any residual deformability<sup>o</sup> in the joints (non-erosive arthritis).

### Subcutaneous Nodules

Painless subcutaneous lesions on extensor surface of elbows, shin and occiput<sup>o</sup>.

### Erythema Marginatum

Red macular rash more easily appreciated in fair skinned individuals sparing the face<sup>o</sup>

### Syndenham's Chorea

Involuntary, purposeless movements<sup>o</sup> associated with emotional lability of patient

### Chronic RHD

- Mitral valve is always involved
- Cardinal anatomic changes in mitral valve- leaflet thickening, commissural fusion and shortening, thickening & fusion of the tendinous cords.
- Fibrosis produces Mitral stenosis <sup>o</sup>also known as 'fish-mouth' or 'button-hole' stenosis.<sup>o</sup>



Fish mouth appearance



### High Yield Facts

#### Aschoff Bodies:

- Focal distinctive inflammatory lesions seen during acute rheumatic fever.<sup>o</sup> They are pathognomonic of RHD
  - Consist of foci of swollen eosinophilic collagen<sup>o</sup> surrounded by:
    - Lymphocytes (primary T cells), Occasional plasma cells, Aschoff giant cells<sup>o</sup> (macrophages of rheumatic fever)<sup>o</sup>, Antischkow cells (pathognomonic for RF)<sup>o</sup>
- Antischkow cells** - modified macrophages with abundant cytoplasm and central round to ovoid nuclei with central wavy ribbon like chromatin hence also called caterpillar cells,<sup>o</sup> seen in any of the 3 layers of the heart<sup>o</sup>

## INFECTIVE ENDOCARDITIS (IE)

- Endocarditis-Inflammation of inner layer of heart

Acute Endocarditis	Subacute Endocarditis
<ul style="list-style-type: none"> <li>Infection of Previously normal<sup>o</sup> valve.</li> <li>Highly virulent<sup>o</sup> organisms</li> <li>Most common caused by <b>Staph. aureus</b><sup>o</sup></li> <li>Produce necrotic and destructive lesions</li> <li>Death of the patient within days to weeks</li> </ul>	<ul style="list-style-type: none"> <li>Infection of Previously damaged valve<sup>o</sup></li> <li>Low virulence<sup>o</sup> organisms</li> <li>Most common <b>Hemolytic (viridans), Streptococcus</b><sup>o</sup></li> <li>Produce less destruction of valves</li> <li><b>Protracted cause of weeks and months-Recover</b> after antibiotic therapy</li> </ul>

### Morphology

- The friable, bulky destructive vegetations contain- **fibrin, bacteria and inflammatory cells** <sup>o</sup>
- Found on the valve cusps, can also extend on to chordae.
- Ring abscess**<sup>o</sup>-When the vegetations erode into myocardium, they can form an abscess
- Most commonly infected valve**<sup>o</sup> - aortic valve and the mitral valve
- Intravenous drug abusers**<sup>o</sup> - **right side**<sup>o</sup> of heart is affected




Image shows friable bulky vegetations



## Clinical Features

- Fever is the **most consistent sign** of IE.
- The other features include weight loss, flu-like syndrome, microthromboemboli (manifest as **splinter or subungual hemorrhages**), erythematous or hemorrhagic non-tender lesions on palms or soles (**Janeway lesions**), subcutaneous nodules in the pulp of digits (**Osler nodes**), and retinal hemorrhages in the eyes (**Roth spots**).
- The disease is diagnosed by **Dukes criteria**.<sup>Q</sup>
- **Blood culture is the investigation of choice**.<sup>Q</sup>



## High Yield Facts

### Most common site for vegetation in:

- **Libman Sack's endocarditis** are the **A-V valves**<sup>Q</sup>, mitral and tricuspid.
- **NBTE** is **mitral**<sup>Q</sup> and less often aortic and Tricuspid.
- **RF** in **mitral**<sup>Q</sup> followed by combined mitral and aortic

### Most frequent causes of the functional valvular lesions

- Aortic stenosis: **Calcification** of anatomically normal and **congenitally bicuspid aortic valves**
- Aortic regurgitation: Dilation of the ascending aorta due to **hypertension**<sup>Q</sup> and **aging**<sup>Q</sup>
- Mitral stenosis: **Rheumatic heart disease**.<sup>Q</sup>
- Mitral regurgitation: **Myxomatous degeneration** (mitral valve prolapse).<sup>Q</sup>

## NONINFECTED VEGETATIONS

### Nonbacterial Thrombotic Endocarditis (NBTE)/ Marantic Endocarditis

- Vegetations on heart valves are sterile and **do not contain microorganisms**.<sup>Q</sup>
- Small, present **along the line of closure, single or multiple**.<sup>Q</sup>
- **Not invasive, do not elicit any inflammatory reaction**<sup>Q</sup>
- **Loosely attached**, so can cause **systemic emboli**<sup>Q</sup> that produce significant infarcts in brain, heart, or elsewhere.

### Predisposing Conditions

- Debilitated patients, such as those with cancer (Ca pancreas, mucinous adenoCa, APML) or sepsis—hence the previous term **marantic endocarditis**<sup>Q</sup>
- **Systemic hypercoagulable state**<sup>Q</sup>
- **Endocardial trauma**<sup>Q</sup>, as from an indwelling catheter

### Libman-sacks Endocarditis (seen in SLE)

- **Small** (1-4 mm), single or multiple, **sterile**<sup>Q</sup>, vegetations with a **warty (verrucous) appearance**<sup>Q</sup>
- Located on the **either or both sides of valve leaflets especially under surfaces of the atrioventricular valves**
- **Mitral and tricuspid valves** are involved
- Associated with an **intense valvulitis**<sup>Q</sup>, characterized by **fibrinoid necrosis**<sup>Q</sup> of the valve substance
- Vegetations consist of **finely granular, fibrinous eosinophilic** material containing cellular debris & nuclear remnants



## Diffrentiating Features of Various Types of Endocarditis

Rheumatic Fever (RF)	Non Bacterial Thrombotic (Marantic Endocarditis)	Libman Sack's Endocarditis	Infective Endocarditis
1. <b>Small, warty</b> <sup>Q</sup> 2. Firm 3. Friable	1. <b>Small, warty</b> <sup>Q</sup> 2. Friable	1. Medium sized (small) 2. Flat, Verrucous 3. Irregular	1. <b>Large</b> <sup>Q</sup> 2. <b>Bulky</b> <sup>Q</sup> 3. Irregular
<b>Along line of closure</b> <sup>Q</sup>	<b>Along line of closure</b> <sup>Q</sup>	1. On surface of cusps 2. (both surfaces may be involved but the <b>undersurface is more likely affected</b> , <sup>Q</sup> less commonly mural endocardium is involved) 3. <b>In pockets of valves</b> <sup>Q</sup>	1. Vegetations on the <b>valve cusps</b> <sup>Q</sup> 2. Less often on mural endocardium
Sterile (no organism)	Sterile	Sterile	<b>Non-sterile (bacteria)</b> <sup>Q</sup>
Embolization is uncommon	<b>Embolization is common</b> <sup>Q</sup>	Embolization is uncommon	<b>Embolization is very common (max chances)</b> <sup>Q</sup>
In <b>rheumatic heart disease</b> <sup>Q</sup>	In <b>cancers (like M3-AML, Ca pancreas), DVT, Trousseau's syndrome</b> <sup>Q</sup>	In SLE	In infective endocarditis





## CARDIOMYOPATHIES




HOCM: Myofiber hypertrophy & disarray




Gross: HOCM-hypertrophic cardiomyopathy

## PERICARDITIS

- Normally, visceral pericardium is separated from the parietal pericardium by a **small quantity (15–50 mL) of fluid<sup>Q</sup>**
- Inflammation of pericardium is called pericarditis<sup>Q</sup>**





Fibrinous pericarditis

### R10<sup>th</sup> Latest Update

- Dilated cardiomyopathy is **most common**<sup>Q</sup> (90% of cases). Most cases are due to mutations in titin (largest known human protein)
- Restrictive cardiomyopathy is the **least frequent**<sup>Q</sup>
- **Naxos syndrome**<sup>Q</sup>: characterized by **arrhythmogenic right ventricular cardiomyopathy**<sup>Q</sup> & **hyperkeratosis**<sup>Q</sup> of plantar palmar skin, associated with mutation in gene encoding the desmosome-associated protein **plakoglobin**<sup>Q</sup>
- **HOCM** is the **leading cause of unexplained left ventricular hypertrophy**
- **HOCM** is **Most common cause of sudden death in young athletes**<sup>Q</sup>
- **Most common genetic mutation in HOCM**<sup>Q</sup> - gene encoding **β-myosin heavy chain (β-MHC)**<sup>Q</sup>
- **Loeffler's endomyocarditis**: Peripheral eosinophilia & eosinophilic infiltrates in multiple organs, including heart<sup>Q</sup>
- **Endocardial fibroelastosis**: Most common in the **first 2 years**<sup>Q</sup> of life
- **Endocardial fibroelastosis** morphologic end-point of different insults including viral infections (e.g., **intrauterine exposure to mumps**<sup>Q</sup>) or mutations in the gene for **tafazzin**<sup>Q</sup>, which affects **mitochondrial inner membrane integrity**<sup>Q</sup>

## MYOCARDITIS

- Myocarditis, i.e., cardiac inflammation, is most commonly caused by viral infections.
- **Coxsackieviruses A<sup>Q</sup> and B<sup>Q</sup> and other enteroviruses<sup>Q</sup> account for most of the cases.**
- **Most common helminth causing myocarditis-Trichinella<sup>Q</sup>**
- **Most common bacterial infection associated with myocarditis- Diphtheria<sup>Q</sup>**



### High Yield Facts

- **Most common cause of pericarditis is viral causes**<sup>Q</sup>
- **Mulibrey nanism-autosomal recessive**<sup>Q</sup> syndrome is characterized by **growth failure, muscle hypotonia, hepatomegaly, ocular changes, enlarged cerebral ventricles, mental retardation, ventricular hypertrophy & chronic constrictive pericarditis**<sup>Q</sup>
- **Most frequent type of pericarditis is fibrinous pericarditis**


### Giant-cell Myocarditis

- Characterized by inflammatory infiltrate containing **multi-nucleate giant cells** with lymphocytes, eosinophils, plasma cells, and macrophages.
- Focal to frequently extensive **necrosis** is present.
- This variant carries a **poor prognosis**.<sup>Q</sup>

## CARDIAC TUMORS

### Myxoma

- **Most common primary tumor** of the heart in adults.<sup>Q</sup>
- **Benign** neoplasms



Myxoma

- Arise from **primitive multipotent mesenchymal cells** 90% are located in the atria, with a **left-to-right ratio of approximately 4:1 (atrial myxomas)**.
- Major clinical manifestations are due to valvular "ball-valve" obstruction, embolization, or a syndrome of constitutional symptoms, such as fever and malaise the latter **most commonly due to cytokine interleukin-6**.





## Morphology

- Always **single**
- Fossa ovalis** in the **atrial septum** is the favored site of origin.
- Histologically, myxomas are composed of stellate or globular myxoma (“**lepidic**”) cells, embedded within an abundant acid **mucopolysaccharide ground substance** and covered on the surface by endothelium


## Papillary Fibroelastoma

- Incidental, **sea-anemone-like lesions**, most often identified at autopsy<sup>Q</sup>
- >80% located on **valves<sup>Q</sup> (ventricular surfaces of semilunar valves & atrial surfaces of AV valves)<sup>Q</sup>**
- Resemble the much smaller, usually trivial, lambl excrescences

## Rhabdomyoma

- Most frequent primary tumor of the pediatric heart<sup>Q</sup>**
- Most common discovered in **first years** of life because of **obstruction of a valvular orifice**
- Actually hamartomas<sup>Q</sup>** rather than true neoplasms as they **regress spontaneously<sup>Q</sup>**
- 50% cases associated with **tuberous sclerosis<sup>Q</sup>**

- Usually **multiple** and involve the **ventricles**, protruding into the lumen
- Histologically**: large, rounded, or polygonal cells containing numerous **glycogen-laden vacuoles** separated by strands of cytoplasm (**spider cells<sup>Q</sup>**)



Large, polygonal cells (“spider cells”) with glycogen vacuoles separated by strands of cytoplasm extending between cell membrane and nucleus

## High Yield Facts

- The most common cardiac tumor is the **secondaries or metastasis<sup>Q</sup>**
- The most common primary cardiac tumor in the adults is the **myxoma<sup>Q</sup>**
- The most common cardiac tumor in the children is the **rhabdomyoma<sup>Q</sup>**
- Carney syndrome**: multiple cardiac & extracardiac myxomas, spotty pigmentation & endocrine overactivity
- Carcinoid heart disease<sup>Q</sup>** typically causes abnormalities of the **right side of the heart<sup>Q</sup>**
- Familial syndromes associated with myxomas**
  - Activating mutations in the GNAS1 gene (in association with McCune-Albright syndrome)
  - Null mutations in PRKAR1A (Carney complex)<sup>Q</sup>

## High Yield Facts

MC cause of <b>aortic aneurysm</b>	Atherosclerosis <sup>Q</sup>	MC site of <b>Tuberculous aneurysms</b>	Thoracic aorta. (T for T) <sup>Q</sup>
MC site of <b>aneurysm caused by atherosclerosis</b>	Abdominal aorta (below renal artery and above bifurcation) <sup>Q</sup>	MC cause of <b>descending aortic aneurysm</b>	Atherosclerosis <sup>Q</sup>
MC cause of <b>Thoracic Aortic Aneurysm</b>	Hypertension <sup>Q</sup>	MC site of <b>Syphilitic aneurysm</b>	Ascending aorta <sup>Q</sup>
MC cause of <b>ascending aortic aneurysm</b>	Cystic medial degeneration <sup>Q</sup> /Systemic hypertension <sup>Q</sup>	MC cause of <b>mycotic abdominal aneurysm</b>	Salmonella gastroenteritis <sup>Q</sup>
MC cause of <b>aortic arch aneurysm</b>	Atherosclerosis <sup>Q</sup>	MC site of <b>Traumatic aneurysms</b>	Descending thoracic aorta just below the site of insertion of ligamentum arteriosum. <sup>Q</sup>
MC site of <b>takayasu arteritis aneurysm</b>	Aortic arch <sup>Q</sup>	<b>Nutritional basis for aneurysm formation</b>	Scurvy <sup>Q</sup> Due to altered collagen cross-linking <sup>Q</sup> True aneurysm <sup>Q</sup>



R10<sup>th</sup>

**Latest Update**

### c-ANCA and p-ANCA Disease Associations

Disease	c-ANCA (PR3) (%)	p-ANCA (MPO) (%)
GPA (WG)	66	24
MPA	26	58
EGPA (CSS)	<5	40

c-ANCA, cytoplasmic antineutrophil cytoplasmic antibody, EGPA (CSS), eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome); GPA (WG), granulomatosis with polyangiitis (Wegener's granulomatosis); MPA, microscopic polyangiitis; MPO, myeloperoxidase; PR3, peroxidase-3.




## NEXT Pattern Questions



Q's

1. A 32-year-old patient underwent renal Biopsy for nephrotic syndrome. Along with glomerular pathology, arterioles show the lesions as depicted in the histopathological picture. Identify the lesion.



- a. Hyperplastic arteriolosclerosis
- b. Mönckeberg medial Sclerosis
- c. Hyaline arteriolosclerosis
- d. Atherosclerosis


**Ans. (c) Hyaline arteriolosclerosis**

- Renal arterioles in diabetes show homogeneous pink deposits suggestive of **Hyaline arteriolosclerosis**



Q's

2. A 40-year-old male patient presented with hypertension due to renal artery involvement, histological examination of the vessel is shown in the diagram. False statement regarding histological features is:




- a. Segmental transmural necrotizing inflammation of small- to medium-sized arteries
- b. In acute phase Fibrinoid necrosis can be seen
- c. Late stage show fibrosis, thrombosis
- d. All vessel or single vessel will be in same stage of inflammation

**Ans. (d) All vessel or single vessel will be in same stage of inflammation**


- Here all stages of inflammation is seen, suggestive of PAN single stage of activity is seen in microscopic polyangitis.





Q's

3. At what duration of myocardial infarction light microscopy demonstrate myocyte necrosis, hypereosinophilia, contraction band necrosis and intense neutrophil infiltrate




- a. 4-6 hours  
b. 12-24 hours  
c. 1-3 days  
d. 3-7 days

Ans. (c) 1-3 days



Q's

4. A 30-year-old male presented with severe dyspnea and fatigue. X-ray showed left atrial enlargement. Physician suspects the patient of having mitral stenosis and gets a histopath examination done, the image of which is shown, it shows?



- a. Sarcoidosis  
b. Tuberculosis  
c. Aschoff bodies  
d. Fungal granuloma

Ans. (c) Aschoff bodies

- As you see the history of severe dyspnea and fatigue, X-ray showed left atrial enlargement, patient of having mitral stenosis. The gross image of valve shows fish mouth appearance and histopathology of the same shows Aschoff bodies.



Q's


5. Vegetations of the following endocarditis has the maximum chances of embolization:
- a. Rheumatic heart disease  
b. Infective endocarditis  
c. Libman-Sacks endocarditis  
d. Subacute bacterial endocarditis

Ans. (b) Infective endocarditis



Q's

6. The swelling of size 1.5 cm is seen over chest of 20-year-old, soft to firm reddish swelling, progressing in size. Histopathological examination showed the following. What could be your possible diagnosis?



- a. Hemangioma  
b. Fibroadenoma  
c. Lipoma  
d. Dermatofibroma


Ans. (a) Hemangioma

- Soft to firm reddish swelling, progressing in size suggestive of hemangioma.



Q's

7. A 55-year-old female patient having large adnexal mass with raised inhibin level. Histopathology shows cord and sheets of tumor cell, often showing nuclear groove as shown in the diagram, based on the above clinical features identify the type of tumor.



- a. Choriocarcinoma  
b. Granulosa cell tumor  
c. Yolk sac tumor  
d. Leydig cell tumor



Ans. (b) Granulosa cell tumor

- Large adnexal mass with raised inhibin level is suggestive of granulosa cell tumor. Histopathology shows cord and sheets of tumor cell, often showing nuclear groove suggestive of Call-Exner body seen in Granulosa cell tumor.



Q's

8. Post surgery image of a tumor in scrotum is shown below along with its histology. What could be your possible diagnosis?




- a. Teratoma      b. Seminoma      c. Yolk sac tumor      d. Lymphoma

Ans. (b) **Seminoma**




## Image-Based Questions

1. A 45-year-old male presented with pulseless disease. On examination, following histopathological finding will be seen




- a. Granulomatous vasculitis      b. Fibrinoid necrosis  
c. Leucocytoclastic vasculitis      d. Thromboangitis obliterans

2. A patient is a known case of polyarteritis nodosa. On examination of biopsy, accumulation of amorphous, basic, proteinaceous material in the vessel wall was seen. This finding is suggestive of:



- a. Fibrinoid necrosis      b. Leucocytoclastic vasculitis  
c. Hyaline arteriosclerosis      d. Caseous necrosis






3. A 34-year-old male patient presented with following clinical feature. Diagnosis



- |                     |                             |
|---------------------|-----------------------------|
| a. Kawasaki disease | b. PAN                      |
| c. MicroPAN         | d. Wegener's granulomatosis |

4. A 15-year-old boy presented with pancarditis, on myocardial biopsy, following finding was seen. Diagnosis?



- |                   |         |
|-------------------|---------|
| a. Aschoff nodule | b. TB   |
| c. FB giant cells | d. None |

5. A 25-year-old male presented with growth in left atrium. Diagnosis?



- |                |                        |
|----------------|------------------------|
| a. Rhabdomyoma | b. Myxoma              |
| c. Metastasis  | d. Papillary elastosis |

## Answers of Image-Based Questions

1. Ans. (a) **Granulomatous vasculitis**

- Here we can see epithelioid cells and giant cells s/o granulomatous vasculitis
- Granulomatous vasculitis is seen in : Giant cell (temporal) arteritis, Takayasu arteritis, Wegener's granulomatosis, Churg-Strauss syndrome, Buerger's disease

2. Ans. (a) **Fibrinoid necrosis**

- Fibrinoid necrosis is a form of necrosis, or tissue death, in which there is accumulation of amorphous, basic, proteinaceous material in the tissue matrix with a staining pattern reminiscent of fibrin.
- How to differentiate from hyaline arteriosclerosis? - please remember in fibrinoid necrosis, pink material has fibrin like quality as opposed to glassy homogenous hyaline in hyaline arteriosclerosis. (refer to image 2)

3. Ans. (d) **Wegener's granulomatosis**

- This is strawberry gums, seen in Wegener's granulomatosis

4. Ans. (a) **Aschoff nodule**

- Here one can see collection of foci of swollen eosinophilic collagen surrounded by : Lymphocytes (primary T cells), Occasional plasma cells, Aschoff giant cells<sup>Q</sup> (macrophages of rheumatic fever)<sup>Q</sup> and Antischkow cells (cells with caterpillar like chromatin marked with an arrow)

5. Ans. (b) **Myxoma**

- The left atrium has been opened to reveal the most common primary cardiac neoplasm-an atrial myxoma. These benign masses are most often attached to the atrial wall. They can produce a "ball valve" effect by intermittently occluding the atrioventricular valve orifice.




## Multiple Choice Questions

### BLOOD VESSELS

#### SCLEROSIS INCLUDING ATHEROSCLEROSIS

1. **Cleft like space in atheromatous plaque mainly contains?** (Recent Question 2016)
  - a. Smooth muscle cell
  - b. Fibrous tissue
  - c. Cholesterol
  - d. Macrophages



2. **Atherosclerosis causes fibroblast plaque formation by injury to?** (Recent Question 2015)
  - a. Endothelium
  - b. Fibroblast
  - c. Macrophage
  - d. Smooth muscle cells
3. **The following arteries are usually spared from extensive atherosclerosis** (Recent Question 2015)
  - a. Popliteal artery
  - b. Internal carotid artery
  - c. Arteries of circle of willis
  - d. Mesenteric arteries
4. **The necrotic core of an atherosclerotic plaque contains**
  - a. T cells
  - b. Collagen
  - c. Lipid
  - d. None of the above
5. **True about atherosclerosis** (Recent Question 2014-15)
  - a. Chronic inflammatory disorder of vessel wall
  - b. Not lead to complications of vessel wall
  - c. Thoracic aorta more than abdominal aorta
  - d. Atherosclerotic plaques do not demonstrate extra-cellular matrix deposition
6. **The coronary artery most commonly involved in atherosclerosis** (Recent Question 2015)
  - a. Left anterior descending artery
  - b. Left main coronary artery
  - c. Right coronary artery
  - d. Circumflex coronary artery
7. **Not true about monckeberg medial sclerosis** (Recent Question 2015)
  - a. Calcification of the walls of muscular arteries
  - b. Typically involving the internal elastic membrane
  - c. Persons older than age 50 are most commonly affected
  - d. Calcifications cause significant narrowing of vessel lumen
8. **Foam cells in atherosclerosis contain lipid in the form of** (Recent Question 2014-15)
  - a. Oxidized LDL
  - b. Reduced LDL
  - c. Oxidized VLDL
  - d. Reduced VLDL

9. **Following are the modifiable risk factor of atherosclerosis except** (Recent Question 2014-15)
  - a. Physical inactivity
  - b. Family history
  - c. Diabetes
  - d. Hypertension
10. **True about the basic structure of artherosclerosis plaque is-** (Recent Question 2014-15)
  - a. Concave part formed by fibrous cap
  - b. Convex part formed by tunica media of the vessel
  - c. Convex part formed by fibrous cap
  - d. Necrotic core contains collagen, elastin and proteoglycans
11. **Atherosclerosis initiation by fibroblast plaque is mediated by injury to** (Recent Question 2014-15)
  - a. Smooth muscle
  - b. Media
  - c. Adventitia
  - d. Endothelium
12. **Atheromatous changes of blood vessels affects early in:** (Recent Question 2015)
  - a. Kidney
  - b. Heart
  - c. Liver
  - d. Spleen
13. **Medial calcification is seen in:** (Recent Question 2015)
  - a. Atherosclerosis
  - b. Arteriolosclerosis
  - c. Monckebergs sclerosis
  - d. Dissecting aneurysm
14. **Changes seen in-atherosclerotic plaque at the time of rupture, are all except:** (Recent Question 2014)
  - a. Thin fibrotic cap
  - b. Multiple foam cap
  - c. Smooth muscle cell hypertrophy
  - d. Cell debris
15. **Infective agent causing atherosclerosis** (Recent Question 2014)
  - a. M. Pneumoniae
  - b. C. Pneumoniae
  - c. H. Influenza
  - d. C. Diphtheriae
16. **In atherosclerosis, increased LDL in monocyte macrophage due to:** (DPG 10, PGI June 99)
  - a. LDL receptors on macrophage
  - b. LDL receptors on endothelium
  - c. Lipids in LDL get oxidized
  - d. All of the above

#### HYPERTENSION

17. **Characteristic histological finding in benign hypertension** (Recent Question 2015)
  - a. Proliferative end arteritis
  - b. Necrotizing arteriolitis
  - c. Hyaline arteriosclerosis
  - d. Cystic medial necrosis
18. **Which of the following is seen in kidney in malignant hypertension?** (Recent Question 2014-15)
  - a. Hyaline necrosis
  - b. Fibrinoid necrosis
  - c. Medial wall hyperplasia
  - d. Microaneurysm
19. **Onion peeling of renal vessels is seen in:** (Recent Question 2014)
  - a. Benign hypertension
  - b. Malignant hypertension
  - c. Diabetic nephropathy
  - d. SLE





20. In a specimen of kidney, fibrinoid necrosis is seen and onion peel appearance is also present. Most probable pathology is: (AIIMS May 2013)
- Hyaline degeneration
  - Hyperplastic arteriosclerosis
  - Glomerulosclerosis
  - Fibrillary glomerulonephritis
21. Typical histology in benign hypertension? (JIPMER 2013)
- Intimal proliferation and hyalinization of media of medium arteries
  - Fibrinoid necrosis of small arteries
  - Loss of endothelial cells of arterioles
  - Formation of new vessels
22. Onion skin thickening of arteriolar wall is seen in:
- Atherosclerosis (Recent Question 2013, DNB 10)
  - Median calcific sclerosis
  - Hyaline arteriosclerosis
  - Hyperplastic arteriosclerosis

### ANEURYSMS


23. Syphilitic aneurysm is seen most commonly in? (Recent Question 2016-17)
- Ascending aorta
  - Arch of aorta
  - Descending aorta
  - Abdominal aorta
24. Which of the following causes pseudoaneurysm? (Recent Question 2015)
- Trauma
  - Atherosclerosis
  - Congenital disease
  - Infection
25. In Marfan syndrome, rupture of aortic aneurysm usually occurs at (Recent Question 2015)
- Ascending aorta
  - Descending aorta
  - Arch of aorta
  - Abdominal aorta
26. For asymptomatic abdominal aortic aneurysm surgery is indicated if the size is greater than (Recent Question 2015)
- 4 cm
  - 4.5 cm
  - 5 cm
  - 5.5 cm
27. Risk of aneurysm rupture is > 25% per year when the size is greater than (Recent Question 2015)
- 4 cm
  - 6 cm
  - 7 cm
  - 8 cm
28. Most common cause of aneurysm (Recent Question 2015)
- Syphillis
  - Atherosclerosis
  - Cystic medial necrosis
  - Hypertension
29. Which of the following is not a cause of aneurysm? (Recent Question 2014)
- Atherosclerosis
  - Cystic medial necrosis
  - Syphillis
  - Mockenbergs sclerosis
30. Visceral aneurysm is most commonly seen in: (Recent Question 2013)
- Splenic
  - Renal
  - Hepatic
  - Coronary

### DISSECTIONS

31. IOC for acute aortic dissection (Recent Question 2015)
- MRI
  - MD- CT scan
  - Transesophageal echocardiography
  - X-ray
32. Most common predisposing factor for aortic dissection (Recent Question 2015)
- Atherosclerosis
  - Syphilis
  - Hypertension
  - Smoking
33. Aortic dissection is not common in the following disease (Recent Question 2015)
- Marfan syndrome
  - Hypertension
  - Cystic medial necrosis
  - Secondary syphilis
34. Classification of aortic dissection depends upon- (Recent Question 2014)
- Cause of dissection
  - Level of aorta affected
  - Percentage of aorta affected
  - None
35. Cystic medial necrosis is seen in- (Recent Question 2013)
- Marfans
  - Friedrichs ataxia Pattern
  - Downs
  - Kawasaki
36. Most common cause of dissecting hematoma is because of- (Recent Question 2016-17)
- Hypertension
  - Marfan's
  - Iatrogenic
  - Kawasaki

### VASCULITIS


37. Small vessel vasculitis is/are? (PGI May 2017)
- Anti GBM ab
  - Takayasu arteritis
  - Kawasaki disease
  - Carcinoma induced vasculitis
  - IgA vasculitis
38. True about Takayasu is all except? (PGI Nov 2016)
- Causes transmural granuloma
  - Granulomatous vasculitis
  - Medium to small vessel
  - Called aortoarteritis
39. All are true about Kawasaki disease except? (Recent Question 2016)
- Thrombocytopenia
  - Dequamation rashes on trunk
  - Fever
  - Conjunctival congestion
40. Berger disease does not involve (Recent Question 2016)
- Artery
  - Veins
  - Nerve
  - Lymphatics
41. Which one is found in HSP? (Recent Question 2016)
- IgA
  - IgM
  - Ig G
  - IgE
42. Which of the following is non-granulomatous arteritis? (Recent Question 2016)
- Takayasu
  - Wegeners
  - Churgstrauss
  - Classical PAN



- 43. HSP is characterized by all except? (APPGMEE 2015)**  
 a. Glomerulonephritis b. Hematochezia  
 c. Thrombocytopenia d. Palpable purpura
- 44. c-ANCA is positive in (Recent Question 2015)**  
 a. Microscopic Polyangitis  
 b. Wegener's Granulomatosis  
 c. Churg Strauss Syndrome  
 d. Behcet's syndrome
- 45. Arterial biopsy of elderly male shows fragmentation of elastic lamina, lymphocyte infiltration and giant cells (Recent Question 2014-15)**  
 a. Temporal arteritis b. Takayasu disease  
 c. Polyarteritis nodosa d. Kawasaki disease
- 46. Temporal arteritis all are associated except (Recent Question 2014-15)**  
 a. Elderly patient  
 b. Low ESR  
 c. Giant cells  
 d. Polymyalgia rheumatica
- 47. Fibrinoid necrosis is seen in (Recent Question 2014-15)**  
 a. Polyarteritis nodosa b. SLE  
 c. HIV d. Sarcoidosis
- 48. Silk Road disease is (Recent Question 2015)**  
 a. Behçet's syndrome  
 b. Giant cell arteritis  
 c. Henoch schonlein purpura  
 d. Wegener's granulomatosis
- 49. Which is not a characteristic of wegers granulomatosis: (Recent Question 2015)**  
 a. Granuloma in vessel wall  
 b. Focal necrotising glomerulonephritis  
 c. Positive for cANCA  
 d. Involves large vessels
- 50. Frequency of renal involvement in HSP (Recent Question 2015)**  
 a. 20-40% b. >80%  
 c. 40-60% d. 10%
- 51. ANCA positive vasculitis – (Recent Question 2014)**  
 a. Henoch schonlein purpura  
 b. Behcet's syndrome  
 c. Wegener's granulomatosis  
 d. None
- 52. Fibrinoid necrosis with neutrophilic infiltration is seen in: (Recent Question 2014)**  
 a. PAN  
 b. Giant cell arteritis  
 c. Takayasu arteritis  
 d. Wegener's granulomatosis
- 53. Necrotizing arterioritis with fibrinoid necrosis is: (Recent Question 2014)**  
 a. Immediate hypersensitivity  
 b. Cell mediated immunity  
 c. Antigen-antibody complex mediated  
 d. Cytotoxic cell mediated
- 54. Which is associated with vasculitis of medium size vessels (Recent Question 2014)**  
 a. Temporal arteritis  
 b. Wegners granulomatosis  
 c. Classic PAN  
 d. Tuberous sclerosis
- 55. All is true about Giant cell arteritis except (Recent Question 2014)**  
 a. Involves large to small sized areteries  
 b. Granulomatous inflammation  
 c. Most commonly involved artery is abdominal aorta  
 d. Segmental nature of the involvement
- 56. In Wegener's granulomatosis cytoplasmic anti neutrophilic antibodies are directed against (Recent Question 2014)**  
 a. Proteinase 1  
 b. Proteinase 2 June 08  
 c. Proteinase 3  
 d. Proteinase 4
- 57. Microscopic polyangiitis is characterized by the following features EXCEPT (APPGMEE 14)**  
 a. Involves small and medium sized arteries and veins  
 b. 75% cases are associated with ANCA positivity  
 c. Palpable purpura, ulcers and vesiculo bullous lesion  
 d. Unlikely to cause pulmonary renal syndrome
- 58. In PAN, cysts are seen in all except: (Recent Question 13, AI 00,95)**  
 a. Lung b. Pancreas  
 c. Liver d. Heart
- 59. In Wegeners glomerulonephritis characteristic feature seen is: (AIIMS Nov 10, Nov 09)**  
 a. Granuloma in the vessel wall  
 b. Focal necrotizing glomerulonephritis  
 c. Nodular glomerulosclerosis  
 d. Interstitial granuloma
- 60. Which of the following is not a common cause of vasculitis in adults: (DNB June 10)**  
 a. Giant cell arteritis  
 b. Kawasaki disease  
 c. Henoch schonlein purpura  
 d. Polyarteritis nodosa

#### VASCULAR TUMORS

- 61. The swelling of size 1.5 cm is seen over chest of 20 year old, soft to firm reddish swelling, progressing in size. Histopathological examination showed the following. What could be your possible diagnosis? (Recent exam 2018)**



- a. Hemangioma b. Fibroadenoma  
 c. Lipoma d. Dermatofibroma



- 62. Kaposi sarcoma true is?** (PGI Nov 2015)  
 a. Causative agent HHV 8  
 b. Can be Seen in depressed cell mediated immunity  
 c. Treatment is chemotherapy and surgery  
 d. Vascular tumor
- 63. Portwine stain is** (Recent Question 2015)  
 a. Capillary hemangioma  
 b. Cavernous hemangioma  
 c. Lymphangioma  
 d. Vascular ectasias
- 64. Multifocal tumor of vascular origin in a patient with AIDS:** (Recent Question 2015)  
 a. Astrocytoma                      b. Gastric Carcinoma  
 c. Kaposi sarcoma                  d. Primary CNS lymphoma
- 65. Find the true statement** (Recent Question 2015)  
 a. Classic Kaposi sarcoma is associated with HIV  
 b. HAART therapy has decreased AIDS associated KS  
 c. Transplant associated AIDS-visceral involvement  
 d. Lymphadenopathic KS-extensive skin lesions
- 66. Fish hook pattern of capillaries is seen in** (Recent Question 2015)  
 a. Capillary hemangioma    b. Cavernous hemangioma  
 c. Angiosarcoma                  d. Hemangiopericytoma
- 67. Pathological feature of pyogenic granuloma:** (Recent Question 2015)  
 a. Epithelioid cells  
 b. Capillary hemangioma  
 c. Granulation tissue  
 d. Giant cells
- 68. Most common site of Angiosarcoma is:** (Recent Question 2014)  
 a. Liver                                  b. Lung  
 c. Kidney                                d. Lip
- 69. CD marker of Angiosarcoma is?** (Recent Question 2013)  
 a. CD 10                                b. CD 19  
 c. CD25                                 d. CD 31
- 70. Spontaneous regression can occur with:** (Recent Question 2013)  
 a. Cavernous hemangioma  
 b. Strawberry angioma  
 c. Nevus flemes  
 d. None of the above
- 71. True about Giant aneurysm:** (Recent Question 2013)  
 a. Rarely rupture  
 b. Most common in middle cerebral artery  
 c. Pressure effect is often the presenting symptom  
 d. Thromboembolic phase is present

#### VEINS AND LYMPHATICS

- 72. Milroy's disease is an example of:** (Recent Question 2015)  
 a. Primary lymphedema  
 b. Secondary lymphedema  
 c. Both  
 d. None

#### MISCELLANEOUS

- 73. The following is not an aging change in heart** (Recent Question 2015)  
 a. Increased LA cavity size  
 b. Increased LV cavity size  
 c. Sigmoid-shaped ventricular septum  
 d. Lambl excrescences

- 74. Raynaud's phenomenon what change is seen in vessels initial stage:** (Recent Question 2014)  
 a. No change                              b. Thrombosis  
 c. Fibrinoid necrosis                  d. Hyaline sclerosis
- 75. Raynaud's phenomenon is seen in:** (PGI May 10)  
 a. SIB  
 b. Systemic sclerosis  
 c. DM  
 d. Hypertension


### CARDIOVASCULAR SYSTEM

#### HEART FAILURE

- 76. Heart failure cells are seen in** (Recent Question 2015)  
 a. Kidney  
 b. Heart  
 c. Lungs  
 d. Brain
- 77. Commonest cause of right ventricular failure is:** (AIIMS 14)  
 a. Cor pulmonale  
 b. Pulmonary involvement  
 c. Endomyocardial fibrosis  
 d. Left ventricular failure
- 78. Heart failure cells are:** (Recent Question 2013)  
 a. Lipofuscin granules in cardiac cells  
 b. Pigmented alveolar macrophages  
 c. Pigmented pancreatic acinar cells  
 d. Pigment cells seen in liver

#### ISCHEMIC HEART DISEASE

- 79. Cardiac biopsy of a patient who died following myocardial infarction is shown below. What is the finding is a feature of reperfusion injury?** (Recent Pattern Question 2020)




- a. Waviness of fibers  
 b. Neutrophils in cardiac muscle  
 c. Eosinophilic contraction bands  
 d. Swelling of cells
- 80. Autopsy specimen of the heart of a patient who died due to myocardial infarction was stained with triphenyltetrazolium chloride dye. Colour of normal part of the heart will be:-** (AIIMS Nov 16)  
 a. Blue  
 b. White  
 c. Red  
 d. Dark brown





81. Given below is the histology of heart from a patient who died of Myocardial infarction. What is the time elapsed after MI? (AIIMS May 16)



- a. 1-2 days                      b. 7 days  
c. 6 hr                          d. 2-3 weeks
82. A 55 years old male presents with severe chest pain radiating to the left arm. ECG shows ST segment elevation in the V4, V5 and V6 leads. CK-MB and troponin levels are found to be increased. The most likely cause for the increase in enzyme in serum is (Recent Question 2016-17, JIPMER May 2015)
- a. Clumping of nuclear chromatin  
b. Lysosomal Autophagy  
c. Mitochondrial swelling   d. Cell membrane defects
83. Irreversible change in ischemia of heart occurs in? (Recent Question 2016-17)
- a. 10 min                      b. 30 min  
c. 60 min                      d. 90 min
84. Recurrent ischaemic events following thrombosis have been pathophysiologically linked to (Recent Question 2016)
- a. Antibodies to thrombolytic agents  
b. Fibrinopeptide A  
c. Lipoprotein A              d. Triglycerides
85. Post mortem finding in a case of death due to myocardial infarction is? (Recent Question 2015)
- a. Fat necrosis                      b. Caseous necrosis  
c. Liquefactive necrosis      d. Coagulative necrosis
86. Autopsy diagnosis of myocardial infarction can be done by immersion of tissue slices in a solution of (Recent Question 2015)
- a. Triphenyl tetrazolium chloride  
b. 100% alcohol  
c. Orcein stain  
d. Crystal violet
87. Due to ischemia, irreversible cell injury to cardiac myocytes occur in (Recent Question 2015)
- a. <2 minutes                      b. 10-20 minutes  
c. 20-40 minutes                  d. >1 hour
88. Earliest light microscopic change in myocardial infarction (Recent Question 2015)
- a. Waviness of fibres              b. Neutrophilic infiltration  
c. Coagulation necrosis          d. Contraction band necrosis
89. The cells seen after 14 hours in the infarcted area in MI are: (Recent Question 2015)
- a. Neutrophils                      b. Lymphocytes  
c. Macrophages                      d. Monocytes

90. The type of necrosis in myocardial infarction is (APPGMEE 14)

- a. Caseous                      b. Coagulative  
c. Liquefactive                  d. Fibrinoid

91. Which of the following is a non-modifiable risk factor for CHD: (Recent Question 2013)

- a. Diabetes                      b. Smoking  
c. Hypertension                  d. Old age

92. Most common site of artery of atherosclerosis: (Recent Question 2013)

- a. LAD                              b. RCA  
c. LCX                              d. Diagonal branch of LAD

93. Fatal arrhythmias are seen if myocardial infarction is: (Recent Question 2013)

- a. Posterior                      b. Inferior  
c. Anterolateral                  d. Subendocardial

94. A 45 years old male had severe chest pain and was admitted to the hospital with a diagnosis of acute myocardial infarction. Four days later he died and autopsy showed transmural coagulative necrosis. Which of the following microscopic features will be seen on further examination? (AIIMS May 11, AI 09)

- a. Fibroblast and collagen  
b. Granulation tissue  
c. Neutrophilic infiltration surrounding coagulatives  
d. Granulomatous inflammation

95. Approximate time, at the end of which the quantity of ATP within ischemic cardiac myocytes is reduced to 10% of original is: (Karn 11)


- a. <2 minutes                      b. 10 minutes  
c. 20 minutes                      d. 40 minutes

96. Autopsy finding after 12 hrs in a case of death due to M.I. (MAHE 05, DPG 10)

- a. Caseous necrosis  
b. Coagulative necrosis  
c. Fat necrosis  
d. Liquefactive necrosis

## RHEUMATIC FEVER AND RHEUMATIC HEART DISEASE

97. 30 years old male presented with severe dyspnoea and fatigue. X-ray showed left atrial enlargement. Physician suspects the patient of having mitral stenosis and gets a histopath examination done, the image of which is shown, it shows? (AIIMS Nov 2017)



- a. Sarcoidosis                      b. Tuberculosis  
c. Aschoff bodies                  d. Fungal granuloma






**98. Pathological feature(s) of rheumatic heart disease is/ are?** (PGI Nov 2017)

- a. Widened mitral annulus
- b. McCallum plaques
- c. Rupture of papillary muscle
- d. Aschoff nodules

**99. Gross findings of heart form a 18 yrs /F presented with history of sore throat 3m back & joint pains has been shown below . On auscultatory finding a murmur was noted. What is your diagnosis.** (AIIMS May 16)



- a. Libman sack endocarditis
- b. Infective endocarditis
- c. Rheumatic carditis
- d. Marantic carditis

**100. In Rheumatic carditis, Mc callums patch is seen in sub-endothelium of?** (Recent Question 2015)

- a. Right atrium
- b. Right Ventricle
- c. Left atrium
- d. Left ventricle

**101. Cells forming aschoff nodules are all the following except** (Recent Question 2015)

- a. T cells
- b. B cells
- c. Plasma cells
- d. Macrophages

**102. Causative organism of rheumatic fever** (Recent Question 2014-15)

- a. Group A Streptococci
- b. Staphylococci
- c. Group B Streptococci
- d. Group D Streptococci

**103. Anitschkow cells are found in?** (Recent Question 2014)

- a. Rheumatoid nodule
- b. Rheumatic myocarditis
- c. Bacterial Endocarditis
- d. Libman sacks endocarditis

**104. McCallum's patch is diagnostic of** (Recent Question 2013)

- a. Infective endocarditis
- b. Rheumatic endocarditis
- c. Myocardial infarction
- d. Tetralogy of Fallot (ToF)

**105. Feature of acute rheumatic fever includes:** (PGI May12)

- a. Carey coombs murmur
- b. Pancarditis
- c. Always cause residual it disease
- d. Chorea
- e. Streptococcal infection

**106. Aschoff's bodies are seen in:** (Jipmer 11)

- a. Rheumatic myocarditis
- b. Rheumatic arthritis
- c. Bacterial endocarditis
- d. Marantic endocarditis Endocarditis

**107. What is the mechanism of acute rheumatic fever** (AIIMS May 10)

- a. Cross reactivity with endogenous antigen
- b. Innocent by slender effect
- c. Due to toxin secretion by streptococci
- d. Release of pyrogenic cytokines

## ENDOCARDITIS

**108. Bulky friable vegetations are seen in:** (Recent Pattern Question 2020)

- a. Rheumatic carditis
- b. Infective endocarditis
- c. Libman-Sacks endocarditis
- d. Nonbacterial thrombotic endocarditis

**109. Libman-Sacks endocarditis most commonly causes** (Recent Question 2015)

- a. Mitral regurgitation
- b. Mitral stenosis
- c. Tricuspid regurgitation
- d. Aortic regurgitation

**110. Small warty vegetations seen on the under surfaces of AV valves, valvular endocardium, chords or mural endocardium of atria or ventricles is characteristic of**

- a. Libman sack endocarditis (Recent Question 2015)
- b. Nonbacterial thrombotic endocarditis
- c. Infective endocarditis
- d. Rheumatic fever

**111. Vegetations of the following endocarditis has the maximum chances of embolization**

- a. Rheumatic heart disease (Recent Question 2015)
- b. Infective endocarditis
- c. Libman-sacks endocarditis
- d. Subacute bacterial endocarditis

**112. Non-bacterial thrombotic endocarditis is seen in** (Recent Question 2015)

- a. Rheumatic fever
- b. Systemic lupus erythematosus
- c. Rheumatoid arthritis
- d. Mucinous adenocarcinoma of pancreas

**113. Non sterile vegetation is seen in** (Recent Question 2015)

- a. Libmann sack's endocarditis
- b. Marantic endocarditis
- c. Infective endocarditis
- d. Rheumatic heart disease

**114. Libmann sack endocarditis is seen in** (Recent Question 2015)

- a. Rheumatic fever
- b. SLE
- c. AML M3
- d. Mucinous adenocarcinoma of pancreas

**115. Which type of endocarditis has vegetation on both sides of the valves:** (Recent Question 2015)

- a. Infective endocarditis
- b. Libman Sack'endocarditis
- c. RF
- d. None

**116. Sterile vegetations are seen in all except-** (DNB Nov. 12 Pattern)

- a. SLE
- b. Infective endocarditis
- c. Rheumatic fever
- d. Marantic endocarditis



**117. Which of the following is associated with destruction of valves?** (Recent Question 2016-17)

- Acute infective endocarditis
- Libman sach's endocarditis
- Rheumatic Heart disease
- All

**118. Flat vegetations in pockets of valves are due to**

- Rheumatic heart disease (DNB Dec 11)
- Libman sacks Endocarditis
- NBTE
- Infective endocarditis

**119. In which of the following vegetation are friable and easily detachable from the cardiac valves:** (AI 10)

- Rheumatic fever
- Rheumatoid heart
- SIB
- Infective endocarditi

### CARDIOMYOPATHY

**120. Match the followings:** (AIIMS Nov 2019)

Column A	Column B
a. Box car nuclei	1. HOCM
b. Myocyte disarray	2. Hypertension
c. Vacuolation in myocytes	3. DCM
d. Myocyte hypertrophy	4. Subendothelial ischemia
	5. Hypersensitive myocarditis

**121. A patient had a quarrel with his brother with heightened emotion and falls suddenly and died. Most likely Cause of death in this case is ?** (AIIMS May 18)

- Arrhythmogenic right ventricle cardiomyopathy
- Takotsubo cardiomyopathy
- Dilated cardiomyopathy
- Chronic ischemic cardiomyopathy.

**122. Dilated cardiomyopathy, gene altered is?** (WBPGE 2016, MHPGME 2016)

- Dystrophin
- Titin
- Sarcomere
- Mitochondrial genes

**123. Pathological features of hypertrophic cardiomyopathy?**

- Death occurs in young athletes (PGI Nov 2015)
- Mostly are genetic
- LV septa is involved
- RA is involved
- Treatment is ablation of septa

**124. Histological finding of hypertrophic cardiomyopathy includes:** (PGI May 2015)

- Myocyte disarray
- Interstitial fibrosis
- Amyloid deposition in muscle
- Myocyte hypertrophy
- Myocardial fibers are arranged in parallel pattern

**125. True about features of Hypertrophic cardiomyopathy:** (PGI Nov 2011)

- Hypertrophy of ventricles without dilatation
- Myocytolysis
- Irregular arrangement of fibers
- Asymmetrical septal hypertrophy
- Myocarditis

**126. Dilated cardiomyopathy is/are seen in infection with:** (PGI May 2011)

- Ischemic heart disease
- Amyloidosis
- Viral myocarditis
- Alcoholic liver disease
- Thyroid disease

**127. Which one of the following is not a cause for restrictive cardiomyopathy:** (DNB Dec 11, AIIMS May 04)

- Alcohol
- Hemochromatosis
- Amyloidosis
- Sarcoidosis

### CARDIAC TUMORS

**128. Two most common tumors of heart in adult are:** (PGI May 2019)

- Myxoma
- Fibroma
- Angiosarcoma
- Metastatic tumors
- Rhabdomyoma

**129. Most common primary tumor of heart is?** (Recent Question 2016)

- Myxoma
- Liposarcoma
- Rhabdomyoma
- Lipoma

**130. Carcinoid of heart involves?** (Recent Question 2015)

- Valvular endocardium of right atrium
- Valvular endocardium of left atrium
- Mural endocardium
- Myocardium

**131. False statement regarding cardiac myxoma** (Recent Question 2015)

- Most common primary tumor of heart
- Most common in left atrium
- More common in females
- 90% familial

**132. Most common primary tumor of heart in children** (Recent Question 2015)

- Myxoma
- Rhabdomyoma
- Lipoma
- Fibroma

**133. Spider cells are seen in** (Recent Question 2015)

- Papillary fibroelastoma
- Tako-tsubo cardiomyopathy
- Rhabdomyoma
- Myxoma

**134. Carcinoid heart disease affects which part:** (Recent Question 2014-15)

- Valvular endocardium
- Pericardium
- Myocardium
- Epicardium

**135. Lepidic cells are characteristic of?** (AP PGME 14)

- Bronchio loaveolar carcinoma
- Mesothelioma
- small cell cancer of lung
- Myxoma of heart

**136. Characteristic pathological finding in carcinoid of heart: (AI 10)**

- Fibrous endocardial thickening of right ventricle and tricuspid valve
- Collagen deposition in wall of right ventricle
- Interstitial fibrous thickening of right ventricles
- Mononuclear inflammatory infiltrate in the wall

**137. Which carcinoma metastasizes to heart?**

(PGI May 2010)

- CA breast
- CA stomach
- CA lung
- CA urinary bladder
- Osteosarcoma

**MYOCARDITIS AND PERICARDITIS****138. Hemopericardium is seen in- (Recent Question 2014)**

- Chest injury
- MI
- Ruptured Aortic aneurysm
- All

**139. The causes of pericarditis are: (Recent Question 2013)**

- Infection
- Trauma
- Neoplasia
- Acute myocardial infarction
- Any of the above

**140. Which worm causes myocarditis:**

(Recent Question 2013)

- |                |                  |
|----------------|------------------|
| a. Trichomonas | b. Trichinella   |
| c. Enterobius  | d. Strongyloides |

**MISCELLANEOUS****141. Tigered effect in myocardium is due to:**

(Recent Question 2015)

- Malignant change
- Fatty change in heart
- Seen in rheumatic fever
- Associated with myocarditis

**142. Commonest complication of prosthetic valve is:**

(Recent Question 2015)

- Embolism
- Subacute bacterial endocarditis
- Rejection
- Infarction CNS

**143. A young female patient came for routine examination. On examination a mid systolic click was found. There is no history of RHD. The histopathological examination is most likely to show- (AIIMS May 12)**

- Myxomatous degeneration and prolapse of the valve
- Fibrinous deposition on the tip of papillary muscle
- Rupture of chordae tendinae
- Aschoff nodule on the mitral valve

**144. What does "cardiac polyp" mean? (AIIMS May 11)**

- |                  |                     |
|------------------|---------------------|
| a. Acute infarct | b. Cardiac aneurysm |
| c. Benign tumor  | d. Fibrinous clot   |

**145. True about subendocardial hemorrhage is all except-**

(AIIMS Nov 10)

- May be seen after head injury
- Involves RV wall
- Continuous pattern of sheetlike
- Flame shaped hemorrhages

**146. Factor responsible for Cardiac Hypertrophy is?**

(DNB Dec 10)

- |          |              |
|----------|--------------|
| a. ANF   | b. TNF alpha |
| c. c-myc | d. TGF beta  |

**Answers with Explanations****1. Ans. (c) Cholesterol**

(Ref: Robbins 9th/pg 496; 8th/pg 500)

Cholesterol clefts: space caused by the dissolving out of cholesterol crystals in sections of tissue embedded in paraffin.

**2. Ans. (a) Endothelium**

(Ref: Robbins 9th/pg 498-499; 8th/pg 505; Harrison 18th/pg 1503)

**Pathogenesis of atherosclerosis** best explained by the 'Response to Injury hypothesis to endothelium'

**3. Ans. (d) Mesenteric arteries**

(Ref: Robbins 9th/pg 498-499; 8th/pg 502)

Spared vessels are:

- Upper extremities vessels
- Mesenteric and renal arteries except at their ostia

**4. Ans. (c) Lipid (Ref: Robbins 9th/pg 496; 8th/pg 500)****5. Ans. (a) Chronic inflammatory disorder of vessel wall**

(Ref: Robbins 9th/pg 494; 8th/pg 500)

**6. Ans. (a) Left anterior descending artery**

(Ref: Robbins 9th/pg 498-499; 8th/pg 502)

**7. Ans. (d) Calcifications cause significant narrowing of vessel lumen**

(Ref: Robbins 9th/pg 491-492; 8th/pg 496-497)

**Mönckeberg medial sclerosis**

- Calcification of the walls (media)**<sup>Q</sup> of muscular arteries, typically involving the internal elastic membrane<sup>Q</sup>
- Do not encroach**<sup>Q</sup> on the vessel lumen

**8. Ans. (a) Oxidized LDL**

(Ref: Robbins 9th/pg 496; 8th/pg 500)

With chronic hyperlipidemia, lipoproteins accumulate within the intima, there they aggregate and become **oxidized** by **free radicals** produced by **inflammatory cells**.

These cannot be completely degraded, hence their ingestion by **macrophages/smooth muscle cells** leads to formation of **foam cells**




9. Ans. (a) **Physical inactivity**

(Ref: Robbins 9th/pg 496; 8th/pg 500)

**Physical inactivity**

10. Ans. (c) **Convex part formed by fibrous cap**

(Ref: Robbins 9th/pg 496; 8th/pg 500)



11. Ans. (d) **Endothelium**

(Ref: Robbins 9th/pg 498-499; 8th/pg 505; Harrison 18th/pg 1503)

12. Ans. (b) **Heart** (Ref: Robbins 9th/pg 498-499; 8th/pg 502)

13. Ans. (c) **Monckebergs sclerosis**

(Ref: Robbins 9th/pg 491-492)

14. Ans. (c) **Smooth muscle cell hypertrophy**

(Ref: Robbins 9th/pg 498-499; 8th/pg 505; Harrison 18th/pg 1503)

15. Ans. (b) **C. Pneumoniae** (Ref: R 9th/pg 496; 8th/pg 1182)

**Infection.** Although circumstantial evidence has been presented linking atherosclerosis to **herpesvirus, cytomegalovirus, and Chlamydothila pneumoniae**, there is no established causal role for infection.

16. Ans. (c) **Lipids in LDL get oxidized**

(Ref: Robbins 9th/pg 496; 8th/pg 500)

With chronic hyperlipidemia, lipoproteins accumulate within the intima, there they aggregate and become **oxidized** by **free radicals** produced by **inflammatory cells**.

17. Ans. (c) **Hyaline arteriosclerosis**

(Ref: Robbins 9th/pg 490, 938-939; 8th/pg 495)

- Also called obliterative endarteritis seen in syphilis/TB/radiation, poisoning, etc
- Malignant MT
- Benign HT
- MC in marfan syndrome

18. Ans. (b) **Fibrinoid necrosis**

(Ref: Robbins 9th/pg 490, 938-939; 8th/pg 495)

19. Ans. (b) **Malignant hypertension**

(Ref: Robbins 9th/pg 490; 8th/pg 495)

20. Ans. (b) **Hyperplastic arteriosclerosis**

(Ref: Robbins 9th/pg 490)

21. Ans. (a) **Intimal proliferation and hyalinization of media of medium arteries**

(Ref: Robbins 9th/pg 490; 8th/pg 495)

There is increased myofibroblastic tissue in the intima along with hyaline deposits in benign hypertension. Intimal proliferation is usually the universal response of a vessel to injury.

22. Ans. (d) **Hyperplastic arteriosclerosis**

(Ref: Robbins 9th/pg 490)

23. Ans. (a) **Ascending aorta**

(Ref: Robbins 9th/pg 503)

Syphilis usually affects proximal ascending aorta (esp aortic ring) > aortic arch.

24. Ans. (a) **Trauma**

(Ref: Robbins 9th/pg 501-502; 8th/pg 506-507)

25. Ans. (a) **Ascending aorta** (Ref: Robbins 9th/pg 501-502)

MC cause of  
**ascending aortic  
aneurysm**

**Marfan's Syndrome**  
Cystic medial degeneration<sup>a</sup> / Systemic  
hypertension<sup>a</sup>

26. Ans. (d) **5.5 cm**

(Ref: Guidelines for the treatment of abdominal aortic aneurysms. J Vasc Surg. 2003;37(5):1106)

Aneurysm repair is the primary treatment for aneurysms that are symptomatic or at a high risk for rupture. Most people with an aneurysm less than 4.0 cm (1.6 inches) in diameter are advised not to have immediate surgery, but rather to follow the aneurysm over time; this is known as watchful waiting.

On the other hand, most patients with an **asymptomatic aneurysm greater than 5.5 cm (2.2 inches) in diameter or that expands more than 0.5 cm within a six-month period** are advised to have repair.

27. Ans. (c) **7 cm**

(Ref: Guidelines for the treatment of abdominal aortic aneurysms. J Vasc Surg. 2003;37(5):1106)

The annual risk of rupture based upon aneurysm size is estimated as follows:

- Less than 4.0 cm in diameter = ~0%
- Between 4.0 to 5 cm in diameter = 1%
- Between 5.0 to 6 cm in diameter = 11%
- > 6.0 cm in diameter = 25%





28. Ans. (b) **Atherosclerosis** (Ref: Robbins 9th/pg 501-502)

29. Ans. (d) **Mockenbergs sclerosis**

(Ref: Robbins 9th/pg 501-502; 8th/pg 506-507)

30. Ans. (a) **Splenic**

(Ref: Semin Intervent Radiol. 2009 Sep; 26(3): 196-206)

- Visceral artery aneurysms (VAAs) and visceral artery pseudoaneurysms (VAPAs) frequently present as life-threatening emergencies
- **Splenic artery aneurysms- most common VAA**
- **Hepatic artery aneurysms (HAAs)- second most common VAA**
- Clinically, the patients with VAPAs, typically present with an antecedent history of arterial trauma or surgical manipulation

31. Ans. (c) **Transesophageal echocardiography**

(Ref: Radiographics.rsna.org MARCH-APRIL 2010, Circulation: Cardiovascular Imaging (November 2009 vol. 2 no. 6 499-506, Harrison 17th ed table 242A))

The dissection is termed *acute* when it is diagnosed **within 14 days** after the first symptoms appear

	Unstable/Critical Conditions	Stable Clinical Condition
Diagnostic modality	TEE with color Doppler	MD-CT with CTA (CT angiography) or MRI with MRA

32. Ans. (c) **Hypertension** (Ref: Robbins 9th/pg 504; 8th/pg 509)

**MC cause of dissection is hypertension<sup>Q</sup>**

33. Ans. (d) **Secondary syphilis**

(Ref: Robbins 9th/pg 504; 8th/pg 509)

**MC cause of dissection is hypertension, seen in men between 40-60 yrs of age.**

**2<sup>nd</sup> MC cause of dissection is cystic medial necrosis (CMN) Marfan's syndrome causes CMN hence dissection**

**Syphilis- Aneurysm of the aorta was the most common complication of syphilitic aortitis.**

**Syphilis will only potentially cause aortic dissection in its tertiary stage**

34. Ans. (b) **Level of aorta affected**

(Ref: Robbins 9th/pg 504)

35. Ans. (a) **Marfans** (Ref: Harrison 18th pg 206)

**Cystic medial necrosis (CMN)**

- Disorder of **large arteries, in particular the aorta<sup>Q</sup>**
- Characterized by an accumulation of **basophilic ground substance in the media<sup>Q</sup>** with cyst-like lesions.
- Diseases causing CMN
  - **Marfan's syndrome<sup>Q</sup>, Chronic aortic dissection<sup>Q</sup>, Congenital heart disease especially bicuspid aortic valve<sup>Q</sup>, Scurvy<sup>Q</sup>, Aortic aneurysm<sup>Q</sup>, Atherosclerotic disease<sup>Q</sup>, Hypertension<sup>Q</sup>, Ehler danlos syndrome (type IV)<sup>Q</sup>**

36. Ans. (a) **Hypertension**

(Ref: Robbins 9th/pg 504; 8th/pg 509)

**MC cause of dissection is hypertension, seen in men between 40-60 yrs of age.**

**2<sup>nd</sup> MC cause of dissection is cystic medial necrosis. (option B).**

37. Ans. (e) **IgA vasculitis**

38. Ans. (c) **Medium to small vessel**

Takayasu arteritis involve large vessels

39. Ans. (a) **Thrombocytopenia**

(Ref: Robbins 9th/pg 510; 8th/pg 515)

40. Ans. (d) **Lymphatics**

(Ref: Robbins 9th/pg 512; 8th/pg 517)

- **Buerger's disease: Segmental, thrombosing, acute and chronic inflammation of medium-sized and small arteries, principally the tibial and radial arteries, with occasional secondary extension into the veins and nerves of the extremities**

41. Ans. (a) **IgA** (Ref: Robbins 9th/pg 926; 8th/pg 934)

- Most common antibody seen in these immune complexes is **IgA<sup>Q</sup>** (IgA levels are elevated)
- The **microscopic hallmark** of HSP is the deposition of **IgA in the walls<sup>Q</sup>** of involved blood vessels.

42. Ans. (d) **Classical PAN**

(Ref: Robbins 9th/pg 506; 8th/pg 511)

43. Ans. (c) **Thrombocytopenia**

(Ref: Robbins 9th/pg 926; 8th/pg 934)

- **HSP: Purpura not due to a low platelet count but due to vasculitis<sup>Q</sup>**
- Platelet count is normal or elevated<sup>Q</sup>

44. Ans. (b) **Wegener's Granulomatosis**

(Ref: Robbins 9th/pg 507, Harrison 18th ed p-2786-87)

- **C-ANCA (proteinase 3<sup>Q</sup> is the target antigen)**
- Typically seen in Wegner's Granulomatosis<sup>Q</sup>

45. Ans. (a) **Temporal arteritis**

(Ref: Robbins 9th/pg 507-8,510)

- **Biopsy and histological confirmation** of temporal artery is the **investigation of choice.**
- It shows **Granulomatous inflammation with giant cells & fragmentation of internal elastic lamina.<sup>Q</sup>**

46. Ans. (b) **Low ESR** (Ref: R 9th/pg 507-8,510; 8th/pg 512-13)

47. Ans. (d) **Sarcoidosis**

(Ref: Robbins 9th/pg 507-8,510; 8th/pg 512-13)

**Fibrinoid necrosis is seen in Pan**

- Aschoff Nodule, SLE, HIV and Malignant Hypertension



**48. Ans. (a) Behçet's disease**

(Ref: Robbins 9th/pg 511, www.behcets.com/american behcet disease association)

Behçet's disease is considered more prevalent in the areas surrounding the old silk trading routes in the Middle East and in Central Asia. Thus, it is sometimes known as **Silk Road Disease**

Linkage between the disease and **HLA-B51** is seen.

**49. Ans. (d) Involves large vessels** (Ref: R 9th/pg 511-512)

**50. Ans. (c) 40-60%**

(Ref: Hepinstall pathology of kidney volume 1- pg 463, JASN May 1, 2002 vol. 13 no. 5 1271-1278)

- **Incidence of Renal Involment in HSP (different in various studies, depends on data of patients and the definition of renal involvement)**
- **Adults : 45 to 85%**
- **Children-20-56%**

**51. Ans. (c) Wegener's granulomatosis**

(Ref: Robbins 9th/pg 507, Harrison 18th ed p-2786-87)

Option a and b are **immune-mediated small-vessel systemic vasculitis**. **C-ANCA positive vasculitis**

**52. Ans. (a) PAN** (Ref: Robbins 9th/pg 509; 8th/pg 511)

**PAN** is characterized by **segmental transmural necrotizing inflammation**<sup>Q</sup> frequently accompanied by **fibrinoid necrosis**.<sup>Q</sup>

OPTION B, C, D are granulomatous vasculitis.

**53. Ans. (c) Antigen-antibody complex mediated**

(Ref: Robbins 9th/pg 509; 8th/pg 511)

PAN-Necrotizing vasculitis with fibrinoid necrosis which is a type of **immune complex mediated vasculitis**<sup>Q</sup>

**54. Ans. (c) Classic PAN** (Ref: Robbins 9th/pg 506-7,510)

**55. Ans. (c) Most commonly involved artery is abdominal aorta** (Ref: Robbins 9th/pg 507-8,510; 8th/pg 512-13)

**56. Ans. (c) Proteinase 3**

(Ref: Robbins 9th/pg 507, Harrison 18th ed p-2786-87)

- **C-ANCA (proteinase 3<sup>Q</sup> is the target antigen)**
- Typically seen in Wegner's Granulomatosis<sup>Q</sup>

**57. Ans. (d) Unlikely to cause pulmonary renal syndrome**

(Ref: Robbins 9th/pg 510-11; 8th/pg 515)

Microscopic polyangiitis (MP)

- **Option a true- MP is small vessel vasculitis**
- Typically **spare medium-sized and larger arteries**
- **Option b is true- P-ANCA** is present in majority of the patients.
- **Option c is true- major clinical features include hemoptysis, hematuria and proteinuria, bowel pain or**

**bleeding, muscle pain or weakness, and palpable cutaneous purpura.**

**Skin findings are as follows:**

- **Palpable purpura** (41%), Leukocytoclastic angitis, livedo reticularis (12%), skin ulcerations, necrosis and gangrene, necrotizing nodules and digital ischemia (7%)
- **Urticaria** - Vasculitis-associated urticaria that lasts longer than 24 hours **Option d is false- Necrotizing glomerulonephritis** (90% of patients) and **pulmonary capillaritis** are common.

**58. Ans. (a) Lung** (Ref: Robbins 9th/pg 509-510; 8th/pg 514-5)

PAN Typically involving renal and visceral vessels but **sparing the pulmonary circulation.**

**59. Ans. (b) Focal necrotizing glomerulonephritis**

(Ref: Robbins 9th/pg 511-512; 8th/pg 516)

- **Wegner's granulomatosis** now called **Granulomatosis with Polyangiitis**
- Focal necrotizing, often crescentic, glomerulonephritis along with Necrotizing or granulomatous vasculitis affecting small to medium-sized vessels.

Option a- granulomas in vessel wall alone is not characteristic of Wegner's. It should be granulomatous or necrotizing vasculitis

**60. Ans. (b, c) b. Kawasaki disease; c. Henoch schonlein purpura** (Ref: Robbins 9th/pg 508,510-12; 8th/pg 511-13)

- Option a - Giant cell arteritis-vasculitis in a patient over the age of 50 years
- Option b - **Kawasaki disease**-vasculitis in **infancy and childhood**
- Option c - **Henoch schonlein purpura**-vasculitis in children
- Option d- Polyarteritis nodosa-vasculitis in 4th or 5th decade

**61. Ans. (a) Hemangioma** (Ref: Robbins 9th ed p 516)

Hemangiomas are very common tumors characterized by increased numbers of normal or abnormal vessels filled with blood. These lesions constitute 7% of all benign tumors of infancy and childhood; most are present from birth and initially increase in size, but many eventually regress spontaneously. While hemangiomas typically are localized lesions confined to the head and neck, they can occasionally be more extensive (angiomatosis) and can occur internally.

**62. Ans. (a, b, c, d); a. Causative agent HHV 8, b. Can be Seen in depressed cell mediated immunity, c. Treatment is chemotherapy and surgery, d. Vascular tumor**

(Ref: Robbins 9th/pg 518.; 8th/pg 523)

**KAPOSI SARCOMA**

- It is caused by KS Herpes virus or **Human herpes virus 8 (HHV8)**-**option a is true**
- Characterized by proliferation of **spindle cells**<sup>Q</sup> which are of **vascular origin**<sup>Q</sup> (Option d is correct)



Transplant and HIV associated Kaposi usually occur in immunosuppressed individuals. (Option b is correct)

Treatment - (Option c is correct)

- HAART is the best way to treat HIV associated Kaposi's sarcoma
- **Topical retinoid treatment**
- Cryosurgery (cryotherapy)
- Surgery; Radiation therapy
- Intralesional chemotherapy

63. Ans. (d) **Vascular ectasias** (Ref: Robbins 9th/pg 515-516)

**Port-wine stain or nevus flammeus**

- It is the **most common form** of **Vascular ectasias**; it persist throughout life<sup>Q</sup>
- Caused by a somatic activating c.548G→A mutation in the **GNAQ<sup>Q</sup>** gene
- Part of Sturge-Weber syndrome<sup>Q</sup> or Klippel-Trénaunay-Weber syndrome

64. Ans. (c) **Kaposi sarcoma** (Ref: Robbins 9th/pg 518)

65. Ans. (b) **HAART therapy has decreased AIDS associated KS**

(Ref: Robbins 9th/pg 518.; 8th/pg 523)

66. Ans. (d) **Hemangiopericytoma**

(Ref: Sternberg's Diagnostic Surgical Pathology, 5th Edition, table 5 11)

**Hemangiopericytoma**

- Tumor derived from **pericytes<sup>Q</sup>**- **perivascular cells** that wrap around blood capillaries
- These tumors most commonly arise from **pelvic retroperitoneum<sup>Q</sup>** or the **limbs<sup>Q</sup>** (particularly thighs).
- Capillaries are arranged in '**fish-hook pattern**;<sup>Q</sup> seen best with silver stains<sup>Q</sup>

67. Ans. (b) **Capillary hemangioma**

(Ref: Robbins/pg 515-516)

**PYOGENIC GRANULOMAS**-lobular capillary hemangioma

**Capillary hemangiomas<sup>Q</sup>** that grows rapidly

Presents as red pedunculated lesions on the **skin, gingival, or oral mucosa<sup>Q</sup>**

Lesions **bleed easily<sup>Q</sup>**

68. Ans. (a) **Liver**

(Ref: Robbins 9th/pg 519; 8th/pg 523)

**Angiosarcoma**

- Malignant endothelial cell neoplasm most commonly seen in **skin, soft tissue, breast and liver<sup>Q</sup>**
- **Hepatic angiosarcoma** is associated with carcinogens including arsenic, thorotrast (a radioactive contrast) and polyvinyl chloride (PVC; a plastic)

69. Ans. (d) **CD 31** (Ref: Robbins 9th/pg 591,519.; 8th/pg 523)

- Endothelial cell origin of angiosarcoma is demonstrated by staining for **CD31, CD34 or VWF<sup>Q</sup>**.
- CD10 - CALLA (common acute lymphoblastic leukemia antigen)

- CD19- pan B marker
- CD25- alpha chain of the IL-2 receptor

70. Ans. (b) **Strawberry angioma**

(Ref: Robbins 9th/pg 515-516)

Lesion	Features
Capillary hemangiomas	<ul style="list-style-type: none"> <li>• <b>Strawberry hemanigioma</b> or <b>juvenile hemangioma</b> is a type</li> <li>• <b>Completely regress</b></li> </ul>
<b>Port-wine stain or nevus flammeus</b>	<ul style="list-style-type: none"> <li>• Caused by a somatic activating c.548G→A mutation in the <b>GNAQ<sup>Q</sup></b> gene</li> <li>• <b>Persist throughout life<sup>Q</sup></b></li> </ul>
Cavernous hemangiomas	<ul style="list-style-type: none"> <li>• <b>Do not<sup>Q</sup></b> spontaneously regress.</li> </ul>

71. Ans. (c, d); c. **Pressure effect is often the presenting symptom**; d. **Thromboembolic phase is present**


(Ref: Neurosurgery [SHC Suppl 3]:SHC1289-SHC1299, 2008, Harrison 18th ed: 2262)

**GIANT ANEURYSMS**

- Definition
  - Diameter of greater than 25 mm
- Most common clinical presentation
  - Mass effect- most common
  - Subarachnoid hemorrhage, intracerebral hemorrhage, or both -
- Most common site
  - Internal carotid artery segments vertebrobasilar region- OPTION b is false
- Other features
  - Seizures
  - High risk for rupture- OPTION A is false
  - **Intraluminal thrombus-occlusion of perforating vessels**
  - **Distal embolic events are common**- OPTION d is true.

72. Ans. (a) **Primary lymphedema**

(Ref: Harrison 17th ed table 243-3)



73. Ans. (b) **Increased LV cavity size**

(Ref: Robbins 9th/pg 526)



- 74. Ans. (a) No change**  
(Ref: Harrison 17th ed Table 243-1)  
Most common cause of Raynaud's Phenomenon -Primary or idiopathic.  
In initial stages of Raynaud's phenomenon no changes is seen late stages may show atherosclerosis or thrombosis
- 75. Ans. (b) Systemic sclerosis**  
(Ref: Harrison 17th ed Table 243-1, Robbins 9th/pg 513)
- 76. Ans. (c) Lungs**  
(Ref: Robbins 9th/pg 529; 8th/pg 535)  
Morphology in Lungs in LVF  
**Lungs: Most commonly** affected organ  
**Heart failure cells**<sup>Q</sup> -hemosiderin containing macrophages.  
Edematous widening of alveolar septa and -Edema fluid in the alveolar spaces.
- 77. Ans. (a) Cor pulmonale** (Ref: R 9th/pg 530; 8th/pg 536)  
**Isolated right-sided heart failure**<sup>Q</sup> is very rare,, occurs with lung disorders. This is referred to as cor pulmonale<sup>Q</sup>  
The **common feature** of these disorders is **pulmonary hypertension**<sup>Q</sup>
- 78. Ans. (b) Pigmented alveolar macrophages**  
(Ref: Robbins 9th/pg 529; 8th/pg 535)
- 79. Ans. (c) Eosinophilic contraction bands**  
(Ref: R 9th pg 544)
- 80. Ans. (b) White**  
(Ref: Robbins 9th/pg 544; 8th/pg 550)  
  - **Necrotic area** can be **visualized after 2-3 hours**<sup>Q</sup> by immersion in **triphenyltetrazolium chloride (TTC)**<sup>Q</sup>
  - **Infarcted area** is revealed as **unstained pale zone**<sup>Q</sup>
  - TTC imparts brick red colour to **non-infarcted myocardium**<sup>Q</sup> where **dehydrogenase enzymes**<sup>Q</sup> are preserved.
- 81. Ans. (a) 1-2 days**  
(Ref: Robbins 9th/pg 544)  
Image is neutrophil infiltrate which is seen 2-24 hr after infarction
- 82. Ans. (d) Cell membrane defects**  
Cell membrane damage leads to leakage of enzymes from the cell; which can be detected in serum.
- 83. Ans. (b) 30 min**  
(Ref: Robbins 9th/pg 544)  
  - **Time for reversible injury in heart is 30 mins**<sup>Q</sup>
- 84. Ans. (c) Lipoprotein A**  
(Ref: Circulation June 29, 2004 vol. 109)  
**Novel Biomarkers in the Prediction of Future Cardiovascular Events**  
Lipoprotein(a)

Homocysteine  
High-sensitivity C-reactive protein (hsCRP)

- 85. Ans. (d) Coagulative necrosis** (Ref: Robbins 9th/pg 544)  
The type of necrosis in myocardial infarction is - **Coagulative**
- 86. Ans. (a) Triphenyl tetrazolium chloride**  
(Ref: Robbins 9th/pg 544)  
(Refer Answer 121)
- 87. Ans. (c) 20-40 minutes** (Ref: R 9th/pg 544; 8th/pg 550)
- 88. Ans. (a) Waviness of fibres**  
(Ref: Robbins 9th/pg 544; 8th/pg 550)  
**Waviness of fibers at border**<sup>Q</sup> (earliest microscopic change)
- 89. Ans. (a) Neutrophils**  
(Ref: Robbins 9th/pg 544; 8th/pg 550)
- 90. Ans. (b) Coagulative** (Ref: Robbins 9th/pg 539-40)
- 91. Ans. (d) Old age** (Ref: Robbins 9th/pg 492; 8th/pg 497)
- 92. Ans. (a) LAD** (Ref: Robbins 9th/pg 542; 8th/pg 549)  
Most common site of artery of atherosclerosis: LAD
- 93. Ans. (b) Inferior** (Ref: Robbins 9th/pg 549; 8th/pg 555)  
"Location of portions of the atrioventricular conduction system (bundle of His) in the inferoseptal myocardium, infarcts of this region may also be associated with heart block."
- 94. Ans. (b) Granulation tissue**  
(Ref: Robbins 9th/pg 544; 8th/pg 550)
- 95. Ans. (d) 40 minutes** (Ref: Robbins 9th/pg 541; 8th/pg 550)
- 96. Ans. (b) Coagulative necrosis**  
(Ref: Robbins 9th/pg 544)
- 97. Ans. (c) Aschoff bodies**
- 98. Ans. (b, d) b. McCallum plaques; d. Aschoff nodules**  
(Ref: R 559)  
  - **Aortic valve thickening**
The cardinal anatomic changes of the mitral valve in chronic RHD are leaflet thickening, commissural fusion and shortening, and thickening and fusion of the tendinous cords along with increase in calcific aortic stenosis.
- 99. Ans. (c) Rheumatic carditis**





100. Ans. (c) **Left atrium**

(Ref: Robbins 9th/pg 557-559; 8th/pg 565-66)

**Mac-callum patches**-map-like areas of **thickened and wrinkled part of the endocardium in the left atrium**<sup>Q</sup>

- Caused by **regurgitant jets of blood flow**<sup>Q</sup>, due to incompetence of the mitral valve
- Seen in **Rheumatic endocarditis**

101. Ans. (b) **B cells**

(Ref: Robbins 9th/pg 557-559; 8th/pg 565-66)

102. Ans. (a) **Group A Streptococci** (Ref: R 9th/pg 557-559)

*Rheumatic heart disease is a* An acute **immunologically mediated multisystem inflammatory disease**<sup>Q</sup> that occurs **few weeks** after an attack of **group A β-hemolytic streptococcal pharyngitis**

103. Ans. (b) **Rheumatic myocarditis** (Ref: R 9th/pg 557-559)

104. Ans. (b) **Rheumatic endocarditis** (Ref: R 9th/pg 557-559)

105. (a, b, d, e); a. **Carey coombs murmur**; b. **Pancarditis**; d. **Chorea**; e. **Streptococcal infection**

(Ref: Robbins 9th/pg 557-559)

**Option a-true: Carey coombs murmur**- occurs in patients with mitral valvulitis due to acute rheumatic fever

**Option b-true Pancarditis**

**Option c-false: Migratory polyarthritides**

- Subsides **spontaneously without any residual deformability**<sup>Q</sup> in the joints (non-erosive arthritis).

**Option d-true: Sydenham's chorea**

**Late manifestation of the disease**

106. Ans. (a) **Rheumatic myocarditis** (Ref: R 9th/pg 557-559)

107. Ans. (a) **Cross reactivity with endogenous antigen**

(Ref: Harrison 18th ed:2552; Robbins 9th/pg 557-559; 8th/pg 565-66)

108. Ans. (b) **Infective endocarditis** (Ref: R 9th pg 560)

109. Ans. (a) **Mitral regurgitation**

(Ref: Robbins 9th/pg 560; 8th/pg 567)

110. Ans. (a) **Libman sack endocarditis** (Ref: R 9th/pg 560)

111. Ans. (b) **Infective endocarditis** (Ref: Robbins 9th/pg 560)

112. Ans. (d) **Mucinous adenocarcinoma of pancreas**

(Ref: Robbins 9th/pg 560; 8th/pg 567)

113. Ans. (c) **Infective endocarditis** (Ref: Robbins 9th/pg 560)

114. Ans. (b) **SLE** (Ref: Robbins 9th/pg 560; 8th/pg 567)

115. Ans. (b) **Libman-Sacks endocarditis** (Ref: R 9th/pg 560)

116. Ans. (b) **Infective endocarditis** (Ref: Robbins 9th/pg 560)

117. Ans. (d) **All** (Ref: Robbins 9th/pg 560,562; 8th/pg 567,569)

- Maximum valve damage:** Acute infective endocarditis
- Libman sack's endocarditis - Associated with an **intense valvulitis**<sup>Q</sup>, characterized by **fibrinoid necrosis**<sup>Q</sup> of the **valve substance**
- Rheumatic Heart disease - The valves show leaflet thickening, commissural fusion and shortening, thickening & fusion of the tendinous cords seen in chronic RHD.

118. Ans. (b) **Libman sacks Endocarditis** (Ref: R 9th/pg 560)

119. Ans. (d) **Infective endocarditis** (Ref: Robbins 9th/pg 560)

120. Ans. (a) 2, (b) 1, (c) 4, (d) 3 (Ref: R 9th pg/568-569)

Myocyte hypertrophy seen in hypertension is best evaluated in correlation with heart size. The classic histologic description is rectangular, hyperchromatic nuclei, often called "box-car" nuclei. In HOCM, histological features include cardiomegaly with left ventricular hypertrophy, hypertrophic myocytes, and myocyte disarray and/or myofiber bundle disorder. Myocyte degeneration with vacuolation is a feature of Arrhythmogenic cardiomyopathy and subendocardial ischemia. The diagnosis of dilated cardiomyopathy is not made histologically. The microscopic findings are very nonspecific and consist of myocyte hypertrophy and myocardial fibrosis.

121. Ans. (b) **Takotsubo cardiomyopathy**

122. Ans. (b) **Titin**

(Ref: Robbins 9th/pg 565-566; 8th/pg 573-574)

- DCM is familial in at least 30% to 50% of cases
- Autosomal dominant inheritance is the predominant pattern;
- Most common- mutations in TTN, a gene that encodes titin (20% of all cases of DCM)

123. Ans. (a, b, c, e) a. **Death occurs in young athletes**; b. **mostly are genetic**; c. **LV septa is involved**; e. **treatment is ablation of septa**

(Ref: Robbins 9th/pg 568-569; 8th/pg 575-576)

124. Ans. (a, b, d); a. **Myocyte disarray**; b. **Interstitial fibrosis**; d. **Myocyte hypertrophy**

(Ref: Robbins 9th/pg 568-569)

The **histologic features** of HOCM myocardium are

- Massive myocyte hypertrophy, transverse myocyte diameters **greater than 40 μm**<sup>Q</sup> (normal, approximately 15 μm);
- Myofiber disarray**<sup>Q</sup>
- Interstitial and replacement **fibrosis**<sup>Q</sup>



**125. Ans. (a, c, d); a. Hypertrophy of ventricles without dilatation; c. Irregular arrangement of fibers; d. Asymmetrical septal hypertrophy**

(Ref: Robbins 9th/pg 568-569)

**126. Ans. (c, d, e); c. Viral myocarditis; d. Alcoholic liver disease; e. Thyroid disease** (Ref: Robbins 9th/pg 565-566)

**Causes of Phenotype**

- Genetic; **alcoholic**<sup>o</sup>; **peripartum**<sup>o</sup>; **myocarditis**<sup>o</sup>; **hemochromatosis**<sup>o</sup>; chronic anemia; doxorubicin (Adriamycin) toxicity; sarcoidosis<sup>o</sup>; idiopathic

**127. Ans. (a) Alcohol** (Ref: R 9th/pg 565-566; 8th/pg 573-574)

**128. Ans. (a) Myxoma; (d) Metastatic tumors**

(Ref: R 9th pg 575)

**129. Ans. (a) Myxoma** (Ref: Robbins 9th/pg 575; 8th/pg 583)

**130. Ans. (a) Valvular endocardium of right atrium**

(Ref: Robbins 9th/pg 562; Heart 2004;90:1224-1228 doi:10.1136/hrt.2004.040329)

- **Carcinoid heart disease**<sup>o</sup> typically causes abnormalities of the **right side of the heart**.<sup>o</sup>
- Preferential **right heart involvement**<sup>o</sup> is most likely related to **inactivation of the vasoactive substances by the lungs**<sup>o</sup>
- The two key investigations for the diagnosis of carcinoid heart disease are **24 hour urinary excretion of 5-hydroxyindole acetic acid (5-HIAA)** and **transthoracic echocardiography**.

**131. Ans. (d) 90% familial**

(Ref: Robbins 9th/pg 575; 8th/pg 583)

Option a –true- Myxomas are the most common primary tumor of the adult heart

Option b –true- About 90% of myxomas arise in the atria, with a left-to-right ratio of approximately 4 : 1.

Option c –true- Approximately 75% of sporadic myxomas occur in females

Option d-false- Most cardiac myxomas are sporadic and arise as isolated masses in the left atrium.

Familial syndromes associated with myxomas have activating mutations in the GNAS1 gene or null mutations in PRKARIA, encoding a regulatory subunit of a cyclic-AMP-dependent protein kinase (Carney complex).

**132. Ans. (b) Rhabdomyoma** (Ref: Robbins 9th/pg 575)

**133. Ans. (c) Rhabdomyoma** (Ref: R 9th/pg 575; 8th/pg 583)

**134. Ans. (a) Valvular endocardium**

(Ref: Robbins 9th/pg 562; Heart 2004;90:1224-1228 doi:10.1136/hrt.2004.040329)

**135. Ans. (d) Myxoma of heart**

(Ref: Robbins 9th/pg 575; 8th/pg 583)

**136. Ans. (a) Fibrous endocardial thickening of right ventricle and tricuspid valve**

(Ref: Robbins 9th/pg 562; Heart 2004;90:1224-1228 doi:10.1136/hrt.2004.040329)

**Carcinoid heart disease:**

- **Characteristic pathological findings** are **endocardial plaques of fibrous tissue** that may involve the **tricuspid valve**<sup>o</sup>, **pulmonary valve**<sup>o</sup>, cardiac chambers, venae cavae, pulmonary artery, and coronary sinus.

**137. Ans. (a, c) a. CA breast; c. CA lung**

(Ref: Cancers and heart by Reynolds 2<sup>nd</sup> pg 316)

**Tumors that are likely to involve heart and pericardium**

- |                |             |
|----------------|-------------|
| • Ca lung      | • Ca breast |
| • Ca esophagus | • Melanoma  |
| • Lymphoma     | • Leukemia  |

**138. Ans. (d) All**

(Ref: Robbins 9th/pg 573-574; 8th/pg 581-82)

**Hemopericardium: Causes**

- Cardiac rupture after transmural myocardial infarction (especially day 5)
- Aortic aneurysm rupture, chest trauma, anticoagulation, leukaemia and TB Pericarditis

**139. Ans. (e) Any of the above**

(Ref: Robbins 9th/pg 573-574)

**140. Ans. (b) Trichinella**

(Ref: Robbins 9th/pg 570-71; 8th/pg 578)

**MC helminth causing myocarditis- trichinella<sup>o</sup>**

**141. Ans. (b) Fatty change in heart**

Fatty change of heart: band of yellow (fatty) myocardium along with red (normal myocardium)

**142. Ans. (a) Embolism** (Ref: Robbins 9th/pg 563; 8th/pg 570)

Thromboembolism is the commonest complication of prosthetic valves.

**143. Ans. (a) Myxomatous degeneration and prolapsed of valve**

(Ref: Robbins 9th/pg 556; 8th/pg 563)

**Mitral valve prolapse (MVP)**

- One or both mitral valve leaflets are “floppy” and prolapse, or balloon back, into the left atrium during systole
- Associated with Marfan syndrome, caused by fibrillin-1 (FBN-1) mutations
- Key histologic change in the tissue is marked thickening of the spongiosa layer with deposition of mucoid (myxomatous) material, called myxomatous degeneration
- Most individuals diagnosed with MVP are asymptomatic
- Discovered incidentally by auscultation of mid-systolic clicks, sometimes followed by a mid to late systolic murmur.



**144. Ans. (d) Fibrinous clot**

(Ref: Textbook of Practical Histology 2<sup>nd</sup> pg 28-29)

**Post mortem fibrinous clot of heart is called cardiac polyp**

**145. Ans. (b) Involves RV wall**

(Ref: Forensic Pathology Reviews Volume 2, 2005, pp 293-306)

**Subendocardial haemorrhage**

<b>Causes</b>	<ul style="list-style-type: none"> <li>• Cardiac injuries and resuscitation</li> <li>• Secondary to noncardiac injuries comprising <b>head injuries</b>, infectious diseases, intoxications, hemorrhagic diathesis, abdominal trauma, asthma, and <b>hypovolemic shock</b><sup>a</sup>.- <b>option a -true</b></li> </ul>
<b>Pathophysiology</b>	<ul style="list-style-type: none"> <li>• Mediated by the autonomic nervous system via hypersecretion of catecholamines<sup>a</sup></li> </ul>
<b>MC sites</b>	<ul style="list-style-type: none"> <li>• Located in the <b>upper part of the interventricular septum</b>, the opposing papillary muscles, and adjacent trabeculae carneae of the <b>free wall of the left ventricle</b><sup>a</sup></li> </ul>
<b>Pattern</b>	<ul style="list-style-type: none"> <li>• <b>Flame shaped, confluent. Sheet like</b><sup>a</sup>.- <b>option c and d -true</b></li> </ul>

**146. Ans. (c) c-myc** (Ref: Robbins 9th/pg 527; 8th/pg 532)

**Cardiac Hypertrophy: Pathophysiology**

**Molecular changes include the expression of immediate-early genes (e.g., FOS, JUN, MYC, and EGR1)**

# Respiratory System and its Disorders

## Key Points

- » **Respiratory system develops** from: **ventral wall of foregut**<sup>o</sup>
- » Sequence of division: Trachea → 2 bronchi → **Bronchioles** → **Terminal Bronchiole** → **Respiratory Bronchiole** → **Alveolar duct** → **Alveolar Sac**<sup>o</sup> (**B-T-R-S** – “**Bu T teRS**”)
- » **Bronchi** have **cartilage** and abundant **subepithelial glands** that produce mucus
- » **Bronchioles: Do not** have **cartilage**<sup>o</sup>, **goblet cells** and **submucosal glands**<sup>o</sup>
- » **Acinus**: Part of the lung **distal to terminal bronchiole**<sup>o</sup> (composed of respiratory bronchioles, alveolar ducts, and alveolar sacs)
- » **Pulmonary lobule**<sup>o</sup>: Cluster of **3 to 5 terminal bronchioles**
- » Entire respiratory tree (larynx, trachea & bronchioles), is lined by **pseudostratified**<sup>o</sup>, **tall, columnar and ciliated**<sup>o</sup> **epithelial cells**
- » Exception to above rule are **vocal cords**: lined by **stratified squamous epithelium**.<sup>o</sup>
- » Bronchial mucosa also contains neuroendocrine cells that have neurosecretory-type granules releasing: **serotonin**<sup>o</sup>, **calcitonin**<sup>o</sup> and **gastrin-releasing peptide (bombesin)**<sup>o</sup>

## Key Recent Updates

- » Basaloid carcinoma is new subtype added to SCC lung
- » NUT carcinoma of lung is very aggressive neoplasm of lung.






## NORMAL ANATOMY

### Microscopic Structure of the Alveolar Wall

- Alveolar epithelium:
  - Type I pneumocytes** (95%)<sup>Q</sup> flattened **respiratory cells**
  - Type II pneumocytes**: Synthesize **surfactant**,<sup>Q</sup> **repair type I cells**
- Alveolar macrophages**: Loosely attached or lying free within the alveolar spaces
- Pores of Kohn**<sup>Q</sup>: Permit the passage of **bacteria** and **exudate** between adjacent alveoli.
- Basement membrane and interstitial tissue
- Network of anastomosing capillaries lined with **endothelial cells**



Alveolar structure

## CONGENITAL MALFORMATIONS OF LUNGS

<b>Pulmonary hypoplasia</b>	Defective development of <b>one or both lungs</b>				
<b>Foregut cysts</b>	<ul style="list-style-type: none"> <li><b>Abnormal detachments of primitive foregut</b></li> <li>Most often located in the <b>hilum</b><sup>Q</sup> or middle <b>mediastinum</b>.</li> <li>3 types: <b>Bronchogenic (most common)</b>,<sup>Q</sup> Esophageal or Enteric</li> </ul>				
<b>Sequestration</b>	<ul style="list-style-type: none"> <li>Discrete area of <b>lung tissue that lacks any connection</b><sup>Q</sup> to the airway system</li> <li>Abnormal <b>blood supply arising from aorta</b><sup>Q</sup></li> <li><b>2 types of Pulmonary sequestration</b>:               <table border="1"> <thead> <tr> <th>Extralobar sequestration</th><th>Intralobar sequestration</th></tr> </thead> <tbody> <tr> <td> <ul style="list-style-type: none"> <li><b>External</b> to lungs</li> <li>With pleural cover</li> <li>Causing mass effect.</li> <li><b>Venous return</b> to <b>right</b> side of heart through <b>IVC</b></li> </ul> </td><td> <ul style="list-style-type: none"> <li><b>More common</b><sup>Q</sup></li> <li>Occur <b>within</b> the lung <b>without pleural cover</b></li> <li>Localized <b>infection</b> or <b>bronchiectasis</b><sup>Q</sup></li> <li>Associated with diaphragmatic hernia, colonic duplication, vertebral abnormalities, and pulmonary hypoplasia</li> <li>Venous drainage through pulmonary veins</li> </ul> </td></tr> </tbody> </table> </li> </ul>	Extralobar sequestration	Intralobar sequestration	<ul style="list-style-type: none"> <li><b>External</b> to lungs</li> <li>With pleural cover</li> <li>Causing mass effect.</li> <li><b>Venous return</b> to <b>right</b> side of heart through <b>IVC</b></li> </ul>	<ul style="list-style-type: none"> <li><b>More common</b><sup>Q</sup></li> <li>Occur <b>within</b> the lung <b>without pleural cover</b></li> <li>Localized <b>infection</b> or <b>bronchiectasis</b><sup>Q</sup></li> <li>Associated with diaphragmatic hernia, colonic duplication, vertebral abnormalities, and pulmonary hypoplasia</li> <li>Venous drainage through pulmonary veins</li> </ul>
Extralobar sequestration	Intralobar sequestration				
<ul style="list-style-type: none"> <li><b>External</b> to lungs</li> <li>With pleural cover</li> <li>Causing mass effect.</li> <li><b>Venous return</b> to <b>right</b> side of heart through <b>IVC</b></li> </ul>	<ul style="list-style-type: none"> <li><b>More common</b><sup>Q</sup></li> <li>Occur <b>within</b> the lung <b>without pleural cover</b></li> <li>Localized <b>infection</b> or <b>bronchiectasis</b><sup>Q</sup></li> <li>Associated with diaphragmatic hernia, colonic duplication, vertebral abnormalities, and pulmonary hypoplasia</li> <li>Venous drainage through pulmonary veins</li> </ul>				
<b>Congenital cystic adenomatoid malformation (CCAM)</b>	<b>Hamartomatous</b> <sup>Q</sup> or <b>dysplastic</b> lung tissue, usually confined to one lobe.				

## ATELECTASIS

**Incomplete expansion**<sup>Q</sup> of the lungs (neonatal atelectasis) or **collapse of previously inflated lung**, producing areas of relatively **airless pulmonary parenchyma**.<sup>Q</sup>

**Three types:**

Type	Pathophysiology
<b>Resorption atelectasis</b>	Due to complete <b>obstruction</b> <sup>Q</sup> of an airway.
<b>Compression atelectasis</b>	Results when <b>fluid</b> (transudate, exudate or blood), <b>tumor</b> , or <b>air</b> <sup>Q</sup> (pneumothorax) accumulate within the pleural cavity.
<b>Contraction atelectasis</b>	Occurs when focal or generalized pulmonary or pleural <b>fibrosis</b> <sup>Q</sup> prevents full lung expansion

## PULMONARY EDEMA

- Results from **increased hydrostatic pressure**<sup>Q</sup> (**left-sided congestive heart failure**)<sup>Q</sup>
- Histologically:
  - Alveolar **capillaries** are **engorged**
  - Intra-alveolar **transudate**<sup>Q</sup> appears as finely **granular pale pink**<sup>Q</sup> material.
  - Alveolar **micro-hemorrhages** & **hemosiderin-laden macrophages**<sup>Q</sup> ("heart failure" cells)<sup>Q</sup> may be seen

## ACUTE LUNG INJURY (ALI) & ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

**Acute lung injury (ALI)** (also called **non-cardiogenic pulmonary edema**) is characterized by the **sudden onset** of **significant hypoxemia** and **bilateral pulmonary infiltrates** on CXR, **without cardiac failure**.<sup>Q</sup>




## Diagnostic Criteria for ALI and ARDS

Oxygenation	Features	Absence of Left Atrial Hypertension
ALI: $\text{PaO}_2/\text{FIO}_2 < 300 \text{ mm Hg}^Q$	<ul style="list-style-type: none"> <li>Bilateral alveolar or interstitial infiltrates on CXR</li> </ul>	PCWP $< 18 \text{ mm Hg}$ or no clinical evidence of increased left atrial pressure
ARDS: $\text{PaO}_2/\text{FIO}_2 < 200 \text{ mm Hg}^Q$	<ul style="list-style-type: none"> <li>Lung Biopsy: Diffuse alveolar damage with hyaline membrane disease</li> </ul>	

## Etiology of ARDS & ALI


Infections	Physical/Injury	Chemical injury	Hematologic conditions
<ul style="list-style-type: none"> <li>Sepsis<sup>Q</sup></li> <li>Gastric aspiration<sup>Q</sup></li> <li>Diffuse pulmonary infections</li> <li>Viral<sup>Q</sup>, <i>Mycoplasma</i>, <i>Pneumocystis</i>,<sup>Q</sup></li> <li>Miliary tuberculosis<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Mechanical trauma, including <b>head injury</b><sup>Q</sup></li> <li>Pulmonary contusions,</li> <li>Fractures with fat embolism</li> <li><b>Near-drowning</b><sup>Q</sup>, <b>Burns</b><sup>Q</sup></li> <li>Ionizing radiation</li> <li><b>Hypothermia</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Oral:</b> Heroin / Barbiturate overdose, Acetylsalicylic acid</li> <li><b>Inhaled:</b> O<sub>2</sub> toxicity, <b>Smoke</b>,<sup>Q</sup> Irritant gases &amp; chemicals</li> <li>Pancreatitis</li> <li>Uremia</li> </ul>	<ul style="list-style-type: none"> <li><b>Multiple transfusions (TRALI)</b><sup>Q</sup></li> <li><b>DIC</b><sup>Q</sup></li> <li>Cardiopulmonary bypass</li> <li>Hypersensitivity reactions</li> </ul>

## Pathogenesis



## Lung Morphology in ARDS & ALI

- **Early stage:**
  - Interstitial and intraalveolar **edema<sup>Q</sup> & inflammation<sup>Q</sup>**
  - **Diffuse alveolar damage or necrosis<sup>Q</sup>**
  - **Fibrin deposition** → Alveoli become lined by **waxy hyaline membranes** (fibrin-rich edema fluid with **necrotic epithelial cells**).<sup>Q</sup>
- **Late organizing stage:**
  - **Type II pneumocytes proliferate**
  - **Granulation tissue<sup>Q</sup>** forms in the alveolar walls and spaces.





## High Yield Facts



- ARDS is also called “**shock lung**”<sup>Q</sup>
- **Earliest event** of ARDS is: diffuse **damage of alveolar capillary wall**<sup>Q</sup>
- Most important **cellular mediator** of ARDS is **Neutrophil**<sup>Q</sup>
- Most important **cytokine** involved in ARDS is **IL8**<sup>Q</sup>
- Histological **diagnostic hallmark** of ARDS is **Diffuse alveolar damage + Hyaline Membrane**<sup>Q</sup>

## OBSTRUCTIVE LUNG DISEASE

**Definition:** Increase in **resistance**<sup>Q</sup> to **airflow** due to **partial or complete obstruction**<sup>Q</sup> at any level of airway.

### Spectrum of Chronic Obstructive Pulmonary Disease

Clinical Term	Primary Site	Major Pathologic Changes	Etiology
<i>Emphysema (pink puffers)</i> <sup>Q</sup>	Acinus <sup>Q</sup>	Airspace <b>dilatation</b> <sup>Q</sup> & wall <b>destruction</b> <sup>Q</sup>	<b>Tobacco smoke</b> <sup>Q</sup>
<i>Chronic bronchitis (“blue bloaters”)</i> <sup>Q</sup> (B-B)	Bronchus <sup>Q</sup>	<b>Mucous gland hyperplasia</b> , <sup>Q</sup> hypersecretion	<b>Tobacco smoke</b> <sup>Q</sup> , air pollutants
<i>Asthma</i>	Bronchus	<b>Smooth muscle hyperplasia</b> <sup>Q</sup> , <b>excess mucus</b> <sup>Q</sup> , <b>inflammation</b> <sup>Q</sup>	Immunological causes
<i>Bronchiectasis</i>	Bronchus	Airway <b>dilation and scarring</b> <sup>Q</sup>	Persistent or severe <b>infections</b> <sup>Q</sup>
<i>Small-airway disease, Bronchiolitis</i>	<b>Bronchiole</b> <sup>Q</sup>	Inflammatory scarring/obliteration	Tobacco smoke, air pollutants


## EMPHYSEMA

**Irreversible dilatation and destruction**<sup>Q</sup> of the airspaces **distal to the terminal bronchiole (acinus)**<sup>Q</sup>, without **fibrosis**.<sup>Q</sup>


### Four Major Types

Features	Centriacinar	Panacinar	Paraseptal	Irregular
<i>Involves</i>	Proximal acinus <sup>Q</sup>	Proximal & distal acinus <sup>Q</sup>	Distal acinus <sup>Q</sup>	Irregular <sup>Q</sup> involvement
<i>Site</i>	Upper lobes <sup>Q</sup> esp apical segments	Lower zones <sup>Q</sup> at the base of lung	Upper half of lungs, cyst-like structures.	Any part of lung can be involved
<i>Etiology</i>	Smokers <sup>Q</sup> Chronic bronchitis <sup>Q</sup>	$\alpha$ 1-antitrypsin deficiency <sup>Q</sup>	Causes Spontaneous pneumothorax <sup>Q</sup>	Depends on involvement, Mostly found at autopsy

### Pathogenesis of Emphysema: Protease-antiprotease Mechanism<sup>Q</sup>








## Morphology in Emphysema

- **Over inflated**<sup>Q</sup> voluminous lungs, often overlapping the heart
- **Large apical blebs** or **bullae**- more common in **irregular and distal**<sup>Q</sup> emphysema
- **Microscopically**: abnormally **large alveoli**<sup>Q</sup> are **separated by thin septa**<sup>Q</sup> with only **focal centriacinar fibrosis**<sup>Q</sup>
- **Enlarged pores of Kohn**<sup>Q</sup>, with septa appearing **float** or **protrude blindly** into alveolar spaces with a **club-shaped end**.

## CHRONIC BRONCHITIS


- **Definition:**
  - **Persistent cough** with **sputum production**<sup>Q</sup> for at least **3 months**<sup>Q</sup> in at least **2 consecutive years**,<sup>Q</sup> in the **absence of any other identifiable cause**.<sup>Q</sup>
- **Pathogenesis:**
  - **Initiating factor** is **exposure to noxious or irritating inhaled substances**<sup>Q</sup> such as **tobacco smoke**<sup>Q</sup> (90% are smokers) and **dust from grain, cotton, and silica**.<sup>Q</sup>
- **Morphology:**
  - Increase in size of **mucous glands (hyperplasia)** with **mild hypertrophy**<sup>Q</sup>
  - **Reid index**<sup>Q</sup> or ratio of **thickness of mucous gland layer: thickness of wall between epithelium & cartilage** is **increased > 0.4**<sup>Q</sup>
  - **Goblet cell hyperplasia**<sup>Q</sup> and **chronic inflammation**<sup>Q</sup>
  - **Bronchiolar wall fibrosis (bronchiolitis obliterans)**.<sup>Q</sup>




## ASTHMA

- **Definition:**
  - **Reversible bronchoconstriction** of conducting airways, along with **inflammation**<sup>Q</sup> & **increased mucus secretion**<sup>Q</sup> usually caused by an **immunological reaction**<sup>Q</sup>, due to increased **airway sensitivity** to a variety of stimuli;
- **Genetic basis:**
  - **Chr 5q polymorphisms** in the **IL13 gene**<sup>Q</sup> (**strongest & most consistent association**)<sup>Q</sup>
  - Polymorphisms in the gene encoding **ADAM33**<sup>Q</sup>
  - **Class II HLA alleles** → **Increased IgE**
  - **IL-4 receptor gene** variants
  - Increased serum levels and lung expression of **YKL-40** (a chitinase-like glycoprotein) correlate with disease severity, airway remodeling and decreased pulmonary function<sup>Q</sup>
- **Pathogenesis:**
  - Exaggerated **TH2 response**<sup>Q</sup> to normally harmless environmental antigens. Type 1 hypersensitivity
- **Morphology:**
  - **Gross: Occlusion of bronchi and bronchioles** by **thick, tenacious mucus** plugs
- **Sputum or bronchoalveolar lavage ("3-Cs"):**
  - **Curschmann spirals**: extrusion of **mucus plugs** from subepithelial mucous gland ducts or bronchioles.<sup>Q</sup>
  - **Charcot-Leyden crystals**-composed of eosinophil protein called **galectin-10**<sup>Q</sup>
  - **Creola bodies**: ciliated columnar cells sloughed from the bronchial mucosa<sup>Q</sup>
- **Histologic findings:**
  - **Thickening of airway wall**
  - **Sub-basement membrane fibrosis**<sup>Q</sup> (due to deposition of **type I and III collagen**)
  - Increased **vascularity**<sup>Q</sup>
  - Increase in the **size of sub-mucosal glands**<sup>Q</sup> and **number of airway goblet cells**<sup>Q</sup>
  - **Hypertrophy and/or hyperplasia** of the **bronchial wall muscle**<sup>Q</sup>


Sputum findings in Bronchial Asthma



Charcot-Leyden crystals



Curschmann spirals  
Asthma




Creola bodies






## BRONCHIECTASIS

- **Definition:**
  - Destruction of smooth muscle and elastic tissue by chronic necrotizing infections leads to **permanent<sup>Q</sup> dilation of bronchi and bronchioles.**
- **Etiology:**
  - Idiopathic
  - Congenital/hereditary: e.g. **cystic fibrosis<sup>Q</sup>**, **intralobar sequestration<sup>Q</sup>**, primary ciliary dyskinesia & **Kartagener's syndrome<sup>Q</sup>**
  - Infections- Bacterial (Tuberculosis<sup>Q</sup>, *Staph aureus*), viral (*Influenza*) and fungal (*Aspergillus*)
  - Bronchial **obstruction**-tumor, mucus plug, **Foreign body<sup>Q</sup>**
  - Others-Rheumatoid Arthritis<sup>Q</sup>, SLE, IBD, GVHD<sup>Q</sup>
- **Gross Morphology:**
  - Involves **lower<sup>Q</sup> lobes bilaterally<sup>Q</sup>**
  - Airways are **dilated**, sometimes up to **four times normal size<sup>Q</sup>**
- **Microscopy:**
  - **Acute & chronic inflammatory exudates** within walls of bronchi & bronchioles
  - **Fibrosis<sup>Q</sup>** may occur



## RESTRICTIVE LUNG DISEASES

**Reduced expansion<sup>Q</sup>** of lung parenchyma and **decreased total lung capacity<sup>Q</sup>**



### High Yield Facts

- Kartagener's syndrome-bronchiectasis, sinusitis, and situs inversus<sup>Q</sup>
- Kartagener's syndrome is seen in **50% patients** with primary ciliary dyskinesia
- Reid index is increased in chronic bronchitis

## CHRONIC DIFFUSE INTERSTITIAL (RESTRICTIVE) DISEASES

### Major Categories

Fibrosing	Granulomatous
<ul style="list-style-type: none"> <li>• Usual interstitial pneumonia (idiopathic pulmonary fibrosis)<sup>Q</sup></li> <li>• Nonspecific interstitial pneumonia<sup>Q</sup></li> <li>• Cryptogenic organizing pneumonia<sup>Q</sup></li> <li>• Associated with connective tissue diseases-RA, SLE<sup>Q</sup></li> <li>• Pneumoconiosis<sup>Q</sup></li> <li>• Drug reactions</li> <li>• Radiation pneumonitis</li> </ul>	<ul style="list-style-type: none"> <li>• Sarcoidosis<sup>Q</sup></li> <li>• Hypersensitivity pneumonitis<sup>Q</sup></li> </ul>
	Smoking related
	<ul style="list-style-type: none"> <li>• Desquamative interstitial pneumonia<sup>Q</sup></li> <li>• Bronchiolitis-associated interstitial lung disease</li> </ul>
Eosinophilic	Others
<ul style="list-style-type: none"> <li>• Idiopathic chronic eosinophilic pneumonia</li> <li>• Other causes of pulmonary eosinophilia are: Churg-Strauss syndrome, Allergic Bronchopulmonary Aspergillosis</li> </ul>	<ul style="list-style-type: none"> <li>• Pulmonary alveolar proteinosis<sup>Q</sup></li> <li>• Langerhans cell histiocytosis <sup>RG</sup></li> <li>• Lymphoid interstitial pneumonia <sup>RG</sup></li> </ul>

## FIBROSING DISEASES

**A. Idiopathic pulmonary fibrosis /Usual interstitial pneumonia:**  
**Prototype of restrictive lung diseases**

**Morphology:** (Pulmonary Fibrosis ← P F → Patchy Fibroblastic foci)<sup>Q</sup>

- Patchy interstitial fibrosis<sup>Q</sup>
- Fibroblastic foci<sup>Q</sup>
- Formation of **cystic spaces (honeycomb lung)<sup>Q</sup>**

**Etiology:** Increased TGF-β<sup>Q</sup> due to alveolar epithelial damage and abnormal cell signaling

**B. Nonspecific interstitial pneumonia:**

**Idiopathic** or associated with **connective tissue diseases<sup>Q</sup>**


Prognosis: **Good<sup>Q</sup>**

**Morphology: 2 varieties**

- **Cellular pattern:** Uniform/patchy chronic interstitial inflammation
  - **Fibrosing pattern:** Diffuse/patchy interstitial **fibrotic lesions**
- Absent:** Fibroblastic foci, honeycombing, hyaline membranes and granulomas<sup>Q</sup>

**C. Cryptogenic Organizing Pneumonia/Bronchiolitis Obliterans Organizing Pneumonia (BOOP)<sup>Q</sup>**

**Histology:** Presence of **polypoid plugs** of loose organizing connective tissue (**Masson bodies**)<sup>Q</sup> within **alveolar ducts, alveoli, and bronchioles**



## PNEUMOCONIOSIS

**Definition:** Diseases induced by organic as well as inorganic particulates and chemical fumes and vapors.

Characteristics	Anthracosis	Silicosis	Asbestosis
<b>Exposure</b>	Coal mining <sup>a</sup> (particularly hard coal)	Foundry work, sandblasting, hard rock mining, stone cutting <sup>a</sup>	Mining, milling, fabrication, and installation and removal of insulation <sup>a</sup>
<b>Type of Mineral dust</b>	Coal particles	Amorphous forms & crystalline forms (Quartz, cristobalite, and tridymite) –more fibrogenic <sup>a</sup>	Serpentine <sup>a</sup> (M.C) and Amphibole <sup>a</sup> (more pathogenic)
<b>Site of involvement</b>	upper lobes <sup>a</sup> and upper zones of the lower lobes	upper lobes <sup>a</sup> and upper zones of the lower lobes	Lower lobes <sup>a</sup> and subpleurally.
<b>Caplan syndrome<sup>a</sup></b>	+	+	+
<b>Lung lesions</b>	Coal macules <sup>a</sup> (1-2 mm) Larger coal nodules <sup>a</sup> Complicated coal workers' pneumoconiosis <sup>a</sup> Progressive massive fibrosis <sup>a</sup> (1-10cm)	Discrete nodules in hilar nodes & upper zones of lungs <sup>a</sup> Hard, collagenous scars Fibrotic lesions in nodes & pleura (pleural thickening) Eggshell calcification <sup>a</sup> Hallmark: Central collagen with peripheral zone of dust-laden macrophages <sup>a</sup> Progressive massive fibrosis <sup>a</sup> (1-10cm) Lung Ca <sup>a</sup> (2 fold risk)	Asbestos bodies <sup>a</sup> Ferruginous bodies <sup>a</sup> Localized fibrous plaques Diffuse pleural fibrosis <sup>a</sup> Pleural effusions Parenchymal interstitial fibrosis (asbestosis) Lung carcinoma <sup>a</sup> Mesotheliomas <sup>a</sup> Laryngeal, ovarian, colon ca <sup>a</sup>

## OTHER LUNG DISEASES CAUSED BY AIR POLLUTANTS

Agent	Disease	Exposure
<b>Mineral dusts</b>		
<b>Beryllium</b>	Acute berylliosis, Beryllium granulomatosis <sup>a</sup> , Lung carcinoma <sup>a</sup>	Mining, fabrication
<b>Iron oxide</b>	Siderosis <sup>a</sup>	Welding
<b>Barium sulfate</b>	Baritosis <sup>a</sup>	Mining
<b>Tin oxide</b>	Stannosis <sup>a</sup>	Mining
<b>Organic dusts that induce hypersensitivity pneumonitis</b>		
<b>Moldy hay</b>	Farmer's lung <sup>a</sup>	Farming
<b>Bagasse</b>	Bagassosis <sup>a</sup>	Manufacturing wallboard, paper
<b>Bird droppings</b>	Bird-breeder's lung <sup>a</sup>	Bird handling

## GRANULOMATOUS DISEASES

### Sarcoidosis

- **Definition:**
  - A systemic disease characterized by non-caseating granulomas in tissues and organs.
- **Epidemiology:**
  - Females more commonly affected than males
- **Genetic basis:**
  - Associated with HLA-A1 and HLA-B8<sup>o</sup>
- **Etiology and Pathogenesis:**
  - Disordered immune regulation in genetically predisposed individuals exposed to certain environmental agents.
  - Cell-mediated response<sup>o</sup> to an unidentified antigen by



### CD4+ helper T cells.

- Intra-alveolar and interstitial CD4/CD8 T-cell ratios = 5 : 1 to 15 : 1<sup>o</sup>
- Increased T cell-derived TH<sub>1</sub> cytokines: IL-2 and IFN-γ → T-cell expansion and macrophage activation
- Increased levels of IL-8, TNF, macrophage inflammatory protein 1α that favor recruitment of T cells & monocytes and contribute to the formation of granulomas.
- TNF concentration in the bronchoalveolar lavage (BAL) fluid is a marker of disease activity<sup>o</sup>
- **Clinical features:**
  - Most commonly presents with bilateral hilar lymphadenopathy or lung involvement<sup>o</sup> (90% cases) followed by Skin > Extrathoracic lymph nodes > Eye > Liver > Spleen > Neurologic > Heart > Kidney (least common)<sup>o</sup>



### Histology:

- All involved tissues show **well-formed non-caseating granulomas**<sup>Q</sup> composed of **epithelioid cells**<sup>Q</sup>, with **Langhans or foreign body-type giant cells**<sup>Q</sup>
- Central necrosis is **unusual**.<sup>Q</sup>
- **Schaumann bodies**<sup>Q</sup>: laminated concretions composed of **calcium and proteins**<sup>Q</sup>
- **Asteroid bodies**<sup>Q</sup>: **stellate inclusions** enclosed within **giant cells**<sup>Q</sup>



### Diagnosis:

- EXCLUDE infections and malignancy
- **Elevated ACE level**<sup>Q</sup> (elevated in other granulomatous diseases but not in malignancy).
- **Positive gallium scan**: Increased activity in
  - Parotids and lacrimal glands (**panda sign**)<sup>Q</sup>
  - Right paratracheal and left hilar area (**lambda sign**)<sup>Q</sup>.
- **Bronchoalveolar lavage (BAL)** shows **increase in lymphocytes**
- **CD4/CD8 ratio**<sup>Q</sup> of lymphocytes in BAL > **3.5:1**<sup>Q</sup> is **strongly supportive** of Sarcoidosis

- **Kviem-Siltzbach procedure** (specific diagnostic test, no longer used): **Non-caseating granulomas** seen **4–6 weeks after Intradermal injection** of splenic tissue extract of a known sarcoidosis patient, is **highly specific** for the diagnosis of Sarcoidosis.
- **Hypercalcemia and/or hypercalciuria** (10% cases) due to increased production of **1,25-dihydroxyvitamin D<sub>3</sub>** by the granuloma.



### High Yield Facts

- **Caplan Syndrome (Rheumatoid pneumoconiosis)**<sup>Q</sup> is a combination of **rheumatoid arthritis (RA)**<sup>Q</sup> and **pneumoconiosis** that manifests as **intrapulmonary nodules**<sup>Q</sup>
- **Silicosis** is **M.C pneumoconiosis** in the world<sup>Q</sup>
- **Crystalline silica** (e.g., quartz) is **most dangerous** silica particle<sup>Q</sup>
- In patients with **Silicosis**, there is increased susceptibility to **tuberculosis**<sup>Q</sup>
- **Silicosis** is **progressive** even after exposure stops<sup>Q</sup>
- **Asbestos bodies**: **golden brown, fusiform or beaded rods** with a translucent center and consists of **asbestos fibers** coated with an **iron-containing proteinaceous material**<sup>Q</sup>
- **Ferruginous bodies**: **Inorganic particulates** coated with **iron-protein complexes**<sup>Q</sup>
- **Lung Ca** and **mesotheliomas (pleural and peritoneal)** develop in workers exposed to asbestos.
- **Amphibole**<sup>Q</sup> variety of asbestos though less prevalent, are **more pathogenic** than chrysotiles, to cause **mesothelioma**
- **Risk of Lung Ca** is increased about **five-fold**<sup>Q</sup>, while that of **mesothelioma** is **1000-fold**<sup>Q</sup> greater in **Asbestos** exposure

## HYPERSENSITIVITY PNEUMONITIS

- **Also called:**
  - Also called **Extrinsic allergic alveolitis**<sup>Q</sup>
- **Definition:**
  - **Immunologically mediated**, predominantly **interstitial** lung disorder due to prolonged exposure to inhaled organic antigens.
- **Pathogenesis:**
  - Involves **type IV**<sup>Q</sup> > **type III**<sup>Q</sup> hypersensitivity reaction (*Harrison's 18<sup>th</sup> ed, Chapter 255*)
- **Morphology:**
  - **Interstitial pneumonitis**:<sup>Q</sup> lymphocytes, plasma cells & macrophages (**eosinophils rare**)
  - **Non-caseating granulomas**<sup>Q</sup> in 2/3<sup>rd</sup> of patients
  - **Interstitial fibrosis, fibroblastic foci, honeycombing & obliterative bronchiolitis** (in late stages)
  - Intra-alveolar infiltrates in more than half of patients

## PULMONARY INFECTIONS


### Pneumonia

- **Definition:** Infection of the **lung parenchyma**;<sup>Q</sup>
- Can be **Typical (airway involvement)** or **Atypical (interstitial involvement)**





## Typical Pneumonia



- **2 patterns**
  - **Bronchopneumonia** (**Patchy consolidation** of the lung)<sup>Q</sup>
  - **Lobar pneumonia** (fibrino-suppurative consolidation of a part of lobe or entire lobe)<sup>Q</sup>

### Stages of Lobar Pneumonia<sup>Q</sup>

Stage	Characteristics
<b>Congestion</b> (1-2 days) <sup>Q</sup>	<ul style="list-style-type: none"> <li>• <b>Grossly:</b> Lung is heavy, boggy &amp; red.</li> <li>• <b>Microscopically:</b> <b>Vascular engorgement</b>,<sup>Q</sup> intra-alveolar fluid with <b>neutrophils</b><sup>Q</sup> &amp; bacteria.</li> </ul>
<b>Red hepatization</b> (2-4 days) <sup>Q</sup>	<ul style="list-style-type: none"> <li>• <b>Grossly:</b> Red, firm &amp; airless, with a <b>liver-like consistency</b><sup>Q</sup>, hence the term <b>hepatization</b>.</li> <li>• <b>Microscopically:</b> Confluent <b>exudation</b> with neutrophils, RBCs, <b>fibrin</b><sup>Q</sup> filling alveoli;</li> </ul>
<b>Gray hepatization</b> (5-8 days) <sup>Q</sup>	<ul style="list-style-type: none"> <li>• <b>Grossly:</b> Grayish brown, dry surface.</li> <li>• <b>Microscopically:</b> <b>Disintegration of RBCs</b> &amp; persistence of a <b>fibrinosuppurative exudate</b></li> </ul>
<b>Resolution</b> (8-9 days)	<ul style="list-style-type: none"> <li>• Exudates within alveoli broken down by enzymatic digestion to produce <b>granular debris</b><sup>Q</sup></li> <li>• Debris may be reabsorbed/ingested by macrophages/expectorated/<b>organized by fibroblasts</b><sup>Q</sup></li> </ul>



### High Yield Facts

- MC cause of **Community-acquired pneumonia** is ***Streptococcus pneumoniae***<sup>Q</sup> followed by *H. influenza*
- Most common cause of **Community-acquired atypical pneumonia** is ***Mycoplasma pneumoniae***<sup>Q</sup> > ***Chlamydia***
- Most common cause of **Hospital-acquired pneumonia** are **Gram-negative rods**<sup>Q</sup>, **Enterobacteriaceae** (*Klebsiella spp.*, *Serratia marcescens*, *Escherichia coli*) and *Pseudomonas spp.*
- Most common cause of **Aspiration pneumonia** are **Anaerobic**<sup>Q</sup> oral flora (*Bacteroides*<sup>Q</sup>, *Prevotella*,

- **Etiology**
  - **Bacterial:**<sup>Q</sup> ***Streptococcus pneumoniae***,<sup>Q</sup> *H. influenzae*, *Moraxella catarrhalis*, *S. aureus*, *K. pneumoniae*, *Pseudomonas spp.*
- **Morphology**
  - In **lobar pneumonia**, 4 stages have been described; see below
  - In **Bronchopneumonia**:
  - Scattered **areas of acute suppurative inflammation**,<sup>Q</sup> usually **multilobar**, **frequently bilateral**<sup>Q</sup> & **basal**<sup>Q</sup> because of the tendency of secretions to gravitate to lower lobes;
  - **Histologically: neutrophil-rich exudate**<sup>Q</sup> that fills the bronchi, bronchioles and adjacent alveolar spaces.
- **Complications**
  - **Abscess formation**<sup>Q</sup> (common with **type 3 *Pneumococci*** or ***Klebsiella*** infections)
  - **Empyema**: Due to spread of infection to **pleural cavity**<sup>Q</sup>
  - **Bacteremic dissemination**<sup>Q</sup> to the heart valves, pericardium, brain, kidneys, spleen, or joints, causing **metastatic abscesses**, endocarditis, meningitis, or suppurative arthritis.

## Atypical Pneumonia

- **Definition:**
  - Acute febrile respiratory disease characterized by **patchy inflammatory changes** in the lungs, largely confined to the **alveolar septa**<sup>Q</sup> & **pulmonary interstitium**<sup>Q</sup>.
- **Differences of atypical from typical pneumonia:**
  - **Moderate** amount of sputum<sup>Q</sup>
  - **No physical findings** of consolidation<sup>Q</sup>
  - Only **moderate elevation of WBCs**<sup>Q</sup>
  - **Lack of alveolar exudate**<sup>Q</sup>

- **Etiology:**
  - **Atypical organisms:** ***Mycoplasma pneumoniae*** (MC)<sup>Q</sup>, *Chlamydia pneumoniae*, *C burnetii* (Q-fever)
  - **Viruses:** ***Influenza virus types A and B***, *Respiratory syncytial viruses (RSV)*, *Parainfluenza (children)*, *Human metapneumovirus*, *Adenovirus*, *Rhinoviruses*, *Rubeola*, *Varicella viruses*
- **Risk Factors:**
  - Extremes of age<sup>Q</sup>, malnutrition<sup>Q</sup>, alcoholism<sup>Q</sup> & underlying debilitating illnesses
- **Morphology:**
  - **Interstitial**<sup>Q</sup> **inflammatory reaction**, virtually **localized within the walls**<sup>Q</sup> of alveoli.



- Alveolar septa are widened and edematous<sup>Q</sup> and usually have a **mononuclear inflammatory infiltrate<sup>Q</sup>** of lymphocytes, macrophages, and occasionally plasma cells.
- Superimposed bacterial infection → **ulcerative bronchitis, bronchiolitis & bacterial pneumonia.**
- Herpes simplex, Varicella & Adenovirus*, may be associated with **necrosis of bronchial and alveolar epithelium<sup>Q</sup>** and **acute inflammation<sup>Q</sup>**.

## Tuberculosis

Causative agent: *Mycobacterium tuberculosis*

- Weakly **Gram positive<sup>Q</sup>** bacilli
- Acid fast (resists decoloration with acid & acid-alcohol)<sup>Q</sup>** on **Ziehl Neelsen (ZN)<sup>Q</sup> staining**, due to a cell wall composed of glycolipids & **mycolic acid<sup>Q</sup>**
- Group specificity** is due to **polysaccharide<sup>Q</sup>**
- Type specificity** is due to **protein antigen<sup>Q</sup>**.

**Main source of transmission:** Person to person transmission of **air-borne<sup>Q</sup>** organisms

## High Yield Facts

- Mycobacterium tuberculosis* enters into macrophages with the help of **mannose binding lectin<sup>Q</sup>** and **CR3<sup>Q</sup>**.
- Macrophages<sup>Q</sup>** are the **primary cells infected** by *M. tuberculosis*.
- IFN- $\gamma$ <sup>Q</sup>** is the **critical mediator** that enables macrophages to contain the *M. tuberculosis* infection.
- NK T cells<sup>Q</sup>** &  **$\gamma\delta$  T-cells<sup>Q</sup>** also produce IFN- $\gamma$ .
- People with genetic **deficiencies in IL-12 & IFN- $\gamma$  pathway**, including STAT1 a signal transducer for IFN- $\gamma$ , are **vulnerable to severe Mycobacterial infections**.
- Polymorphisms in genes for HLA, IFN- $\gamma$ , IFN- $\gamma$  receptor & TLR2** are associated with **increased susceptibility** to TB
- Factors contributing in pathogenesis of TB: **Cord factor, Lipoarabinomannan<sup>Q</sup>, Complement system, M. tuberculosis heat shock proteins.**
- Risk of acquiring TB infection** is determined mainly by **exogenous factors<sup>Q</sup>** while **risk of developing TB disease** depends largely on **endogenous factors<sup>Q</sup>**.

## Pathogenesis

- Entry & replication in macrophages:**
  - By **inhibition of phagolysosome formation<sup>Q</sup>**
- TH1 response:**
  - Alveolar macrophages** that **present TB antigen** to T cells, also secrete **IL-12** which activates T cells to differentiate to **TH1 cells**.
- Macrophage activation & bacteria killing:**
  - IFN  $\gamma$ <sup>Q</sup>** produced by **TH1 cells** – Stimulates: **Maturation of phago-lysosome**
  - Stimulates production of **NO** → **reactive nitrogen intermediates**
  - Mobilizes antimicrobial peptides (**defensins**) against *M. tuberculosis*
  - Stimulates **autophagy** to destroy *M. tuberculosis*.

## Granulomatous inflammation & tissue damage:

- Activated macrophages differentiate into '**epithelioid histiocytes**'
- Some epithelioid cells may fuse to form **giant cells<sup>Q</sup>**
- Combination of **epithelioid cells & giant cells forms granulomas<sup>Q</sup>**

## Morphology

**A. Primary Tuberculosis:** in previously unexposed (unsensitized)<sup>Q</sup> person.

- Primary organ** involved is mostly **lungs<sup>Q</sup>**
- Usually involves **lower part of upper lobe** or **upper part of lower lobe<sup>Q</sup>** (close to pleura)<sup>Q</sup>
- Ghon's focus<sup>Q</sup>** : Grey white area of inflammation with consolidation
- Ghon's complex:<sup>Q</sup>** Ghon's focus + inflamed regional lymph nodes
- Histology:** **Granulomatous inflammatory reaction** with both **caseating & non-caseating** tubercles
- Simon's Focus<sup>Q</sup>** : Occult hematogenous dissemination to apex of lung

**Fate of primary TB:**

- 95% cases:** controlled by immunity Calcified healed lesions & hilar lymph nodes (**Ranke complex**)
- 5% cases:** Primary tuberculosis is progressive (**Progressive Primary TB**)
  - resembles acute bacterial pneumonia
  - **Lymphohematogenous dissemination** may cause **tuberculous meningitis &/or miliary TB**

**B. Secondary Tuberculosis:** in a previously sensitized host<sup>Q</sup>

Also called 'adult type' or '**reactivation tuberculosis**' or 'chronic pulmonary TB'

- Involves **apical & posterior segments of upper lobe** due to high O<sub>2</sub> concentration (**Puhl's lesion<sup>Q</sup>**)
- Infraclavicular lesion** is called **Assman's Focus<sup>Q</sup>**
- Regional lymph node involvement is late<sup>Q</sup>**
- Cavitation occurs readily<sup>Q</sup>** with erosion of the cavities into an airway
- Histology:** Active lesions show **coalescent tubercles** with central caseation (**Caseous Necrosis<sup>Q</sup>**)
- AFB** can be seen in **early phase** of granuloma formation but usually not seen in late fibro-calcific stage

**C. Extrapulmonary TB:**


- MC site is lymph node<sup>Q</sup>** MC cervical & supraclavicular ('**Scrofula**')<sup>Q</sup>
- Pleural TB:** **Exudative<sup>Q</sup>** Pleural effusion, tuberculous empyema, or obliterative fibrous pleuritic;
- Renal TB:** **sterile pyuria<sup>Q</sup>**
- Genital TB:** Preferentially involves **fallopian tube<sup>Q</sup>** in females & **epididymis<sup>Q</sup>** in males
- Skeletal TB:** Most common site **spine<sup>Q</sup>** (**Pott's disease**)<sup>Q</sup> > hip > knee
- Paravertebral cold abscess** may form;
- TB meningitis** (paresis of cranial nerves especially ocular, is frequent finding)<sup>Q</sup>
- GI TB (MC site terminal ileum and caecum)<sup>Q</sup>**



- **Tuberculous pericarditis** (MC cause of chronic constrictive pericarditis)<sup>Q</sup>
- **Endobronchial, endotracheal & laryngeal tuberculosis** may develop by spread through lymphatic channels or from expectorated infectious material.


## RECENT EXAM<sup>Q</sup>

- **Miliary/Disseminated TB:** When bacteria disseminate through systemic **arterial system** & involve **lungs** ± multiple organs. Most prominent in the **liver**, bone marrow, spleen, adrenals and meninges.




Nodules  
2–4 mm  
in size

Here one can see multitude of small tan(-) yellow granulomas, about 2 to 4 mm in size, scattered throughout the lung parenchyma. The miliary pattern gets its name from the resemblance of the granulomas to millet seeds. Diagnosis miliary-TB.



Necrosis

TB Granuloma



Ghon's focus

AFB staining showing mycobacteria

- In **congenital TB**, primary organ involved is **liver**<sup>Q</sup>. Cantwell revised criteria for congenital TB.

### Cantwell criteria

- **Presence of proven Tuberculous disease with atleast 1 of the following:**
  1. Lesions in the newborn baby during the first week of life.
  2. Primary hepatic complex or caseating hepatic granuloma.
  3. Tuberculous infection of the placenta or maternal genital tract.
  4. Exclusion of possibility of postnatal transmission by investigation of contacts, including hospital staff.

## Diagnosis

- **Mantoux (Tuberculin test)**<sup>Q</sup> & **IFN-γ release assay (IGRA)**<sup>Q</sup> indicate **infection with TB & not TB disease**<sup>Q</sup>
- **Sputum Microscopy** by **ZN staining**<sup>Q</sup>
- **Petroff's method**<sup>Q</sup> is best suited for decontamination of sputum
- **Auramine Rhodamine**<sup>Q</sup> stain **more sensitive** than ZN staining
- **Culture** media for TB are: **LJ media, Middlebrook media**
- **Bactec/MGIT** method may be used for early diagnosis
- **Rapid** diagnostic tests for TB include: **PCR, Line probe assay**<sup>Q</sup>, **GeneXpert**<sup>Q</sup>
- **FNAC/Biopsy** of involved organ shows **caseating/non-caseating granuloma with/without AFB**<sup>Q</sup>

## High Yield Facts

- **"Primary"** TB occurs in **non-immune** host & **"secondary"** TB occurs in a **host immune** to *M. tuberculosis*.
- **Immunocompromised** people **do not form granulomas** & their macrophages **contain many AFB**<sup>Q</sup>
- **Hemoptysis** (in 20–30% cases) may result from rupture of a dilated vessel in a cavity (**Rasmussen's aneurysm**)<sup>Q</sup> or from **Aspergilloma** formation in an old cavity
- **Adenosine deaminase (ADA)** level in pleural fluid > **40 IU/L**<sup>Q</sup> indicates Tuberculosis.
- **Causes of Necrotizing epithelioid cell granulomas:**

- |                         |                       |   |
|-------------------------|-----------------------|---|
| • Tuberculosis          | • Tuberculoid leprosy | • Wegener's Granulomatosis <sup>Q</sup> |
| • Cat's scratch disease | • Syphilis            |   |



## OTHER TYPES OF PNEUMONIA

### Cryptogenic Organizing Pneumonia (COP)

- Noninfectious pneumonia characterized by inflammation of bronchioles and surrounding structure
- Formerly known as: Bronchiolitis obliterans organizing pneumonia or BOOP
- Micrograph shows a Masson body (pale circular and paucicellular), as may be seen in cryptogenic organizing pneumonia. The Masson body plugs the airway
- Cultures: Sputum and blood cultures are negative and has no response to antibiotics




Image showing Masson bodies

## LUNG ABSCESS

- Definition:**
  - Local suppurative** process within the lung, characterized by **necrosis** of lung tissue.
- Etiology:**
  - Bacteroides**, **Fusobacterium**, & **Peptococcus** (3 most common bacteria), Aerobic and anaerobic *Streptococci*, *S. aureus* & gram -ve organisms.
- Risk factors:**
  - Aspiration of infective material** (Most common cause)

## TUMORS OF LUNGS & PLEURA


### Lung Carcinoma

#### Etiology and Pathogenesis

- Tobacco Smoking:** 60 times<sup>Q</sup> greater among habitual heavy smokers (2 packs/day for 20 years)
- Industrial Hazards:** Exposure to **asbestos**<sup>Q</sup>, **arsenic**<sup>Q</sup>, **chromium**<sup>Q</sup>, **uranium**, **nickel**, **vinyl chloride** and **mustard gas**, increase the risk of developing lung cancer. High-dose **ionizing radiation** is carcinogenic.
- Air Pollution:** Radon

R10<sup>th</sup>

Latest Update








## Types of Lung Ca

### ■ Adenocarcinoma:


- **Peripherally** located<sup>Q</sup>
- **Well-differentiated tumors** with obvious glandular element
- Express thyroid transcription factor-1 (TTF-1)<sup>Q</sup> and Napsin A
- Electron Microscopy shows **short, plump microvilli**<sup>Q</sup>
- 2 subtypes:
  - **Microinvasive:** ≤3 cm in size and ≤5 mm invasion<sup>Q</sup>
  - **Mucinous:** Tend to **spread aerogenously**<sup>Q</sup> forming **satellite tumors**<sup>Q</sup>



Adenocarcinoma

### ■ Squamous cell carcinoma:

- **Strongly associated with smoking**<sup>Q</sup>
- **Central in location**<sup>Q</sup>
- Precursor lesions- **Squamous metaplasia** or **dysplasia** of bronchial epithelium
- Mass like lesion which may infiltrate surrounding areas
- **Hemorrhage** or **necrosis** which may also form **cavity lesions**<sup>Q</sup>
- **Keratinization** ("squamous pearls")<sup>Q</sup> on **light microscopy** and/or **Intercellular bridges**<sup>Q</sup> on **electron microscopy** is diagnostic




Squamous cell Ca

### ■ Small cell carcinoma:


- **Most malignant Lung Ca**<sup>Q</sup>, most **aggressive**<sup>Q</sup>, **wide metastasis**<sup>Q</sup> and **fatal**<sup>Q</sup>

- **Strongest relationship to cigarette smoking**<sup>Q</sup>
- **Central > peripheral** in location<sup>Q</sup>
- No preinvasive phase.
- **Light Microscopy:**
  - Small cells with **salt and pepper pattern**<sup>Q</sup>, **nuclear molding**<sup>Q</sup> is prominent



Small cell Ca

- **Basophilic staining** of vascular walls due to encrustation by DNA from necrotic tumor cells (**Azzopardi effect**)<sup>Q</sup>



Azzopardi effect

- **Electron microscopy:** **Dense-core neurosecretory granules**<sup>Q</sup> releasing neuroendocrine markers such as **chromogranin**<sup>Q</sup>, **synaptophysin**<sup>Q</sup> and **CD57**<sup>Q</sup>, **parathormone-related protein**<sup>Q</sup>
- **Most common lung Ca** associated with **ectopic hormone production**<sup>Q</sup>
- Immunohistochemistry: **BCL2**<sup>Q</sup> in 90% of tumors
- **Large cell carcinoma:**
  - Diagnosis of exclusion
  - **Large nuclei**<sup>Q</sup>, **prominent nucleoli**<sup>Q</sup> and a **moderate amount of cytoplasm**





## Differences between Small Cell and Non-small Cell Lung Carcinoma

Features	Small cell lung carcinoma	Non-Small cell lung carcinoma
<b>Tumor Suppressor gene abnormalities</b>		
3p deletions	>90%	>80%
RB mutations	~90%	~20%
p(6KDN2A) mutation	~10%	>50%
TP53 mutation	>90%	>50%
<b>Dominant oncogene abnormalities</b>		
KRAS mutations	Rare	~30% (adenocarcinoma)
EQR mutations	Absent	~20% (adenocarcinoma, non-smokers, woman)
ALK rearrangements	Absent	4%–6% adenocarcinoma, non-smokers, often have signet ring morphology
Responses to chemotherapy and radiotherapy	Often complete response but recur invariably	Incomplete



### High Yield Facts

- **Lung Carcinoma is most frequently diagnosed<sup>Q</sup>** major cancer in the world
- **Lung Carcinoma is most common cause of cancer mortality<sup>Q</sup>** worldwide
- Most common Lung Ca **world-wide is Adenocarcinoma<sup>Q</sup>**
- Most common lung Ca in **India is Squamous cell Ca<sup>Q</sup>**
- Most common cancer associated with **smoking** is squamous cell carcinoma
- Most specifically associated with smoking: small cell carcinoma
- Most common cancer in **non-smokers<sup>Q</sup>** and **women<sup>Q</sup>**, **young age<sup>Q</sup>** is **Adenocarcinoma**
- Lung Ca with **best prognosis- Adenocarcinoma<sup>Q</sup>**
- Lung Ca with **worst prognosis- Small Cell Ca<sup>Q</sup>**
- **Lung Carcinoids may occur in patients with MEN type I**
- Most common paraneoplastic manifestations of Bronchoalveolar carcinoma is systemic sclerosis

## Metastasis from Lung Ca to Other Organs

- Lymphatic and hematogenous pathways<sup>Q</sup>
- **Sq cell Ca shows late metastasis<sup>Q</sup>**
- **Brain > Adrenals, Liver, and bone**


## Paraneoplastic Syndromes in Lung Carcinoma


Hormone Secreted	Clinical Manifestations
<b>Antidiuretic hormone (ADH)<sup>Q</sup></b>	<b>Hyponatremia</b> due to <b>SIADH<sup>Q</sup></b>
<b>Adrenocorticotrophic hormone (ACTH)<sup>Q</sup></b>	<b>Cushing syndrome<sup>Q</sup></b>
<b>Parathormone, PTH -related peptide, PGE</b>	<b>Hypercalcemia<sup>Q</sup></b> [Most commonly seen with SCC lung]
<b>Calcitonin</b>	<b>Hypocalcemia</b>
<b>Gonadotropins</b>	<b>Gynecomastia</b> [Most commonly seen with large cell carcinoma lung]
<b>Serotonin and bradykinin</b>	<b>Carcinoid syndrome<sup>Q</sup></b>

## Other Systemic Manifestations of Lung Carcinoma

- **Lambert-Eaton myasthenic syndrome<sup>Q</sup>**
- **Peripheral neuropathy<sup>Q</sup>**
- **Acanthosis nigricans<sup>Q</sup>**
- Leukemoid reactions
- **Trousseau syndrome** (deep vein thrombosis and thromboembolism)<sup>Q</sup>


- **Hypertrophic pulmonary osteoarthropathy**, associated with **clubbing of the fingers.<sup>Q</sup>**
- Apical lung cancers (**Pancoast tumors**): Severe pain along ulnar nerve
- **Horner syndrome<sup>Q</sup>** (enophthalmos, ptosis, miosis, and anhidrosis) on the same side as the lesion





## Carcinoid Tumors of Lungs

- **Epidemiology:**
  - 1% -5% of all lung tumors<sup>Q</sup>
  - Age of Onset: < 40 years<sup>Q</sup> of age, Male is equally involved to females<sup>Q</sup>
  - 20% - 40% of patients are non-smokers<sup>Q</sup>
- **Classification:**
  - **Classical carcinoid:** <2 mitosis/10 hpf (high power fields) and lack necrosis<sup>Q</sup>
  - **Atypical carcinoid:** 2-10 mitosis/hpf with increased pleomorphism, necrosis, prominent nucleoli<sup>Q</sup>, may cause Carcinoid syndrome<sup>Q</sup>
- **Clinical Features:**
  - Obstructive line bronchiectasis, emphysema & atelectasis<sup>Q</sup>
  - Elaboration of vasoactive amines: Carcinoid syndrome characterized by diarrhea, flushing & cyanosis.<sup>Q</sup>
- **Gross Morphology:**
  - Carcinoids may arise centrally or may be peripheral
  - Spherical polypoid masses<sup>Q</sup> (3-4 cm) that project into the lumen of bronchus
  - Some tumors penetrate bronchial wall (invasive)<sup>Q</sup> to fan out in the peribronchial tissue, producing 'collar-button lesion'.<sup>Q</sup>
- **Light Microscopy:**
  - Nest/rosette-like arrangements of cells separated by a delicate fibrovascular stroma<sup>Q</sup>
- **IHC:**
  - Stains positive for Serotonin<sup>Q</sup>, Neuron-specific enolase<sup>Q</sup>, Bombesin<sup>Q</sup>, Calcitonin<sup>Q</sup>
- **Electron microscopy:**
  - Dense-core granules,<sup>Q</sup> characteristic of all neuroendocrine tumors
- **Reported 5-year survival rates:**
  - Typical carcinoids (Best Prognosis)<sup>Q</sup>,
  - Small cell carcinoma (Worst prognosis)<sup>Q</sup>




Lung carcinoid showing tumor nests

## Metastatic Tumors

Lung is the most common site of metastatic neoplasms<sup>Q</sup>


### Pleural Tumors

#### Solitary Fibrous Tumor

- **Definition:**
  - Soft-tissue tumor of the pleura<sup>Q</sup> and, less commonly, in the lung
- **Epidemiology:**
  - No relationship to asbestos exposure.<sup>Q</sup>
- **Genetics:**
  - Cryptic inversion of *chr 12* involving *NAB2-STAT6 fusion gene*<sup>Q</sup> 
- **Gross Morphology:**
  - May be attached to the pleural surface by a pedicle, dense fibrous tissue with occasional cysts filled with viscid fluid
- **Microscopy:**
  - Tumor shows whorls of reticulin<sup>Q</sup> and collagen fibers<sup>Q</sup> with interspersed spindle cells<sup>Q</sup> resembling fibroblasts
  - Tumor cells are CD34+ve<sup>Q</sup> and keratin -ve<sup>Q</sup> by immunostaining

#### Malignant Mesothelioma

- **Epidemiology:**
  - Rare tumor but risk of mesothelioma in heavily asbestos exposed individuals (25-45 yrs).
  - No increased risk of mesothelioma in asbestos workers who smoke<sup>Q</sup>
  - SV40 (simian virus 40) is often associated<sup>Q</sup>
- **Gross Morphology:**
  - Diffuse lesion arising either from visceral or parietal pleura
  - May cause extensive pleural effusion & direct invasion of thoracic structures
- **Light Microscopy:**
  - Epithelioid (60%) Most common variety<sup>Q</sup>, Sarcomatoid (20%), Mixed (20%)
- **IHC:**
  - Strong positivity for keratin proteins, calretinin,<sup>Q</sup> WT-1,<sup>Q</sup> cytokeratin 5/6<sup>Q</sup> & D2-40<sup>Q</sup>
- **Electron microscopy:**
  - Long microvilli & abundant tonofilaments but absent microvillous rootlets and lamellar bodies<sup>Q</sup>
- MC gene mutated is BAP<sub>1</sub>



Asbestos body



Electron microscopy Image shows long and thin microvilli

### R10<sup>th</sup> Latest Update

		TTF-1	
		+	–
p63	+	NSCLC, favor adenocarcinoma	NSCLC, favor squamous cell carcinoma
	–	NSCLC, favor adenocarcinoma	NSCLC, NOS

### R10<sup>th</sup> Latest Update

#### Squamous cell carcinoma - 2015 UPDATES

Number of subtypes have been reduced to three, which makes the diagnosis easier and avoids rare subtypes with confusing names

- Keratinizing
- Non-keratinizing
- Basaloid squamous cell carcinoma (**new category added**)

#### Note The Terminologies:

- Basaloid squamous cell carcinoma -if this component is >50% of the tumor, regardless of the presence of any keratinization.
- In tumors with <50% basaloid component, this can be acknowledged in the diagnosis “with basaloid features”.

#### Neuroendocrine Tumors-2015 Updates

- They are grouped into **high grade tumors (small cell and large cell neuroendocrine carcinoma)**, **intermediate and low grade tumors (atypical and typical carcinoids)**, and the **preinvasive diffuse idiopathic pulmonary neuroendocrine cell hyperplasia (DIPNECH)**, and for each of these, there are characteristic molecular alterations

Small cell carcinoma
Large cell neuroendocrine carcinoma
Carcinoid tumor
Diffuse idiopathic pulmonary neuroendocrine cell hyperplasia

The following entities are listed according to the new WHO classification:

- NUT CARCINOMA
- Lymphoepithelioma-like carcinoma

#### Nut Carcinoma

- Chromosomal translocation between the **NUT gene (NUTM1) on chromosome 15q14** and other genes: BRD4 on chromosome 19p13.1 (70%), BRD3 on chromosome 9q34.2 (6%), or an unknown partner gene (24%)
- Very aggressive tumor with a median survival of 7 months






## NEXT Pattern Questions



Q's

1. A 45-year-old male presented with severe respiratory distress. O/E he had pedal edema, bilateral crepitation on auscultation. He was admitted to emergency department and expired 2 days of admission. Lung biopsy done suggested the following. What is your diagnosis?



- a. CMV pneumonitis      b. Small cell Ca lung  
c. Tuberculosis      d. Heart failure cells


**Ans. (d) Heart failure cells**

- With the history of heart failure symptoms, notice the brown colored macrophages with hemosiderin called heart failure cells.



Q's

3. A middle aged immunocompromised male came with fever and breathlessness. HRCT showed a middle lobe lesion with infiltration. Lung biopsy from the lesion is shown in image. Most likely Diagnosis is:



- a. Tuberculosis Pneumonia  
b. Cryptococcus pneumonia  
c. Small cell carcinoma lung  
d. CMV pneumonia


**Ans. (b) Cryptococcus pneumonia**

- Cryptococcus pneumonia is common in HIV immunocompromised patients. Notice the clear capsulated cryptococcus and the PAS + stained capsule of the same.



Q's

2. A Factory worker was working in a factory from past 20 years, and now presenting with pleural thickening and fibrosis. Histopathology of lesion is shown in below image. Most likely diagnosis is:



- a. Asbestosis  
b. Byssinosis  
c. Coal Worker Pneumoconiosis  
d. Silicosis


**Ans. (a) Asbestosis**

- Note the brown colored the alveoli suggestive of Asbestosis.



Q's

4. A 58-year-old female presents with difficulty of breathing. CT scan show peripherally located mass lesion. Histopathological diagram of lung biopsy is shown below, based on it what is your diagnosis?




- a. Adenocarcinoma  
b. Squamous cell carcinoma  
c. Carcinoid tumor  
d. Small cell carcinoma


**Ans. (a) Adenocarcinoma**

- The mass in the bronchus which is peripherally located, shows malignant invasive glands suggestive of adenocarcinoma





5. Which of the following disease is classically associated with the following?



- a. Bronchiectasis      b. Interstitial lung disease      c. Chronic Bronchitis      d. Emphysema


**Ans. (c) Chronic Bronchitis**

- In chronic bronchitis, there is increase in Reid index which means the ratio of submucosal mucin gland layer by bronchial wall layer is more than 0.4.




## Image-Based Questions

1. A 45-year-old male presented with severe respiratory distress. O/E he had pedal edema, bilateral crepitation on auscultation. He was admitted to emergency department and expired 2 days of admission. Lung biopsy done suggested the following. What is your diagnosis?




- a. CMV pneumonitis  
b. Small cell Ca lung  
c. Tuberculosis  
d. Heart failure cells

2. A 50-year-old male chronic smoker who smokes 20 cigarettes/ day for last 15 years died of severe respiratory distress. His lung autopsy shows the following. Identify the lesion?



- a. Centriacinar emphysema  
b. Panacinar Emphysema  
c. Chronic bronchitis  
d. Bronchiectasis



3. Autopsy gross specimen from lungs suggested the following, What is your interpretation?




- a. Chronic bronchitis
- b. Bronchiectasis
- c. Lung Ca
- d. Pleural mesothelioma

5. Identify the given test being shown and its importance?




- a. Schick Test, Confirmed TB diagnosis
- b. Mantoux test, latent Tb diagnosis
- c. Mantoux test, TB disease
- d. Mantoux test, Prior TB exposure

4. A full term neonate rapidly develops progressive respiratory distress shortly after birth. lung biopsy done post-mortem reveals the following, what is your diagnosis?




- a. Pulmonary hemorrhage
- b. Pulmonary alveolar proteinosis
- c. Pneumonia
- d. Pulmonary edema

6. A 40-year-old female was admitted to hospital with fever, loss of appetite and weight, cough and weakness of 3 months duration. A chest X-ray showed consolidation at the upper part of lung. She died a few days after admission. Autopsy findings of gross lung and its biopsy findings have been shown. What is your diagnosis?



- a. Sarcoidosis
- b. Ca lung
- c. Tuberculosis Lung
- d. Bronchiectasis






7. A 56-year-old male who presented with severe respiratory distress, high grade fever and streaks of blood in sputum, died 7 days after hospitalization in AIIMS ICU. The lung autopsy has been shown, Identify the pathology?



- a. Lobar pneumonia  
b. Bronchopneumonia  
c. Tuberculosis  
d. Bronchogenic Carcinoma

8. A 56-year-old male presented with lesion at the lung apex. He was working in asbestos factory for last 20 years. The lung biopsy was seen under electron microscope which revealed the following. What is your diagnosis?



- a. Adenocarcinoma Lung  
b. Mesothelioma  
c. Lung metastasis  
d. Benign pleural fibroma



## Answers of Image-Based Questions

1. Ans. (d) **Heart failure cells**
  - The lung biopsy shows intra-alveolar **transudate granular pale pink** material along with **alveolar micro-hemorrhages & hemosiderin-laden macrophages ("heart failure" cells)** seen in left-sided congestive heart failure.
2. Ans. (a) **Centriacinar emphysema**
  - Central areas in the lung specimen shows marked emphysematous damage surrounded by relatively spared alveolar spaces.
3. Ans. (b) **Bronchiectasis**
  - Cut surface of lung shows markedly distended peripheral bronchi upto four times the normal size.
4. Ans. (b) **Pulmonary alveolar proteinosis**
  - The alveoli are filled with a dense, amorphous, protein-lipid granular precipitate, while the alveolar walls are normal.
5. Ans (d) **Mantoux test, Prior TB exposure**
  - Mantoux (Tuberculin test)** indicate **infection with TB and Type IV hypersensitivity to antigens but not TB disease**.
6. Ans. (c) **Tuberculosis lung**
  - Gross lung shows upper lobe cavitations and its biopsy findings shows caseating granuloma suggestive of Tuberculosis.
7. Ans. (a) **Lobar pneumonia**
  - The gross specimen of lung shows consolidation of a large portion of an entire lower lobe .
8. Ans. (b) **Mesothelioma**
  - Electron microscopy shows long microvilli & abundant tonofilaments but absent microvillous rootlets and lamellar bodies.




## Multiple Choice Questions

### RESPIRATORY SYSTEM AND ITS CONGENITAL MALFORMATIONS

1. **Ciliocytophthoria is seen in:** (AIIMS Nov 2019)
  - a. Kartagener
  - b. Situs inversus
  - c. Acute respiratory infection
  - d. Cystic fibrosis
2. **True about pulmonary sequestration:** (PGI May 2019)
  - a. Female preponderance
  - b. Supplied by bronchial supply
  - c. May have independent venous drainage
  - d. Intralobular variety more common
  - e. May be associated with other congenital anomalies
3. **Kartageners syndrome cause of infertility is?** (AIIMS May 18)
  - a. Oligospermia
  - b. Asthenospermia
  - c. Undescended testis
  - d. Epididymis obstruction
4. **Bronchial mucosa secretes all except?** (Recent Question 2015)
  - a. Bombesin
  - b. Calcitonin
  - c. Serotonin
  - d. Bradykinin
5. **Most common type of foregut cysts are?** (Recent Question 2014)
  - a. Bronchogenic
  - b. Esophageal
  - c. Enteric
  - d. Mixed type
6. **Blood supply of bronchogenic sequestration is?** (Recent Question 2014)
  - a. Aorta
  - b. Pulmonary artery
  - c. Pulmonary Vein
  - d. Bronchogenic artery
7. **Hamartomatous lung tissue is?** (Recent Question 2015)
  - a. Hypoplasia of lung
  - b. Congenital cyst
  - c. Lobar sequestration
  - d. Congenital cystic adenomatoid malformation
8. **Respiratory system develops from?** (Recent Question 2013)
  - a. Ventral wall of foregut
  - b. Dorsal wall of foregut
  - c. Ventral wall of midgut
  - d. Dorsal wall of midgut
9. **Bronchogenic cyst is lined by?** (Recent Question 2013)
  - a. Stratified Squamous epithelium
  - b. Squamous epithelium
  - c. Non ciliated pseudostratified columnar epithelium
  - d. Ciliated pseudostratified columnar epithelium
10. **Which cells produce surfactant in conducting part of the lung?** (MAHA 2015)
  - a. Goblet cells
  - b. Brush cells
  - c. Basal cells
  - d. Clara cells
11. **Pores of Kohn are present in?** (Recent Question 2013)
  - a. Bronchioles
  - b. Alveoli
  - c. Bronchus
  - d. Terminal bronchioles

### ARDS AND PNEMONIA

12. **A middle aged immunocompromised male came with fever and breathlessness. HRCT showed a middle lobe lesion with infiltration in the lung as shown in image. Most likely Diagnosis is?** (AIIMS May 18)




- a. Tuberculous Pneumonia
  - b. Cryptogenic organizing pneumonia
  - c. Small cell carcinoma lung
  - d. CMV pneumonia
13. **The following are true regarding hyaline membrane disease except:** (APPGMEE 2015)
  - a. Basic abnormality is deficiency of surfactant
  - b. Prenatal diagnosis by low amniotic fluid L/S ratio
  - c. Intratracheal surfactant helps
  - d. Occurs in babies born post-dates
14. **What is false about ARDS?** (Recent Question 2016)
  - a. Mucus plug in alveoli
  - b. Interstitial edema
  - c. Hyaline membrane present
  - d. Interstitial infiltrates by cells
15. **Etiology of ARDS are all except?** (Recent question 2015)
  - a. Multiple transfusion
  - b. Sepsis
  - c. Aspiration of gastric contents
  - d. Fat embolism
16. **Characteristic feature of best sputum sample is /are?** (PGI Nov 2015)
  - a. Presence of leukocytes
  - b. Respiratory epithelium
  - c. Neutrophils
  - d. Mucus with inflammatory cells
  - e. Presence of alveolar macrophages
17. **Heart failure cells are actually?** (Recent Question 2015)
  - a. Alveolar macrophages
  - b. Type I pneumocytes
  - c. Type 2 pneumocytes
  - d. Pulmonary edema fluid cells
18. **Heart Failure cells are:** (Recent Question 2014, 2013)
  - a. Lipofuscin granules in cardiac cells
  - b. Pigmented alveolar macrophages
  - c. Pigmented pancreatic acinar cells
  - d. Pigment cells seen in liver





19. **Terminal stage of pneumonia is:** (Recent Question 2014)
  - a. Congestion
  - b. Red hepatization
  - c. Gray hepatization
  - d. Resolution
20. **ARDS is due to defect in?** (AIIMS May 2014)
  - a. Type 1 pneumocytes
  - b. Type 2 pneumocytes
  - c. Endothelial cells
  - d. Clara cells
21. **All are recognized causes of Adult Respiratory Distress Syndrome (ARDS) EXCEPT:** (APPGMEE 14)
  - a. Smoke inhalation
  - b. Malignant hypertension
  - c. Hypothermia
  - d. Viral pneumonias
22. **In Hyaline Membrane Disease the pathology in the lung consists of:** (Recent Question 2013)
  - a. Albumin and complement
  - b. Fibrin
  - c. Precipitated surfactant
  - d. Mucus
23. **Heart failure cells contain:** (Recent Question 2013)
  - a. Hemosiderin
  - b. Lipofuscin
  - c. Myoglobin
  - d. Albumin
  - e. Pneumonia
24. **Patient with h/o long standing depressive illness come to emergency with acute breathlessness. The X-ray shows diffuse infiltrates with predominance in right middle lobe and right lower lobe. The patient did not survive and the following picture in the lungs was seen on autopsy?** (AIIMS Nov 2017)




- a. Severe necrosis with fungal hyphae, severe fungal pneumonia
  - b. Coagulation necrosis, Tuberculosis
  - c. Vegetable matter; Aspiration pneumonia
  - d. Severe necrosis; severe necrotizing pneumonia
25. **In the stage of Grey hepatization:** (Recent Question 2013)
    - a. WBCs fill the alveoli
    - b. RBCs fill the alveoli
    - c. Organisms fill the alveoli
    - d. Accumulation of fibrin in alveoli
  26. **The most common fate of lobar pneumonia is:** (Recent Question 2013)
    - a. Consolidation
    - b. Resolution
    - c. Abscess formation
    - d. Empyema


27. **Characteristic histopathological finding in SHOCK Lung:** (AI 12, AIIMS Nov 07, May 08)
  - a. Diffuse alveolar necrosis
  - b. Interstitial pulmonary edema
  - c. Diffuse interstitial inflammation
  - d. Intra-cellular debris

## TUBERCULOSIS

28. **Histological picture of a lesion excised from the right cervical region is shown below. What is your diagnosis?** (Recent Pattern Question 2020)



- a. Necrotizing granulomatous inflammation
  - b. Neurofibroma
  - c. Schwannoma
  - d. Hodgkin lymphoma
29. **A patient underwent lung transplantation. The resected lung from the patient showed following features. What could be your possible diagnosis?** (Recent exam 2018)




- a. Bronchiectasis
  - b. Lung abscess
  - c. Lung carcinoma
  - d. Miliary tuberculosis
30. **True about miliary tuberculosis:** (PGI May 2016)
    - a. Occur primarily due to hematogenous spread
    - b. Miliary lesion is generally of size 1-2 mm
    - c. Diffuse bilateral crepitation is always present
    - d. Onset is generally acute
    - e. Sputum smear microscopy is negative in 80% of cases
  31. **MC site of TB reactivation in lung is?** (Recent Question 2016)
    - a. Apex
    - b. Base
    - c. Subpleural
    - d. Near bronchus
  32. **TB infects which cell** (Recent Question 2016)
    - a. Macrophage
    - b. Lymphocyte
    - c. Neutrophils
    - d. Eosinophils




- 33. Miliary TB is:** (Recent Question 2014)
- Primary
  - Post-primary
  - Extra-pulmonary
  - None
- 34. All of the following are lesions seen in primary tuberculosis except:** (Recent Question 2015)
- Simon's focus
  - Ghon's focus
  - Ranke complex
  - Puhl's lesion
- 35. Primary site of involvement in congenital TB:** (Recent Question 2015)
- Lungs
  - Liver
  - Lymph nodes
  - Stomach
- 36. Most common site of gastrointestinal TB:** (Recent Question 2014)
- Stomach
  - Duodenum
  - Terminal ileum
  - Colon
- 37. Miliary TB occurs due to spread via:** (Recent Question 2014)
- Arteries
  - Veins
  - Lymphatics
  - Direct invasion
- 38. Ghons complex refers to:** (Recent Question 2013)
- Healed parenchymal lesions
  - Necrotic lymph nodes
  - Parenchymal lesion along with inflamed lymph nodes
  - Complication in enlarged hilar lymph nodes
- 39. In TB, cytokine which plays a major role in conversion of macrophage to epithelioid cell is?** (JIPMER 2013)
- IFN- $\gamma$
  - TNF
  - IL-12
  - Macrophage chemoattractant protein
- 40. Infraclavicular lesion of tuberculosis is known as:** (AIIMS 11)
- Ghon's focus
  - Puhl's focus
  - Assmans focus
  - Simmon's focus
- 41. All of the following statements about Interferon gamma release assays are true (IGRA) except?** (DNB June 2012)
- More specific than tuberculin skin testing
  - ESAT-6 and CFP-10 are the antigens used
  - Quantitative and qualitative measurement of Interferon gamma released by mycobacterium tuberculosis in the body
  - Lesser cross-reactivity to BCG than tuberculin testing
- 42. The most important function of epithelioid cells in tuberculosis is:** (DPG 10)
- Phagocytosis
  - Secretory
  - Antigenic
  - Healing

## OBSTRUCTIVE DISEASES OF LUNGS


- 43. A 30-year-old male presented with history of dyspnea, cough and sputum production. The patient died of respiratory failure. Gross image of lung is shown below. What is the likely etiology?** (Recent Pattern Question 2020)



- Cystic fibrosis
  - Mutation in dynein arms
  - Alpha 1 antitrypsin deficiency
  - Antibodies against type IV collagen
- 44. A 75-year-old male, known smoker presented to pulmonology department with history of cough. Biopsy was taken which showed the following. What is the change shown?** (Recent Pattern Question 2020)




- Dysplasia
  - Metaplasia
  - Hyperplasia
  - Atrophy
- 45. The main diagnostic criteria of ABPA are all except?** (APPGMEE 2015)
- Pulmonary infiltrates
  - Bronchial asthma
  - Distal bronchiectasis
  - Eosinophilia
- 46. Which of the following structures in the lung is likely to be affected the most in a patient who smoked a pack and half of cigarettes per day for 30 years and developed centrilobular emphysema?** (AP 2013)
- Alveolar sac
  - Terminal bronchiole
  - Alveolar duct
  - Respiratory bronchiole
- 47. Chronic bronchitis can be a premalignant condition, which involves:** (Recent Question 2015)
- Columnar to squamous
  - Squamous to columnar
  - Squamous to cuboidal
  - Cuboidal to squamous
- 48. Emphysema pathologically involves beyond the:** (Recent Question 2015)
- Bronchi
  - Terminal bronchiole
  - Respiratory bronchiole
  - Alveolar Sac



49. **Commonest type of emphysema is:** (Recent Question 2014)  
a. Centriacinar                      b. Obstructed  
c. Irregular                         d. Panacinar
50. **All are obstructive lung disease except:** (Recent Question 2014)  
a. Emphysema                      b. Interstitial fibrosis  
c. Asthma                            d. Bronchitis
51. **Curshmann's crystals are seen in:** (Recent Question 2014)  
a. Bronchial asthma                b. Bronchiectasis  
c. Chronic bronchitis              d. Wegener's granulomatosis
52. **Which of the following finding, composed of shed epithelium with thick mucus are seen in bronchial asthma?** (JIPMER 2014)  
a. Creola body                      b. Councilman body  
c. Curshmann spirals              d. Charcot leyden crystals
53. **Hyperplasia of smooth muscle of airway is seen in?** (Recent Question 2013)  
a. Emphysema                      b. Asthma  
c. Alveolar proteinosis            d. Bronchiectasis
54. **Creola bodies are seen in:** (Recent Question 2013)  
a. Bronchial asthma                b. Chronic bronchitis  
c. Emphysema                      d. Bronchiectasis
55. **Bronchiectasis means -----of bronchi:** (Recent Question 2013)  
a. Inflammation                    b. Dilatation  
c. Cavitation                        d. All
56. **Reid index is useful in:** (AI 12)  
a. Glomerulonephritis            b. Cirrhosis  
c. Chronic bronchitis              d. ARDS
57. **Centriacinar emphysema primarily involves:** (PGI Nov 2011)  
a. Upper lobe                        b. Middle lobe  
c. Lower lobe                        d. All lobes  
e. Lower part of upper lobe
58. **Difference between bronchial asthma and COPD is:** (Jipmer 11)  
a. Reversible bronchoconstriction  
b. Hyperventilation on chest X ray  
c. Acute exacerbation by URTI  
d. Decreased FEV1/FVC
59. **Which of the following is NOT a complication of bronchiectasis:** (AIIMS Sep 10)  
a. Lung abscess                      b. Lung cancer  
c. Amyloidosis                      d. Empyema
60. **Reid's Index is?** (DNB Dec 10)  
a. Increased in Chronic Bronchitis  
b. Decreased in Chronic Bronchitis  
c. Increased in Bronchial Asthma  
d. Decreased in Bronchial Asthma
61. **Which one of the following is NOT a feature of Kartagener's syndrome:** (UPSC 09), (WBPB 2016)  
a. Bronchiectasis  
b. Ciliary dyskinesia  
c. Dysphagia  
d. Situs inversus

## INTERSTITIAL LUNG DISEASES

62. **A Factory worker was working in a factory from past 20 years , and now presenting with pleural thickening and fibrosis. Histopathology of lesion is shown in below image. Most likely diagnosis is?** (AIIMS May 18)




- a. Asbestosis  
b. Cotton Fiber  
c. Coal Worker Pneumoconiosis  
d. Silicosis
63. **Silicosis biopsy features and radiological correlation?** (AIIMS May 2017)  
a. Lower lobe involved  
b. Dense collagen and calcifications in the lymph nodes seen  
c. Progressive massive fibrosis can be seen as late complication  
d. Immune granuloma can be seen  
e. Macules may be seen
64. **The lung pathology occurring in persons working in cotton- wool industries is** (Recent Question 2016-17)  
a. Asthma like features  
b. Hypersensitivity pneumonitis  
c. Lung Ca  
d. Chronic bronchitis
65. **Bagassosis occurs in people working in which industry?** (Recent Question 2016-17)  
a. Silica                                b. Wallboard paper  
c. Cotton                              d. Asbestosis
66. **Most common cause for lung abscess?** (Recent Question 2016-17)  
a. Staph aureus                      b. Staph pyrogen  
c. Bacteroids                        d. Klebsiella
67. **APBA is associated with?** (Recent Question 2016-17)  
a. Central bronchiectasis  
b. Bronchitis  
c. Midlung bronchiectasis  
d. Peripheral bronchiectasis
68. **All are true about hypersensitivity pneumonitis except:** (Recent Question 2016-17)  
a. Type IV Hypersensitivity reaction  
b. More common in smoker  
c. Bronchoalveolar lavage shows CD4+ and CD8+ T lymphocytes  
d. May presents with cough, dyspnea & breathlessness





69. Cavitary lesion in right lower lung with dyspnoea with following histopathological appearance



**Most likely diagnosis:**

(AIIMS Nov 2015)

- a. Echinococcus with 2 layers
  - b. Strongyloides with 2 layers
  - c. Paragonimus with 2 layers
  - d. Cysticercosis with 3 layers
70. Causative agent of Farmer's lung is:
- a. Thermophilus actinomycetes (Recent Question 2015)
- b. Aspergillus
- c. Penicillium glabrum
- d. Rhizopus
71. Causative particle in asbestosis is?
- (Recent Question 2015)
- a. Amphibole                      b. Crysolite
- c. Tridymite                      d. Gristobalite
72. All are true about silicosis except? (PGI Nov 2015)
- a. Bifringent crystals seen
- b. Pleural plaque seen
- c. Lower lobe is usually involved
- d. Most common pneumoconiosis
73. Asbestosis causes? (Recent Question 2014)
- a. Lymphoma                      b. Leukaemia
- c. Renal cell carcinoma                      d. Mesothelioma
74. Anthracosis is due to inhalation of:
- (Recent Question 2013)
- a. Coal dust                      b. Asbestos dust
- c. Silica dust                      d. Beryllium dust
75. Ferruginous bodies are seen in? (DNB Aug 12)
- a. Sarcoidosis
- b. Silicosis
- c. Asbestosis
- d. Coal worker's pneumoconiosis
76. A female presents with history of progressive breathlessness. Histology shows heterogenous patchy fibrosis with several fibroblastic foci. The most likely diagnosis is: (AI 11)
- a. Cryptogenic organizing pneumonia
- b. Non specific interstitial pneumonia
- c. Usual interstitial pneumonia
- d. Desquamative interstitial pneumonia
77. Schaumann bodies are seen in: (AIIMS 11)
- a. Sarcoidosis                      b. Chronic bronchitis
- c. Asthma                      d. Syphilis
78. Asteroid bodies are seen in? (DNB Dec 11, June 10)
- a. Sarcoidosis                      b. Syphilis
- c. Chromoblastomycosis                      d. Sporotrichosis

79. The following does not occur with asbestosis: (DPG 11, DNB Dec 08)

- a. Methemoglobinemia
- b. Pneumoconiosis
- c. Pleural mesothelioma
- d. Pleural calcification

80. Laminated concretions of calcium and proteins are: (Maharashtra 10)

- a. Schaumanns bodies
- b. Ferruginous bodies
- c. Asteroid bodies
- d. Gamma Gandy bodies

81. The dangerous particle size causing pneumoconiosis varies from:

- a. 100-150 μm
- b. 50-100 μm
- c. 10-50 μm
- d. 1-5 μm

## TUMORS OF LUNGS & PLEURA


82. WHO 2015 new inclusion in lung squamous cell carcinoma is/are? (AIIMS May 2017)

- a. Basaloid type
- b. Lymphoepithelioid type
- c. Papillary type
- d. Clear cell variety
- e. Small cell variety

83. Tumor marker for lung adenocarcinoma?

- a. ck7
- b. ck20 (AIIMS May 2017)
- c. TTF1
- d. CK 11
- e. Berep-4

84. A 34 yr woman presented with coughing, dyspnea, flushing, diarrhea, hypotension, hematemesis for 3 months, Bronchoscopy shows large intraductal mass. Histopathology of the mass is given below. Mitosis was 5/hpf and is chromogranin positive. What is your diagnosis? (AIIMS May 2016)



- a. Small cell CA
  - b. Large cell CA
  - c. Typical carcinoid tumor stage IV
  - d. Atypical carcinoid Stage IV
85. Most common lung malignancy in woman and with smokers less than 10 packet cigarette per year? (Recent Question 2016-17)
- a. Small cell Ca- b. Sq cell Ca
- c. Adenoca
- d. Carcinoids

86. Mutations associated with nonsmall cell Lung ca is? (Recent Question 2016-17)

a. p 53- b. EGFR
- c. Rb
- d. Myc

87. Lung Ca metastasize early is? (Recent Question 2016-17)

a. Adeno ca- b. Sq cell Ca
- c. Small cell Ca
- d. Large cell ca

88. Carcinoid of lung (bronchial adenoma) arise from (JIPMER 2015)

a. Ciliated cell- b. Kulchitsky cell
- c. Type 2 pneumocytes
- d. Clara cell





89. **Lymphoma like picture in lung cancer is seen in which subtype?** (Recent Question 2015)
  - a. Squamous cell carcinoma
  - b. Adenocarcinoma
  - c. Small cell carcinoma
  - d. Large cell carcinoma
90. **Which of the following is the marker for mesothelioma:** (Recent Question 2015)
  - a. ck7
  - b. ck22
  - c. ck 5/6
  - d. TTF-1
91. **Which of these is used as a marker in mesothelioma?** (Recent Question 2015)
  - a. Calretinin
  - b. TTF-1
  - c. CK-8/9
  - d. Glypican
92. **Most common posterior mediastinal tumor is?** (Recent Question 2015)
  - a. Thymoma
  - b. Neuroma
  - c. Chordoma
  - d. Pleuroma
93. **Which malignancy is most commonly associated with asbestos?** (Recent Question 2015)
  - a. Malignant mesothelioma
  - b. Benign Pleural fibroma
  - c. Squamous Cell Ca Lung
  - d. Carcinoids
94. **Ectopic ACTH production is seen in:**
  - a. Small cell carcinoma is lung (Recent Question 2014)
  - b. Anaplastic carcinoma of lung
  - c. Squamous cell carcinoma of lung
  - d. Adenocarcinoma of cerebellum
95. **Primary pleural tumor is:** (Recent Question 2014)
  - a. Mesothelioma
  - b. Myxoma
  - c. Lipoma
  - d. Fibroma
96. **Most common mediastinal tumor is:** (Recent Question 2014)
  - a. Neurogenic tumor
  - b. Pericardial cyst
  - c. Hernia
  - d. Teratoma
97. **A 60 yr old person presents with a mass located at central bronchus causing distal bronchiectasis and recurrent pneumonia. Which of the following findings is expected from biopsy of the mass?** (AIIMS Nov 14)
  - a. Abundant osteoid matrix formation
  - b. Contains all three germ layers
  - c. Spindle cells with abundant stromal matrix
  - d. Small round cells and hyperchromatic nuclei with nuclear moulding
98. **A hyperplastic mass containing neuroendocrine cells in an area of chronic inflammation and scarred tissue of lung is called?** (JIPMER 2014)
  - a. Carcinoid
  - b. Tumorlet
  - c. Hamartoma
  - d. Teratoma
99. **Small cell cancer commonly metastasizes to:** (Recent Question 2014)
  - a. Brain
  - b. Liver
  - c. Adrenal
  - d. Kidney
100. **Most common type of carcinoma lung is:** (Recent Question 2013)
  - a. Small cell carcinoma
  - b. Adenocarcinoma
  - c. Squamous cell carcinoma
  - d. Large cell carcinoma
101. **Marker of small cell cancer of lung is:** (Recent Question 2013)
  - a. Chromogranin
  - b. Cytokeratin
  - c. Desmin
  - d. Vimentin
102. **Which paraneoplastic syndrome is not seen with Small Cell Ca Lung:** (Recent Question 2013)
  - a. PTH
  - b. ACTH
  - c. ADH
  - d. Carcinoid syndrome
103. **Carcinoid tumor develops from:** (Recent Question 2013)
  - a. Enterochromaffin cells
  - b. Neuroectoderm
  - c. J cells
  - d. Goblet cells
104. **A 60-year-old male had a chronic history of exposure of asbestosis. He now presents with a mass in the apex of right lung. Which of the following would be seen on electron microscopy of a biopsy from the lesion?** (AIIMS Nov 2013)
  - a. Melanosomes
  - b. Neurosecretory granules
  - c. Numerous long slender microvilli
  - d. Desmosomes with secretory endoplasmic reticulum
105. **True about lung carcinoma:** (PGI Dec 13)
  - a. Squamous cell Ca is most common carcinoma
  - b. Squamous cell Ca cause myopathy
  - c. Small cell Ca has best prognosis
  - d. BronchoalveolarCa involves proximal airways
  - e. Hypercalcemia is common with Small cell Carcinoma
106. **Immunohistochemical marker used for detection of AdenoCa lung?** (JIPMER 2013)
  - a. TTF
  - b. GFAP
  - c. Progesterone Receptors
  - d. AFP
107. **Paraneoplastic syndromes are most commonly associated with?** (JIPMER 2013)
  - a. Bronchial adenoCa
  - b. BronchoalveolarCa
  - c. Small cell Ca
  - d. Bronchial Carcinoid
108. **Incorrect statement about Small cell Ca of lung:** (PGI May 12)
  - a. Not associated with smoking
  - b. Surgical resection alone is the treatment of choice
  - c. Associated with paraneoplastic syndrome
  - d. Most patients have distant metastases on diagnosis
  - e. Contain neurosecretory granules
109. **Which of the following causes malignant mesothelioma?** (DNB Aug 12)
  - a. Smoking
  - b. Asbestosis
  - c. Silicosis
  - d. Pneumoconiosis
110. **Histological findings in bronchoalveolar carcinoma includes:** (PGI Nov 2011)
  - a. Clara cells
  - b. Adenosquamous
  - c. Mucin secreting cells
  - d. Type II pneumocytes
  - e. Neuroendocrine cells
111. **Following is true about bronchial carcinoids:** (JIPMER 11)
  - a. Highly radiosensitive
  - b. Metastasis common
  - c. Carcinoid syndrome does not manifest
  - d. Commonly arise from terminal bronchioles



**112. PTH like substance is produced by which type of lung malignancy:** (JIPMER 11)

- Squamous cell carcinoma
- Oat cell carcinoma
- Adeno carcinoma
- Large cell carcinoma

**113. True about lung carcinoma:** (AI 10)

- More than 75% of lung cancers are squamous cell type
- Oat cell carcinoma frequently show cavitation
- Lung calcification is characteristically seen in oat cell carcinoma
- Oat cell carcinoma is commonly associated with bilateral hilar lymphadenopathy

**114. On biopsy, characteristic finding of malignant mesothelioma is:** (AIIMS May 10)

- Myelin
- Desmin
- Weibel-palade bodies
- Branching microvilli
- Fibrosis

**115. Which of the following is not true about Bronchoalveolar carcinoma:** (Maharashtra 10)

- Adenocarcinoma
- Stromal invasion with desmoplasia
- Preservation of alveolar structure
- Grows along pre-existing anatomical structures



## Answers with Explanations

**1. Ans. (c) Acute Respiratory infection**

Ciliocytophthoria (CCP) defines a degenerative process of the ciliated cells consequent to viral infections after acute respiratory infections, and it is characterized by typical morphological changes.

**2. Ans. (c) May have independent venous drainage; (d) Intralobular variety more common; (e) May be associated with other congenital anomalies** (Ref: Robbins 9th/pg 670)

### Sequestration

- Discrete area of lung tissue that lacks any connection to the airway system
- Abnormal blood supply arising from the aorta
- Venous return to the right side of the heart through IVC (extralobar) or pulmonary veins (intralobar)

**3. Ans. (b) Asthenospermia**

Asthenozoospermia (or asthenospermia) is the medical term for reduced sperm motility due to defect in ciliary movement

**4. Ans. (d) Bradykinin** (Ref: Robbins 9th/pg 669-670)

Bronchial mucosa contains **neuroendocrine cells** that have neurosecretory-type granules releasing: **serotonin<sup>Q</sup>, calcitonin<sup>Q</sup>, and gastrin-releasing peptide (bombesin)<sup>Q</sup>**

**5. Ans. (a) Bronchogenic** (Ref: Robbins 9th/pg 670; 8th/pg 679)

**6. Ans. (a) Aorta** (Ref: Robbins 9th/pg 670; 8th/pg 679)

**7. Ans. (d) Congenital cystic adenomatoid malformation**

(Ref: Robbins 9th/pg 670; 8th/pg 679)

**Congenital cystic adenomatoid malformation (CCAM): Hamartomatous<sup>Q</sup> or dysplastic lung tissue, usually confined to one lobe.**

**8. Ans (a) Ventral wall of foregut** (Ref: R 9th/pg 669-670)

- Respiratory system develops from: ventral wall of foregut (formed when embryo is 4 weeks old) while** Cartilagenous, muscular & connective tissue of trachea and lungs are derived from splanchnic mesoderm.

**9. Ans. (d) Ciliated pseudostratified columnar epithelium**

(Ref: Robbins 9th/pg 670; 8th/pg 679)

- A **bronchogenic cyst is rarely connected to the tracheobronchial tree.**
- Microscopically, the cyst is lined by ciliated pseudostratified columnar epithelium with squamous metaplasia occurring in areas of inflammation. The wall contains bronchial glands, cartilage, and smooth muscle.

**10. Ans. (d) Clara cells** (Ref: Robbins 9th/pg 670; 8th/pg 679)

**Club cells**, also known as **bronchiolar exocrine cells**, originally known as **Clara cells**, are **dome-shaped cells with short microvilli**, found in the **small airways (bronchioles)** of the lungs, produce substances similar to surfactant.

**11. Ans. (b) Alveoli** (Ref: Robbins 9th/pg 669-670; 8th/pg 678)

**Pores of Kohn<sup>Q</sup>** permit the passage of **bacteria** and **exudate** between adjacent alveoli.

**12. Ans. (d) CMV pneumonia**

Large cell with basophilic intranuclear inclusion is the clue

**13. Ans. (d) Occurs in babies born post-dates**

(Ref: Robbins 9th/pg 672)

**14. Ans. (a) Mucus plug in alveoli** (Ref: R 9th/pg 672-673)

**15. Ans. (d) Fat embolism** (Ref: Robbins 9th/pg 672-673)

**Etiology of ARDS & ALI**



16. Ans. (a, d) **a. Presence of leukocytes; d. Mucus with inflammatory cells**

(Ref: Mayo Clin Proc. 1975 Jun; 50(6):339-44)

Good quality sputum samples have:

- Should not be contaminated by oropharyngeal flora
- $\geq 10$  leukocytes with mucus, but  $< 25$  squamous epithelial cells per low-power field (LPE,  $\times 100$ ),

17. Ans. (a) **Alveolar macrophages** (Ref: R 9th/pg 669-670)

18. Ans. (b) **Pigmented alveolar macrophages**

(Ref: Robbins 9th/pg 669-670; 8th/pg 678)

19. Ans. (d) **Resolution** (Ref: Robbins 9th/pg 704-705)

20. Ans. (a) **Type I pneumocytes**

(Ref: Robbins 9th/pg 672-673)

**In ARDS**

- Initially, there is destruction of **type I pneumocytes**<sup>Q</sup> → **hypoxemia**
- Later damage and necrosis of **type II alveolar pneumocytes**<sup>Q</sup> → leads to **surfactant deficiency**<sup>Q</sup>

21. Ans. (b) **Malignant hypertension** (Ref: R 9th/pg 672-673)

22. Ans. (b) **Fibrin** (Ref: Robbins 9th/pg 672-673)

**Hyaline membrane consists of** fibrin-rich edema fluid with **necrotic epithelial cells**.

23. Ans. (a) **Hemosiderin** (Ref: Robbins 9th/pg 669-670)

24. Ans. (c) **Vegetable matter; Aspiration pneumonia**

25. Ans. (d) **Accumulation of fibrin in alveoli**

(Ref: Robbins 9th/pg 704-705; 8th/pg 712-713)

26. Ans. (b) **Resolution** (Ref: Robbins 9th/pg 704-705)

- In **most cases of pneumonia, resolution** occurs;
- $< 10\%$  of patients have pneumonia severe enough to merit hospitalization & in most such instances death results from a complication, as stated below;

27. Ans. (a) **Diffuse alveolar necrosis**

(Ref: R 9th/pg 672-673)

ARDS is also called "**shock lung**"

28. Ans. (a) **Necrotizing granulomatous inflammation**

(Ref: Robbins 9th/pg 376)

29. Ans. (d) **Miliary tuberculosis**

(Ref: Robbins 9th ed p376)

Miliary pulmonary disease occurs when organisms draining through lymphatics enter the venous blood and circulate back to the lung. Individual lesions are either microscopic or small, visible (2-mm) foci of yellow-white consolidation scattered through the lung parenchyma (the adjective "miliary" is derived from the resemblance

of these foci to millet seeds). Miliary lesions may expand and coalesce, resulting in consolidation of large regions or even whole lobes of the lung. With progressive pulmonary tuberculosis, the pleural cavity.

30. Ans. (a, b, e) **a. Occur primarily due to hematogenous spread, b. Miliary lesion is generally of size 1-2 mm, e. Sputum smear microscopy is negative in 80% of cases**

(Ref: Robbins 9th/pg 375-376; 8th/pg 370-371)

31. Ans. (a) **Apex**

### Postprimary (Adult-Type) Disease

- Also referred to as **reactivation** or **secondary TB**, postprimary TB is probably most accurately termed **adult-type TB**, since it may result from endogenous reactivation of distant latent infection or recent infection (primary infection or reinfection).
- It is usually localized to the apical and posterior segments of the upper lobes, where the substantially higher mean oxygen tension (compared with that in the lower zones) favors mycobacterial growth

32. Ans. (a) **Macrophage**

(Ref: Robbins 9th/pg 375; 8th/pg 370)


- **Mycobacterium tuberculosis** enters into macrophages with the help of **mannose binding lectin**<sup>Q</sup> and **CR3**<sup>Q</sup>.
- **Macrophages**<sup>Q</sup> are the **primary cells infected** by *M. tuberculosis*

33. Ans. (b) **Post-primary** (Ref: Robbins 9th/pg 375-376)

Fate of primary TB:

- **95% cases:** Cell mediated immunity controls infection → Calcified healed lesions & hilar lymph nodes formed (**Ranke complex**).
- **5% cases:** Primary tuberculosis is progressive (**Progressive Primary TB**) → resembles acute bacterial pneumonia → **Lympho-hematogenous dissemination** may cause **tuberculous meningitis** &/or **miliary tuberculosis**.

34. Ans. (d) **Puhl's lesion** (Ref: Robbins 9th/pg 375-376)





35. Ans. (b) **Liver** (Ref: Nelson 19 ed Chap 212)

In congenital TB, primary organ involved is liver.

- Congenital tuberculosis is rare because the most common result of female genital tract tuberculosis is infertility.
- Congenital transmission usually **occurs from a lesion in the placenta** through the umbilical vein.
- **Primary infection in the mother** just before or during pregnancy is more likely to cause congenital infection than is reactivation of a previous infection.
- The tubercle bacilli **first reach the fetal liver**, where a primary focus with periportal lymph node involvement may occur.

Organisms pass through the liver into the main fetal circulation and infect many organs.

36. Ans. (c) **Terminal ileum** (Ref: Robbins 9th/pg 375-376)

Extrapulmonary TB:

- **MC site is lymph node<sup>o</sup>**, MC cervical & supraclavicular ("Scrofula")<sup>o</sup>
- **Pleural involvement: Exudative<sup>o</sup> Pleural effusion**, tuberculous empyema, or obliterative fibrous pleuritic

Genitourinary TB: sterile pyuria<sup>o</sup>; Preferentially involves fallopian tube<sup>o</sup> in females & epididymis<sup>o</sup> in males

- **Skeletal TB**: Most common site **spine<sup>o</sup> (Pott's disease)** <sup>o</sup>> hip > knee
- **TB meningitis** (paresis of cranial nerves especially ocular, is frequent finding)<sup>o</sup>
- **In Gastro-intestinal (GI) TB**: MC site **terminal ileum and caecum<sup>o</sup>**

37. Ans. (a) **Arteries** (Ref: Robbins 9th/pg 375-376)

- Miliary TB is due to **hematogenous spread** of tubercle bacilli.
- Although in **children** it is often the consequence of **primary infection**, in **adults** it may be due to either **recent infection or reactivation of old disseminated foci**.
- Lesions are usually yellowish **1-2 mm granulomas** that resemble millet seeds (thus the term *military*)
- **Miliary pulmonary disease** occurs when organisms draining through lymphatics enter the venous blood and circulate back to the lung
- **Systemic miliary tuberculosis** occurs when bacteria disseminate through the **systemic arterial system**.
- Miliary tuberculosis is most prominent in the **liver, bone marrow, spleen, adrenals, meninges, kidneys, fallopian tubes, and epididymis**, but could involve any organ

38. Ans. (c) **Parenchymal lesion along with inflamed lymph nodes** (Ref: Robbins 9th/pg 375-376; 8th/pg 370-371)

- **Ghon's focus<sup>o</sup>**: Grey white area of **inflammation with consolidation**
- **Ghon's complex<sup>o</sup>**: Ghon's focus + **inflamed regional lymph nodes**

39. Ans. (a) **IFN-g** (Ref: Robbins 9th/pg 371-372; 8th/pg 368)

40. Ans. (c) **Assmans focus** (Ref: Robbins 9th/pg 375-376)

- Classically involves **apical and posterior segments of upper lobe** due to high O<sub>2</sub> concentration (**Puhl's lesion<sup>o</sup>**)
- Infraclavicular lesion** is called **Assman's Focus**; Refer to Ans 44 above;

41. Ans. (c) **Quantitative and qualitative measurement of Interferon gamma released by mycobacterium tuberculosis in the body** (Ref: Harrison 18th/chapter 165)

- **IGRAs are more specific** than the TST as a result of **less cross-reactivity due to BCG vaccination** and sensitization by **non-tuberculous mycobacteria**
- Measure **T cell release of IFN-** in response to stimulation with the highly **TB-specific antigens ESAT-6 and CFP-10**
- This test is **performed in vitro** & not in vivo, hence the answer is option C

42. Ans. (b) **Secretory**

(Ref: Immunology, 8th edition, by David Male; pg 427)

- **Epithelioid cells** are transformed macrophages that have **lost their phagocytic function** but **retained their secretory activity**.
- Epithelioid cells contain **numerous endoplasmic reticulum & golgi bodies**.
- They **secrete IL10, TNF alpha, TGF beta**;

43. Ans. (c) **Alpha 1 antitrypsin deficiency** (Ref: R 9th pg 675)

Features are suggestive of Emphysema

44. Ans. (b) **Metaplasia** (Ref: Robbins 9th/pg 675)

45. Ans. (c) **Distal bronchiectasis** (Ref: Harrison 18th/pg 2120)

46. Ans. (d) **Respiratory bronchiole**

(Ref: Robbins 9th/pg 675)

Cigarette smoking is most likely to cause damage to smaller segments of airway like acinus

47. Ans. (a) **Columnar to squamous**

(Ref: Robbins 9th/pg 679)

Chronic bronchitis is an example of squamous metaplasia in which normal columnar epithelium of respiratory tract is replaced by squamous epithelium;

48. Ans. (b) **Terminal bronchiole**

(Ref: Robbins 9th/pg 675-676)

Emphysema is defined as **irreversible dilatation and destruction<sup>o</sup>** of the airspaces **distal to the terminal bronchiole (acinus)<sup>o</sup>**, without fibrosis.<sup>o</sup>

49. Ans. (a) **Centriacinar** (Ref: Robbins 9th/pg 675-676)

Centriacinar emphysema is the most common form, constituting more than 95% of clinically significant cases.

50. Ans. (b) **Interstitial fibrosis**

(Ref: Robbins 9th/pg 674-675)

Obstructive lung diseases include: Emphysema, Chronic Bronchitis, Asthma, Bronchiectasis,





51. **Ans. (a) Bronchial asthma** (Ref: Robbins 9th/pg 679-680)
- Curschmann spirals: extrusion of **mucusplugs** from subepithelial mucous gland ducts or bronchioles.<sup>Q</sup>
52. **Ans. (c) Curshmann spirals**
- (Ref: Robbins 9th/pg 679-680)
53. **Ans. (b) Asthma** (Ref: Robbins 9th/pg 679-680)
- Characteristic histologic findings ("airway remodeling") of asthma include: Hypertrophy and/or hyperplasia of the bronchial wall muscle<sup>Q</sup>
54. **Ans. (a) Bronchial asthma** (Ref: Robbins 9th/pg 679-680)
55. **Ans. (b) Dilatation** (Ref: Robbins 9th/pg 683-684)
- Bronchiectasis refers to **destruction of smooth muscle and elastic tissue** by chronic necrotizing infections leading to permanent<sup>Q</sup> dilation of bronchi and bronchioles.<sup>Q</sup>
56. **Ans. (c) Chronic bronchitis** (Ref: Robbins 9th/pg 679)
57. **Ans. (a) Upper lobe** (Ref: Robbins 9th/pg 675-676)
58. **Ans. (a) Reversible bronchoconstriction**
- (Ref: Robbins 9th/pg 674-675; 8th/pg 683-684)
59. **Ans. (b) Lung cancer** (Ref: Robbins 9th/pg 683-684)
60. **Ans. (a) Increased in Chronic Bronchitis**
- (Ref: Robbins 9th/pg 679; 8th/pg 687-688)
61. **Ans. (c) Dysphagia** (Ref: Robbins 9th/pg 683-684)
- Kartagener syndrome refers to **bronchiectasis, sinusitis, and situs inversus**;<sup>Q</sup>
  - It is seen in **50% patients** with primary ciliary dyskinesia
62. **Ans. (a) Asbestosis**
- Look at knobbed ends and brown colour, this is suggestive of asbestos bodies. As they also contain iron, so they take blue colour on perls stain
63. **Ans. (b, c, d) b. Dense collagen and calcifications in the lymph nodes seen; c. Progressive massive fibrosis can be seen as late complication; d. Immune granuloma can be seen**
64. **Ans. (a) Asthma like features** (Ref: Robbins 9th 689)
- In 10-25% cases, disease may be progressive, with chest tightness recurring or persisting throughout the workweek.
  - After >10 years of exposure, workers with recurrent symptoms are more likely to have an **obstructive pattern** on pulmonary function testing.
65. **(b) Wallboard paper** (Ref: Robbins 9th/pg 688)

66. **Ans. (a) Staph aureus** (Ref: Robbins 9th pg)
67. **Ans. (a) Central bronchiectasis**
- (Ref: Robbins 9/683)
- Predominant involvement of the central airways is reported in association with allergic bronchopulmonary aspergillosis (ABPA), in which an immune-mediated reaction to *Aspergillus* damages the bronchial wall.
68. **Ans. (b) More common in smoker**
- (Ref: Robbins 9th/694-95; Davidson 22nd/719-20; Harrison 19th/ 1681-83)
69. **Ans. (c) Paragonimus with 2 layers**
- (Ref Harrison's 19th ed/1429; CDC website)
- The figure shows: **Eggs of Paragonimus sp. taken from a lung biopsy stained with hematoxylin and eosin**
70. **Ans. (a) Thermophilus actinomycetes**
- (Ref: Robbins 9th/pg 689; 8th/pg 687)
- In *farmer's lung*, inhalation of proteins, such as thermophilic actinomyces bacteria and fungal spores that are present in moldy bedding and feed, are most commonly responsible for the development of Hypersensitivity pneumonitis.
71. **Ans. (a) Amphibole** (Ref: Robbins 9th/pg 689)
- Serpentine<sup>Q</sup> (M.C)** and **Amphibole<sup>Q</sup> (more pathogenic)** as particles are asbestor particle associated with asbestosis.
72. **Ans. (c) Lower lobe is usually involved**
- (Ref: Robbins 9th/pg 689; 8th/pg 687)
- Silicosis involves upper lobe more commonly than lower lobe. Examination of the nodules by polarized microscopy reveals the birefringent silicate particles (silica is weakly birefringent).
73. **Ans. (d) Mesothelioma** (Ref: Robbins 9th/pg 691-692)
74. **Ans. (a) Coal dust** (Ref: Robbins 9th/pg 689; 8th/pg 697)
75. **Ans. (c) Asbestosis** (Ref: Robbins 9th/pg 691-692)
76. **Ans. (c) Usual interstitial pneumonia** (Ref: R 9th/pg 685)
77. **Ans. (a) Sarcoidosis** (Ref: Robbins 9th/pg 693-694)
- Histology in Sarcoidosis:
- Well-formed non-caseating granulomas<sup>Q</sup> composed of an aggregate of tightly clustered epithelioid cells<sup>Q</sup>, with Langhans or foreign body-type giant cells; Central necrosis is unusual.<sup>Q</sup>
  - Schaumann bodies: laminated concretions composed of calcium and proteins<sup>Q</sup>
  - Asteroid bodies: Stellate inclusions enclosed within giant cells<sup>Q</sup>



78. Ans. (a) **Sarcoidosis** (Ref: Robbins 9th/pg 693-694)

79. Ans. (a) **Methemoglobinemia**

(Ref: Robbins 9th/pg 691-692)

80. Ans. (a) **Schaumanns bodies**

(Ref: Robbins 9th/pg 693-694)

81. Ans. (d) **1-5 µm** (Ref: Robbins 9th/pg 689; 8th/pg 697)

For pneumoconiosis, the most dangerous particles are from **1 to 5 µm** in diameter, because particles of this size may reach the **terminal small airways and air sacs and settle in their linings**.

82. Ans. (a) **Basaloid type**

### Squamous Cell Carcinoma - 2015 UPDATES

Number of subtypes have been reduced to three, which makes the diagnosis

- Keratinizing
- Non-keratinizing
- Basaloid squamous cell carcinoma (new category added)

83. Ans. (a, c, e) a. **ck7; b. TTF1; c. BerEP-4**

Adenocarcinoma lung is positive for TTF-1, BerEP-4, CK 7+/ CK 20-. They have EGFR, K-RAS & ALK gene mutation.

84. Ans. (d) **Atypical carcinoid Stage IV**

(Ref: Robbins 9th / pg 231-234; 8th / pg 221-229)

- In the given question features like **coughing, dyspnea, flushing, diarrhea, hypotension, haematemesis** for 3 months is suggestive of a **neu roendocrine tumor** most likely to be **carcinoid**.
- **Morphology of a typical carcinoid tumor:** Nests or trabeculae of medium sized polygonal cells with lightly eosinophilic cytoplasm, low nuclear grade, round to oval finely granular nuclei; may have rosettes or small acinar structures with variable mucin.
- **Also remember:**
  - **Typical- <2 mitosis/10 hpf** (high power fields) and lack Necrosis; **Atypical carcinoids- 2-10 mitosis/hpf** with increased pleomorphism, necrosis, prominent nucleoli<sup>Q</sup>, may cause Carcinoid syndrome<sup>Q</sup>

85. Ans. (c) **Adenoca** (Ref: Robbins 9th/pg 701; 8th/pg 710)

86. Ans. (b) **EGFR**

Mutations in small cell Ca lung include TP53 (75% -90%), RB (~100%), chr 3p deletions and MYC family.

87. Ans. (c) **Small cell Ca (Ref: Robbins 9th/716)**

88. Ans. (b) **Kulchitsky cell** (Ref: Robbins 9th/pg 719)

- Enterochromaffin (EC) cells, or "Kulchitsky cells", are a type of enteroendocrine and neuroendocrine

cell occurring in the epithelia lining the lumen of the digestive tract and the respiratory tract that release serotonin.

- Tumors from these cells results in carcinoid.

89. Ans. (d) **Large cell carcinoma** (Ref: Robbins 9th/pg 715)

**Large cell carcinoma** is an undifferentiated malignant epithelial tumor that lacks the cytologic features of other forms of lung cancer. The cells typically have large nuclei, prominent nucleoli, and a moderate amount of cytoplasm. This gives a lymphoma like picture.

90. Ans. (c) **ck 5/6** (Ref: Robbins 9th/pg 723-724)

Morphology of Malignant mesothelioma:

IHC Shows: Strong positivity for keratin protein, **calretinin<sup>Q</sup>**, **Wilmstum or 1 (WT-1)<sup>Q</sup>**, **cytokeratin 5/6<sup>Q</sup>**, and **D2-40<sup>Q</sup>**.

91. Ans. (a) **Calretinin** (Ref: Robbins 9th/pg 723-724)

92. Ans. (b) **Neuroma** (Ref: Robbins 9th/pg 721; 8th/pg 731)

93. Ans. (a) **Malignant mesothelioma**

(Ref: R 9th/pg 723-724)

**Rare tumor** but risk of developing mesothelioma in heavily asbestos exposed individuals is as high as **7% to 10%**.

94. Ans. (a) **Small cell carcinoma is lung**

(Ref: R 9th/pg 715-717)

**Small cell lung ca is the most common lung Ca** associated with **ectopic hormone production** like **ADH, ACTH, PTH**.

95. Ans. (a) **Mesothelioma** (Ref: Robbins 9th/pg 723-724)

Pleural tumors:

- Solitary fibrous tumor
- Malignant mesothelioma

96. Ans. (a) **Neurogenic tumor** (Ref: Robbins 9th/pg 721)

Neurogenic tumor is the most common mediastinal tumor.

97. Ans. (d) **Small round cells and hyperchromatic nuclei with nuclear moulding** (Ref: Robbins 9th/pg 715-717)

This typical presentation of 60 yr/M presenting with a **mass located at central bronchus** causing distal **bronchiectasis and recurrent pneumonia** is suggestive of small cell Carcinoma.

In Small cell Ca lung, location in lungs are most commonly central and usually presents with mass within the bronchus. Light Microscopy feature of Small cell Ca is:

- Small cells with **salt and pepper pattern, hyperchromatic nuclei<sup>Q</sup>**, **nuclear molding<sup>Q</sup>** is prominent
- **Basophilic staining** of vascular walls due to encrustation by DNA from necrotic tumor cells (**Azzopardi effect**)<sup>Q</sup>

About other options:

- Abundant **osteoid matrix** formation points towards **metastatic Osteosarcoma**;
- Contains **all three germ layers**: is consistent with a **Teratoma**;
- Spindle cells with abundant stromal matrix** is suggestive of **Sarcoma**;



98. Ans. (b) **Tumorlet**

(Ref: Robbins 9th/pg 719-720; 8th/pg 729)

**Benign tumorlets** of lungs are **small**, insignificant, **hyperplastic nests**<sup>Q</sup> of neuroendocrine cells seen in areas of **scarring or chronic inflammation**<sup>Q</sup>

99. Ans. (c) **Adrenal**

(Ref: Robbins 9th/pg 717; 8th/pg 725)

**Metastasis from Lung Ca to other organs:**

- Lymphatic and hematogenous pathways<sup>Q</sup>
- Sq cell Ca shows late metastasis<sup>Q</sup>
- **Adrenals** (>50%), Liver (30%- 50%), Brain (20%) and bone (20%)

100. Ans. (b) **Adenocarcinoma** (Ref: Robbins 9th/pg 715)

101. Ans. (a) **Chromogranin** (Ref: Robbins 9th/pg 717)

**Electron microscopy of Small cell Carcinoma lung:** dense-core neurosecretory granules<sup>Q</sup> releasing neuroendocrine markers such as **chromogranin**<sup>Q</sup>, **synaptophysin**<sup>Q</sup>, and **CD57**<sup>Q</sup>, **parathormone-related protein**<sup>Q</sup>

102. Ans. (a) **PTH** (Ref: Robbins 9th/pg 715-717)

Most common variety causing **hypercalcemia** is **Squamous cell Ca**, while all other hormones are usually secreted by **small cell Ca**

103. Ans. (a) **Enterochromaffin cells** (Ref: R 9th/pg 719-720)

Carcinoid tumor develops from **Enterochromaffin cells** which contains neuroendocrine hormones.

104. Ans. (c) **Numerous long slender microvilli**

(Ref: Robbins 9th/pg 723-724; 8th/pg 733-734)

Prolonged asbestos exposure increases the risk of **mesothelioma**, which presents as a localized gray-white pleural mass.

Features	Adenocarcinoma	Malignant Mesothelioma
<b>Immunohistochemistry</b>	Carcinoembryonic antigen (CEA), CD15, Ber-EP4, MUC4, thyroid transcription factor 1 (TTF-1), Napsin A	Calretinin, WT1, keratin 5/6, claudin-4, thrombomodulin, D2-40/podoplanin, h-caldesmon, caveolin-1, vimentin
<b>Electron microscopy</b>	Short & plump microvilli	longer and more slender microvilli

About other options,

**A. Melanosomes:** Seen in **Malignant Melanoma**

**B. Neurosecretory granules:** Seen in **Small Cell Carcinoma of lung**

**D. Desmosomes with secretory endoplasmic reticulum:** can suggest Adenocarcinoma

105. Ans. (a) **Squamous cell Ca is most common carcinoma**

(Ref: Harrison 18th/chapter 89 Robbins 9th/pg 715-717)

A.	True	Most common Lung Ca in India is Squamous cell ca, while Adenocarcinoma world wide
B.	False	<b>Lambert-Eaton myasthenic syndrome</b> is a <b>paraneoplastic syndrome</b> associated with <b>Small cell Lung Ca</b>
C.	False	Small cell Ca has the poorest prognosis
D.	False	Bronchoalveolar Ca mostly involves distal airways while Small cell & Squamous cell Carcinoma involve proximal airways
E.	False	Hypercalcemia is common with Squamous cell Carcinoma

106. Ans. (a) **TTF**

(Ref: Robbins 9th/pg 715-717; 8th/pg 724-725)

Adenocarcinoma expresses thyroid transcription factor-1 (**TTF-1**); Refer to Ans 136 above;

107. Ans. (c) **Small cell Ca** (Ref: Robbins 9th/pg 715-717)

108. Ans. (a,b) **a. Not associated with smoking; b. Surgical resection alone is the treatment of choice**

(Ref: Robbins 9th/pg 715-717; 8th/pg 724-725)

**Small Cell Carcinoma (SCC)** is the most common lung Ca associated with smoking

Since, SCC is highly malignant, Surgery with chemotherapy is the treatment of choice

109. Ans. (b) **Asbestosis** (Ref: Robbins 9th/pg 723-724)

Malignant mesothelioma is associated with **Asbestosis**. **Smoking does NOT increase the risk of malignant mesothelioma.**

110. Ans. (a,c,d) **a. Clara cells; c. Mucin secreting cells; d. Type II pneumocytes** (Ref: Robbins 9th/pg 715-717)

Bronchoalveolar Ca

- Previously a subtype of Adenocarcinoma is now classified as **pre-malignant lesion of Lung Ca**
- Has **no invasion** and produces **pneumonia like consolidation**, hence called "**Lepidic**"<sup>Q</sup>
- Consists of mucin secreting **bronchiolar cell**<sup>Q</sup>, **Clara cells**<sup>Q</sup>, **Type II pneumocytes**<sup>Q</sup>

111. Ans. (c) **Carcinoid syndrome does not manifest**

(Ref: Robbins 9th/pg 719-720; 8th/pg 729)

112. Ans. (a) **Squamous cell carcinoma**

(Ref: R 9th/pg 715-717)



**113. Ans. (d) Oat cell carcinoma is commonly associated with bilateral hilar lymphadenopathy**

(Ref: R 9th/pg 715-717)

Oat cell Ca is the other name for small cell ca

a.	False, as <b>most common lung Ca is Adenoca</b>
b.	False as <b>cavitations are most common in squamous cell Ca</b>
c.	False as <b>Squamous cell ca is most often associated with calcification</b>
d.	True as <b>metastasis in small cell ca is common to hilar nodes &amp; distant sites like adrenal &amp; CNS</b>

**114. Ans. (d) Branching microvilli** (Ref: R 9th/pg 715-717)

Short Branching microvilli is the finding in electron microscopy

**115. Ans. (b) Stromal invasion with desmoplasia**

(Ref: Robbins 9th/pg 715-717; 8th/pg 724-725)



[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines, typical of notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Gastrointestinal Tract and its Disorders

## Key Points

- » Esophagus does not have serosa
- » Most common fungal organism causing Esophageal infections is Candida
- » Barret esophagus is characterized by intestinal metaplasia within the esophageal squamous mucosa
- » Most common esophageal cancer in India is Squamous cell Carcinoma
- » Most important risk factors for Adenocarcinoma is Barrett's esophagus
- » Curling ulcers-Ulcers occurring in the proximal duodenum and associated with severe burns or trauma
- » Hallmark of Crohn disease-non-caseating granulomas
- » Most common site of carcinoid tumor: Tracheobronchial tree followed by ileum, followed by rectum
- » Classic FAP at least 100 polyps are necessary for a diagnosis
- » Squamous cell carcinoma most common tumor of anal canal

## Key Recent Updates

- » Dysbiosis is seen in pseudomembranous colitis
- » Combination of strong crypt CG3 staining and loss of DAS 1 stain is seen in 45% UC. Hence, these are new markers of UC.



### The GI tract contains four layers:-

- **Mucosa** consisting of lining epithelium, **lamina propria** and muscularis mucosae
- **Submucosa**-mucous secreting glands, **Meissner's plexus**
- **Muscularis propia** (inner circular layer, outer longitudinal layer)-**Auerbach's plexus** in between these two layers.
- **Adventitia or Serosa**

## CONGENITAL ABNORMALITIES<sup>Q</sup>

### Diverticulum

#### Meckel's Diverticulum

**Most common true diverticulum** which occurs in the **ileum**. (antimesenteric side)

True diverticulum is defined by the presence of **all three layers** of the bowel wall.<sup>Q</sup>

- Due to failed involution of the **vitelline duct**<sup>Q</sup>
- **Common site** of gastric ectopia, can cause **occult bleeding**.<sup>Q</sup>

### Mnemonic

#### Meckel's diverticulum Rule of 2s<sup>Q</sup>



- Occur in **2%** population
- **Within 2 feet** of ileocecal valve
- **2 inches** long
- 2 times more common in males
- Symptomatic by **age 2**

#### Acquired Diverticulum

Most common in sigmoid colon

### Hirschsprung Disease (Congenital Aganglionic Megacolon)<sup>Q</sup>

- Normal migration of **neural crest cells from cecum to rectum is arrested prematurely**
- Distal intestinal segment lacks **both the Meissner's submucosal and the Auerbach myenteric plexus**.<sup>Q</sup>
- **Proximal to aganglionic segment, colon undergoes progressive dilation**<sup>Q</sup>
- Defect always **begins at the rectum**,<sup>Q</sup> but extends proximally for variable lengths.
- **Aganglionic region** may have a grossly **normal or contracted appearance**.<sup>Q</sup>
- In contrast, **the normally innervated proximal colon** may undergo **progressive dilation (megacolon)**<sup>Q</sup>
- Heterozygous **loss of function mutations in RET gene** causes **most of familial cases & 15% of sporadic cases**<sup>Q</sup>



Aganglionosis of the colon (Hirschsprung's Disease)

### Omphalocele

- Abdominal viscera herniating into a ventral membranous sac.

### Gastroschisis

- Herniation of **all layers**<sup>Q</sup> of the abdominal wall, from peritoneum to skin.


### Ectopia

- **Most frequent site** of ectopic gastric mucosa is the **upper third of the esophagus**


## ESOPHAGUS

### ACHALASIA CARDIA

- Occurs due to selective **loss of function of inhibitory neurons** like those secreting vasoactive intestinal peptide and nitric oxide which causes relaxation of LES whereas cholinergic innervations is intact



### Screening Test





### High Yield Facts

- Achalasia differs from Hirschsprung's disease since dilated esophagus contains less ganglion cells whereas dilated colon contains normal ganglion cells proximal to constricted aganglionic segment in Hirschsprungs.
- Inlet patch:<sup>Q</sup> Ectopic gastric mucosa is the upper third of the esophagus

### Diagnosis

- Barium swallow shows **bird beak appearance** of the esophagus
- Method of choice-Manometry<sup>Q</sup>

## LACERATIONS

Mallory-Weiss Tears	Boerhaave Syndrome
<ul style="list-style-type: none"> <li>Longitudinal <b>mucosal</b> tears</li> <li>These tears usually <b>cross the gastroesophageal junction</b></li> <li>Associated with <b>severe retching or vomiting</b><sup>Q</sup> secondary to acute alcohol intoxication.</li> <li>Generally require <b>surgical intervention</b><sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Transmural</b> tearing<sup>Q</sup></li> <li>Most common location: <b>left posterolateral part 3–5 cm above the gastroesophageal junction.</b></li> <li><b>Require surgical intervention</b></li> </ul>

## ESOPHAGITIS

- Inflammation of the esophageal mucosa is known as esophagitis
- Most common cause**<sup>Q</sup>-Reflux of gastric contents into the lower esophagus due to **transient lower esophageal sphincter relaxation**<sup>Q</sup>
- Gold standard** for the diagnosis of reflux esophagitis is **24 hours pH study**<sup>Q</sup>.




### High Yield Facts

- Esophageal infection
- Herpes viruses** typically cause **punched-out ulcers**<sup>Q</sup>
  - CMV** causes **shallower ulcerations**<sup>Q</sup> with nuclear & cytoplasmic inclusions **within capillary endothelium and stromal cells**<sup>Q</sup>
  - M.C fungal** organism causing Esophageal infections is **Candida**<sup>Q</sup> followed by mucormycosis and Aspergillus.


## BARRETT'S ESOPHAGUS

- Characterized by **intestinal metaplasia** within the esophageal squamous mucosa.<sup>Q</sup>
- Occurs due to **chronic gastroesophageal reflux disease (GERD)**.<sup>Q</sup>
- Risk of **dysplasia** correlates with **length of esophagus affected. Long segment has the higher risk**<sup>Q</sup>
- Confers an **increased risk of esophageal adenocarcinoma**.<sup>Q</sup>
- Classified as long segment (if >3 cm is involved) or short segment (if <3 cm is involved).

- Diagnosis: Endoscopy and biopsy.
- Microscopically, **lower** esophageal squamous epithelium is replaced by columnar epithelium
- Definite diagnosis is made only when columnar mucosa contains the intestinal goblet cells which show distinct mucous vacuoles that stain pale blue by hematoxylin and eosin.**<sup>Q</sup>



Barrett's Esophagus



Alcian Blue Positivity in Barretts Esophagus




### Latest Update

Dysplasia is detected in 0.2%–2% of persons with Barrett's esophagus each year

- Barrett's ulcer** is the ulcer in the columnar lined portion of Barrett's esophagus.<sup>Q</sup>

## ESOPHAGEAL TUMORS







## Adenocarcinoma

Risk factors for Adenocarcinoma	Risk is reduced by
<ul style="list-style-type: none"> <li><b>Barrett's esophagus (Most important)</b>, Tobacco exposure, Radiation, Obesity, Gastroesophageal reflux disease (GERD), <b>Scleroderma<sup>Q</sup></b>, <b>Alcohol<sup>Q</sup></b>, Medications : Long term use (&gt; 5 yr) of Theophylline &amp; Beta-agonists</li> </ul>	<ul style="list-style-type: none"> <li>Fresh fruits and vegetables, <b>Helicobacter pylori<sup>Q</sup></b> (reduces incidence of Barrett esophagus)</li> </ul>

- Most frequently in **Caucasians**, M > F, Distal 1/3 rd of esophagus<sup>Q</sup>
- By the time symptoms appear, the tumor has usually spread to **submucosal lymphatic vessels**;
- Overall 5-year survival is less than 25%.
- Microscopically, most cancers are mucin producing glandular tumors with intestinal type features




### Latest Update

Progression of Barrett's esophagus to adenocarcinoma occurs through the stepwise acquisition of genetic and epigenetic changes


- Early stages:** Mutation of TP53 and downregulation of p16/INK4a
- Late stages:** Amplification of EGFR, ERBB2, MET, cyclin D1, and cyclin E genes.

## Squamous Cell Carcinoma

- Most common site:** middle third<sup>Q</sup> of the esophagus
- More common in **African Americans<sup>Q</sup>**
- Foci of **dysplastic epithelium as well as in situ carcinoma are present adjacent<sup>Q</sup>** to the mucosa



Squamous cell carcinoma



Adeno carcinoma



### Latest Update

#### Mutations with SCC esophagus<sup>Q</sup>

- Amplification of the transcription factor gene SOX2<sup>Q</sup>
- Overexpression of the cell cycle regulator cyclin D1; and loss-of-function mutations in the tumor suppressors TP53, E-cadherin, and NOTCH1.<sup>Q</sup>

### Risk Factors for Squamous Cell Carcinoma

- Tobacco and alcohol consumption<sup>Q</sup>
- Poverty
- Caustic esophageal injury
- Chronic achalasia<sup>Q</sup>
- Tylosis et plamaris
- Plummer Vinson syndrome
- Hot beverages or food
- Radiation-tumor occurs 5 to 10 or more years<sup>Q</sup> after exposure
- Long-standing esophagitis
- Human papillomavirus (HPV) infection (in high-risk areas but not in low-risk regions)<sup>Q</sup>
- Polycyclic hydrocarbons, nitrosamines
- Nutritional deficiency of vitamin A, vitamin C, riboflavin, zinc, molybdenum
- Long-standing celiac disease<sup>Q</sup>
- Ectodermal dysplasia<sup>Q</sup> and epidermolysis bullosa



### High Yield Facts

- Most Common benign tumor of esophagus leiomyomas<sup>Q</sup>**
- Most Common esophageal cancer worldwide is squamous cell carcinoma<sup>Q</sup> (Robbins 9th ed pg 758)<sup>pg<sup>Q</sup></sup>
- Most Common esophageal cancer in India is Squamous cell Carcinoma<sup>Q</sup>**
- Most prevalent esophageal cancer worldwide (**old + new cases**) is **squamous cell carcinoma<sup>Q</sup>**

Most Common esophageal cancer worldwide (**new cases**) is **Adenocarcinoma<sup>Q</sup>**


- Most important risk factors for Adenocarcinoma is **Barrett's esophagus<sup>Q</sup>**
- Most Common site of esophageal carcinoma- middle 1/3<sup>rd</sup> of the esophagus**
- Most Common site of esophageal carcinoma in India is middle one-third of the esophagus**
- Most Common type of esophageal cancer in **upper 1/3<sup>rd</sup> of esophagus**: Squamous cell cancer<sup>Q</sup>
- Most Common type of esophageal cancer in **middle 1/3<sup>rd</sup> of esophagus**: Squamous cell cancer<sup>Q</sup>
- Most Common type of esophageal cancer in **lower 1/3<sup>rd</sup> of esophagus**: Adenocarcinoma<sup>Q</sup>
- Milk, termed mursik, which contains the carcinogen acetaldehyde**: Esophageal squamous cell carcinoma
- Plummer Vinson syndrome (also known as Patterson Kelly syndrome)**: Triad of iron-deficiency anemia, esophageal webs and glossitis
- Tylosis et plamaris**: Hyperkeratosis and pitting of palms and soles



## STOMACH

Anatomic regions of Stomach and types of cells in them


Cardia	Fundus	Body	Antrum
<ul style="list-style-type: none"> <li>Mucin-secreting<sup>Q</sup> Foveolar cells<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Chief cells<sup>Q</sup></li> <li>Parietal cells<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Chief cells<sup>Q</sup></li> <li>Parietal cells<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Mucin-secreting foveolar cells<sup>Q</sup></li> <li>Endocrine cells, such as G cells<sup>Q</sup></li> </ul>



- **Gastropathy**-When inflammatory cells are rare or absent.<sup>Q</sup>
  - 2 types:

Ménétrier disease	Zollinger-ellison syndrome
<ul style="list-style-type: none"> <li>Characterised by the hypertrophy of gastric mucosa and not by <b>exophytic growth</b>.<sup>Q</sup></li> <li>Associated with excessive secretion of transforming growth factor <math>\alpha</math> (TGF-<math>\alpha</math>) a ligand for the tyrosine kinase epidermal growth factor receptor, resulting in selective expansion of surface foveolar mucous cells and hypersecretion of mucus</li> <li>Risk of gastric <b>adenocarcinoma</b> is <b>increased</b> in adults with Ménétrier disease<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Caused by <b>gastrin-secreting tumors</b>.<sup>Q</sup></li> <li>These gastrinomas are most commonly found in the <b>small intestine or pancreas</b><sup>Q</sup></li> <li><b>Most remarkable feature is a doubling of oxyntic mucosal thickness due to a five-fold increase in the number of parietal cells.</b><sup>Q</sup></li> </ul>



## Gastropathy




## GASTRITIS

Gastritis is the **inflammation of the gastric mucosa**.


It can be divided into:







Chronic gastritis



Lymphocytes and plasma cells infiltrating gastric mucosa  
S/o chronic gastric

#### Risk factors of

##### Acute gastritis

- Heavy smoking
- Excessive Alcohol
- Excessive NSAID-aspirin  
Ibuprofen, Naproxen
- Uremia
- Ischemia and shock
- Stress (Major surgery,  
Burns, severe infections)

##### Chronic gastritis

- Drugs-NSAIDS
- **H. pylori**
- Alcohol and smoking
- Radiation
- **Gastrectomy with gastroenterostomy**
- Uremia
- Pernicious anemia

### Chronic Gastritis

- Most common cause of chronic gastritis is *H.pylori* infection and Autoimmune gastritis

#### Chronic gastritis

##### With H. Pylori (90%) [Type B]

- **Most common type**
- Not associated with intestinal metaplasia / duodenal epithelium

H.pylori causes gastritis in 2 patterns

It causes gastritis in two patterns

a. Antral predominant gastritis: Seen in individuals having lower proinflammatory cytokines, tumor necrosis factor (TNF) and interleukin-1 $\beta$  (IL-1 $\beta$ ); Associated with high acid production and increased risk of duodenal ulcer.

a. Pangastritis followed by multifocal atrophic gastritis: Seen in individuals having higher proinflammatory cytokines tumor necrosis factor (TNF) and interleukin-1 $\beta$  (IL-1 $\beta$ ) production; Associated with lower gastric acid production and increased risk of adenocarcinoma.



##### Without H. Pylori (10%) [Type A]

- **Autoimmune gastritis** Most common cause of chronic gastritis without H.Pylori
- Body-Fundic Predominant (SPARES ANTRUM)
- Associated with other autoimmune disorders e.g pernicious anemia, Hashimoto's Thyroiditis, type 1 diabetes
- Characterized by diffuse atrophic gastritis (mucosal damage of the oxyntic (acid-producing) mucosa within the body and fundus.)

### Chronic Gastritis Associated with H. pylori (in 90% Patients)

- Intraepithelial neutrophils and subepithelial plasma are characteristic of H. pylori gastritis

- Special stains **non-silver stains**<sup>o</sup> (like Giemsa, Diff-Quick, Acridine orange) and **silver stain**<sup>o</sup> (like warthin-Starry, Steiner Stain).



### H. pylori Causes

- Gastritis
- Peptic ulcer disease
- Gastric cancer - **Iron deficiency may also be a risk factor for H. pylori-associated gastric cancer.**<sup>Q</sup>
- Gastric lymphoma especially MALT lymphoma

### Virulence Factors of H. pylori

- H. pylori is a Gram-negative flagellated bacteria

- **Flagella:** Allows the bacteria to be motile in viscous mucus
- **Urease:** Produces ammonia from endogenous urea-elevates local gastric pH and enhances bacterial survival
- **Adhesins:** Like **BabA** (responsible for enhanced binding in people with blood group O)
- **Toxins:** cytotoxin-associated gene A (CagA) and VacA

### Diagnosis

- **Screening test:** Serum ELISA for antibodies against *H. pylori*
- **Urea breath test** (radiolabeled urea is broken down to radiolabeled CO<sub>2</sub> by urease enzyme which is detected, thus suggesting presence of *H. pylori* infection)
- **Gold standard:** antral biopsy showing the bacilli<sup>Q</sup>, highlighted by Warthin-Starry silver stain.
- **Most specific investigation:** Culture of bacteria (done on Skirrow's medium)<sup>Q</sup>

### Autoimmune Gastritis

- **Most common** form of chronic gastritis in patients **without H. pylori infection.**<sup>Q</sup>
- **Most common cause of diffuse atrophic gastritis**<sup>Q</sup>
- **No link to HLA alleles.**
- **Median age at diagnosis is 60.**
- Autoimmune gastritis is characterized by: Antibodies to **parietal cells** (most **prominently the H<sup>+</sup>, K<sup>+</sup>- ATPase, or proton pump,**<sup>Q</sup>) and **intrinsic factor**<sup>Q</sup> -present in 80% patients, detected in serum and gastric secretions
- **CD4<sup>+</sup> T cells** directed against parietal cell components, including the H<sup>+</sup>, K<sup>+</sup>-ATPase- **principal agents** of injury
- **Defective gastric acid secretion (achlorhydria)**<sup>Q</sup>**hypergastrinemia**
- **Hyperplasia of gastrin producing G cells** in the antral mucosa may result in **gastric carcinoid tumor**<sup>Q</sup>

### STRESS-RELATED MUCOSAL DISEASE

Stress-related mucosal disease occurs in patients with severe trauma, extensive burns, intracranial disease, major surgery, other forms of severe physiologic stress.

- Gastric lesions usually develop during the **first 3 days** of their illness
  - **Stress ulcers:** Most common in individuals with shock, sepsis, or severe trauma.
  - **Curling ulcers:** Ulcers occurring in the proximal duodenum and associated with severe burns or trauma
  - **Cushing ulcers:** Gastric, duodenal, and esophageal ulcers arising in persons with intracranial disease. These have high incidence of perforation
    - **Most commonly**<sup>Q</sup> occurs due to local ischemia
    - **Most common**<sup>Q</sup> complication- **Bleeding** followed by perforation
    - **Recover completely** with treatment





### Non-stress-related causes of gastric bleeding:

- **Dieulafoy lesion:** submucosal artery that **does not branch properly** within the wall of the stomach and thus has diameter of up to 3 mm, or 10 times the size of mucosal capillaries.
  - Found along the lesser curvature, near the gastroesophageal junction
  - Bleeding is often associated with NSAID use and may be recurrent
- **GAVE (watermelon stomach):** Longitudinal stripes of edematous erythematous mucosa alternate with less severely injured and paler mucosa. The erythematous stripes are created by **ectatic mucosal vessels**. Most common cause- idiopathic, others being cirrhosis and systemic sclerosis.<sup>Q</sup>

## COMPLICATIONS OF CHRONIC GASTRITIS

1. Peptic Ulcer Disease
2. Gastritis Cystica


## PEPTIC ULCER DISEASE (PUD)

The location of the peptic ulcer (in decreasing order of frequency) is:

- **Proximal Duodenum** : near pyloric valve and involve the **anterior duodenal wall**.<sup>Q</sup>
- Stomach (**lesser curvature** near the junction of body and antrum)
- Gastroesophageal junction in **GERD or Barrett's esophagus**
- **Ileal Meckel's diverticulum** containing ectopic gastric mucosa.

**ACTIVE peptic ulcers** are made up of following histological layers:



- **Necrotic zone**<sup>Q</sup>-Base of peptic ulcers has thin layer of fibrinoid debris<sup>Q</sup>
- **Superficial exudative zone**<sup>Q</sup> -Zone of neutrophil predominant infiltrate<sup>Q</sup>
- **Granulation tissue zone**<sup>Q</sup> -Base having active granulation tissue with mononuclear leukocytes<sup>Q</sup>
- **Zone of cicatrization**<sup>Q</sup>-Zone of fibrous or collagenous scar<sup>Q</sup>
- **Gastric ulcer can be 2 types:** benign and malignant



### High Yield Facts

- PUD does not impart an increased risk of gastric cancer<sup>Q</sup>, but patients who have had partial gastrectomies for PUD have a slightly higher risk of developing cancer in the residual gastric stump, possibly due to hypochlorhydria, bile reflux, and chronic gastritis.<sup>Q</sup>
- **Most common cause of PUD older than 60 years of age** is increased NSAID use<sup>Q</sup> (especially low-dose aspirin combined with other NSAIDs.)




Benign Gastric Ulcer	Malignant Gastric Ulcer
<ul style="list-style-type: none"> <li>• Generally at lesser curvature<sup>Q</sup></li> <li>• Smooth radiating folds<sup>Q</sup> with Hampton line &amp; collar<sup>Q</sup></li> <li>• Overhanging margins<sup>Q</sup> showing regeneration</li> <li>• Mucosal rugae projects outwards from the margins of ulcer<sup>Q</sup></li> <li>• Huge base<sup>Q</sup></li> <li>• Preserved peristalsis<sup>Q</sup></li> <li>• Heals within 8-10 weeks<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• At greater curvature<sup>Q</sup></li> <li>• Interrupted nodular, clubbed folds with Lasman Kirklin complex<sup>Q</sup> (malignant ulcer with no mass)</li> <li>• Eccentric with heaped up and everted margins<sup>Q</sup></li> <li>• Mucosal rugae stop far of the ulcer<sup>Q</sup></li> <li>• Necrotic base<sup>Q</sup></li> <li>• No peristalsis<sup>Q</sup></li> <li>• No healing<sup>Q</sup></li> </ul>
 <p>Benign gastric ulcer</p>	 <p>Malignant gastric ulcer</p>

### Gastritis Cystica

- Exuberant reactive epithelial proliferation associated with entrapment of epithelial-lined cysts
- May be found within the **submucosa (gastritis cystica polyposa)<sup>Q</sup>** or deeper layers of the gastric wall (**gastritis cystica profunda<sup>Q</sup>**)

### GASTRIC POLYPS



### High Yield Facts

- **Hyperplastic gastric polyps** are called inflammatory polyps<sup>Q</sup> & usually associated with H. pylori<sup>Q</sup>
- **Fundic gastric polyp** usually is associated with familial adenomatous polyposis (FAP)<sup>Q</sup>
- **Fundic gastric polyp** has increased due to proton pump inhibitor therapy<sup>Q</sup>. It is glandular hyperplasia due to gastrin oversecretion in response to reduced acidity<sup>Q</sup>.
- Gastric dysplasia and adenomas<sup>Q</sup> are recognizable precursor lesions associated with gastric adenocarcinoma.
- Most common site of Gastric Adenocarcinoma secondary to H. pylori infection is **Antrum<sup>Q</sup>**
- Most common site of Gastric Adenocarcinoma secondary to Pernicious anemia is **Fundus and Body<sup>Q</sup>**
- **Linitis plastica**- leather bottle appearance-seen in diffuse gastric cancer<sup>Q</sup>, it is also seen in metastasis from cancers of breast and lung<sup>Q</sup>.





## GASTRIC MALIGNANCIES


### Gastric Adenocarcinoma<sup>Q</sup>: Most Common Gastric Malignancy

#### Classification of Gastric Adenocarcinoma

Based on Depth of invasion	Based on Lauren's histological classification
<ul style="list-style-type: none"> <li>• <b>Early gastric cancer (superficial spreading type<sup>Q</sup>):</b> <ul style="list-style-type: none"> <li>▪ Involvement of <b>mucosa and the submucosa<sup>Q</sup></b> irrespective of the involvement of perigastric lymph nodes</li> <li>▪ <b>Associated with best prognosis.<sup>Q</sup></b></li> </ul> </li> <li>• <b>Late gastric cancer:</b> <ul style="list-style-type: none"> <li>▪ Involvement of the <b>muscle layer</b> of the stomach</li> <li>▪ <b>Poor prognosis</b></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• <b>Intestinal type:</b> composed of the <b>neoplastic intestinal glands<sup>Q</sup></b></li> <li>• <b>Diffuse type: <i>linitis plastica</i></b> <ul style="list-style-type: none"> <li>▪ Non-cohesive cells which do not form glands</li> <li>▪ "<b>Signet ring</b>"<sup>Q</sup> appearance (because mucin in the cell pushes the nucleus to the periphery)</li> <li>▪ <b>Worst prognosis<sup>Q</sup></b></li> <li>▪ <b>E-cadherin mutation frequently associated<sup>Q</sup></b></li> </ul> </li> </ul>



Intestinal type adeno Ca



Diffuse gastric Ca

#### Clinical Features

- The most common location of the gastric cancer is the **antrum of the stomach.<sup>Q</sup>**
- Cancer of the **gastric cardia is on the rise especially due to Barrett's esophagus<sup>Q</sup>**
- **Lesser curvature<sup>Q</sup>** is involved more often than the greater curvature.
- **Diffuse gastric cancer**-no definite lump, strong **desmoplastic reaction** that stiffens the gastric wall. These tumors show diffuse rugal flattening and a rigid, thickened wall-**leather bottle appearance termed linitis plastica<sup>Q</sup>.**

#### Investigation of Choice

Endoscopy with Biopsy of Lesion<sup>Q</sup>


#### Metastasis

- Occurs to the liver (**first organ to be affected**) followed by lungs, bone, ovary (where it is known as **Krukenberg's tumor**), periumbilical lymph nodes (**Sister Mary Joseph nodule**), peritoneal cul-de-sac (**Blumer's shelf** palpable on rectal or vaginal examination) and **left supraclavicular lymph node (Virchow's lymph node).<sup>Q</sup>**


#### Prognostic Factors

The **depth of invasion and the extent of nodal and distant metastases** at the time of diagnosis are the **most powerful prognostic indicators** in gastric cancer<sup>Q</sup>

#### Genes in Gastric Cancer







## GI Lymphomas


- Gastrointestinal tract is the most common extranodal site involved by lymphoma
- Most common lymphomas are **NHL**
- Most frequent sites in order of its occurrence are the **stomach followed by small intestine and ileocecal region**<sup>Q</sup>
- 90% of the primary gastrointestinal lymphomas are of B- cell lineage with very few T-cell lymphomas and Hodgkin's lymphoma

### Gastric Lymphoma

- Stomach is a **common site** for extranodal lymphomas
- Accounts for 5% of all gastric malignancies
- MC: Indolent extranodal marginal zone B-cell lymphomas** (mucosa-associated lymphoid tissue (MALT))
- Most common** inducer: **H. pylori**<sup>Q</sup>
- Most common presenting symptoms** are **dyspepsia and epigastric pain**<sup>Q</sup>
- Diagnostic lesions** on biopsy-**lymphoepithelial lesions**<sup>Q</sup>
- Translocations** are associated with gastric MALToma, **t(11;18)(q21;q21)**<sup>Q</sup> & **t(1;14)(p22;q32)** and **t(14;18)(q32;q21)**.

## Gastrointestinal Stromal Tumor (GIST)

- Most common mesenchymal tumor** of the abdomen with mean age of 60 yrs
- Most common site (> 50%)** is **stomach**; > **small intestine (30%)** **colon and rectum**
- Most common mutation** is **KIT (80%) > PDGFRA (8%) > SDH mutations (SDH deficient)**
- Arise from **interstitial cells of Cajal**, or pacemaker cells, of the gastrointestinal muscularis propria
- Increased incidence of GIST is seen in **NF-1**
- Most useful diagnostic marker** is **c-kit (CD117)** detectable in 95% of the patients.
- Microscopically the tumor may show either epithelioid cells, spindle cells or mixed (both the epithelioid cells and spindle cells).
- Size and mitotic rate** are predictive of behavior<sup>Q</sup>
- Those with mutations in **KIT or PDGFRA** often respond to the tyrosine kinase inhibitor **imatinib**.



GIST—Microscopy shows spindle cells

SDHB deficient (pediatric type) GIST	Usual GIST, (SDHB POSITIVE)
<ul style="list-style-type: none"> <li>Predominantly pediatric and young adult</li> <li>F:M ratio as high as 9:1</li> <li>All are gastric, most in antrum</li> <li>Frequently multiple, simultaneous or metachronous<sup>Q</sup></li> <li>Lymph node metastases common<sup>Q</sup></li> <li>Poor response to imatinib</li> <li>No CKIT or PDGFRA mutations<sup>Q</sup></li> <li>Protracted course (e.g. 15 years), even if metastatic</li> </ul>	<ul style="list-style-type: none"> <li>Predominantly older adults</li> <li>M = F</li> <li>May occur throughout gastrointestinal tract</li> <li>Usually solitary<sup>Q</sup></li> <li>Lymph node metastases rare</li> <li>Responsive to imatinib</li> <li>CKIT or PDGFRA mutations (90%)<sup>Q</sup></li> <li>Poor prognosis if metastatic</li> </ul>

**Latest Update**

SDH-deficient GISTs are located exclusively in the stomach, showing predilection for children and young adults with female preponderance.

- The tumor generally pursues an indolent course and exhibits primary resistance to imatinib therapy in most cases.

**Carney-Stratakis Syndrome**

- Germline mutations in **succinate dehydrogenase genes SDHB, SDHC or SDHD**<sup>Q</sup>
- No germline or somatic KIT or PDGFRA mutation**<sup>Q</sup>
- Familial paraganglioma and GIST**<sup>Q</sup>
- Associated with Carney's triad:** Carney's triad is gastric GIST + paraganglioma + pulmonary chondroma.

## SMALL INTESTINE AND COLON

### ANGIODYSPLASIA

- Occurs in 6th decade, in cecum/right colon
- Important cause of **major episodes of lower intestinal bleeding**<sup>Q</sup>
- Malformed **submucosal and mucosal blood vessels**,<sup>Q</sup>

### MALABSORPTION SYNDROMES

- Presents **most commonly as chronic diarrhea**<sup>Q</sup>
- Hallmark: Steatorrhea**, characterized by **excessive fecal fat and bulky, frothy, greasy, yellow or clay-colored stools**.<sup>Q</sup>

## Celiac Disease (Celiac Sprue or Gluten Sensitive Enteropathy)

- Fundamental disorder is **sensitivity to gluten**<sup>Q</sup>
- Gluten is protein component (gliadin) of **wheat**<sup>Q</sup> related grains (**Oat, barley & rye**)<sup>Q</sup>.
- Hallmark: T-cell mediated chronic inflammatory reaction**<sup>Q</sup> with an autoimmune component
- Almost all individuals share **HLA-DQ2** or **HLA-DQ8**<sup>Q</sup> haplotype.
- The **epithelial cells secrete** excess **IL-5** that **activates CD8+ T cells** & increases the **risk of lymphoma**<sup>Q</sup>



## Mnemonic

### Gluten

- B** – Barley
- R** – Rye
- O** – Oat
- W** – Wheat

## High Yield Facts

### Serology of Celiac disease

- **Latent celiac disease**-in which positive serology is not accompanied by villous atrophy
- Most sensitive tests are the measurement of IgA antibodies against tissue transglutaminase (tTG) or IgA or IgG antibodies to deamidated gliadin peptide (DGP)<sup>Q</sup>
- **Specific test** but not sensitive-IgA anti-endomysial antibodies<sup>Q</sup>


## High Yield Facts

- Most common **chronic malabsorptive disorders**: **Pancreatic insufficiency, Celiac disease, Crohn disease**<sup>Q</sup>
- Intestinal GVHD is an important cause of malabsorption and diarrhea after allogeneic hematopoietic stem cell transplantation.<sup>Q</sup>
- In **celiac disease**, **proximal intestine**<sup>Q</sup> is involved whereas in **tropical sprue**, **whole of the intestine**<sup>Q</sup> is involved

## Clinical Features

- Classic presentation includes **diarrhea, flatulence, weight loss and fatigue**<sup>Q</sup>
- **Associations**: **Dermatitis herpetiformis**<sup>Q</sup>, Auto immune diseases, Down's & Turner syndrome, IgA, nephropathy
- **Dramatic improvement** in features of malabsorption **after withdrawal of gluten**<sup>Q</sup> containing substances from diet
- On long term, increased risk of **malignancy**<sup>Q</sup> - **enteropathy-associated T-cell lymphoma**<sup>Q</sup> > **Small intestinal adenocarcinoma**


## Diagnosis



- **Morphology Biopsy**: from **second portion of the duodenum or proximal jejunum** are **diagnostic**
- Overall mucosal thickness remains same in celiac sprue
- **Diffuse enteritis**, with **marked atrophy or total loss of villi**<sup>Q</sup>
- **Hyperplastic, elongated, tortuous crypts** with **increased mitotic activity**<sup>Q</sup>
- Increased numbers of **intraepithelial CD8+ T lymphocytes**<sup>Q</sup>



Normal





Celiac disease

## Environmental Enteropathy <sup>Q</sup>

- Also called as tropical enteropathy or tropical sprue.
- No accepted criteria for diagnosis of environmental enteropathy.
- **No single infectious agent** has been linked.
- Biopsy of the intestine shows the **diffuse enteritis with atrophy of the villi**.<sup>Q</sup>

## Whipple's Disease

- **Multisystem illness** caused by Gram +ve actinomycete "**Tropheryma whippeli**"<sup>Q</sup>
- May involve any organ of the body but principally affects the **intestine, CNS and joints**.<sup>Q</sup>



### High Yield Facts

- **Hallmark** of Whipple's disease had been presence of **PAS positive macrophages** containing the characteristic bacilli. But, similar picture (PAS +ve macrophages with bacilli) can also be seen with **M. avium complex** (cause of diarrhea in AIDS)
- Acid-fast stain are helpful, since **mycobacteria stain positively while T. whippelii do not**<sup>Q</sup>.
- Patients with sickle cell disease are particularly susceptible to Salmonella osteomyelitis.<sup>Q</sup>
- The development of dementia is a relatively late symptom and is extremely poor prognostic sign<sup>Q</sup> of Whipple's disease.

### Clinical Features

- Malabsorption<sup>Q</sup>
- Multisystem involvement along with fever, lymphadenopathy and arthralgias<sup>Q</sup>
- CNS involvement (10%) dementia, seizures, coma, myoclonus<sup>Q</sup>

### Diagnosis

- **Morphologic hallmark:** Duodenal biopsy showing PAS positive diastase resistant macrophages showing characteristic granules (**lysosome stuffed with partially digested microorganisms**<sup>Q</sup>).
- Election microscopy shows **rod shaped microorganisms**<sup>Q</sup>.
- Diagnosis is **confirmed** by identification of **T. whippelii** by polymerase chain reaction (PCR).<sup>Q</sup>

### Treatment

- **Prolonged use** of **double-strength trimethoprim/sulfamethoxazole** for approximately **one year**<sup>Q</sup>.
- PAS positive macrophages can persist following successful treatment, and the **presence of bacilli outside of macrophages** is indicative of **persistent infection** or an early sign of **recurrence**<sup>Q</sup>.

## MICROSCOPIC COLITIS

### Two Types

- **Collagenous colitis:** Presence of a **dense subepithelial collagen layer**, increased numbers of intraepithelial lymphocytes, and a mixed inflammatory infiltrate within the lamina propria
- **Lymphocytic colitis:**
  - Increase in intraepithelial lymphocytes > **T lymphocyte/5 colonocyte**<sup>Q</sup>
  - **Strong association with celiac disease and autoimmune diseases**<sup>Q</sup>.

### Latest Update

- **Autoimmune enteropathy**<sup>Q</sup> is an X-linked disorder characterized by severe persistent diarrhea and autoimmune disease that is caused by mutation in **FOXP3 gene**,<sup>Q</sup> resulting in defective function of regulatory T cells.


## INFECTIOUS DISEASES

The important causes of infections in the intestine are as follows:

### Enteric Fever (Typhoid)


- It is caused by infection with **Salmonella enterica** & its two subtypes, typhi (most common in endemic countries) and paratyphi (most common travellers)
- Clinically-**step-ladder pyrexia**, rose spots (erythematous macular lesions on chest and abdomen), abdominal pain, vomiting
- **Characteristic: Hypertrophy of Peyer patches in the terminal ileum and their ulceration, presence of longitudinal ulcers (oval ulcers with long axis along the long axis of the ileum)**<sup>Q</sup>.
- Microscopic-macrophages having bacteria and red blood cells (**erythrophagocytosis**)
- **Gallbladder colonization** with **S. typhi** or **S. paratyphi** is associated with gallstones and the **chronic carrier state**.
- Liver, Bone marrow and lymph node shows small, randomly scattered foci of parenchymal necrosis in which hepatocytes are replaced by macrophage aggregates, called **typhoid nodules**<sup>Q</sup>;
- Complications include hemorrhage and perforation.
- Extraintestinal complications - encephalopathy, meningitis, seizures, endocarditis, myocarditis, pneumonia & cholecystitis
- **Blood culture is the diagnosis of choice**<sup>Q</sup>
- **Widal test** is used for measuring the antibody titer





## Mnemonic

Tie Typhoid → (Tie) → Longitudinal ulcers  
TB – Transverse ulcers  
Amoebiasis – flask shaped ulcers



## Intestinal Tuberculosis (ITB)

It can present itself in two of the following forms:

- Primary infection
- Secondary following reactivation, usually from a primary pulmonary focus



Distribution of tuberculous lesions

Ileum > cecum > ascending colon > jejunum > appendix > sigmoid > rectum > duodenum > stomach > esophagus

## Routes of Infection

- Primary:** Unpasteurized milk and milk product are regarded as the main route of transmission of zoonotic. TB caused by *Mycobacterium bovis* in countries where there are no effective eradication programs.
- Secondary:** Ingestion, sputum from an active focus in the lung, Hematogenous, Direct spread from adjacent organs



## Pathology

- The **ileocecal region** is the **most common site**<sup>o</sup> of involvement
- ITB usually has one of three forms: ulcerative, hypertrophic or ulcerohypertrophic or fibrous
- Tuberculous granulomas initially form in the **mucosa or Peyer's patches**<sup>o</sup>
- Transverse Ulcers**<sup>o</sup> are seen, they are relatively superficial, with a different appearance from those in Crohn's disease
- ITB progresses slowly and presents late with complications, acute or subacute obstruction due to mass (tuberculoma), stricture formation in the ileocecal region or perforation leading to peritonitis
- The mesenteric lymph nodes are enlarged; matted and caseous. **Tabes mesenterica**.<sup>o</sup>
- Diagnosis – widening of the ileocecal angel (known as “**pulled up cecum**”<sup>o</sup> on barium radiography,
  - Histological evidence of caseating epithelioid cell granulomas along with langhan giant cells
  - Confirmation of acid fast bacilli on **Ziehl-Neelsen** and **culture/PCR positivity**<sup>o</sup>

## Amoebiasis

- It is caused by infection with an anaerobic protozoa *E. histolytica*<sup>o</sup>
- Flask-shaped ulcers**<sup>o</sup> (ulcer with a broad base but narrow neck).
- Cecum and ascending colon**<sup>o</sup>
- The ulcers usually involve the mucosa and the submucosa (not the muscle layer)
- Microscopy shows presence of **trophozoites**<sup>o</sup> in the ulcer
- Liver** shows hepatic abscess (called as “**anchovy sauce pus**”).<sup>o</sup>





## INFLAMMATORY BOWEL DISEASE (IBD)

Inflammatory bowel disease (IBD) is a chronic condition resulting from **inappropriate mucosal immune** activation.

It is primarily of two types: Crohn's disease and ulcerative colitis.

### Crohn's Disease

- Sharply delineated and transmural involvement of bowel by inflammatory process
  - Any portion of intestine can be involved (most common site: ileum)
  - Associated with HLA-DR1/DQw5 and an abnormal T-cell response **TH1 cells**.

**Hallmark of crohn's disease- non-caseating granulomas.**



### High Yield Facts

- Earliest change in **Crohn's disease** is **Apathoid ulceration**<sup>Q</sup>.
- Earliest change in **Ulcerative colitis** is **Blurring of mucosal stripe and granular appearance**<sup>Q</sup>.

#### Metastatic Crohn's disease

- A misnomer since there is no cancer
- Described when cutaneous granulomas form nodules

## ULCERATIVE COLITIS


- Ulcerative colitis extends only into the mucosa and submucosa.
- Mural thickening is not present, the serosal surface is normal, and strictures do not occur**
- Associated with **HLA-DR2, polymorphism in IL-10 gene** and an **abnormal T-cell response particularly of Th2 cells**<sup>Q</sup>
- Always involves the rectum** and extends proximally in a **continuous fashion** to involve part or all of the colon. Disease of the entire colon is termed **pancolitis**<sup>Q</sup>
- In severe cases of pancolitis, mild mucosal inflammation of the distal ileum is termed as backwash Ileitis**<sup>Q</sup>

### Mnemonic

#### Extraintestinal manifestations of UC-SEAS




- S**kin manifestations: erythema nodosum, pyoderma gangrenosum
- E**ye inflammation: iritis, episcleritis
- A**rthritis
- S**clerosing cholangitis


- Approximately **2.5% to 7.5%** of individuals with ulcerative colitis also have **primary sclerosing cholangitis**<sup>Q</sup>







Feature	Crohn's Disease	Ulcerative Colitis
<b>A. Macroscopic features</b>		
• Distribution	Segmental with <b>skip areas</b> <sup>q</sup>	<b>Continuous</b> without skip areas <sup>q</sup>
• Location	Commonly <b>terminal ileum (most common) and/or ascending colon</b>	Commonly <b>rectum sigmoid colon and extending upwards</b>
• Extent	Usually involves the <b>entire thickness of the affected segment of bowel wall</b>	Usually superficial, <b>confined to mucosal layers</b>
• Ulcers	<b>Serpiginous ulcers</b> , that develop into deep <b>fissures</b> <sup>q</sup>	Superficial mucosal ulcers without fissures
• Pseudopolyps	Rarely seen	<b>Commonly present</b> <sup>q</sup>
• Fibrosis	<b>Common</b>	Rare
• Shortening	Due to fibrosis	Due to contraction of muscularis
<b>B. Microscopic features</b>		
• Depth of inflammation	Typically <b>transmural</b> <sup>q</sup>	<b>Mucosal<sup>q</sup> and Submucosal</b>
• Type of inflammation	<b>Non-caseating granulomas</b> <sup>q</sup> and infiltrate of mononuclear cells (lymphocytes, plasma cells and macrophage)	Crypt abscess and non-specific acute and chronic inflammatory cells (lymphocytes, plasma cells neutrophils, eosinophils, mast cells)
• Mucosa	Patchy ulceration	Hemorrhagic mucosa with ulceration
• Submucosa	<b>Widened</b> due to edema and lymphoid aggregates	Normal or reduced in width
• Muscularis	Infiltrated by inflammatory cells	Usually spared, except in cases of <b>Toxic Megacolon</b> <sup>q</sup>
• Fibrosis	Present	Usually absent
<b>C. Complications</b>		
• Fistula formation	Internal and external fistulae in 10% case	Extremely <b>rare</b> <sup>q</sup>
• Malignant changes	Less common but present	May occur in disease of more than 10 years duration ( <b>more common</b> <sup>q</sup> )
• Fibrous strictures	<b>Common</b> <sup>q</sup>	Never <sup>q</sup>
• Toxic megacolon	–	<b>Risk present</b> <sup>q</sup>
• Features	<ul style="list-style-type: none"> <li>Hose pipe appearance<sup>q</sup></li> <li>Cobble-stone appearance<sup>q</sup></li> <li>Halo sign on CT<sup>q</sup></li> <li>String sign of cantor<sup>q</sup></li> <li>Raspberry/rosethorn appearance<sup>q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Garden hose appearance<sup>q</sup></li> <li><b>Pseudopolyps</b><sup>q</sup></li> <li><b>Pipestem colon (Ahastral)</b><sup>q</sup></li> </ul>
<div>  <p>Histological findings are suggestive of non-caseating granuloma. It is a hallmark of Crohn's disease</p> </div> <div>  <p>Cobblestone appearance Crohn's disease</p> </div> <div>  <p>Crypt abscess (ulcerative colitis) &gt; Crohns disease</p> </div>		



### High Yield Facts

- Extraintestinal manifestations of Crohn's disease include uveitis, migratory polyarthritis, sacroiliitis, ankylosing spondylitis, erythema nodosum, and clubbing of the fingertips, any of which may develop before intestinal disease is recognized.<sup>a</sup>
- Approximately 2.5–7.5% of individuals with ulcerative colitis also have primary sclerosis cholangitis.
- Polymorphism of the IL-23 receptor is **protective** in both the types of inflammatory bowel disease.
- Autosomal recessive mutations of the **IL-10 and IL-10 receptor genes** are linked to severe, early onset IBD
- Smoking is a strong exogenous risk factor for development of CD whereas smoking partly relieves symptoms in UC.<sup>a</sup>
- Anti-flagellin antibodies are common in Crohn's disease and uncommon in ulcerative colitis patients.<sup>a</sup>
- Gene associated with Crohn's disease NOD2 > ATG16L1 and IRGM <sup>RG</sup>



### Latest Update

#### Colitis-Associated Neoplasia

Depends on following parameters:

- Duration of the disease:** Risk increases 8 to 10 years after disease onset.
- Extent of disease: Pancolitis > Left side
- Nature of inflammation - ↑ neutrophils more risk
- Duration of the disease:** Risk increases 8 to 10 years after disease onset.




### High Yield Facts

Antibody	GI Disorder
<i>Antiendomysial antibody</i> <sup>a</sup>	<i>Celiac sprue</i>
<i>Antisaccharomyces cerevisiae antibody</i> <sup>a</sup>	<i>Crohn's disease</i>
<i>p-Antineutrophil cytoplasmic antibody</i> <sup>a</sup>	<i>Ulcerative colitis</i>







## COLORECTAL POLYPS

Polyps are most common in the colorectal region but may occur in the esophagus, stomach, or small intestine.





### Non-Neoplastic Polyp

Features	Hamartomatous Polyps	Hyperplastic Polyps	Inflammatory Polyps
<b>Cause</b>	Caused by <b>germline mutations in tumor suppressor genes or proto-oncogenes</b> . <sup>a</sup>	<p>↓ <b>Epithelial turnover</b>      Delayed surface epithelial shedding</p> <p><b>Piling of goblet cells &amp; absorptive cells</b></p>	Result of <b>chronic cycles of injury and healing</b> . <sup>a</sup>
<b>Clinical Presentation</b>	Types: <ul style="list-style-type: none"> <li>Juvenile polyp</li> <li>Peutz Jegher's syndrome (Discussed below)</li> </ul>	<ul style="list-style-type: none"> <li>Most commonly found in the <b>left colon</b></li> <li>Sixth and seventh decades of life</li> <li>Typically less than 5 mm in diameter</li> </ul>	<p>Clinical triad of rectal bleeding, mucus discharge, and an inflammatory lesion of the anterior rectal wall.</p> <p><b>Solitary rectal ulcer syndrome is an inflammatory polyp</b><sup>a</sup></p>
<b>Hallmark</b>	discussed below separately	<p><b>Hallmark: Serrated surface architecture. Restricted to the upper third, or less, of the crypt.</b><sup>a</sup></p> <p>Piling goblet cells</p>  <p>Hyperplastic polyp. (60-70years) Due to ↓ epithelial turnover</p>  <p>Serrated surface in upper third of crypt</p> <p>Lumen is not round</p> 	<p><b>Hallmark: Mixed inflammatory infiltrates, erosion.</b></p>  <p><b>10X- surface mucosal ulceration</b></p>
<b>Malignant potential</b>	Associated with increased cancer risk, <b>either within the polyps or at other intestinal or extra-intestinal sites</b> <sup>a</sup>	<b>Non-neoplastic</b> <sup>a</sup>	<b>Non-neoplastic</b> <sup>a</sup>





## Types of Hamartomatous Polyps

Features	Juvenile Polyps	Peutz Jegher's Syndrome
		
Age	Children younger than 5 years of age	Median age of 11 years
Site	Rectum	Jejunum
Symptoms	<b>Rectal bleeding<sup>Q</sup></b> Most juvenile polyps are less than 3 cm in diameter	They come to attention for following <ul style="list-style-type: none"> <li>Intussusception, Mucocutaneous pigmentation, Secondary cancers</li> </ul>
Morphology	<ul style="list-style-type: none"> <li>Sporadic single, called <b>retention polyps</b>.</li> <li>Juvenile polyposis syndrome (JPS): autosomal dominant syndrome, 3 to as many as 100 polyps</li> </ul>	<ul style="list-style-type: none"> <li>Multiple GI hamartomatous, <b>polyps and mucocutaneous hyperpigmentation</b> (especially lips, buccal mucosa,<sup>Q</sup></li> <li><b>Hallmark histology: The arborization and presence of smooth muscle intermixed with lamina propria.</b></li> </ul>
Mutation	<b>SMAD4<sup>Q</sup></b>	Heterozygous loss-of-function of <b>STK11<sup>Q</sup></b>
Malignant potential	<ul style="list-style-type: none"> <li><b>Dysplasia is extremely rare in sporadic juvenile polyps<sup>Q</sup></b></li> <li>Juvenile polyposis syndrome is associated with dysplasia and 30 to 50% develop colonic adenocarcinoma by age 45.</li> </ul>	<ul style="list-style-type: none"> <li>Markedly increased risk of several malignancies</li> <li>Surveillance is recommended <ul style="list-style-type: none"> <li><b>At birth, for sex cord tumors of the testes;</b></li> <li><b>Late childhood for gastric and small intestinal cancers;<sup>Q</sup></b></li> <li><b>Second and third decades of life for colon, pancreatic, breast, lung, ovarian, and uterine cancers.<sup>Q</sup></b></li> </ul> </li> </ul>

## Other Syndromes with Hamartomatous Polyps<sup>Q</sup>


- **Cowden Syndrome and Bannayan-Ruvalcaba-Riley Syndrome:** Autosomal dominant hamartomatous polyp syndromes associated with loss-of-function mutations in **PTEN** hence also known as **PTEN hamartoma syndrome**.
- **Cronkhite-Canada Syndrome:** Nonhereditary and develops in individuals over 50 years of age

## Neoplastic Polyps

- Any tumor mass lesion in the GI tract like this includes adenocarcinomas, neuroendocrine (carcinoid) tumors, stromal tumors, lymphomas, and even metastatic cancers from distant sites) can produce a mucosal protrusion or polyp.

- The most common neoplastic polyps are colonic adenomas. Adenomas are intraepithelial neoplasms like small pedunculated, polyps to large sessile lesions
- Colorectal adenomas are characterized by the presence of epithelial dysplasia
- Colonic adenomas are classified as tubular, tubulovillous, or villous based on their architecture
- Tubular adenomas- **most common<sup>Q</sup>** adenoma
- Villous adenomas contain foci of invasion more frequently
- **The size of the adenoma is the most important characteristic which correlates with the risk of malignancy.<sup>Q</sup>**





### Sessile Serrated Adenomas

- **Lack cytologic dysplasia<sup>o</sup>** and share morphologic features with hyperplastic polyps, more at right colon.
- Differs from **hyperplastic polyps** by serrated architecture throughout the **full length<sup>o</sup>** of the glands
- **Intramucosal carcinoma:** When dysplastic epithelial cells breach the basement membrane to invade the lamina propria or muscularis mucosae.
- **Intramucosal carcinomas** have little or no metastatic potential as functional lymphatic channels are absent in the colonic mucosa.

### FAMILIAL ADENOMATOUS POLYPOSIS (FAP)


- **Inheritance:**
  - AD
- **Cause:**
  - Adenomatous polyposis coli (APC; 5q21) mutation
- **Predominant Site(s):**
  - None

- **Syndromes associated:**
  - Gardner syndrome
  - Turcot syndrome
- **Extraintestinal manifestations**
  - Congenital hypertrophy of retinal pigment epithelium<sup>o</sup>
- **MYH-associated polyposis**
  - Autosomal recessive
  - Some polyposis patients without APC loss have mutations of the base-excision repair gene MYH (also referred to as MUTYH). In these cases, polys are similar to attenuated FAP, with polyp development at later ages, the presence of fewer than 100 adenomas, and the delayed appearance of colon cancer, often at ages of 50 or older.
- **Lynch syndrome OR Hereditary Non-Polyposis Colorectal Cancer**
  - **Inheritance:** AD
  - **Cause:** Mutations in **DNA repair genes**. MSH2, MLH1 leading to **microsatellite instability**
  - Predominant site: Right colon



### High Yield Facts

- Most common extra intestinal manifestation of juvenile polyposis is pulmonary A-V malformation
- **Classic FAP:** At least **100 polyps** are necessary for a diagnosis<sup>Q</sup>
- **Attenuated FAP:** Lower number of adenomatous polyps (around 30)
- **Gardner syndrome:** osteomas of mandible, skull, and long bones, epidermal cysts, desmoid tumors, thyroid tumors, and dental abnormalities, including **unerupted and supernumerary teeth**
- **Turcot syndrome:** Intestinal adenomas and tumors of the central nervous system (**Most common-Medulloblastoma or Glioblastoma**)
- **Squamous cell carcinoma-** most common tumor of anal canal, most common HPV



Adenomatous Polyposis

## COLORECTAL MALIGNANCY

### Adenocarcinoma

- **Most common malignancy of the GI tract<sup>Q</sup>**
- Colorectal cancer incidence peaks at 60 to 70 years of age.
- The **risk factors** for the colon cancer are:
- **Etiology (MNEMONIC: CRAPS)**
  - Chronic ulcerative colitis
  - Ratio of animal fat: fibre diet
  - Adenomatous polyps
  - Familial Polyposis
  - Strong family history of colon cancer.



### High Yield Facts


- **Intramucosal carcinoma-** when dysplastic epithelial cells breach the basement membrane to invade the **lamina propria or muscularis mucosae**.<sup>Q</sup>
- Have **no metastatic potential<sup>Q</sup>** as **functional lymphatic channels are absent** in the colonic mucosa
- NSAIDS ↓ risk of colon cancer c pharmacologic chemoprevention

### Genetic Mutations in Colon Carcinoma


- **APC/β-catenin pathway (mc;80%)** followed by late event mutations like activating mutations in KRAS then SMAD2 and SMAD4 and later in 70% to 80% of colon cancers. Telomerase re-activation can further progress the tumor
- **Microsatellite instability pathway:** Occurs due to DNA mismatch repair deficiency. Mutations accumulate in microsatellite repeats, referred to as microsatellite instability (MSI). Tumors with this pathway - MSI high, or MSI-H, tumors. Common genes where this occurs are : type II TGF-β receptor. pro-apoptotic protein BAX<sup>Q</sup>
- **Microsatellite instability** without mutations in DNA mismatch repair enzymes<sup>Q</sup> with CpG island hypermethylation phenotype (CIMP). Often show BRAF gene mutation
- **Increased CpG island methylation (CIMP)** in the absence of microsatellite instability. Often they have K-RAS mutation
- **No CpG island methylation (CIMP)** show p53 mutation

### Prognosis

Depth of invasion and the presence or absence of lymph node metastases<sup>Q</sup>



Colon Carcinoma (Gross)



Colon adenocarcinoma





## TUMORS OF THE APPENDIX

- Most common tumor of the appendix is the **well differentiated neuroendocrine (carcinoid) tumor**.<sup>Q</sup>
- Forms a solid bulbous swelling at the **distal tip**<sup>Q</sup> of the appendix.
- Nodal metastases are very infrequent<sup>Q</sup>, and distant spread is exceptionally rare.

## UPDATE ON CARCINOID TUMOR


### Carcinoid Tumor

- Arises from the **endocrine cells called as argentaffin tissue** (also called as Kulchitzky cells of crypts of Lieberkuhn) now properly referred to as **well-differentiated neuroendocrine tumors**<sup>Q</sup>
- The carcinoid tumors can be of the following types:
  - Foregut carcinoid tumors:** Arise from the esophagus, stomach and the duodenum proximal to the ligament of Trietz, these are usually **benign**.
  - Midgut carcinoid tumors:** Arise from the jejunum and ileum; these are **aggressive and metastasize frequently**.<sup>Q</sup>
  - Hindgut carcinoid tumors:** Arise from the **appendix, colon and rectum; usually benign**.<sup>Q</sup>


- Grossly, carcinoids are **intramural or submucosal** masses that create small polypoid lesions, which on cut section gives yellow tan color.
- On electron microscopy, the tumor cells show **dense core granules** in the cytoplasm
- Immunohistochemistry-these granules stain positively with **chromogranin A, neuron-specific enolase and synaptophysin**

### Carcinoid Syndrome (5% Cases of Carcinoid Tumor)

- Manifest when vasoactive substances from the tumor enters systemic circulation escaping hepatic degradation.
- It is strongly associated with **metastatic disease**.
- Clinical features** are **Intestinal hypermotility** (Vomiting, diarrhea), **Vasomotor symptoms** like flushing and cyanosis of the skin, **Systemic fibrosis** (Affect cardiac valves, endocardium, retroperitoneal and pelvic fibrosis)
- Cardiac lesions are present in **50% of the patients** with the carcinoid syndrome. Most common is **tricuspid regurgitation (not tricuspid stenosis) followed by pulmonary stenosis**. (TIPS- Tricuspid insufficiency and pulmonary stenosis)<sup>Q</sup>



Carcinoid



Dense core granules

### Diagnosis

- Serum chromogranin A** levels ( $\uparrow$  in 56–100% of carcinoid tumors) and the level correlates with tumor bulk
- The levels of 5-HT and its metabolite 5-hydroxyindoleacetic acid (5-HIAA) is elevated in the urine. Measurement of **5HIAA is most frequently used**.



## High Yield Facts


- Most common site of carcinoid tumor: Tracheobronchial tree followed by ileum followed by rectum<sup>Q</sup>
- Most common site of carcinoid tumor in foregut<sup>Q</sup>-Bronchus, lung, trachea<sup>Q</sup> (27.9%) > Stomach (4.6 %) > Duodenum (2.0%)
- Most common site of carcinoid tumor in midgut<sup>Q</sup> - Appendix (4.8%) > Ileum<sup>Q</sup> (14.9%) > Colon (8.6%)
- Most common site of carcinoid tumor in hindgut<sup>Q</sup> – rectum<sup>Q</sup> (13.6%)
- At Presentation, most common symptom of carcinoid syndrome is diarrhea and flushing and during the course of disease most common symptom is flushing<sup>Q</sup>
- The cardiac changes are largely **right sided** due to inactivation of both serotonin and bradykinin in the blood during passage through the lungs by the **monoamine oxidase present in the pulmonary vascular endothelium**.<sup>Q</sup>

R10<sup>th</sup>


## Latest Update

### Pseudomembranous Colitis Antibiotic-associated colitis or antibiotic-associated diarrhea

- Most commonly caused by **Clostridium difficile**<sup>Q</sup>
- **Hallmark: Adherent layer of inflammatory cells and necrotic debris at sites of colonic mucosal injury (pseudomembrane)**<sup>Q</sup>
- Occurs due to disruption of the normal colonic microbiota due to antibiotics, which allows **C. difficile** overgrowth to produce large amounts of **two toxins, toxin A and toxin B**<sup>Q</sup>.
- Toxins released by **C. difficile** cause the **ribosylation of small GTPases**
- **Diagnosis: detection of C. difficile toxin (Not Culture)**<sup>Q</sup> and is supported by the characteristic histopathology



Pseudomembranous enterocolitis



Mucopurulent exudate erupting from crypts to form mushroom like cloud on surface

R10<sup>th</sup>

## Latest Update

### Microbiome and Dysbiosis

- The microbiome is the diverse microbial population of bacteria, fungi, and viruses found in or on the human body (e.g., in the intestinal tract, skin, upper airway, and vagina)
- Dysbiosis refers to changes in composition of the microbiome that are associated with disease. These changes may result from therapies or various pathophysiologic conditions.
- Use of some antibiotics is an important risk factor for intestinal infections caused by toxin-producing *Clostridium difficile*. These antibiotics kill or inhibit normal commensal bacteria, allowing overgrowth of *C. difficile*.
- Restoration of the microbiome by duodenal infusion of stool containing commensal flora from healthy donors successfully treats *C. difficile* infection in many individuals who have relapsed after antibiotic therapy






## NEXT Pattern Questions



Q's

1. A 45/m presented with abdominal pain. Endoscopy shows a mass in stomach following which a biopsy was taken as shown below. Which of the following statement is not true about the case?



- GISTs without mutated KIT or PDGFRA have mutations in succinate dehydrogenase complex
- Mutation of KIT or PDGFRA is an early event in sporadic GISTs
- Prognosis correlates with tumor size, mitotic index, and location
- Patients having mutations in KIT are Imatinib resistant


Ans. (d) **Patients having mutations in KIT are Imatinib resistant**

- The given image shows the histopath of spindle cells and with the biopsy taken from stomach, its GIST. Now, the cases of GIST have mutation in KIT, PDGFR- $\alpha$ . Otherwise, they have SDH loss mutation. Kit mutation is important as these cases are sensitive to Imatinib.



Q's

2. A 50-year-old male presented with recurrent bloody diarrhea. Colonoscopy was done followed by histopathology as shown below. What is your diagnosis?



- Pseudomembranous colitis
- Non-Hodgkin lymphoma colon
- Ulcerative colitis
- Crohn's disease


Ans. (d) **Crohn's disease**

- Non caseating granuloma as seen in the histopathology is virtually the hallmark of Crohn's disease.



Q's

3. A 30-year-old presented with malabsorption with diarrhea, weight loss, abdominal pain, Occasionally polyarthritis, CNS complaints. Duodenal biopsy shows the following. Which is the correct statement?



- Infiltration of histiocytes with RBCs in the lamina propria
- Granuloma in the lamina, can be intestinal TB
- Macrophages with PAS (+) ve material inside the lamina propria representing Whipple's disease
- Eosinophils in the lamina propria


Ans. (c) **Macrophages with PAS (+) ve material inside the lamina propria representing Whipple's disease**

- Malabsorption with diarrhea, weight loss, abdominal pain and occasionally polyarthritis, CNS complaints are seen in Whipple's disease. Remember, these cases show PAS (+) ve material inside macrophages in the lamina propria layer.



Q's


4. All of the following statements are true regarding colonic polyposis shown below except ?



- FAP without mutation in APC gene have another target of MYH gene
- Untreated case develop malignancy
- Prophylactic colectomy prevent the risk of development of cancer in other organ as well
- Gardner syndrome encompasses the subset of FAP patients with brain tumor medulloblastoma

Ans. (d) **Gardner syndrome encompasses the subset of FAP patients with brain tumor medulloblastoma**

- The colonic polyposis syndrome in FAP is Turcot syndrome which has FAP with brain tumor and not Gardner syndrome.



Q's

5. Intestinal biopsy from a patient A 25-year-old male presenting with symptoms of fatigueness, chronic diarrhea, bloating, or chronic fatigue. What is not true about the given condition?



- Characteristic itchy, blistering skin lesion, is seen in 10% cases
- Most sensitive tests are the measurement of IgA antibodies against tissue transglutaminase
- Most common celiac disease-associated cancer is adenocarcinoma colon
- Hla-DQ 2 and HLA DQ-8/ B8 is useful for its high negative predictive value


Ans. (c) **Most common celiac disease-associated cancer is adenocarcinoma colon**

- The symptoms of fatigueness, chronic diarrhea, bloating, or chronic fatigue are suggestive of malabsorption. Histopath shows loss of villi and so is Celiac disease. Remember, Celiac disease causes T cell lymphoma and not adenoma as the most common tumor.




## Image-Based Questions

1. A 45-year-old male complained of dysphagia. On investigation, he was HIV positive. He underwent endoscopy and biopsy. Endoscopy findings and histological findings are suggestive of?




- Herpes
- CMV
- Candida
- Pseudomonas

2. A 50-year-old male with Gastroesophageal reflux disease—Diagnosis?



- Squamous metaplasia
- Columnar metaplasia
- Dysplasia
- Malignancy






3. Old male with H/O long intake of antibiotics. Diagnosis:




- a. Pseudomembranous Colitis
- b. Chrons
- c. Ulcerative colitis
- d. Amoebic colitis

4. Histological findings are suggestive of non-caseating granuloma. It's a hallmark of:



- a. Crohn disease
- b. Ulcerative colitis
- c. Salmonella
- d. Amoebiasis

5. Gross findings are suggestive of:



- a. Pseudopolyps
- b. Pseudopipe
- c. Cobblestone
- d. Ulcers

6. A 20-year-old male with osteomas of the skull.




- a. Adenomatous Polyposis
- b. Crohn's disease
- c. Ulcerative colitis
- d. None




## Answers of Image-Based Questions

1. Ans. (b) **CMV esophagitis**
  - Endoscopy showed shallow ulcers at lower end of esophagus
  - Histopathology shows large eosinophilic intranuclear inclusion and multiple small cytoplasmic inclusions
2. Ans. (b) **Columnar metaplasia Histological findings of barrett esophagus**
  - Metaplastic columnar epithelium (specialized) with goblet cells is seen adjacent to squamous epithelium of esophagus hence suggestive of columnar metaplasia
  - Long-segment: Barrett's mucosa extends 3 cm or more. Short-segment: Barrett's mucosa extends less than 3 cm
3. Ans. (a) **Pseudomembranous enterocolitis**
  - The mucosal surface of the colon seen here is hyperemic and is partially covered by a yellow-green exudate.
4. Ans. (a) **Crohn's disease**
  - Non caseating epithelioid cell granuloma. Look at the presence of giant cells and lot of lymphocytes
5. Ans. (a) **Pseudopolyps suggestive of ulcerative colitis**
  - Pseudopolyps can be seen clearly as raised red islands of inflamed mucosa.
6. Ans. (a) **Adenomatous Polyposis**
  - Gross picture shows multiple polyposis with numerous small polyps covering the colonic mucosa.
  - A 20-year-old male should be suspected as having faulty APC gene as he is having multiple polyps and also having osteomas of skull. Other extra colonic manifestations should be looked for.






## Multiple Choice Questions


1. A 40-year-old immunocompromized patient presented with complaints of dysphagia. UGI scopy showed multiple ulcers in the distal esophagus. Biopsy from the esophagus showed the following. What is the diagnosis?  
(Recent exam 2018)




- a. Candida  
b. Cytomegalovirus  
c. Herpes  
d. Eosinophilic esophagitis
2. On endoscopy and barium swallow the following findings are seen, which of the following will be seen on histopathology?  
(AIIMS Nov 2017)



3. A 40-year-old patient presented with heart burn and increased salivation. UGE scopy was done and biopsy was taken and is as shown below. Which of the is the diagnosis?  
(AIIMS May 2017)



- a. Barret esophagus  
b. Adenocarcinoma  
c. Esophagitis  
d. Squamous cell carcinoma
4. Prognostic factors for carcinoma esophagus is/are:  
(PGI May 16)
- a. Depth of invasion  
b. Lymph node status  
c. Tumour grading  
d. Stage of the disease
5. Layer absent in esophagus (Recent Question 2016)
- a. Mucosa  
b. Serosa  
c. Muscularis  
d. lamina propria
6. A 30-year-old software engineer came to OPD with chief complains of heartburn. On endoscopic biopsy, the lesion shows the following (figure below). Identify the lesion, stain has been done for what and what additional features should be looked for?  
(AIIMS Nov 2015)



- a. Adenocarcinoma; PAS; malignancy  
b. Barretts oesophagus; mucin stain; dysplasia  
c. Squamous cell carcinoma; cytokeratin, squamous pearls  
d. Infection; fungal stain; inclusion body
7. Most frequent site of ectopic gastric mucosa is:  
(Recent Question 2015)
- a. Upper third of esophagus  
b. Middle third of esophagus  
c. Lower third of esophagus  
d. Duodenum
8. Achalasia cardia: (Recent Question 2015)
- a. Absence of nerves  
b. Absence of muscles  
c. Hypertrophy of nerves  
d. None





9. **Predisposing factors for Esophagus Ca:** (PGI JUNE 13)  
 a. Tylosis  
 b. Achalasia  
 c. Barrett's esophagus  
 d. Scleroderma  
 e. Plummer-Vinson syndrome
10. **M.C. site of Ca esophagus is:** (Recent Question 2013)  
 a. Middle 1/3  
 b. Upper 1/3  
 c. Lower 1/3  
 d. Lower end of esophagus

### STOMACH

11. **A 60-year-old person presented with some stomach tumor with following features: mesenchymal solitary mass below mucosa of stomach with intact mucous, Spindle shaped cells and epithelioid cell on biopsy. This tumor is positive for:** (PGI May 2019)  
 a. DOG 1  
 b. CD117  
 c. CD 34  
 d. KIT  
 e. CD 99
12. **What is true about Succinate dehydrogenase deficient GIST?** (AIIMS May 2017)  
 a. Negative for C-KIT & CD117  
 b. YOUNG age  
 c. MC site stomach  
 d. Sensitive to Imatinib  
 e. Aggressive clinical course
13. **Tumor most commonly associated with H pylori:** (Recent Question 2016)  
 a. MALTOMA  
 b. Adenocarcinoma  
 c. Squamous cell carcinoma  
 d. None
14. **Krukenberg tumour of ovary is due to carcinoma of** (MH PG 2014)  
 a. Stomach  
 b. Lung  
 c. Central nervous system  
 d. Thyroid
15. **Chronic gastritis is caused by all except:** (Recent Question 2015)  
 a. H. Pylori  
 b. Pernicious anaemia  
 c. Gastrectomy with gastroenterostomy  
 d. Overuse of salicylates
16. **Which of the following is the most outermost histological layer of peptic ulcer** (Recent Question 2015)  
 a. Necrotic zone  
 b. Superficial exudative zone  
 c. Granulation tissue zone  
 d. Zone of cicatrization
17. **Most common type of gastric polyp is:** (Recent Question 2015)  
 a. Hyperplastic polyp  
 b. Hamartomatous polyp  
 c. Malignant polyp  
 d. Familial polyposis
18. **Not true about GIST:** (Recent Question 2015)  
 a. Stomach is the most common site  
 b. High propensity of malignant change  
 c. Associated with c-KIT mutation  
 d. Histology shows epithelioid and spindle shaped cells
19. **MC site for stomach Ca:** (Recent Question 2015)  
 a. Lesser curvature  
 b. Antrum  
 c. Greater curvature  
 d. Pylorus

20. **Most common site of curling's ulcer?** (Recent Question 2015)  
 a. Proximal Duodenum  
 b. Esophagus  
 c. Distal duodenum  
 d. D. jujenum
21. **Most common site of GIST is** (Recent Question 2014)  
 a. Ileum  
 b. Esophagus  
 c. Colon  
 d. Stomach
22. **Best prognosis in Carcinoma stomach is seen in** (APPGMEE 14)  
 a. Superficial spreading type  
 b. Ulcerative type  
 c. Linitis plastica type  
 d. Polypoidal type
23. **Endoscopic biopsy from a case of H.pylori related duodenal ulcer is most likely to reveal:** (Recent Question 2013)  
 a. Antral predominant gastritis  
 b. Multifocal atrophic gastritis  
 c. Acute erosive gastritis  
 d. Gastric atrophy
24. **Most common complication of gastric ulcer:** (AIIMS June 13)  
 a. Tea pot stomach  
 b. Scirrhus carcinoma  
 c. Perforation  
 d. Massive haematemesis
25. **Histologic examination of the lesion in stomach reveal fat-laden cells, likely cause is:** (AIIMS Nov 11)  
 a. Lymphoma  
 b. Postgastrectomy  
 c. Signet-cell carcinoma  
 d. Atrophic gastritis
26. **Gastric carcinoma is associated with all EXCEPT:** (DNB Dec 11)  
 a. Inactivation of p53  
 b. Over expression of C-erb  
 c. Over expression of C-met  
 d. Activation of RAS
27. **True about autonomic atrophic gastritis:** (PGI Nov 2011, 2009)  
 a. Loss of parietal cells  
 b. Hypertrophy of G cells  
 c. Apoptosis of gland epithelial cells  
 d. Hypertrophy of ECL cells  
 e. Active inflammation to neuroendocrine gland
28. **The most common site of a benign (peptic) gastric ulcer is** (AIIMS June 11, 04)  
 a. Upper third of lesser curvature  
 b. Greater curvature  
 c. Pyloric antrum  
 d. Lesser curvature near incisura angularis
29. **Hour glass deformity is seen in:** (DNB Dec 11)  
 a. Carcinoma stomach  
 b. Peptic ulcer  
 c. Duodenal atresia  
 d. CHPS
30. **False about the malignant ulcer of stomach is:** (AI 2010)  
 a. The mucosal folds don't reach the edge of the ulcer  
 b. Mucosal folds are thickened and fused  
 c. Ulcer crater is eccentric  
 d. Margins of the ulcer are overhanging
31. **Which of the following markers is specific for gastrointestinal stromal tumor (GIST):** (AI 10, 09)  
 a. CD 117  
 b. CD 34  
 c. CD123  
 d. S-100




32. When carcinoma of stomach develops secondarily to pernicious anemia, it is usually situated in the:  
 a. Prepyloric region      b. Pylorus (DNB 2010)  
 c. Body      d. Fundus
33. True about *H. pylori* infection:  
 a. Gram +ve aerobe (WB PGME 2016, PGI Nov 2009)  
 b. Invade gastric mucosa and cause ulcer  
 c. Rapid urease test on endoscopy is diagnostic  
 d. Serology confirms eradication  
 e. Causes MALT lymphoma of stomach

### SMALL INTESTINE

34. A 23-year-old lady presented with diarrhea, vomiting and poor appetite. Biopsy showed crypt hyperplasia, villous atrophy and CD8+ cells in the lamina propria. Skin manifestations have been shown. What could be the diagnosis? (Recent Pattern Question 2020)




- a. Whipple's disease  
 b. Chronic pancreatitis  
 c. Environmental enteropathy  
 d. Celiac disease
35. Which of the following is associated with PAS positive macrophages? (Recent exam 2018)  
 a. Whipple disease      b. Abetalipoproteinemia  
 c. Crohn's disease      d. Ulcerative colitis
36. A patient presented with complains of chronic constipation and diarrhoea. There was excessive associated weight loss. Intestinal biopsy was obtained and it showed the following findings. What is your diagnosis? (AIIMS Nov 16)



- a. Giardia      b. Entamoeba  
 c. Whipple's disease      d. CMV

37. Which sugar is used to diagnose intestinal malabsorption? (Recent Question 2016-17)  
 a. Xylose      b. Glucose  
 c. Amylase      d. Lactose
38. Duodenal villous atrophy is seen in (Recent Question 2016-17)  
 a. Crohn's disease  
 b. Ulcerative colitis  
 c. Celiac disease  
 d. Cystic fibrosis
39. All are true about Peutz Jegher's syndrome except: (WBPGME 2016, Recent Question 2015)  
 a. Autosomal dominant  
 b. Hamartomatous polyps do not develop into adenocarcinoma  
 c. Gain of function mutation in LKB1/STK11  
 d. Chance of fatal intussusceptions
40. True about gluten sensitive enteropathy: (PGI May 2015)  
 a. Diet should exclude barley, wheat and rye  
 b. Intestinal biopsy is diagnostic  
 c. Anti IgA endomycial antibody is specific  
 d. Mucosal hyperplasia
41. Pathologist examines biopsy from a patient presenting with bleeding per rectum with a past history of intussusception for the past 6 months. Histopathology obtained has been shown below. Identify the Pathology? (AIIMS NOV 2015)



- a. Tubule villous adenoma  
 b. Adeno carcinoma  
 c. Hamartoma  
 d. Juvenile polyposis syndrome
42. Celiac sprue is associated with (Recent Question 2014-15)  
 a. HLA DQ1  
 b. HLA DQ2  
 c. HLA DQ3  
 d. HLA DQ4
43. Characteristic histopathology finding in Whipples disease is: (Recent Question 2014-15)  
 a. PAS positive macrophages and rod shaped bacilli in lamina propria  
 b. Shortened thickened villi with increased crypt depth  
 c. Blunting and flattening of mucosal surface and absent villi  
 d. Mononuclear infiltration at base of crypts




44. A 2-year-old male presented with abdominal distension, chronic diarrhea, severe anemia and failure to thrive? Which of the following is the investigation of choice?  
 a. Anti milk protein antibody (Recent Question 2015)  
 b. Anti endomysial antibody  
 c. Antinuclear antibody d. Intestinal biopsy
45. True about Peutz-jeghers syndrome: (PGI May 2013)  
 a. Pigmentary changes in skin and mucous membrane around mouth  
 b. Adenomatous polyp in intestine  
 c. Most common of pattern inheritance is autosomal recessive  
 d. 20-30% premalignant  
 e. May presents as anemia in children
46. Whipple's disease is characterized by?  
 a. Foamy macrophages (Recent Question 13)  
 b. AFB positive  
 c. Papillary projections  
 d. Villous atrophy
47. True about abdominal lymphoma: (PGI Nov 2011)  
 a. GIT lymphoma most commonly has polypoid appearance  
 b. Primary small intestinal lymphoma are most common  
 c. Lymphoma is most common primary malignant neoplasm of colon  
 d. Stomach is most common site for extranodal lymphoma  
 e. MALT lymphoma is associated with H. pylori infection
48. True about intestinal lymphoma: (PGI Nov 2010)  
 a. Involves in Non-Hodgkin's lymphoma  
 b. Made up of predominantly T cells  
 c. C-kit positive  
 d. Most common at ileocecal junction  
 e. Lymphadenectomy is always done
49. Paneth cells contain: (DNB June 11)  
 a. Zinc  
 b. Copper  
 c. Molybdenum  
 d. Selenium


### LARGE INTESTINE

50. Which of the following is/are features of typical Ulcerative colitis: (PGI May 2019)  
 a. Crypt distortion  
 b. Chronic inflammatory cells in the lamina propria  
 c. Crypt abscess  
 d. Granuloma  
 e. Crypt branching
51. Which of the following is a feature of Crohn's disease?  
 a. Pseudopolyps can be seen (Recent exam 2018)  
 b. Non-caseating granulomas are present  
 c. Backwash ileitis may be associated with Crohn's disease  
 d. Both b and c
52. Which of the following is the earliest change in intestine which occurs in Crohn's disease? (Recent exam 2018)  
 a. Cobblestone appearance  
 b. Aphthous ulcer  
 c. Perforation  
 d. Stricture


53. A 50-year-old male presented with recurrent bloody diarrhea. Colonoscopy showed geographical ulcers. Histopathology is shown below. What is your diagnosis?



- a. Pseudomebranous colitis (AIIMS May 2017)  
 b. Non-hodgkin lymphoma colon  
 c. Adenocarcinoma colon  
 d. Crohn disease
54. About Crohn's disease true is? (AIIMS May 2017)  
 a. Loss of haustration b. Linear fissure  
 c. Cobblestone colon d. String sign of Kantor  
 e. Pipe stem colon
55. Identify the parasite in the intestinal biopsy of a HIV positive patient. (Recent Pattern Question 2020)





- a. Giardia  
 b. CMV  
 c. Amoebic colitis  
 d. Cryptosporidium
56. A 26-year-old male presented with abdominal pain and bloody diarrhea of one week duration. The following colonoscopic biopsy is diagnostic of infection with (AIIMS May 16)



- a. Giardiasis b. Amoebiasis  
 c. Enterobius d. Severe bacterial infection





- 57. Extraintestinal manifestations of Crohn disease include ALL EXCEPT (JIPMER 2016)**
- Uveitis
  - Migratory polyarthritis
  - Sacroiliitis
  - Pericholangitis
- 58. Which gene involved in colonic adenoma to colonic carcinoma? (Recent Question 2016-17)**
- TP 53 (Protoncogene mutation)
  - Rb
  - K RAS (Protoncogene mutation)
  - EGFR
- 59. Which of the following is NOT associated with an increased risk of Gastrointestinal malignancy? (Recent Question 2015)**
- Cowden's syndrome
  - Lynch syndrome
  - Gardner's syndrome
  - HNPCC
- 60. In carcinoma of unknown primary, if the tissue marker CDX-2 is positive, it indicates: (Recent Question 2015)**
- Bladder cancer
  - Gastrointestinal cancer
  - Lung cancer
  - Thyroid cancer
- 61. All the following conditions are characterized by neoplastic polyps except: (MHPGMEE 2016, Recent Question 2015)**
- Peutz-Jeghers syndrome
  - Gardner syndrome
  - Turcot syndrome
  - Lynch syndrome
- 62. An 11-year-old girl presents with abdominal pain, no diarrhea, freckles lips, nostrils, buccal mucosa, palmar surfaces of the hands. Likely diagnosis: (Recent Question 2015)**
- Gardner syndrome
  - Cowden syndrome
  - Peutz-Jeghers syndrome
  - Cronkhite Canada syndrome
- 63. Not true about Cowden syndrome: (Recent Question 2015)**
- Mutation of PTEN gene
  - Multiple hamartomas
  - Increased risk of GI malignancy
  - Risk of follicular thyroid cancer
- 64. Most common site of carcinoid tumor in hindgut: (Recent Question 2015)**
- Caecum
  - Rectum
  - Transverse colon
  - Descending colon
- 65. Pseudopolyps are features of: (Recent Question 2015)**
- Crohn's disease
  - Ulcerative colitis
  - Celiac sprue
  - Whipple's disease
- 66. True about ulcerative colitis, all except: (Recent Question 14)**
- Rectum involved
  - Pseudopolyps
  - Pancolitis
  - Noncaseating granuloma
- 67. Inheritance of Gardner syndrome is: (Recent Question 2014)**
- Autosomal recessive
  - Autosomal dominant
  - X linked
  - None of the above
- 68. Aganglionic segment is encountered in which part of colon in case of Hirschsprung disease: (AIIMS Nov 14)**
- Distal to dilated segment
  - In Whole colon
  - Proximal to dilated segment
  - In the dilated segment
- 69. Acquired diverticulum most common site is: (Recent Question 2015)**
- Sigmoid colon
  - Ileum
  - Ascending colon
  - Transverse colon
- 70. Regarding FAP all true except- (JIPMER 2017)**
- Autosomal recessive
  - Duodenal polyp
  - More than 100 polyps
  - Extraintestinal manifestations
- 71. A 5 year-old boy presented with bleeding per rectum. PR showed rectal polyp, biopsy showed the following. What is your diagnosis? (Recent Pattern Question 2020)**
- 
- Villous adenoma
  - Peutz-Jeghers polyp
  - Juvenile polyp
  - Serrated adenoma
- 72. Which polyp has got the maximum risk of turning into malignancy: (Recent Question 2014)**
- Pseudopolyp
  - Hyperplastic polyp
  - Tubular villous adenomas (multiple)
  - Adenomatous polyps
- 73. Gene involved in HNPCC is: (Recent Question 2014)**
- APC
  - PTEN
  - HLH1
  - SKTH
- 74. FALSE statement regarding Hirschsprung disease is (AP-PGMEE 14)**
- Aganglionosis always involves distal rectum
  - Non passage of meconium in first 24 hours is a cardinal feature
  - Diagnosis is established by Suction rectal biopsy
  - No passage of stools after per rectal examination
- 75. The minimum number of polyps necessary for a diagnosis of Familial Adenomatous Polyposis (FAP) is: (APPGMEE 14)**
- 05
  - 10
  - 50
  - 100
- 76. Premalignant conditions of the GIT are: (PGI 13)**
- Ileocaecal TB
  - Familial polyposis
  - Villous adenomas
  - Ulcerative colitis
- 77. True about Carcinoembryonic antigen (CEA): (PGI May 2013)**
- Used for monitoring of recurrence of colon cancer
  - Specific for colon cancer
  - Increased in smokers
  - Increased in colon cancer





78. **In Hirschsprung's disease, staining used for diagnosis is?** (Recent Question 2013)  
 a. Fontana stain                      b. Trichome stain  
 c. AChE                                  d. Auramine Rhodamine stain
79. **Osteomas, adenomatous polyps of intestine and periampullary carcinomas are seen in:** (Recent Question 2013)  
 a. Cowden syndrome                  b. Peutz Jeghers syndrome  
 c. FAP                                      d. Gardner syndrome
80. **Dietary factors associated with colon carcinoma:**  
 a. High fiber                              b. Low fiber (PGI Dec 12)  
 c. Smoked fish                           d. High fat intake  
 e. Japanese are common to develop Ca colon
81. **Anti-Saccharomyces cerevisiae antibodies are seen in?** (PGI Dec 12, DNB June 11)  
 a. Crohn's disease                      b. Scleroderma  
 c. SLE                                      d. Sjogren's syndrome
82. **Inflammatory bowel disease with transmural involvement and skip lesions is?** (DNB June 11)  
 a. Crohn's disease                      b. Ulcerative colitis  
 c. Shigella infection                    d. Clostridium infection
83. **Not true about familial polyposis colon cancer syndrome?** (Jipmer 11)  
 a. Autosomal recessive  
 b. Associated with fibromas and osteomas  
 c. Associated with brain tumors  
 d. 100% incidence of colon ca
84. **True about intestinal lymphoma:** (PGI Nov 10)  
 a. Involved in non-Hodgkins lymphoma  
 b. Made up of predominantly T cells  
 c. C-kit positive  
 d. Most common site is ileocaecal junction  
 e. Lymphadenopathy always seen

### MISCELLANEOUS

85. **Least predilection for distal ileum:** (Recent Question 2014)  
 a. Carcinoid syndrome  
 b. Meckel diverticulum  
 c. Chrons disease  
 d. Zollinger Ellison syndrome
86. **Mesenteric tumours are:** (Recent Question 2013, Karnat 05)  
 a. Usually solid                              b. Usually cystic  
 c. Highly malignant                        d. Highly vascular



## Answers with Explanations

1. **Ans. (b) Cytomegalovirus** (Ref: Robbins 9th ed p 755)  
 CMV causes shallower ulcerations and characteristic nuclear and cytoplasmic inclusions within capillary endothelium and stromal cells. Infected cells are strikingly enlarged, often to a diameter of 40 µm, and show cellular and nuclear pleomorphism. Prominent intranuclear basophilic inclusions spanning half the nuclear diameter are usually set off from the nuclear membrane by a clear halo
2. **Ans. (a) A**  
 • The gross morphology in the lower part of esophagus shows tumor masses that may be polypoid, or exophytic, and protrude into and obstruct the lumen suggestive of carcinoma A.
3. **Ans. (a) Barret esophagus** (Ref: R 9/ p 757)  
 • Barrett's esophagus is the metaplastic change in the esophageal lining in which the normal squamous epithelium is changed to columnar epithelium due to prolonged gastroesophageal reflux (GERD)  
 • Columnar metaplasia, glandular metaplasia, goblet cell metaplasia.  
 • Definite diagnosis is made only when columnar mucosa contains **the intestinal goblet cells, which will be positive for PAS, Alcian blue (mucin stain)**
4. **Ans. (a, b, c, d) a. Depth of invasion; b. Lymph node status c. Tumour grading d. Stage of the disease**
5. **Ans. (b) Serosa** (Ref: Robbins 9th/pg ; 8th/pg 768)
6. **Ans. (b) Barretts oesophagus; mucin stain; dysplasia**  
 (Ref: Robbin's 9th/ 757)  
 • There are two figures given in this question.  
 • The first figure shows **lower oesophagus, in which normal squamous epithelium has been replaced by columnar epithelium.**  
 • This type of replacement of one mature tissue by another is called as **Metaplasia.**  
 • Now since in this case, **squamous epithelium is getting replaced by columnar epithelium; this is referred to as Columnar metaplasia** referred to as **Barrett's esophagus.**  
 • This can be diagnosed by H& E stain (1<sup>st</sup> picture) which shows **when columnar mucosa contains the intestinal goblet cells which show distinct mucous vacuoles that stain pale blue by hematoxylin and eosin.**  
 • **For specific staining of mucus, mucin stain can be used (2<sup>nd</sup> picture)**
7. **Ans. (a) Upper third of esophagus**  
 (Ref: R 9th/pg ; 8th/pg 768)  
 Ectopic gastric mucosa is seen most commonly in the **upper third of the esophagus**  
 This is known as inlet patch.



**8. Ans. (a) Absence of nerves**

(Ref: Robbins 9th/pg; 8th/pg 768)

Achalasia cardia is caused by selective **loss of function of inhibitory neurons** like those secreting vasoactive intestinal peptide and nitric oxide which causes relaxation of LES whereas cholinergic innervations is intact

**9. Ans. (a, b, c, d, e); a. Tylosis; b. Achalasia; c. Barrett's esophagus; d. Scleroderma; e. Plummer-Vinson syndrome** (Ref: Robbins 9th/pg 758-59; 8th/pg 773)

**10. Ans. (c) Lower 1/3**

(Ref: Robbins 9th/pg 759, Bailey & love 24th ed: 1009)

**11. Ans. (a) DOG 1; (b) CD117; (c) CD 34; (d) KIT**

(Ref: R 9th pg 785)

The findings are suggestive of GIST, IHC for GIST are DOG 1, CD117, CD 34, KIT

**12. Ans. (a, b, c) a. Negative for C-KIT & CD117; b. YOUNG age; c. MC site stomach**

Most gastrointestinal stromal tumors (GISTs) are characterized by KIT or platelet-derived growth factor alpha (PDGFRA) activating mutations. However, there are still 10%-15% of GISTs lacking KIT and PDGFRA mutations, called wild-type GISTs (WT GISTs). Among these so-called WT GISTs, a small subset is associated with succinate dehydrogenase (SDH) deficiency, known as SDH-deficient GISTs. In addition, GISTs that occur in Carney triad and Carney-Stratakis syndrome represent specific examples of SDH-deficient GISTs. SDH-deficient GISTs locate exclusively in the stomach, showing predilection for children and young adults with female preponderance. The tumor generally pursues an indolent course and exhibits primary resistance to imatinib therapy in most cases.

**13. Ans. (a) MALTOMA** (Ref: Robbins 9th/pg 773; 8th/pg 786)

- H. pylori gastritis induces mucosa-associated lymphoid tissue (MALT) that can give rise to B cell lymphomas (MALTomas).
- Histologically, gastric MALToma takes the form of a dense lymphocytic infiltrate in the lamina propria. Characteristically, the neoplastic lymphocytes infiltrate the gastric glands focally to create diagnostic **lymphoepithelial lesions**

**14. Ans. (a) Stomach** (Ref: Robbins 9th/pg 771; 8th/pg 785)

**Metastasis** from stomach cancer Occurs to the liver (**first organ to be affected**) followed by lungs, bone, ovary (**where it is known as Krukenberg's tumor**), periumbilical lymph nodes (Sister Mary Joseph nodule), peritoneal cul-de-sac (**Blumer's shelf** palpable on rectal or vaginal examination) and **left supraclavicular lymph node (Virchow's lymph node)**

**15. Ans. (d) Overuse of salicylates** (Ref: R 9th/pg; 8th/pg 775)

**16. Ans. (d) Zone of cicatrization** (Ref: Robbins 9th/pg 766)

**17. Ans. (a) Hyperplastic polyp**

(Ref: Robbins 9th/pg 769; 8th/pg 783)

**18. Ans. (b) High propensity of malignant change**

(Ref: Robbins 9th/pg 775; 8th/pg 789)

**19. Ans. (b) Antrum** (Ref: Robbins 9th/pg 771; 8th/pg 785)

- The most common location of the gastric cancer is the **antrum of the stomach**.<sup>Q</sup>
- Cancer of the **gastric cardia is on the rise especially due to Barrett esophagus**<sup>Q</sup>
- **Lesser curvature**<sup>Q</sup> is involved more often than the greater curvature.
- **Hence the most common site is lesser curvature of anteropyloric region**<sup>Q</sup>

**20. Ans. (a) Proximal duodenum** (Ref: Robbins 9th/pg 762)

**Curling ulcers**-Ulcers occurring in the proximal duodenum and associated with severe burns or trauma

**21. Ans. (d) Stomach** (Ref: Robbins 9th/pg 775; 8th/pg 789)

**22. Ans. (a) Superficial spreading type** (Ref: R 9th/pg 771)

The **depth of invasion and the extent of nodal and distant metastases** at the time of diagnosis remain the **most powerful prognostic indicators** in gastric cancer.

**23. Ans. (a) Antral predominant gastritis**

(Ref: Harrison 18th ed: 2458, 17th ed:1870)

- The **most common cause** of chronic gastritis- **H. pylori**<sup>Q</sup>
- Within the stomach, H. pylori are **most often found in the antrum – antral predominant gastritis**<sup>Q</sup> or **type B gastritis**<sup>Q</sup>

**24. Ans. (d) Massive haematemesis**

(Ref: Robbins 9th/pg 767)

**25. Ans. (d) Atrophic gastritis**

(Ref: Odze RD, Goldblum JR (2009). *Surgical Pathology of the GI Tract, Liver, Biliary Tract, and Pancreas*. Philadelphia, PA: Saunders, <http://www.histopathology-india.net/gaxan.htm>, *Gastrointestinal Pathology: an atlas and text*. Philadelphia, PA: Lippincott Williams & Wilkins)

Collection of lipid laden macrophages within the lamina propria is defined as **gastric xanthoma**

Associated with pathological lesions such as **chronic gastritis** & intestinal metaplasia, **atrophic gastritis**, and gastric ulcer. **Rarely these are also seen in duodenogastric reflux after gastric surgery hence d> b**

Usually located at the antral and the lesser curvature Positive with Sudan black, oil red O (on frozen section), and CD68

**Differential diagnosis**

Signet ring carcinoma-mucin laden tumor cells infiltrate gastric wall. PAS/alcan blue positive



26. **Ans. (d) Activation of RAS** (Ref: Robbins 9th/pg 771)

- MC mutations seen in gastric carcinoma- **p53**
- Least common mutation seen in gastric carcinoma- K-Ras
- MC mutations seen in diffuse type gastric carcinoma- CDH1 gene mutations
- MC mutations seen in intestinal type gastric carcinoma- **Wnt pathway, APC,  $\beta$ -catenin, e-erb 2 amplification**
- Mutations seen in both intestinal and diffuse type gastric cancers- **p53, Cmet and cyclin E genes amplification**

27. **Ans. (a, b, d); a. Loss of parietal cells; b. Hypertrophy of G cells; d. Hypertrophy of ECL cells** (Ref: R 9th/pg 764)

28. **Ans. (d) Lesser curvature near incisura angularis**

(Ref: Robbins 9th/pg 766; 8th/pg 776)

29. **Ans. (b) Peptic ulcer**

(Ref: Robbins 9th/pg 766; 8th/pg 776)

30. **Ans. (d) Margins of the ulcer are overhanging**

(Ref: Chandrasoma Taylor 3ed: 582,587) Read Pretext

31. **Ans. (a) CD 117** (Ref: Robbins 9th/pg 775; 8th/pg 789)

**Most useful diagnostic marker is c-kit (CD117)** detectable in 95% of the patients.

32. **Ans. (d) Fundus** (Ref: Robbins 9th/pg 771; 8th/pg 785)

- **Most common site of Gastric Adenocarcinoma secondary to H. pylori infection is Antrum<sup>Q</sup>**
- **Most common site of Gastric Adenocarcinoma secondary to Pernicious anemia is Fundus and Body<sup>Q</sup>**

33. **Ans. (e) Causes MALT lymphoma of stomach**

(Ref: Robbins 9th/pg 764; 8th/pg 777)

- Option a false-H.pylori is a gram-negative flagellated bacteria
- Option b false-The organism is concentrated within the **superficial mucus<sup>Q</sup>** overlying epithelial cells
- Option c-true, rapid urease test has high sensitivity and specificity, can be **false negative<sup>Q</sup>** with recent use of PPIs. **Gold standard test is : antral biopsy showing the bacilli<sup>Q</sup>**
- Option d: Serology- Cannot be used for early follow-up  
Urea breath test- useful for follow-up after treatment
- Option e-true- **Most common inducer of gastric MALToma -H Pylori**

34. **Ans. (d) Celiac disease** (Ref: R 9th pg 782)

35. **Ans. (a) Whipple disease** (Ref: Robbins 9th ed p 792)

36. **Ans. (a) Giardia** (Ref: www.ncbi.nlm.nih.gov/pubmed/104699)

This image shows Giardia lamblia infection of the small intestine. The small pear-shaped trophozoites live in the duodenum and become infective cysts that are excreted. They produce a watery diarrhea. A useful test for diagnosis of infectious diarrheas is stool examination for ova and parasites.

37. **Ans. (a) Xylose**

(Ref: [https://en.wikipedia.org/wiki/D-xylose\\_absorption\\_test](https://en.wikipedia.org/wiki/D-xylose_absorption_test))

D-xylose is a monosaccharide, that does not require enzymes for digestion prior to absorption. Its absorption requires an intact mucosa only. In contrast, polysaccharides require enzymes, such as amylase, to break them down so that they can eventually be absorbed as monosaccharides. This test was previously in use but has been made redundant by antibody tests.

In normal individuals, a 25 g oral dose of D-xylose will be absorbed and excreted in the urine at approximately 4.5 g in 5 hours. A decreased urinary excretion of D-xylose is seen in conditions involving the GI mucosa, such as small intestinal bacterial overgrowth and Whipple's disease if the D-xylose urinary excretion is not normal after a course of antibiotics, then small intestinal bacterial overgrowth is ruled out and non-infectious cause of malabsorption (i.e., celiac disease) is suggested.

38. **Ans. (c) Celiac disease** (Ref: R9/ 1782)

D Xylose absorption test is a medical test to diagnose condition that cause malabsorption of proximal small intestine.

39. **Ans. (c) Gain of function mutation in LKB1/STK11**

(Ref: Robbins 9th/pg 806; 8th/pg 817)

40. **Ans. (a, c) a. Diet should exclude barley, wheat & rye; c. Anti IgA endomysial antibody is specific**

(Ref: Robbins 9th/pg 782; 8th/pg 795)

41. **Ans. (c) Hamartoma**

(Ref: Robbins 9th/pg 806; SEE ans 57 Above)

**Clinically** history of intussusception and pathologically we see **arborization and presence of smooth muscle intermixed with lamina propria- marked with an arrow, both are suggestive of peutz jegher polyp which is a hamartomatous polyp.**

42. **Ans. (b) HLA DQ2** (Ref: Robbins 9th/pg 782; 8th/pg 795)

43. **Ans. (a) PAS positive macrophages and rod shaped bacilli in lamina propria**

(Ref: Robbins 9th/pg 783; 8th/pg 796)

44. **Ans. (b) Anti endomysial antibody**

(Ref: <http://emedicine.medscape.com/article/932104-clinical>, Robbins 9th/pg 782; 8th/pg 795)

45. **Ans. (a, e); a. Pigmentary changes in skin and mucous membrane around mouth; e. May presents as anemia in children** (Ref: Robbins 9th/pg 806; 8th/pg 817)

PJ Syndrome come to attention for following

- Intussusception
- Mucocutaneous pigmentation
- Secondary cancers
- Hematochezia can present in 14% cases – can present as anemia



46. Ans. (a) **Foamy macrophages** (Ref: Robbins 9th/pg 783)

47. Ans. (d, e); **d. Stomach is most common site for extranodal lymphoma; e. MALT lymphoma is associated with H. pylori infection**

(Ref: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3042647/>)

48. Ans. (a) **Involves in Non-Hodgkin's lymphoma**

(Ref: <http://www.ncbi.nlm.nih.gov/pmc/articles/C30426L47/>;  
See Ans 66)

49. Ans. (a) **Zinc**

(Ref: Ross histology; 4th ed, Pathology of stomach and duodenum: 320)

50. Ans. (a) **Crypt distortion; (b) Chronic inflammatory cells in the lamina propria; (c) Crypt abscess; (e) Crypt branching**

(Ref: R 9th pg 799)

Noncaseating granuloma is the hallmark of Crohn's disease.

51. Ans. (c) **Non-caseating granulomas are present**

(Ref: Robbins 9th ed p 799)

52. Ans. (b) **Aphthous ulcer** (Ref: Robbins 9th ed p 799)

The earliest lesion in Crohn disease, the aphthous ulcer, may progress, and multiple lesions often coalesce into elongated, serpentine ulcers oriented along the axis of the bowel.

53. Ans. (d) **Crohn disease** (Ref: R 9/ 798)

The picture shows – transmural inflammation (blue part) and an overlying ulcer. The best option here is Crohn's disease.

54. Ans. (c, d) **c. Cobblestone colon; d. String sign of Kantor**

Aphthoid ulcers develop into linear ulcers and fissures to produce an ulceronodular or “cobblestone” appearance.

55. Ans. (d) **Cryptosporidium** (Ref: R 9th pg 800)

56. Ans. (b) **Amoebiasis** (Ref: <https://msu.edu>)

57. Ans. (d) **Severe bacterial infection**

(Ref: Gut liver 2010, 4 (3): 338-344)

Uveitis and polyarthritis are common extra intestinal manifestations of Crohn's disease.

Sacroiliitis is more common in Crohn's disease (21%) as compared to U.C (12.2%) especially in patients with upper GI or perianal involvement.

Pre cholangitis is common extra intestinal manifestation of U.C.

58. Ans. (c) **K RAS (Protooncogene mutation)**

<b>Normal colon</b>	<b>APC at 5q21</b>	Germline (inherited) or somatic (acquired) mutations of cancer suppressor genes (“first hit”)
<b>Mucosa at risk</b>	<b>APC b-catenin</b>	Methylation abnormalities Inactivation of normal alleles (“second hit”)
<b>Adenomas</b>	<b>K-RAS at 12p12</b>	Protooncogene mutations
<b>Adenomas</b>	<b>TP53 at 17p13 LOH at 18q21 (SMAD 2 and 4)</b>	Homozygous loss of additional cancer suppressor genes
<b>Carcinoma</b>	<b>Telomerase,</b>	Additional mutations Gross chromosomal alterations

Remember both p53 & KRAS mutations occur, However p53 to tumor suppressor gene (not oncogene)

59. Ans. (a) **Cowden's syndrome** (Ref: Robbins 9th/pg 1316)

All are neoplastic polyps except Cowden syndrome  
In Cowden syndrome – there is no increased risk of GI malignancy but other organ malignancies can develop  
**Cowden Syndrome and Bannayan-Ruvalcaba-Riley Syndrome** – autosomal dominant hamartomatous polyp syndromes associated with loss-of-function mutations in **PTEN** hence also known as **PTEN hamartoma syndrome**.

60. Ans. (b) **Gastrointestinal cancer**

(Ref: Am J Surg Pathol 2003;27:303)

- CDX2: Also called caudal-related homeobox gene 2, caudal type homeobox transcription factor 2
- Fairly specific marker of GI origin for adenocarcinomas

61. Ans. (a) **Peutz-Jeghers syndrome** (Ref: Robbins 9th/pg 806)

Syndromes with hamartomatous polyps –  
**Cowden Syndrome and Bannayan-Ruvalcaba-Riley Syndrome** – autosomal dominant hamartomatous polyp syndromes associated with loss-of-function mutations in **PTEN** hence also known as **PTEN hamartoma syndrome**.  
**Cronkhite-Canada Syndrome** – contrasts sharply with other hamartomatous polyposis syndromes as it is nonhereditary and develops in individuals over 50 years of age

62. Ans. (c) **Peutz-Jeghers syndrome**

(Ref: Robbins 9th/pg 806)

63. Ans. (c) **Increased risk of GI malignancy**

(Ref: Robbins 9th/pg 1316)

64. Ans. (b) **Rectum** (Ref: Harrison 18th ed: 350-3)

65. Ans. (b) **Ulcerative colitis**

(Ref: Robbins 9th/pg 798; 8th/pg 810, Harsh Mohan 4th ed: 543)



**66. Ans. (d) Noncaseating granuloma**

(Ref: Robbins 9th/pg 798-800; 8th/pg 810-811)

Noncaseating granuloma is seen in chrons

**67. Ans. (b) Autosomal dominant** (Ref: Robbins 9th/pg 809)**68. Ans. (a) Distal to dilated segment**

(Ref: Robbins 9th/pg 751)

**69. Ans. (a) Sigmoid colon** (Ref: Robbins 9th/pg 751)

- True diverticulum is defined by the presence of **all three layers** of the bowel wall.<sup>o</sup>
- **Most common true diverticulum** is the Meckel diverticulum, which occurs in the **ileum**.
- **Acquired diverticula:** Most common site: **sigmoid colon**<sup>o</sup>

**70. Ans. (a) Autosomal recessive****71. Ans. (c) Juvenile polyp** (Ref: R 9th pg 808)**72. Ans. (d) Adenomatous polyps**

(Ref: Robbins 9th/pg 808,809)

Option a is pseudopolyp in inflammatory condition-are not premalignant condition

Option b- hyperplastic polyp is **Non-neoplastic polyp**

Here the doubt arises between 2 options: option c and d

Option d-adenomatous polyposis (FAP) is an autosomal dominant disorder in which patients develop numerous colorectal adenomas as teenagers

Colorectal adenocarcinoma develops in 100% of untreated FAP patients, often before age 30 and nearly always by age 50

**Option c- adenomas-** The most common neoplastic polyps are colonic adenomas, which are precursors to the majority of colorectal adenocarcinomas.

**73. Ans. (a) APC** (Ref: Robbins 9th/pg 809; 8th/pg 820-824)**74. Ans. (d) No passage of stools after per rectal examination**

(Ref: <http://www.aafp.org/afp/2006/1015/p1319.html>)

- Most cases of Hirschsprung disease are diagnosed in the newborn period. Hirschsprung disease should be considered in any newborn that fails to pass meconium within 24-48 hours of birth-option b true
- Distal intestinal segment that lacks both the Meissner submucosal and the Auerbach myenteric plexus. - option a is true
- Proximal to aganglionic segment, colon undergoes progressive dilation
- Rectal examination may demonstrate a tight anal sphincter and explosive discharge of stool and gas - option d is false
- A rectal suction biopsy can detect hypertrophic nerve trunks and the absence of ganglion cells in the colonic submucosa, confirming the diagnosis.- option c is true

- Down syndrome (trisomy 21) is the most common chromosomal abnormality associated with the disease, accounting for approximately 10 percent of patients

**75. Ans. (d) 100** (Ref: Robbins 9th/pg 809; 8th/pg 820-824)

- **Classic FAP -At least 100 polyps are necessary for a diagnosis**<sup>o</sup>
- **Attenuated FAP-lower number of adenomatous polyps (around 30)**

**76. Ans. (b, c); b. Familial polyposis; c. Villous adenomas**

(Ref: Robbins 9th/pg 808,809)

**77. Ans. (a, c, d); a. Used for monitoring of recurrence of colon cancer; c. Increased in smokers; d. Increased in colon cancer**

(Ref: De Mais, Daniel. ASCP Quick Compendium of Clinical Pathology, 2nd Ed. ASCP Press 2009)

CEA-Carcinoembryonic antigen (CEA) is a **glycoprotein**<sup>o</sup>

CEA measurement is mainly used as a **tumor marker** to monitor colorectal carcinoma treatment, to identify **recurrences** after surgical resection, for **staging** or to localize cancer spread through measurement of biological fluids.

**78. Ans. (c) AChE**

(Ref: <http://ajcp.ascpjournals.org/content/126/1/9.full.pdf>)

In addition to absent intrinsic ganglion cells in hirschsprung disease, most striking and diagnostically useful finding is the presence of hypertrophic nerve fibers in the myenteric and submucosal plexuses.

The diagnostic approach in hirschsprung is:

- Identify ganglion cells.- H & E stain
- Acetylcholinesterase histochemistry (AChE staining), -
  - **Hirschsprung disease**-abnormally thick and numerous nerve fibers in the muscularis mucosa and lamina propria
  - **Normal rectal mucosa**-relatively sparse, thin AChE-positive nerve fibers that are limited largely to the deep muscularis mucosa

**79. Ans. (d) Gardener syndrome** (Ref: Robbins 9th/pg 1316)

- **Gardner syndrome:** osteomas of mandible, skull, and long bones, epidermal cysts, desmoid tumors, thyroid tumors, and dental abnormalities, including **unerupted and supernumerary teeth**
- **Turcot syndrome-** intestinal adenomas and tumors of the central nervous system (**MC- Medulloblastoma or Glioblastoma**)
- **Both gardener and turcot syndrome have mutated APC gene and are associated with FAP**

**80. Ans. (b, d); b. Low fiber; d. High fat intake**

(Ref: Robbins 9th/pg 811; 8th/pg 822)

**81. Ans. (a) Crohn's disease** (Ref: Robbins 9th/pg 798)**82. Ans. (a) Crohn's disease** (Ref: Robbins 9th/pg 798)



83. Ans. (a) Autosomal recessive (Ref: Robbins 9th/pg 809)

84. Ans. (a) Involved in non-Hodgkins lymphoma

(Ref: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3042647>)

85. Ans. (d) Zollinger Ellison syndrome

(Ref: Robbins 9th/pg 798; 8th/pg 810)

Option a, b, c occur ileum. Option d -ZES (gastrinoma) does not occur in distal ileum

86. Ans. (b) Usually cystic

(Ref: World J Surg Oncol. May 19 2009;7:47)

Mesenteric tumors have been described as cystic in 40-60% of cases- **option b is true**

Solid primary tumors of the mesentery are rare- **option a is false**

**Option c is false**- Malignant primary mesenteric tumors are extremely uncommon

**Option d is false** -Two thirds of malignant mesenteric tumors are mesenchymatous (most characterized as leiomyosarcoma or liposarcoma), while the remainder are primarily lymphomas.

[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are approximately 20 lines visible. The paper has a slightly aged or off-white appearance. There is no handwriting or other markings on the page.

# Liver, Gallbladder, Pancreas and its Disorders

## Key Points

- » Normal adult liver weighs **1400 to 1600 gm** & has a **dual blood supply** from **portal vein** & **hepatic artery**
- » **Kupffer cells** are **modified macrophages of the liver**
- » **Acetaminophen poisoning** is the most common cause of hepatic toxicity in **Western countries**
- » **Capillarization of Sinusoids** is the **Hallmark of Cirrhosis**
- » **Proliferation & activation** of hepatic stellate cells is caused by **PDGF- $\beta$  and TNF**
- » Most common **mode of Hepatitis B transmission** in **India** is **Horizontal**
- » **20%** of individuals **with chronic HCV** infection progress to cirrhosis
- » HEV infection has the **highest mortality rate** among **pregnant** women
- » Diagnostic hallmark of **Chronic Hepatitis B** is "**ground-glass**" hepatocytes
- » Anti-liver kidney microsome-1 (**anti-LKM-1**) is **seen in type II autoimmune hepatitis**
- » **Alcoholic liver disease begins in acinus zone 3** & extends outwards toward the portal tracts
- » Most common acquired metabolic disorder is non-alcoholic fatty liver disease
- » Peliosis hepatitis is associated with anabolic steroids, Danazol, OCPs and tamoxifen
- » **Hepatoblastoma** is the **most common liver tumor of early childhood**
- » **Fibrolamellar Carcinoma** is variant of HCC, associated with a generally favorable prognosis
- » **Metastasis to Liver** are the **most common malignant tumors in liver**
- » **Pancreas divisum** is the **most common<sup>o</sup>** congenital anomaly of the pancreas

## Key Recent Updates

- » NASH is a component of metabolic syndrome.
- » Pediatric NAFLD shows diffuse steatosis, portal fibrosis and portal and parenchymal mononuclear cells.





## LIVER

### NORMAL ANATOMY OF LIVER

- Normal adult liver weighs **1400 to 1600 g**.
- Liver has a **dual blood supply**, with **portal vein providing 60%-70%** & **hepatic artery supplying 30%-40%**.
- Hepatic micro-architecture is based on the **lobular model**.
- Hepatocytes around central hepatic vein are called "**centrilobular**" (**zone 3**) while those near the portal tract are "**periportal**" (**zone 1**)
- Between the trabecular plates of hepatocytes are **vascular sinusoids**, lined by fenestrated endothelial cells.
- Space** between **sinusoids** and **hepatocytes** is **Space of Disse**.
- Kupffer cells** are **modified macrophages of the liver** that are attached to the sinusoids
- Fat-containing myofibroblastic **stellate cells** are found in the **space of Disse**.
- Flow of bile:** Hepatocytes → **bile canaliculi** → **canals of Hering** → **bile ductules** (periportal region) → **terminal bile ducts** (in portal triad)

### HEPATIC INJURY

Occurs in two forms: Necrosis and Apoptosis

Necrosis	Apoptosis
<ul style="list-style-type: none"> <li>Due to <b>ischemic</b> or <b>hypoxic</b> injury &amp; <b>oxidative stress</b><sup>Q</sup></li> <li>Can be Confluent, Zonal or Bridging Necrosis</li> </ul>	<ul style="list-style-type: none"> <li>Apoptotic hepatocytes in <b>yellow fever</b> are called '<b>Councilman bodies</b>'<sup>Q</sup></li> <li><b>Apoptotic bodies</b> in <b>acute &amp; chronic hepatitis</b> are termed <b>acidophil bodies</b><sup>Q</sup> (due to deep eosinophilic staining)</li> </ul>

### LIVER FAILURE

#### Acute Liver Failure

**Acute liver illness** associated with **encephalopathy** and **coagulopathy** that occurs **within 26 weeks** of the initial liver injury in the absence of pre-existing liver disease.

**Morphology:** Depends on the duration and nature of injury: shows **Massive hepatic necrosis** with **parenchymal loss** with **regenerating hepatocytes**.

#### High Yield Facts

- M.C cause** of acute liver failure is massive hepatic necrosis induced by **drugs or toxins**.
- Acetaminophen poisoning** is the most common cause in **Western countries** followed by **autoimmune hepatitis**
- In Asia, **acute hepatitis B & E** are the M.C cause of acute liver failure

## CIRRHOSIS

Defined histopathologically by:

- Bridging fibrous scars** linking portal tracts with one another and portal tracts with terminal hepatic veins.<sup>Q</sup>
- Fibrosis is the key feature of progressive damage to the liver.**<sup>Q</sup>
- Parenchymal nodules:** due to repeated cycles of **hepatocyte regeneration and scarring**
  - <3 mm – **micronodules**<sup>Q</sup>
  - >3 mm – **macronodules**<sup>Q</sup>
- Disruption of the architecture of the entire liver.**

### Causes

Acquired Causes	Inherited Metabolic Liver Disease
<ul style="list-style-type: none"> <li><b>Alcoholism</b></li> <li><b>Chronic viral hepatitis</b> (Hepatitis B &amp; C)</li> <li><b>Non-Alcoholic Steato-Hepatitis (NASH)</b></li> <li>Autoimmune hepatitis</li> <li>Biliary cirrhosis <ul style="list-style-type: none"> <li>Primary biliary cirrhosis</li> <li>Primary sclerosing cholangitis</li> </ul> </li> <li>Autoimmune cholangiopathy</li> <li>Cardiac cirrhosis</li> </ul>	<ul style="list-style-type: none"> <li>Hemochromatosis</li> <li>Wilson's disease</li> <li><math>\alpha</math>1-Antitrypsin deficiency</li> <li>Cystic fibrosis</li> </ul>
	Cryptogenic Cirrhosis
	Many patients who were thought to have cryptogenic cirrhosis are ultimately found to have nonalcoholic steatohepatitis

### Pathogenesis

The central pathogenic processes in cirrhosis are:


- Death of hepatocytes**
- Extracellular matrix (ECM) deposition**<sup>Q</sup>
- Vascular reorganization**<sup>Q</sup>

R<sup>9th</sup>


#### Latest Update

In cirrhosis, Kupffer cell activation leads to:

Functions performed	Cytokines involved
Proliferation & activation of hepatic stellate cells	PDGF- $\beta$ and TNF
Contraction of myofibroblasts	Endothelin-1 (ET-1)
Fibrosis	TGF- $\beta$ , Metalloproteinase 2 (MMP-2); Tissue inhibitors of MMP 1 & 2 (TIMP-1 & -2)
Chemotaxis to areas of injury	PDGF & monocyte chemoattractant protein-1 (MCP-1).



Gross liver showing nodular surface S/o Cirrhosis




Mic of cirrhotic liver showing nodules

- **Capillarization of Sinusoids is the Hallmark of Cirrhosis<sup>Q</sup>**
  - Normally collagen **types I & III** are concentrated in portal tracts and around central veins while **type IV collagen** are present in the **space of Disse**.
  - In cirrhosis, **types I and III collagen** are deposited in the **space of Disse**
  - Leads to **loss of fenestrations of sinusoids (Capillarization of sinusoids)**, impairing the function of sinusoids as channels that permit the exchange of solutes between hepatocytes and plasma.

### Scar Formation

- Principal cell type involved in scar deposition is the **hepatic stellate cell (ITO cells)<sup>Q</sup>**
- **ITO cells** are **Vitamin-A storage cells<sup>Q</sup>**
- In acute & chronic injury, the **stellate cells** get activated to **highly fibrogenic cells** called **myofibroblasts<sup>Q</sup>**

## HEPATITIS



### Acute Infective Hepatitis

- **Etiology: Hepatotrophic viruses: Hep A, B, C, D and E<sup>Q</sup>.**
  - Other Systemic Viruses which can cause hepatitis: EBV, CMV, Herpes virus and Adenovirus Yellow fever (yellow fever virus)

### Hepatitis A

- **Causative organism:** Hepatitis A virus (HAV)-Non-enveloped 27-nm, heat-, acid- & ether-resistant **RNA virus** in the **Hepatovirus** genus of the **Picornavirus** family<sup>Q</sup>

- **Mode of transmission: Feco-oral<sup>Q</sup> (most common), Sexual ±<sup>Q</sup>, Percutaneous route**
- **Incubation period: 2 to 6 weeks<sup>Q</sup>**
- **Clinical course**
  - **Fulminant hepatitis:** 0.1%
  - **Progression to chronicity:** None<sup>Q</sup>
  - **Carrier state:** None<sup>Q</sup>
  - **Prognosis:** Excellent
- **Serology**
  - **Early fecal shedding** of HAV **2-3 weeks before & 1 week after<sup>Q</sup>** the onset of jaundice.
  - **Diagnosis:** **IgM anti-HAV<sup>Q</sup>**
  - **Previous infection:** **IgG anti-HAV<sup>Q</sup>**

### Hepatitis B


- **Causative organism:**  
**Hepatitis B virus (HBV): Hepadna virus family**
  - 42 nm double-shelled virion with spherical 3.2-kb **DNA**, circular, ss/ds (**incomplete ds**)<sup>Q</sup>
  - Hepatitis B has eight subtypes and eight genotypes (A-H)
  - Genotypes **B (adw)** and **C (adr)** predominate in Asia<sup>Q</sup>
- **HBV Genes & Antigens**

Gene	Antigen produced
<b>C gene</b>	<b>HBcAg</b> , hepatitis B core antigen
<b>C &amp; Pre C genes</b>	<b>HBsAg</b> (hepatitis B e antigen)
<b>S gene</b>	<b>HBsAg</b> , (hepatitis B surface antigen): large, middle, and small HBsAg
<b>P gene</b>	<b>DNA polymerase (pol)</b> and <b>reverse transcriptase</b>
<b>X gene</b>	<b>HBx Ag:</b> virus <b>replication</b> and <b>transcriptional transactivator</b>

- **Mode of transmission**
  - In **High prevalence** regions (**≥8%**): **Perinatal** (most important)<sup>Q</sup>
  - In **Intermediate prevalence** areas (**2-7%; India**)<sup>Q</sup>: **Horizontal-** Blood products, percutaneous- minor breaks in the skin/mucous membranes, perinatal (vertical)
  - In **Low prevalence** areas (**<2%**): Sexual and intravenous drug abuse.



- **Incubation period:** 2 to 26 weeks
- **Clinical course in Hep B can be presented as:**



### Serologic Patterns of Hepatitis B Infection

HBsAg	Anti-HBs	Anti-HBc	HBeAg	Anti-HBe	Interpretation
+	-	IgM	+	-	Acute hepatitis B, high infectivity <sup>Q</sup>
+	-	IgG	+	-	Chronic hepatitis B, high infectivity
+	-	IgG	-	+	"Precore-mutant" (HBeAg -ve)
+	+	+	+/-	+/-	(Surface mutant) <sup>Q</sup> Seroconversion from HBsAg to anti-HBs
-	-	IgM	+/-	+/-	1. Acute hepatitis B 2. Anti-HBc "window" <sup>Q</sup>
-	-	IgG	-	+/-	Hep B in remote past
-	+	IgG	-	+/-	Recovery from hepatitis B
-	+	-	-	-	After Hep B vaccination

### High Yield Facts

- The most common **mode of Hepatitis B transmission in India** is: **Horizontal**<sup>Q</sup>
- **Age** at the time of onset of infection is the **best predictor of chronicity**
- **Younger** the age at the time of HBV infection, **higher is the probability of chronicity**
- **Prognosis** of Hepatitis B worsens with age
- **Pre-core Mutants** are strains of HBV that **do not produce HBeAg** despite HBV DNA.
- **Vaccine-induced escape mutant:** replicate in the presence of vaccine-induced immunity.

### High Yield Facts

#### Incidence of Fulminant Hepatitis in:

- Hep A-0.1%
- Hep-B ~1%
- Hep C-0.1%
- Hep D-5.20%
- Hep E-1-2% (20% in pregnancy)

### Hepatitis C

- **Causative organism:** 40-60 nm enveloped virus, 9.4-kb RNA, linear, ssRNA virus of **Hepacivirus** family
- **Mode of transmission:** IV drug abuse, sexual, needle-stick injury, blood products (horizontal), perinatal ±
- **Incubation period:** 15-160 days
- **Clinical course:**
  - **Fulminant hepatitis:** 0.1%<sup>Q</sup>
  - **Progression to chronicity:** common **upto 85%**<sup>Q</sup>
  - **Cirrhosis:** 20% of individuals **with chronic HCV** infection.<sup>Q</sup>
- **Prognosis:** Moderate

### Hepatitis D

- **Causative agent:** Hepatitis D virus (HDV) or "the delta agent," is a unique RNA virus that is dependent for its life cycle on HBV
- **Mode of transmission:**
  - Percutaneous > Sexual > Perinatal
  - **More common in IV drug abusers & multiple blood transfusions.**<sup>Q</sup>
- **2 types of infection:**
  - **Co-infection:** Due to exposure to serum containing **both HDV & HBV at the same time.**
    - Higher rate of acute hepatic failure in intravenous drug users.
    - Acute co-infection by HDV & HBV is best indicated by **Anti-HDV & Anti-HBcIgM**<sup>Q</sup>
- **Superinfection:** When a **chronic carrier of HBV** is exposed to HDV
  - Disease progresses to **cirrhosis** and **hepatocellular carcinoma**
  - With chronic delta hepatitis, **HBsAg & anti-HDV IgG and IgM persist for months**<sup>Q</sup>
- **Incubation period:** 30-180 days
- **Clinical course:**
  - **Progression to chronicity:** Common<sup>Q</sup>
- **Prognosis:** Acute → good, but Chronic → poor
- **Prevention:** **Vaccination for HBV** also prevents HDV infection<sup>Q</sup>





## High Yield Facts

- **Persistent infection** and **chronic hepatitis** are the **hallmarks** of HCV infection
- Due to changing structure of **HCV RNA polymerase**, **multiple genotypes** of viruses are found in the same patient after some duration of Hep C infection – “**Quasi-species**”
- **E2 protein** of the envelope **most variable region** of the entire viral genome → escape from neutralizing antibodies
- **Genomic instability & antigenic variability** are responsible for **persistent infection & ineffective HCV vaccine**.
- Hepatitis C (especially **HCV genotype 3**) infection is the association with **metabolic syndrome**
- HCV can give rise to **insulin resistance & non-alcoholic fatty liver disease (NAFLD)**

## Hepatitis E

- **Causative organism:** 32–34 nm non-enveloped icosahedral 7.6 kb ss linear RNA of **Hepevirus family**
- **Mode of transmission:** **Faeco-oral**<sup>Q</sup>
- **Incubation period:** 14–60 days
- **Clinical course:**
  - **Progression to chronicity:** None<sup>Q</sup> (Seen in AIDS & immunosuppressed **transplant patients**)<sup>Q</sup>
- **Prognosis:** Good (except in pregnant women)<sup>Q</sup>




## High Yield Facts

- **HEV infection** accounts for more than **30%–60% cases of sporadic acute hepatitis** in India, exceeding the frequency of HAV.<sup>Q</sup>
- Characteristic feature of HEV infection is the **high mortality rate** among **pregnant women**, approaching **20%**.<sup>Q</sup>
- **Carrier state:** Individual who **harbors** and can **transmit** an organism, but has **no symptoms**.


## Morphology of Acute Viral Hepatitis

- **Gross:** Normal or slightly mottled and in **severe cases liver may shrink** greatly.<sup>Q</sup>
- **Microscopically**
  - “**Ballooning degeneration**”:<sup>Q</sup> Diffuse swelling due to hepatocyte injury
  - “**Dropout**” of **hepatocytes**: cytoplasm looks empty & is surrounded by scavenger macrophages
  - **Lymphoplasmacytic** (mononuclear) infiltrate (also seen in chronic hepatitis)
  - **Minimal** or absent **portal-inflammation**
  - “**Spotty necrosis**”/**lobular hepatitis**:<sup>Q</sup> Scattered parenchymal injury throughout hepatic lobule
  - **Councilman body**:<sup>Q</sup> Intensely eosinophilic apoptotic hepatocytes with pyknotic nucleus
  - **Confluent necrosis in severe acute hepatitis**<sup>Q</sup>
  - **Central to portal bridging necrosis**:<sup>Q</sup> with increasing severity
  - **Parenchymal collapse**:<sup>Q</sup> In most severe cases



## Chronic Hepatitis

- **Definition:** Symptomatic, **biochemical**, or **serologic** evidence of **continuing or relapsing** hepatic disease for **more than 6 months**<sup>Q</sup>




- **Morphology**
  - **Defining histologic feature** is **mononuclear portal infiltration**<sup>Q</sup>
  - **Interface hepatitis:** Inflammatory infiltrate **at the interface** between hepatocellular **parenchyma & portal tract stroma**, along with lobular hepatitis
  - **Hallmark** of **progressive chronic liver damage** is deposition of **fibrous tissue (scarring)**<sup>Q</sup>
  - May progress to **bridging fibrosis**
  - **Cirrhosis (scarring with nodule formation)** in most severe cases
- **Histologic grading & staging:** Histologic activity index (HAI) & METAVIR score: Based on Biopsy assessment-
  - Grading is based on inflammation & Necrosis
  - Staging is based on fibrosis
  - **Type of Necrosis:**
    - Periportal necrosis, including piecemeal necrosis and/or bridging necrosis
    - Intralobular necrosis: Confluent/ Focal
- **Type of Inflammatory Activity** (grade)
  - Portal Inflammation: Mild, moderate or Severe
  - **Type of Fibrosis:** Portal, Bridging, Cirrhosis






## High Yield Facts

- Inflammatory cells in both **acute and chronic** viral hepatitis are **mainly T cells**
- In **Acute Hepatitis A**: mononuclear infiltrate is rich in **Plasma cells**
- Diagnostic hallmark of **Chronic Hepatitis B** is “**ground-glass**” **hepatocytes** (cells with **endoplasmic reticulum swollen by HBsAg**) (arrow head)



Ground Glass Hepatocytes

- In Chronic hepatitis C:
  - Lymphoid aggregates
  - Bile duct injury mimicking primary biliary cirrhosis.
  - Focal mild to moderate macrovesicular steatosis (especially in HCV genotype 3 infections)



Lymphoid Aggregates in Portal Tracts


- **Etiology** rather than the histologic pattern is the **most important determinant** of the probability of developing **progressive chronic hepatitis**.
- **Piecemeal necrosis** or **interface hepatitis**- **disruption of the limiting plate** of periportal hepatocytes by inflammatory cells<sup>Q</sup>
- **Piecemeal necrosis** is an important diagnostic criterion in Chronic active hepatitis
- **Bridging necrosis**: confluent necrosis that **bridges vascular structures**—between **portal tract** or between **portal tract and central vein**<sup>Q</sup>

## AUTOIMMUNE HEPATITIS


- **Definition**: Chronic hepatic inflammatory process manifested by **elevated serum AST**, liver-associated serum **autoantibodies** and **hypergammaglobulinemia**<sup>Q</sup>
- **Classification**: (based on serology)

Features	Type I	Type II	Type III
<b>Characteristic auto-antibodies</b>	<ul style="list-style-type: none"> <li>• Antinuclear (<b>ANA</b>),</li> <li>• Anti-smooth muscle actin (<b>SMA</b>)</li> <li>• Anti-soluble liver antigen/liver-pancreas antigen (<b>anti-SLA/LP</b>) antibodies<sup>Q</sup></li> <li>• Anti-mitochondrial (<b>AMA</b>) antibodies<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Anti-liver kidney microsome-1 (<b>anti-LKM-1</b>) (<b>directed against CYP2D6</b>)<sup>Q</sup></li> <li>• Anti-liver cytosol-1 (<b>ACL-1</b>) antibodies</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Absent ANA &amp; anti-LKM1</b> but <b>anti-SLA/LP</b> present</li> </ul>
<b>Age at presentation</b>	Middle-aged to older individuals <sup>Q</sup>	Predominantly childhood and young adulthood	More severe than type I, but is now considered a part of type I
<b>Treatment failure</b>	Infrequent	Frequent; relapse common	

- **Morphology (On Microscopy)**
  - **Parenchymal destruction followed rapidly by scarring**: Extensive interface hepatitis, (**perivenular or bridging necrosis**)
  - **Plasma cell predominance**<sup>Q</sup> in the mononuclear inflammatory infiltrates
  - Hepatocyte “**rosettes**”<sup>Q</sup> in areas of marked activity.



Auto immune hepatitis



Hepatocyte rosettes

- **Prognosis**
  - Better in adults than in children
  - **Immunosuppressive** therapy leads to **remissions in 80%** of patients.<sup>Q</sup>

## DRUG AND TOXIN INDUCED LIVER DISEASE

A diagnosis of drug or toxin-induced liver injury may be made on the basis of:

- **Temporal association** of liver damage with drug toxin exposure
- **Recovery** (usually) **upon removal** of the inciting agent
- **Exclusion** of other potential causes

### Patterns of Injury in Drug- and Toxin-Induced Hepatic Injury: (Refer Answers of this Chapter)




#### High Yield Facts

- **Alcohol** produces more toxic liver injury than any other agent
- Most common hepatotoxin causing **acute liver failure** is **acetaminophen** (centrilobular hepatic necrosis)
- Most common hepatotoxin causing **chronic liver disease** is **alcohol**

## ALCOHOLIC LIVER DISEASE


- **Epidemiology:**
  - Excessive ethanol consumption is the **leading cause of liver disease in the West**
  - Alcohol is the eighth highest risk factor for death
- **Dose dependent severity:**
  - In men, **>60–80 g/d** of alcohol for 10 years produces fatty liver<sup>Q</sup>
  - **160 g/d for 10–20 years** causes **hepatitis or cirrhosis**
  - Only **15%** of alcoholics develop **alcoholic liver disease**.<sup>Q</sup>
  - Risk of developing **HCC** is **1% to 6%** of cases annually<sup>Q</sup>
- **Risk Factors:**

- **Females are more susceptible<sup>Q</sup>** to hepatic injury than men
- **African Americans** are more prone. **ALDH\*2**, a variant of aldehyde-dehydrogenase (ALDH), found in 50% of Asians are more prone
- **Iron overload**,<sup>Q</sup> infections with **HCV and HBV**<sup>Q</sup> synergize with alcohol.
- Concurrent **Hepatitis C** infection is associated with **younger age** for severity, more **advanced histology**, **decreased survival**.<sup>Q</sup>
- **Pathogenesis: Alcohol causes-**
  - Changes in **lipid metabolism** & decreased export of lipoproteins
  - Cell injury by **reactive oxygen species** and cytokines.
- **Clinical & Laboratory features:**
  - **AST:ALT Ratio = 2:1 or 3:1**<sup>Q</sup>
  - Increased GGT (non-specific), Bilirubin, Alkaline phosphatase
- **Morphology: Liver disease begins in acinus zone 3<sup>Q</sup>** & extends outwards toward the portal tracts.
  - **Fatty liver** (Hepatocellular steatosis):
    - Macroscopically, liver is large (4-6 kg), soft, **yellow and greasy**.
    - **Fatty change is completely reversible** if there is abstinence from further intake of alcohol<sup>Q</sup>
    - **Microscopically, Microvesicular** fatty change that coalesces to **Macrovesicular change**.
  - **Alcoholic Steatohepatitis:**
    - Hepatocyte swelling and necrosis
    - **Mallory-Denk bodies** (previously called **Mallory Hyaline bodies**):<sup>Q</sup> Clumped, amorphous, eosinophilic material in ballooned hepatocytes.
    - **Neutrophilic reaction** around hepatocytes.



Mallory-Denk body (eosinophilic inclusions)

- **Morphology: Alcoholic steatofibrosis:**
  - Fibrosis begins with sclerosis of central veins
  - Perisinusoidal scar accumulates in the space of Disse spreading outward, encircling in a **chicken wire fence pattern**<sup>Q</sup>



Fatty change and fibrosis (stained blue) in a characteristic perisinusoidal chicken wire fence pattern (Masson trichrome stain)

- **Perisinusoidal scarring** leads to a classic **micronodular**<sup>Q</sup> or **Laennec cirrhosis**.

### High Yield Facts

- **Mallory-Denk bodies** are composed of Intermediate filaments-keratins 8 and 18 with ubiquitin.
- **Causes of Mallory-Denk bodies: "WAIT in PHC"**
  - W** : Wilson disease
  - A** : Alcoholic liver disease
  - I** : Indian Childhood Cirrhosis
  - T** : Alpha1 anti-Trypsin deficiency
  - P** : Primary Biliary Cirrhosis
  - HC** : Hepatocellular Carcinoma

## METABOLIC LIVER DISEASE

### Non-Alcoholic Fatty Liver Disease (NAFLD)

- **Definition:** **Hepatic steatosis** (fatty liver) in individuals **who do not consume alcohol** or do so in very small quantities (less than 20 g of ethanol/week).<sup>Q</sup>
- **Pathogenesis:**
  - Two hit model for NAFLD.
  - 1. **Insulin resistance**<sup>Q</sup> gives rise to **hepatic steatosis**.
  - 2. Hepatocellular **oxidative injury**<sup>Q</sup> resulting in liver cell necrosis and inflammatory reactions

### Morphology: (microscopy)

#### Non-alcoholic steatohepatitis (NASH)

- Pathologic **steatosis** is defined as involving **more than 5%** of hepatocytes<sup>Q</sup>
- Macrovesicular and microvesicular steatosis seen in hepatocytes
- Cirrhosis may be seen in later stages

### Difference from Alcoholic hepatitis:

#### In NASH:

- **Mononuclear cells** more prominent than neutrophils
- **Portal fibrosis**- more prominent
- Mallory-Denk bodies- less prominent
- Hepatocyte ballooning - less prominent

### High Yield Facts

- Most common cause of **chronic liver disease** in West: **NAFLD**<sup>Q</sup>
- Most common cause of **metabolic liver disease**<sup>Q</sup> is **NAFLD**<sup>Q</sup>
- Most common acquired metabolic disorder is **non-alcoholic fatty liver disease**<sup>Q</sup>
- Histologic hallmarks of **NAFLD**<sup>Q</sup> are most consistently associated with the **metabolic syndrome**
- **Cardiovascular disease** is a frequent cause of **death** in patients with NASH<sup>Q</sup>
- Level of **hedgehog pathway activity**<sup>Q</sup> correlates with stage of **fibrosis** in NAFLD
- NASH has increased risk of **hepatocellular carcinoma**.<sup>Q</sup>
- **>90%** of previously described "**cryptogenic cirrhosis**"<sup>Q</sup> (i.e., cirrhosis of unknown cause) is now thought to represent such "**burned out**" NAFLD<sup>Q</sup>

### Inherited Metabolic Diseases




- Hemochromatosis
- Wilson's disease
- $\alpha_1$ -antitrypsin deficiency

	Hemochromatosis	Wilson's disease	$\alpha_1$ -antitrypsin deficiency
<b>Genetic Defect</b>	HFE <sup>Q</sup> (C282Y) gene mutation, Chr 6p21.3 <sup>Q</sup>	ATP7B gene <sup>Q</sup> mutation on Chr 13 <sup>Q</sup>	<ul style="list-style-type: none"> <li>• Deficiency of <math>\alpha_1</math>-antitrypsin which is a "protease inhibitor" (Pi)</li> <li>• Gene on Chr 14<sup>Q</sup></li> </ul>
<b>Mode of Inheritance</b>	Autosomal Recessive	Autosomal Recessive	Autosomal Recessive
<b>Pathophysiology</b>	Excessive iron absorption	<ul style="list-style-type: none"> <li>• Decrease in copper transport into bile</li> <li>• Impaired incorporation of Cu into ceruloplasmin</li> <li>• Impaired ceruloplasmin secretion into the blood</li> </ul>	Increased protease enzymes (neutrophil elastase, cathepsin G, and proteinase 3) released from neutrophils

Contd...





	Hemochromatosis	Wilson's disease	$\alpha_1$ -antitrypsin deficiency
<b>Clinical Features</b>	<ul style="list-style-type: none"> <li>M:F=7:1</li> <li><b>Deposition of hemosiderin:</b> liver, pancreas, myocardium, pituitary gland, adrenal gland, thyroid, parathyroid, joints (<b>arthritis</b>), and skin; <b>Hypogonadism</b> in both sexes</li> <li>In advanced cases, triad of               <ol style="list-style-type: none"> <li>(1) <b>Micronodular cirrhosis</b><sup>a</sup></li> <li>(2) <b>Bronze diabetes</b><sup>a</sup></li> <li>(3) <b>Skin pigmentation</b><sup>a</sup></li> </ol> </li> <li>Pancreatic fibrosis may also be seen</li> </ul>	<ul style="list-style-type: none"> <li>Acute or chronic liver disease,</li> <li>Neuropsychiatric manifestations, frank psychosis, or a Parkinson disease–like syndrome</li> <li><b>Kayser-Fleischer rings</b><sup>a</sup>, green to brown deposits in Descemet's membrane in the limbus of the cornea</li> </ul>	<ul style="list-style-type: none"> <li><b>Pulmonary emphysema</b></li> <li>Liver disease</li> <li>Cutaneous panniculitis</li> <li>Arterial aneurysm</li> <li>Bronchiectasis</li> <li>Wegener's granulomatosis</li> </ul>
<b>Investigations</b>	Suggestive of Iron Overload: <ul style="list-style-type: none"> <li>Increased Serum Fe &amp; Ferritin</li> <li>Decreased TIBC</li> <li><b>Buccal mucosal biopsy: iron staining</b><sup>a</sup></li> <li><b>Liver Iron = 6000–18,000 mg/gm of liver</b><sup>a</sup></li> </ul>	<ul style="list-style-type: none"> <li>Decrease in serum ceruloplasmin,</li> <li><b>Most Sensitive test:</b> increase in hepatic copper (<b>&gt;200 mg/g of dry liver weight</b>)<sup>a</sup></li> <li><b>Most specific screening test:</b> Increased urinary excretion of copper (<b>&gt;100 mg/day</b>)<sup>a</sup></li> <li>Genetic testing</li> </ul>	<ul style="list-style-type: none"> <li>Liver biopsy</li> <li>Genetic testing:               <ul style="list-style-type: none"> <li>■ <b>Pi MM</b>-Wild type (normal)<sup>a</sup></li> <li>■ <b>Pi MZ</b>-heterozygous</li> <li>■ <b>PiS</b>-Moderate deficiency</li> <li>■ <b>PiZZ</b>- Null phenotype (most severe)<sup>a</sup></li> </ul> </li> </ul>
<b>Liver Biopsy Findings</b>	<ul style="list-style-type: none"> <li>Golden-yellow hemosiderin granules in periportal hepatocytes</li> <li><b>Absent inflammation</b></li> <li><b>Micronodular cirrhosis</b><sup>a</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Fatty change (steatosis)</b></li> <li>Cholestasis</li> <li><b>Acute hepatitis</b></li> <li><b>Chronic hepatitis</b></li> <li>Macrovesicular steatosis</li> <li>Mallory bodies</li> <li><b>Cirrhosis</b></li> <li><b>Massive liver necrosis</b> is rare</li> </ul>	<ul style="list-style-type: none"> <li><b>Hallmark: PAS +ve diastase-resistant round-to-oval cytoplasmic globular inclusions in hepatocytes</b><sup>a</sup></li> <li>Neonatal hepatitis</li> <li><b>Cirrhosis</b></li> <li><b>Fibrosis of portal tract</b></li> <li>Fatty change</li> <li>Mallory bodies</li> </ul>
<b>Stains used</b>	<ul style="list-style-type: none"> <li>Prussian Blue stain</li> </ul>	<ul style="list-style-type: none"> <li><b>Rhodamine stain</b> for Cu</li> <li><b>Orcein stain</b> for copper-associated protein</li> </ul>	<ul style="list-style-type: none"> <li>PAS(+) ve, diastase resistant globules</li> </ul>
<b>Images</b>	 <p>Prussian blue stain</p>	 <p>Rhodamine stain</p>	 <p>PAS +ve globules</p>

## CHOLESTATIC DISEASES

### Jaundice

Yellow discoloration of the sclera, skin, and mucous membranes indicating Hyperbilirubinemia.


Jaundice occurs when there is any one or more of following:

- Bilirubin overproduction
- Hepatitis
- Obstruction to the flow of bile





## Bilirubin Metabolism



### Causes of Jaundice

#### 1. Predominantly Unconjugated Hyperbilirubinemia

##### Excess production of bilirubin

- Hemolytic anemias
- Ineffective erythropoiesis (e.g., pernicious anemia, thalassemia)

##### Reduced hepatic uptake

- Drug interference with membrane carrier systems
- Gilbert syndrome

##### Impaired bilirubin conjugation

- Physiologic jaundice of the newborn (decreased UGT1A1 activity, decreased excretion)
- Breast milk jaundice ( $\beta$ -glucuronidases in milk)
- Crigler-Najjar syndrome types I and II: Genetic deficiency of UGT1A1 activity
- Gilbert syndrome
- Diffuse hepatocellular disease (e.g., viral or drug-induced hepatitis, cirrhosis)

#### 2. Predominantly Conjugated Hyperbilirubinemia

- Defective secretion
  - Rotor syndrome
  - Dubin-Johnson syndrome
- Cholestasis
  - Intrahepatic bile duct obstruction
  - Extrahepatic bile duct obstruction.



Dark stained liver in Dubin Johnson syndrome

### Hereditary Hyperbilirubinemias


Disorder	Inheritance	Defects
<b>Unconjugated Hyperbilirubinemia</b>		
<i>Crigler-Najjar syndrome type I</i>	AR <sup>a</sup>	Absent UGT1A1 activity <sup>a</sup>
<i>Crigler-Najjar syndrome type II</i>	AD <sup>a</sup> with variable penetrance	Decreased UGT1A1 activity <sup>a</sup>
<i>Gilbert syndrome</i>	Promoter mutation: AR Missense mutations: AD	Decreased UGT1A1 activity <sup>a</sup>

Contd...



Conjugated Hyperbilirubinemia		
<b>Dubin-Johnson syndrome</b>	AR; mutation in canalicular multidrug resistance protein 2 ( <b>MRP2</b> ) <sup>a</sup>	Impaired bilirubin glucuronide excretion
<b>Rotor syndrome</b>	AR	Decreased hepatic uptake, storage & biliary excretion
<b>Progressive Familial Intrahepatic Cholestasis (PFIC)</b>	AR	<b>PFIC1- ATP8B1</b> <b>PFIC2- ABCB11</b> <b>PFIC3- ABCB4</b>

## Neonatal Cholestasis




## AUTOIMMUNE CHOLANGIOPATHIES

### Two Types

- Primary biliary cirrhosis
- Primary sclerosing cholangitis

### Intrahepatic Biliary Disease Tract

Features	Primary Biliary Cirrhosis	Secondary Biliary Cirrhosis	Primary Sclerosing Cholangitis
<b>Involvement</b>	Intrahepatic <sup>a</sup> biliary tree	Extrahepatic <sup>a</sup> biliary tree	Both extrahepatic <sup>a</sup> & intrahepatic <sup>a</sup> biliary tree
<b>Etiology</b>	<ul style="list-style-type: none"> <li>Sjögren syndrome<sup>a</sup></li> <li>Scleroderma</li> <li>Thyroid disease</li> </ul>	<ul style="list-style-type: none"> <li>Biliary Atresia</li> <li>Gallstones</li> <li>Stricture</li> <li>Carcinoma of pancreatic head</li> </ul>	<ul style="list-style-type: none"> <li>Inflammatory bowel disease<sup>a</sup></li> <li>Retroperitoneal fibrosis</li> </ul>
<b>Sex predilection</b>	F:M = 6 : 1	None	F:M = 1 : 2 <sup>a</sup>
<b>Clinical features</b>	Pruritus, jaundice, malaise, dark urine, light stools, hepatosplenomegaly		
<b>Lab findings</b>	Conjugated hyperbilirubinemia, increased Alkaline Phosphatase, bile acids, cholesterol		
<b>Autoantibodies</b>	95% AMA-positive <sup>a</sup>	None; as it is not an autoimmune disease	65% Atypical p-ANCA positive <sup>a</sup>
<b>Important pathologic findings</b>	Florid duct lesions (lympho-plasmacytic inflammation & loss of small ducts only)	Prominent bile stasis <sup>a</sup> in bile ducts, bile ductular proliferation with surrounding neutrophils, portal tract edema	Inflammatory destruction; fibrotic obliteration of bile ducts ("onion-skin" fibrosis) <sup>a</sup>





## NODULES AND TUMORS

### Nodular Hyperplasias

Occurs due to focal or diffuse **alterations in hepatic blood supply**

#### Two Types

Features	Focal Nodular Hyperplasia	Nodular Regenerative Hyperplasia
<b>Peak age</b>	30–40 yrs	50–70 yrs
<b>M:F ratio</b>	1:10 <sup>Q</sup>	1:1
<b>Presentation</b>	Asymptomatic <sup>Q</sup>	Portal hypertension
<b>Associated conditions</b>	OCP use <sup>Q</sup> (66%–95%), cavernous hemangioma	Connective tissue disease, CMPN, drugs/toxins, HIV, Post-transplant, vascular disorders, Rheumatoid arthritis
<b>Nodules</b>	Well-demarcated but poorly encapsulated <sup>Q</sup>	Numerous <b>micronodular</b>
<b>Cut surface often</b>	Central gray-white, depressed stellate scar	Tan-white, rarely hemorrhagic
<b>Septa/scar</b>	Usually present	Absent



#### High Yield Facts

- Peliosis hepatitis is associated with **anabolic steroids, Danazol, OCPs and tamoxifen**
- Von Meyenburg Complexes are “bile duct hamartomas,” mimic metastases to the liver
- **Primary Sclerosing Cholangitis is an autoimmune disease that predominantly occurs in males**
- **Primary Sclerosing Cholangitis** is associated with inflammatory bowel disease in 70% cases.
- Prevalence of PSC in persons with ulcerative colitis is about 4%
- **AMA**-Anti-mitochondrial Ab directed against **E2 component of the pyruvate dehydrogenase<sup>Q</sup> complex (PDC-E2)**
- **Atypical p-ANCA (perinuclear staining pattern)<sup>Q</sup>** seen in **Primary Sclerosing Cholangitis**, is directed **against a nuclear envelope protein<sup>Q</sup>**, instead of myeloperoxidase).

### Benign Tumors

- **Most common benign liver tumor is Cavernous hemangioma.<sup>Q</sup>**

#### Hepatocellular Adenomas

- Benign neoplasms developing from hepatocytes.
- **Associated with: Oral contraceptives<sup>Q</sup> (estrogen rich) and anabolic steroids.**
- **Morphology of Hepatic adenoma: Cords of hepatocytes, with an arterial vascular supply without portal tracts.<sup>Q</sup>**

R<sup>9th</sup>

Latest Update

#### Classification of adenomas

- **HNF1- $\alpha$  inactivated adenomas**
  - No risk of malignant transformation
  - Often associated with OCP use
  - In individuals with MODY-3
- **$\beta$ -Catenin activated adenoma:** Mutation in the  $\beta$ -catenin gene
  - Diagnostic hallmark: Nuclear translocation for  $\beta$ -catenin
  - Leading to marked atypia
  - Very high risk for malignant transformation
- **Inflammatory adenomas:**
  - Associated with non-alcoholic fatty liver disease
  - Characterized by activating mutations in gp130, a co-receptor for IL-6 leading to JAK-STAT signaling and overexpression of acute phase reactants (C-reactive protein & serum amyloid-A)
  - 10% have  $\beta$ -catenin activating mutations
  - Risk for malignant transformation is intermediate

### Malignant Tumors

#### Hepatoblastoma

- **Most common liver tumor of early childhood.<sup>Q</sup>**
- Rarely occurs beyond the age of 3 years.
- Characterized by: Activation of the **WNT signaling pathway<sup>Q</sup>** (APC gene)
- Associated with **Beckwith-Wiedemann syndrome<sup>Q</sup>**

#### Hepatocellular Carcinoma (HCC)

- **Epidemiology:**
  - Peak incidence age: 20–40 yrs; M:F = 4–8:1
- Factors associated with increased risk of HCC
  - **Common**
    - **Cirrhosis (M.C)<sup>Q</sup>**
    - **Hepatitis B >>C** chronic infection
    - Chronic Alcoholism
    - **NASH/NAFLD**
    - Aflatoxin B<sub>1</sub> or other mycotoxins
  - **Less common causes**
    - Primary biliary cirrhosis **Hemochromatosis**,  $\alpha$ 1 Antitrypsin deficiency
    - Glycogen storage diseases
    - Citrullinemia, Porphyria cutanea tarda, **Hereditary tyrosinemia**,
    - **Wilson's disease**

R<sup>9th</sup>

Latest Update

#### Genetic Factors:

- **Activation of  $\beta$ -catenin (40% cases) & inactivation of p53 (60% cases)** are the 2 most common early mutational events
- Activation of **IL-6/JAK STAT pathway** suppress hepatocyte differentiation & promote their proliferation by regulating function of transcription factor HNF4- $\alpha$






- **Gross Morphology:**
  - Unifocal (usually large) mass or Multifocal, Diffusely infiltrative
- **Light Microscopy**
  - Pattern of growth can be **trabecular, solid, or tubular with Mallory hyaline bodies**
- **Ultrastructural findings:**
  - Malignant hepatocyte with **numerous mitochondria, microbodies** & abundant **glycogen**.<sup>Q</sup> Cells also contain intracytoplasmic bile products
- **Immunohistochemistry**
  - **HepPar-1<sup>Q</sup>** is a **monoclonal antibody** that reacts to a cytoplasmic marker of normal & neoplastic hepatocytes
- **Glypican-3<sup>Q</sup>** - stains most **hepatocellular carcinomas** (especially those associated with cirrhosis) and **high-grade dysplastic nodules**, but not normal liver (**most specific**)
- **Serum markers**
  - **p-CEA (canalicular staining): 100% specificity;**<sup>Q</sup> often **negative in Poorly Differentiated-HCC**
  - **AFP:** 90-95% specificity, Arginase-1, TTF-1, **PIVKA-2<sup>Q</sup>, DCP (Des CarboxyProthrombin)<sup>Q</sup>**
- **Metastasis from HCC:**
  - **Intrahepatic metastases**, by either vascular invasion or direct extension
  - **Hematogenous metastases** (extrahepatic): occurs to **lung** & portal vein and **right side of heart** via IVC

R10<sup>th</sup>


**Latest Update**

#### Precursor Lesions of Hepatocellular Carcinoma (HCC)

Characteristics	Hepatocellular Adenoma	Small cell change	Large cell change	Low-grade Dysplastic Nodule	High-Grade Dysplastic Nodule
<b>Focality in liver</b>	Single or multiple	Diffuse	Diffuse	Single or multiple	Single or multiple
<b>Premalignant</b>	Yes	Yes	In Some HBV	Uncertain	Yes
<b>Association with cirrhosis</b>	Rare	Common	Common	Usual	Usual




Gross showing gray white growth in liver S/o HCC



Trabecular pattern in HCC

#### Fibrolamellar Variant of HCC



Nests and cords of malignant-appearing, oncocytic hepatocytes separated by dense bundles of collagen

#### Hepatocellular Carcinoma vs Fibrolamellar Variant of Hepatocellular Carcinoma

	Hepatocellular Carcinoma	Fibrolamellar variant of HCC
<b>Age</b>	70 years	20-30 years
<b>Male : Female</b>	3-4:1	1 : 1
<b>Tumour marker</b>	Alphafetoprotein (AFP) very high	AFP Normal, but <b>Neurotensin</b> elevated
<b>Prognosis</b>	Poor	Good





## Cholangiocarcinoma (CCA)

- **Origin:** Biliary tree, arising from bile ducts within and outside liver.
- **Risk Factors**
  - Chronic inflammation, cholestasis, hepatolithiasis & fibropolycystic liver disease and primary sclerosing cholangitis
  - Liver flukes (Opisthorchis and Clonorchis species)
  - Hepatitis B and C

R<sup>9th</sup>

### Latest Update

#### Precursor Lesions of Cholangiocarcinoma

Characteristics	Biliary intraepithelial neoplasia, high grade (BilIN-3)	Mucinous cystic Neoplasm	Intraductal papillary Biliary Neoplasia
<b>Focality in liver</b>	Diffuse or multifocal	Single	Focal or diffuse
<b>Commonly associated diseases</b>	PSC, Hepatolithiasis, Liver flukes	None	None
<b>Association with cirrhosis</b>	Sometimes	No	No

- **Site**
  - **Extrahepatic forms:**
    - 50-60 % **perihilar (Klatskin tumor)**: At the junction of the right and left hepatic ducts.
    - 20% to 30% are distal and posterior to the duodenum.
  - Intrahepatic-10%.
- **Clinical Features:**
  - **Intrahepatic:** Cholestasis, symptomatic liver mass
  - **Extrahepatic** (hilar & distal): symptoms of biliary obstruction, cholangitis, right upper quadrant pain.
- **Morphology**
  - **Gross:** Small to massive lesions
  - **Microscopically:** **Adenocarcinoma** with marked **desmoplasia**
- **Prognosis:** Poor

### High Yield Facts

- Metastasis to Liver are the most common malignant tumors in liver (more common than primary).
- Common primary sites from where Metastasis occurs to Liver are colon, breast, lung, and pancreas.
- Biliary intraepithelial neoplasias (BilIN-1, -2, or -3: low to high grade,) are premalignant for **cholangiocarcinoma**
- **BilIN-3**, the highest grade lesion, incurs the **highest risk** of malignant transformation
- **Cholangiocarcinoma** is the **second** most common **primary** malignant tumor of the liver after HCC

R<sup>9th</sup>

### Latest Update

#### Other Primary Hepatic Malignant Tumors

Tumors	Characteristics
<b>Combined hepatocellular &amp; cholangiocarcinoma</b>	Originates from a multipotent stem cell.
<b>Mucinous cystic neoplasms &amp; Intraductal papillary biliary neoplasia</b>	May occur as in situ lesions or as invasive cholangiocarcinoma.
<b>Angiosarcoma of liver</b>	Associated with <b>vinyl chloride, arsenic, or Thorotrast exposure</b> , poor prognosis
Epithelioid hemangioendothelioma	Endothelial malignancy, better prognosis
<b>Hepatic lymphomas</b>	Most are <b>diffuse large B-cell lymphomas</b> > MALT lymphomas. Seen in middle aged men Associated with Hep B, Hep C, HIV, and PBC.
<b>Hepatosplenic delta-gamma T cell lymphoma</b>	Most common in young adult males, has a predilection for <b>hepatic &amp; splenic sinusoids</b> as well as the marrow.



## GALLBLADDER (GB)

### Acute Cholecystitis

- **Definition:** Acute inflammation of GB
- **Etiology:** 90% cases caused by obstruction (**stone**); 10% **Acalculous**

### Acalculous Cholecystitis

- **Definition:** Inflammation of GB resulting from ischemia (cystic artery is an end artery)
- **Pathophysiology:** **Inflammation and edema** of the GB wall, GB **stasis**, accumulation of biliary sludge & mucus, causing **cystic duct obstruction** in the **absence of stone**.
- **Risk factors**
  - **Sepsis** with hypotension and multisystem organ failure;
  - **Immunosuppression**
  - Major trauma and **burns**
  - **Diabetes mellitus**
  - Infections: **Salmonella, Staphylococci, E.coli**



## Chronic Cholecystitis

- **Definition:** Chronic inflammation of GB which may be a **sequel** to repeated bouts of acute cholecystitis, or in absence of antecedent attacks.
- **Etiology:** Associated with **cholelithiasis in more than 90% of cases**<sup>Q</sup>
- **Morphology:**
  - **Rokitansky-Aschoff sinuses**<sup>Q</sup>: Outpouchings of the mucosal epithelium through the wall
  - **Porcelain gallbladder**<sup>Q</sup>: **Dystrophic calcification** within GB wall, Increased GB cancer.<sup>Q</sup>
  - **Xanthogranulomatous cholecystitis**<sup>\*</sup>
    - Massively thickened, shrunken, nodular wall, chronically inflamed with foci of necrosis & hemorrhage
    - **Triggered by rupture of Rokitansky-Aschoff sinuses** into wall of GB, followed by accumulation of **lipid laden foamy macrophages (xanthoma cells)**<sup>Q</sup>
- **Hydrops of GB:** Atrophic, chronically obstructed gallbladder containing clear secretions

## Adenomyomatosis GB

Benign proliferation of gallbladder surface epithelium with gland-like formation, extramural sinuses, transverse strictures, and/or fundal nodule ("adenoma" or "adenomyoma") formation.



### High Yield Facts

- In the West, 90% of gallstones are cholesterol stones, rest are pigment stones composed of bilirubin calcium salts
- **Carcinoma gallbladder is the most common malignancy of the extrahepatic biliary tract.**<sup>Q</sup>

## Carcinoma Gallbladder

- **Epidemiology**
  - Occurs most frequently in the **seventh decade** of life
  - It is slightly **more common in women** (M:F = 1:2)
- **Risk factors:** Gallstones (cholelithiasis)<sup>Q</sup>
- **Genetics:** ERBB2 (Her-2/neu) mutation<sup>Q</sup>
- **Gross Morphology**
  - Most common sites of involvement are the fundus and the neck;
  - Only 20% involve the lateral walls.
  - **Infiltrating (more common)**<sup>Q</sup> or **exophytic**.
- **Microscopy:** Adenocarcinoma<sup>Q</sup> > Squamous cell Ca > Carcinosarcoma
- **Prognosis:** Papillary tumors have a **better prognosis**<sup>Q</sup>.

## PANCREAS

### CONGENITAL ANOMALIES

- **Pancreas Divisum**
  - **Most common**<sup>Q</sup> congenital anomaly of the pancreas
  - **Failure of fusion** of the fetal duct systems of the dorsal and ventral pancreatic primordia
  - Predisposes to **chronic pancreatitis**<sup>Q</sup>
- **Annular Pancreas:** **Band-like ring** of normal pancreatic tissue that encircles the 2nd part of duodenum
- **Ectopic Pancreas:** In sites like **stomach & duodenum**<sup>Q</sup>, followed by the jejunum, Meckel diverticula & ileum
- **Agenesis of Pancreas:** Due to mutation in **PDX1 gene**, encoding a homeobox transcription factor that is critical for pancreatic development



### High Yield Facts

- **Pancreas is a retroperitoneal organ** extending from the C-loop of the duodenum to the hilum of the spleen.
- Pancreas develops from dorsal & ventral pancreatic primordial buds.
- Pancreas consists of an:
  - Exocrine part (80% to 85%) that secrete enzymes, stored in granules as proenzymes (zymogens)<sup>Q</sup>
  - Endocrine part: that secretes important hormones
- Somatostatin is also secreted by extra-islet neuroendocrine cells

## ACUTE PANCREATITIS

- **Definition:** **Reversible** pancreatic parenchymal injury associated with inflammation.
- **Etiology:**
  - **Excessive alcohol intake (most common)**<sup>Q</sup>
  - Pancreatic duct obstruction (e.g. **gallstones**)
  - Traumatic injuries
  - **Drugs:** Azathioprine, L-Asparaginase, Furosemide, estrogens, Dideoxyinosine
  - Infections e.g. **mumps**<sup>Q</sup>



- Metabolic disorders leading to **hypercalcemia**<sup>Q</sup>
- Ischemia
- Hereditary factors
- Genetic factors:**

Gene (chr)	Protein product	Function / Characteristics
<b>CFTR (7q31)</b>	Cystic fibrosis transmembrane conductance regulator	<b>Loss-of-function mutations</b> limit HCO <sub>3</sub> secretion → inspissation of secreted fluids & <b>duct obstruction</b>
<b>PRSS1 (17q34)</b>	Serine protease 1 (trypsinogen 1)	Cationic trypsin, <b>Gain-of – function mutations prevent self- inactivation of trypsin</b>
<b>SPINK1 (5q32)</b>	Serine protease inhibitor, kazal type 1	Inhibitor of trypsin, <b>Mutations cause loss-of-function, increasing trypsin activity</b>

- Morphology:**
  - Inappropriate activation** of digestive **enzymes within** the substance of the **pancreas** leads to:
    - Microvascular leak and edema, Fat necrosis, Acute inflammation, Destruction of pancreatic parenchyma, Destruction of blood vessels and interstitial hemorrhage

## PANCREATIC CYSTIC NEOPLASMS

- Serous cystic neoplasms: (also known as serous cystadenomas):** **VML gene:** always Benign
- Mucinous Cystic Neoplasms:** 1/3<sup>rd</sup> associated with **invasive adenocarcinoma**<sup>Q</sup>
- Intraductal papillary mucinous neoplasms (IPMNs):** Can progress to an **invasive cancer**.<sup>Q</sup>
- Solid-pseudopapillary neoplasms:** **Wnt signaling pathway (β-catenin)**<sup>Q</sup> oncogene

## PANCREATIC CARCINOMA


- Precursor lesions:** “Pancreatic intraepithelial neoplasias” (PanINs)
- Molecular Alterations:**

Gene	Chr	%	
<b>KRAS</b>	12p	90	<b>Most commonly involved Oncogene</b> <sup>Q</sup>
<b>p16/CDKN2A</b>	9p	95	<b>Most commonly involved Tumor suppressor gene</b> <sup>Q</sup>
<b>TP53</b>	17p	50-70	<b>Involved in Response to DNA damage</b>
<b>SMAD4</b>	18q	55	<b>TGF β pathway</b>
<b>BRCA2</b>	13q	10	<b>Germ-line mutation</b> <sup>Q</sup>

- Etiology:**
  - Cigarette smoking (strongest environmental influence)**<sup>Q</sup>, Fat rich diet
  - Chronic pancreatitis** & diabetes mellitus (new-onset diabetes mellitus in an older patient may be the first sign)
- Inherited Predisposing conditions:**

Disorder	Gene (Chromosome)
Hereditary breast and ovarian cancer	<b>BRCA2</b> (13q)
Familial atypical multiple-mole melanoma syndrome	<b>p16/CDKN2A</b> (9p)
Hereditary pancreatitis	<b>PRSS1</b> (7q) and <b>SPINK1</b>
Peutz-Jeghers syndrome (maximum risk of Pancreatic Ca)	<b>LKB1</b> (19p)
Hereditary Non polyposis Colorectal Cancer (HNPCC)	<b>MLH1, MSH2</b> (2p)

- Site: Pancreatic Head (most common)**<sup>Q</sup> > Body > Tail
- Clinical features:**
  - 50% **Pancreatic Head** Carcinoma develop **jaundice**
- Important Characteristics**
  - Mostly adenocarcinoma**<sup>Q</sup>, **Highly invasive**



Malignant glands infiltrating into stroma

- Elicits “**desmoplastic response**” (dense fibrosis).
- Migratory thrombophlebitis (Trousseau sign)**<sup>Q</sup>, 10%- due to release of **Platelet activating factor** and procoagulants from carcinoma or its necrotic products.



R10<sup>th</sup>

**Latest Update**

### Molecular serum markers of liver fibrosis

Marker	Function
<b>Liver function</b>	
ALT	Metabolic enzymes in the liver
AST	Metabolic enzymes in the liver
<b>ECM formation</b>	
PHINP	Propeptide of collagen type III
PINP	Propeptide of collagen type I
Type IV collagen	Basement membrane formation
P4NP 7S	N-terminal pro-peptides of type IV collagen 7S domain
PVCP	Propeptide of collagen type V
HA	Component of ECM
YKL-40	Glycoprotein involved in ECM turnover
MFAP	Glycoprotein involved in ECM turnover
<b>Fibrinolytic process</b>	
<b>Neo-epitope</b>	
MMP-1/MMP-13	Degrade fibrotic matrix
MMP-2	Degrades basal membranes and fibrotic matrix
MMP-9	Degrades basal membranes
TIMP-1	Inhibits MMP-1 activity

Marker	Function
<b>ECM degradation</b>	
CO3-610	Collagen type III fragment generated by MMP-9
CO6-MMP	Collagen type VI fragment generated by MMP-2,9
CO1-764	Collagen type I fragment generated by MMP-2,9,13
C4M	Collagen type IV fragment generated by MMP-9
<b>Cytokines</b>	
TGF-β	Growth factor stimulates production of ECM by HSC
CTGF	Potent pro-fibrogenic factor
PDGF	Growth factor stimulates proliferation of HSC
TNF-α	Inflammatory cytokine involved in fibrogenesis
IL-4, 6, 8, 18	Inflammatory cytokine involved in fibrogenesis

R10<sup>th</sup>

**Latest Update**

### Laboratory Evaluation of Liver Disease

Test Category	Blood Measurement*
Hepatocyte integrity	Cytosolic hepatocellular enzymes 1. Serum aspartate aminotransferase (AST) 2. Serum alanine aminotransferase (ALT) 3. Serum lactate dehydrogenase (LDH)
Biliary excretory function	Substances normally secreted in bile <sup>+</sup> • Serum bilirubin • Urine bilirubin • Serum bile acids • Plasma membrane enzymes (from damage to bile canaliculus) Serum alkaline phosphatase Serum, γ-glutamyl transpeptidase (GGT)
Hepatocyte function	Proteins secreted into the blood • Serum albumin • Prothrombin time (PT) • Partial thromboplastin time (PTT) Hepatocyte metabolism • Serum ammonia • Aminopyrine breath test (hepatic demethylation)

R10<sup>th</sup>

**Latest Update**

### World Health Organization Criteria for the Metabolic Syndrome

One of	Diabetes mellitus or Impaired glucose tolerance or Impaired fasting glucose or Insulin resistance
and two of:	<ul style="list-style-type: none"> <li><b>Blood pressure:</b> ≥ 140/90 mm Hg</li> <li><b>Dyslipidemia:</b> Triglycerides (TG): ≥1.695 mmol/L and high-density lipoprotein cholesterol (HDL-C) ≤ 0.9 mmol/L (male), ≤1 mmol/L (female)</li> <li><b>Central obesity:</b> waist-hip ratio &gt; 0.90 (male); &gt; 0.85 (female), or body mass index &gt; 30 kg/m<sup>2</sup></li> <li><b>Microalbuminuria:</b> urinary albumin excretion rate of ≥ 20 µg/min of albumin-to-creatinine ≥30 mg/gm</li> </ul>






## NEXT Pattern Questions



Q's

1. A 44-year-old patient presented with jaundice and had needle prick injury 2 years back and liver biopsy shown below. Based on the histological features what classify type of hepatitis?



- a. Hepatitis B virus induced hepatitis
- b. Hepatitis C virus induced hepatitis
- c. Hepatitis A virus induced hepatitis
- d. Hepatitis E virus induced hepatitis


**Ans. (a) Hepatitis B virus induced hepatitis**

- With a history of Jaundice for last 2 years and the liver biopsy showing homogeneous pink color suggestive of ground glass appearance, this is most likely to be the case of Hepatitis B infection.



Q's


2. A 54-year-old patient presented with jaundice (raised direct bilirubin) and presence of pANCA antibody. Liver biopsy shows the following. Which of the following statement are true regarding the condition?



- a. Most common antibody seen is pANCA
- b. It is associated with ulcerative colitis
- c. On histology it shows circumferential onion skin fibrosis around duct
- d. All of the above


**Ans. (d) All of the above**

- Obstructive jaundice (raised direct bilirubin) and presence of pANCA antibody is suggestive of primary sclerosing cholangitis. 70% cases are often seen with ulcerative colitis. Biopsy of liver will show circumferential onion skinning appearance.




## Image-Based Questions

1. Liver biopsy from a patient shows dark brownish-black deposits in hepatocytes, canaliculi, Kupffer cells, and ductules. Polarised microscopy of the same shows maltese cross picture of red birefringence in the larger deposits. What is your diagnosis?




- a. Melanosis
- b. Cholestasis
- c. Erythropoietic protoporphyria
- d. Hepatitis B

3. Identify the areas Labeled 1, 2 and 3?




- a. Periportal, Midzonal, Centrilobular
- b. Periportal, Centrilobular, Midzonal
- c. Midzonal, Centrilobular, Midzonal
- d. Periportal, Centrilobular, Midzonal

2. A 61-year-old man had leg swelling with grade 2 pitting edema upto the knees, prominent jugular venous distention to the level of the mandible & increasing levels of serum AST and ALT. A diagnosis of congestive heart failure was made. The gross appearance of the liver has been shown in the figure. Identify the condition?




- a. Portal vein thrombosis
- b. Cirrhosis
- c. Fatty liver
- d. Chronic venous congestion

4. 45/M chronic alcoholic presented with pain abdomen, USG suggested fatty liver. Liver biopsy done has been shown below. What is your interpretation and likely diagnosis?




- a. Lymphocytic infiltrate, Hep C
- b. Neutrophilic Infiltrate, Hep B
- c. Macrovesicular steatosis, Alcoholic liver disease
- d. Squamous pearls, Metastasis






5. A 6-week-old infant presented with conjugated Hyperbilirubinemia. HIDA scan done was not suggestive of Extrahepatic biliary atresia. Liver biopsy showed the following finding. Identify the underlying Etiology?




- Wilson's disease
- Cystic fibrosis
- $\alpha$ 1 antitrypsin deficiency
- Idiopathic neonatal hepatitis

7. 25/M presented with jaundice, clay coloured stool and pruritus. Bilirubin was 7 gm%, Direct Bilirubin 5gm%, ALP 500 IU/L. Biopsy from biliary tract revealed the following. What is your diagnosis?



- PSC
- SBC
- PBC
- Bile duct stones

6. 45/M, a chronic alcoholic presented to Med OPD of AIIMS with Jaundice. S. Bilirubin was 4.5 mg% with direct Bilirubin being 3mg%. Liver biopsy was done which suggested the following. What is your interpretation and likely diagnosis?




- Malory Hyaline bodies, HCC
- Malory Hyaline bodies, Chronic Hepatitis B
- Ground glass appearance, HCC
- Ground Glass appearance, Chronic Hepatitis B

8. The condition associated with their formation is:



- Hypomotility of gall bladder
- Accelerated cholesterol crystal nucleation
- Hypersecretion of mucus
- All of the above



## Answers of Image-Based Questions

**1. Ans. (c) Erythropoietic protoporphyria**

- Liver biopsy showing dark brownish-black deposits in hepatocytes, canaliculi, Kupffer cells, and ductules with polarised microscopy of the same showing maltese cross picture of red birefringence in the larger deposits, a characteristic of Erythropoietic protoporphyria.

**2. Ans. (d) Chronic venous congestion**

- The cut surface of the liver has a variegated mottled red appearance, representing congestion and hemorrhage in the centrilobular regions of the parenchyma. Also note that on microscopic examination, the centrilobular region is suffused with red blood cells and atrophied hepatocytes are not easily seen. Portal tracts and the periportal parenchyma are intact.

**3. Ans. (a) Periportal, Midzonal, Centrilobular**

- Liver is divided histologically into lobules. The center of the lobule is the central vein. At the periphery of the lobule are portal triads. Functionally, the liver can be divided into three zones, based upon oxygen supply. Zone 1 encircles the portal tracts where the oxygenated blood from hepatic arteries enters. Zone 3 is located around central veins, where oxygenation is poor. Zone 2 is located in between.

**4. Ans. (c) Macrovesicular steatosis, Alcoholic liver disease**

- Liver with mixed small and large fat droplets (white round structures, steatosis) a feature seen in alcoholic liver disease.

**5. Ans. (c)  $\alpha$ 1 antitrypsin deficiency**

- The history given is that of neonatal hepatitis. Liver biopsy here shows Periodic acid–Schiff (PAS) stain after diastase digestion of the liver, characteristic magenta cytoplasmic granules seen in hepatitis due to  $\alpha$ 1 antitrypsin deficiency.

**6. Ans. (d) Ground Glass appearance, Chronic Hepatitis B**

- The microscopic section of liver hepatocytes here show large pale, finely granular pink cytoplasmic inclusions on hematoxylin and eosin staining.

**7. Ans. (a) PSC**

- Clinical feature of jaundice, clay coloured stool and pruritus is suggestive of obstructive jaundice. Biopsy showing fibrotic obliteration of bile ducts ("onionskin" fibrosis) seen in primary biliary cirrhosis.

**8. Ans. (d) All of the above**

- The figure shows gall bladder filled with gall stones. Causes of its formation can include
  - a. Hypomotility of gall bladder
  - b. Accelerated cholesterol crystal nucleation
  - c. Hypersecretion of mucus.






## Multiple Choice Questions

### STRUCTURE OF LIVER

1. **In Obstructive jaundice, which of the following enzyme is elevated?** (Recent Question 2016-17)
  - a. GGT
  - b. AST
  - c. ALT
  - d. LDH
2. **The given figure shows which of the following?** (AIIMS Nov 2015)



- a. Amyloidosis: grey areas are viable; white areas are necrotic
  - b. Nutmeg liver: red areas are viable pericentral areas; white areas are periportal necrotic areas
  - c. Nutmeg liver: Red areas are pericentral necrotic areas, white areas are viable fibrotic periportal area
  - d. Amyloidosis - necrotic white periportal viable grey pericentral areas
3. **Absent urobilinogen in urine with icterus indicates?** (AIIMS Nov 2015)
  - a. Perihepatic obstruction
  - b. Hemolysis
  - c. Hepatitis
  - d. Liver failure
4. **Massive hepatocellular necrosis is seen with** (Recent Question 2015)
  - a. Tetracycline
  - b. Macrolides
  - c. Methyldopa
  - d. Acetaminophen
5. **Liver damage in shock** (Recent Question 2015)
  - a. Centrilobular necrosis
  - b. Diffuse necrosis
  - c. Periportal necrosis
  - d. Spotty necrosis
6. **Nutmeg liver is seen in** (Recent Question 2015)
  - a. Chronic venous congestion of liver
  - b. Portal vein obstruction
  - c. Hepatic artery obstruction
  - d. Non-alcoholic hepatic steatosis
7. **False statement about alagille syndrome** (Recent Question 2015)
  - a. Complete absence of bile ducts
  - b. Normal liver
  - c. Mutations in Jagged 1 gene
  - d. No Risk of hepatocellular carcinoma
8. **Most common hepatotoxin causing acute liver injury** (Recent Question 2015)
  - a. Acetaminophen
  - b. Alcohol
  - c. Paracetamol
  - d. Halothane
9. **Fibrin ring granuloma in liver is caused by** (Recent Question 2015)
  - a. Sulphonamides
  - b. Amlodarone
  - c. Isoniazid
  - d. Allopurinol

10. **Peliosis hepatis is caused by** (Recent Question 2015)
  - a. Contraceptives
  - b. Anabolic steroids
  - c. Erythromycin
  - d. Ezetemibe
11. **Steatohepatitis with mallory-Denk bodies is caused by** (Recent Question 2015)
  - a. Alcohol
  - b. Enalapril
  - c. Vitamin A
  - d. Methotrexate
12. **Periportal fibrosis is caused by?** (Recent Question 2015)
  - a. Alcohol
  - b. Methotrexate
  - c. Rifampicin
  - d. OCPs
13. **The following is not a feature of non-cirrhotic portal fibrosis** (Recent Question 2015)
  - a. Intimal fibroelastosis
  - b. Lymphocytic infiltration
  - c. Portal fibrosis
  - d. Bridging fibrosis
14. **Perivenular fibrosis is caused by** (Recent Question 2015)
  - a. Methotrexate
  - b. Alcohol
  - c. OCPs
  - d. Amiodarone
15. **In cirrhosis, the proliferation and activation of the following cell results in fibrosis** (Recent Question 2015) (WB PG 2016)
  - a. Hepatocytes
  - b. Stellate cells
  - c. Kupffer cells
  - d. Bile duct epithelium
16. **Centrilobular necrosis of liver may be seen with-** (Recent Question 2014)
  - a. Phosphorus
  - b. Arsenic
  - c.  $\text{CCl}_4$
  - d. Ethanol
17. **Mallory bodies contain -** (Recent Question 2013)
  - a. Vimentin
  - b. Cytokeratin
  - c. Keratin
  - d. Collagen
18. **Which substance is/are not deposited in hepatocyte?** (PGI May 10, June 01)
  - a. Lipofuscin
  - b. Pseudomelanin
  - c. Bile pigment
  - d. Iron
  - e. Melanin
19. **On stopping Alcohol, all the following changes are reversible EXCEPT -** (DNB Dec 10)
  - a. Hepatitis
  - b. Cirrhosis
  - c. Microvesicular fatty change
  - d. Macrovesicular fatty change
20. **Which one of the following is not a feature of liver histology in non-cirrhotic portal fibrosis (NCPF)?** (AI 05, DPG 10)
  - a. Fibrosis in and around the portal tracts
  - b. Thrombosis of the medium and small portal vein branches
  - c. Non-specific inflammatory cell infiltrates in the portal tracts
  - d. Bridging fibrosis
21. **Bile infarct is related to:** (PGI Nov 10)
  - a. Hepatitis B
  - b. Dubin Johnson syndrome
  - c. Extrahepatic cholestasis
  - d. Intrahepatic cholestasis
  - e. Occlusion of hepatic artery



## HEPATITIS SEROLOGY & CLINICAL FEATURES

- 22. A patient presented with fibrosis of liver. ALT 40IU/uL. Serology report suggested:** (AIIMS Nov 2016)  
 HbeAg -ve                      Anti HBc Ab +ve  
 Anti HCV Ab +ve  
**Next line of investigation will be?**  
 a. Liver biopsy                      b. HBsAg  
 c. HBV DNA                      d. HCV RNA
- 23. Which hepatitis marker can be used to diagnose Acute Hepatitis B?** (Recent Question 2016-17)  
 a. HBc Ag                      b. Anti HBc Igm  
 c. HBs Ag                      d. HBe Ag
- 24. Which of the following viral markers signifies the ongoing viral replication in the case of Hepatitis-B infection?** (Recent Question 2016-17)  
 a. Anti-HBs                      b. Anti-HBc  
 c. HBe Ag                      d. HBs Ag
- 25. Serology profile of a patient suggested the following, what is your diagnosis?** (AIIMS Nov 2015)  
 • Hbs Ag: non reactive  
 • IgG anti Hbc: reactive  
 • HbeAg: Non reactive  
 • Hep B viral DNA: undetectable  
 a. Window period  
 b. Chronic hepatitis inactive stage  
 c. Recovery from remote infection  
 d. Recovery from acute infection
- 26. Reverse transcriptase is a RNA dependent DNA polymerase. Which of these use it ?** (AIIMS May 2015)  
 a. Hepatitis A virus  
 b. Hepatitis B virus  
 c. Hepatitis E virus  
 d. Hepatitis C virus
- 27. Which hepatitis causes more morbidity in pregnant female?** (Recent Question 2016)  
 a. Hep A                      b. Hep B  
 c. Hep C                      d. Hep E
- 28. Fecooral transmission is seen in?** (Recent Question 2016)  
 a. Hep A                      b. Hep B  
 c. Hep C                      d. Hep E
- 29. HBV DNA polymerase is encoded by which of the following gene?** (Recent Question 2015)  
 a. P                      b. X  
 c. C                      d. S
- 30. Calciviridae is?** (Recent Question 2015)  
 a. Hepatitis A                      b. Hepatitis E  
 c. Hepatitis C                      d. Hepatitis D
- 31. In Chronic Hepatitis B (HBV) infection presence of HBeAg (Hepatitis B e antigen) suggests which of the following?** (MAHA 16)  
 a. Ongoing viral replication  
 b. Resolving infection  
 c. Development of cirrhosis  
 d. Development of Hepatoma
- 32. Most common route hepatitis E transmission is?** (Recent Question 2015)  
 a. Sexual                      b. Faeco-oral  
 c. Horizontal                      d. Vertical

- 33. Gene responsible for mutation of HBV is?** (Recent Question 2015)  
 a. X gene                      b. S gene  
 c. P gene                      d. C gene
- 34. A nurse got a needle prick injury. Which of the following suggests active phase of hepatitis?** (AIIMS Nov 14)  
 a. IgM Ab of HBc                      b. IgG Ab of HBc  
 c. IgG of HBs                      d. Anti HbeAb
- 35. Most common subtype of Hepatitis B in North India is?** (Recent Question 2014)  
 a. adr                      b. adw  
 c. ayw                      d. ayr
- 36. Acute hepatitis 'B' can be diagnosed by:** (PGI May 12)  
 a. HBsAg                      b. IgM anti-HBcAb  
 c. HBeAg                      d. IgG anti-HBcAb  
 e. Core antigen
- 37. Hepatitis virus that causes chronic liver disease is?** (DNB Aug. 12 Pattern) (WBPG 2016)  
 a. Hepatitis A                      b. Hepatitis B  
 c. Hepatitis C                      d. Hepatitis D
- 38. Not a complication of acute viral Hepatitis?** (DNB Aug 12 Pattern)  
 a. Aplastic anemia                      b. Acute pancreatitis  
 c. Autoimmune hepatitis                      d. Hepatocellular carcinoma
- 39. A 25-year-old person presents with mild icterus. His HBsAg +ve, HBeAg -ve with SGOT and SGPT raised 5-6 times the original value. HBV DNA levels were >1,00,000/ml. What is your diagnosis?** (AI 2010)  
 a. Wild type HBV                      b. Surface mutant HBV  
 c. Precore mutant HBV                      d. Active HBV carrier

## HEPATITIS HISTOLOGY

- 40. Which of the following is/are features of acute hepatitis?** (PGI May 18)  
 a. Interface hepatitis  
 b. Mononuclear cell infiltration  
 c. Apoptosis of hepatocyte  
 d. Portal tracts Inflammation  
 e. Bridging fibrous
- 41. Hepatitis infection persists in 3% asymptomatic individuals. Why is there an increased risk of developing liver cancer in these patients?** (AIIMS Nov 2017)  
 a. Inability to induce inflammation to remove organism  
 b. Increased liver transaminases  
 c. High rate of hepatocyte proliferation  
 d. Integration of viral DNA to host DNA
- 42. Hallmark of chronic hepatitis** (Recent Question 2015)  
 a. Interface hepatitis  
 b. Ballooning degeneration of hepatocytes  
 c. Cholestasis  
 d. Periportal fibrosis and bridging fibrosis
- 43. Ground glass hepatocyte is seen in which hepatitis?** (Recent Question 2015)  
 a. Hepatitis A                      b. Hepatitis B  
 c. Hepatitis D                      d. Hepatitis E
- 44. Councilman bodies are seen in-** (Recent Question 2014, DNB Dec 09)  
 a. Alcoholic cirrhosis                      b. Wilson's disease  
 c. Acute viral hepatitis                      d. Autoimmune hepatitis



45. **Microvesicular fatty change in hepatocytes is seen due to infection with:** (Recent Question 2014)  
 a. Hepatitis A                      b. Hepatitis B  
 c. Hepatitis C                      d. Hepatitis D
46. **Histopathology of chronic hepatitis shows -** (Recent Question 2013)  
 a. Ballooning of hepatocytes  
 b. Councilman bodies  
 c. Bridging fibrosis  
 d. All of the above
47. **Chronic persistent hepatitis and chronic active hepatitis are differentiated by -** (DNB June 11)  
 a. Anti-Smith Ab  
 b. C-Reactive Protein  
 c. Arthritis  
 d. Liver biopsy
48. **Which of the following is single most important indicator of likelihood of progression of hepatitis to liver cirrhosis -** (MH 10)  
 a. Etiology  
 b. Associated serological findings  
 c. Presence of bridging necrosis  
 d. Presence of mallory hyaline bodies
49. **Chronic Active Hepatitis is most reliably distinguished from chronic Persistent hepatitis by the presence of -** (UPSC 05, 10)  
 a. Extrahepatic manifestations  
 b. Significant titre of anti-smooth muscle antibody  
 c. Characteristic liver histology  
 d. Hepatitis B surface antigen

#### ALCOHOLIC LIVER DISEASE

50. **The specific marker for alcoholic hepatitis?** (AIIMS Nov 18)  
 a. GGT                                  b. Alanine transaminase  
 c. Alkaline phosphatase          d. LDH
51. **Alcohol is a risk factor for which of the following cancers?** (PGI May 18)  
 a. Liver                                b. Esophagus  
 c. Gastric                              d. Cervical  
 e. Breast
52. **True about morphological feature(s) of alcoholic steatosis?** (PGI Nov 2017)  
 a. Fat droplet in hepatocyte  
 b. Mallory body may be seen  
 c. May cause hepatic necrosis  
 d. Perisinusoidal fibrosis may be seen  
 e. Macro-nodule formation
53. **Malory Hyaline body is seen in?** (Recent Question 2016-17)  
 a. Acute Hep A                      b. Chronic Hep A  
 c. Acute Hep B                      d. Chronic Hep B
54. **Ground glass hepatocytes are seen in ?** (Recent Question 2016-17)  
 a. Hep A                                b. Hep B  
 c. Hep C                                d. Hep D
55. **Malory Denk bodies are not seen in?**  
 a. Non alcoholic liver disease (Recent Question 2016-17)  
 b. Wilsons  
 c. Indian childhood cirrhosis  
 d. Chronic hep B

56. **Focal or confluent periportal necrosis along with ballooning degeneration of hepatocytes with or without Mallory bodies and megamitochondria suggestive of?** (Recent Question 2015/WBPG 2014)  
 a. Acute Hepatitis B                b. Chronic Hepatitis B  
 c. Alcoholic liver injury            d. Primary HCC
57. **Which of the following is not a feature of Alcoholic liver disease?** (WBPG 2015)  
 a. Macrovesicular fat within hepatocytes  
 b. Lipogranuloma  
 c. Lymphocytic infiltration of portal tracts  
 d. Portal & sinusoidal collagen deposits
58. **The sign of reversible injury in a case of alcoholic liver disease -** (Recent question 2014)  
 a. Loss of cell membrane          b. Nuclear karyolysis  
 c. Cytoplasmic vacuole            d. Pyknosis
59. **An obese female presented with features of hepatitis. She is a known diabetic. Which of the following would be the liver biopsy feature?** (AIIMS May 2014)  
 a. NASH                                b. Hepatocyte necrosis  
 c. Cirrhosis                            d. Lipoid necrosis
60. **A child presented with viral fever followed by unconsciousness. CT scan was suggestive of cerebral edema. Which of the following would be a finding on liver biopsy?** (AIIMS May 2014)  
 a. Peacemeal necrosis              b. Microvesicular steatosis  
 c. Bridging fibrosis                d. Ballooning degeneration
61. **Pathological manifestation of chronic alcoholism include all of the following except -** (Recent Question 2013)  
 a. Piecemeal necrosis  
 b. Ballooning degeneration  
 c. Microvesicular fatty changes  
 d. Central hyaline sclerosis
62. **If a patient has bilirubin 20 mg/dl, AST=313 IU/L, ALT=103 IU/L & GGT=44 IU/L. Most probable diagnosis is:** (PGI May 2013)  
 a. Viral hepatitis                    b. Alcoholic hepatitis  
 c. Billiary atresia                    d. Drugs  
 e. Autoimmune hepatitis

#### FATTY LIVER

63. **Microvesicular fatty liver is caused by-** (Recent Question 2014)  
 a. DM                                    b. Valproate  
 c. Starvation                         d. IBD
64. **Which does not cause microvesicular steatosis -** (Recent Question 2013)  
 a. Alcoholic fatty liver  
 b. Tetracycline toxicity  
 c. Acute fatty liver of pregnancy  
 d. Reyes syndrome

#### AUTOIMMUNE HEPATITIS

65. **About autoimmune hepatitis, which of the following is true?** (PGI Nov 2017)  
 a. More common in female  
 b. Anti-liver kidney microsome-1 (anti-LKM-1) antibody is found in type I subtype only  
 c. Oral corticosteroids are given in severe cases  
 d. Associated with other autoimmune disease  
 e. Rarely lead to cirrhosis




66. **Characteristic antibodies of autoimmune hepatitis include all of the following except:** (APPGMEE 2015)
- ANAs
  - Anti-CCP antibodies
  - Smooth muscle antibodies
  - Anti LKM antibodies
67. **Antibody present in autoimmune hepatitis type 2** (Recent Question 2015)
- Anti-LKM1 antibody
  - Anti-nuclear antibody
  - Anti-endomysial antibody
  - Anti-mitochondrial antibody
68. **40/male presented with fever and jaundice. The clinical picture improved on immunosuppressive therapy, what is the most likely diagnosis?** (AIIMS May 2014)
- Autoimmune hepatitis
  - Primary biliary cirrhosis
  - Infectious hepatitis
  - Secondary biliary cirrhosis
69. **Most common antibody in autoimmune hepatitis is?** (DNB Aug 12)
- U1 RNP
  - Anti-Sm
  - ANA
  - Anti-LKM
70. **In adults, most common autoimmune disease of liver is?** (DNB Aug 12)
- Autoimmune hepatitis
  - Sclerosing cholangitis
  - $\alpha$ -1 antitrypsin deficiency
  - Primary biliary cirrhosis
71. **Autoimmune hepatitis has the following Antibodies except?**
- ANA
  - ANCA
  - Anti LKM-1
  - Anti- SLA

#### HYPERBILIRUBINEMIA

72. **Causes of unconjugated hyperbilirubinemia is/are?**
- Hemolysis due to ABO incompatibility (PGI May 18)
  - Criggler-Najar syndrome
  - Gilbert syndrome
  - Sepsis
  - Roter's syndrome
73. **True about surgical jaundice:** (PGI May 2016)
- Increase of serum bilirubin
  - Increase acid phosphatase
  - Increase alkaline phosphatase
  - Urine bilirubin is absent
  - Stool sterocobilinogen absent
74. **Regarding Gilbert's syndrome, which one of the following statements is not correct?** (Recent Question 2016-17)
- Jaundice becomes severe with time.
  - Hyperbilirubinemia increases after fasting.
  - Inheritance of disease is autosomal dominant.
  - Liver histology is normal.
75. **A 55-year-old gentleman presented with history of right upper quadrant discomfort, jaundice, pruritis, fever, fatigue and weight loss. His serum bilirubin and alkaline phosphatase levels are raised and he also gives history of treatment for inflammatory bowel disease. He is most likely to be suffering from:** (Recent Question 2016-17)
- Benign bile duct stricture with cholangitis
  - Biliary worms
  - Bile duct malignancy
  - Primary sclerosing cholangitis
76. **True about primary biliary cirrhosis:** (PGI May 2015)
- More common in female
  - Periportal fibrosis
  - May be associated with Rheumatoid arthritis & crohn's disease
  - Jaundice may be present
  - Autoimmune disease are seen
77. **Flord duct lesions are diagnostic of** (Recent Question 2015)
- Klatskin tumor
  - Primary sclerosing cholangitis
  - Primary biliary cirrhosis
  - Secondary biliary cirrhosis
78. **False statement regarding alagille syndrome** (Recent Question 2015)
- Mutation in jagged-1-gene
  - Portal and bile ducts are completely absent
  - Micronodural cirrhosis of liver
  - All of the above
79. **Onion skin fibrosis is seen in** (Recent Question 2015)
- Primary biliary cirrhosis
  - Secondary biliary cirrhosis
  - Primary sclerosing cholangitis
  - Progressive familial intrahepatic cholestasis
80. **Which of the following does not cause cholestasis in newborn?** (Recent Question 2016)
- ABO incompatibility
  - Sepsis
  - Tyrosenemia
  - Biliary atresia
81. **All of the following are autosomal recessive except:** (Recent Question 2015)
- Gilbert's syndrome
  - CrigglerNajjar type I
  - CrigglerNajjar type II
  - Dubin Johnson syndrome
82. **Antimitochondrial antibodies are positive in** (Recent Question 2015)
- Primary sclerosing cholangitis
  - Secondary biliary cirrhosis
  - Primary biliary cirrhosis
  - Primary hemochromatosis
83. **Which one of the following inherited conditions causes direct hyperbilirubinemia:** (APPGMEE 2015)
- Gilbert syndrome
  - Type I CriglerNajjar syndrome
  - Rotor syndrome
  - Type II CriglerNajjar syndrome
84. **About Gilbert syndrome, true are all except-** (Recent Question 2014)
- Causes cirrhosis
  - Autosomal dominant
  - Normal liver function test
  - Normal histology
85. **Unconjugated hyperbilirubinemia is seen in -** (Recent Question 2014)
- Rotor syndrome
  - Dubin-Johnson syndrome
  - Gilbert syndrome
  - Bile duct obstruction





- 86. Feature of unconjugated bilirubin is/are:** (PGI Nov 2011)  
 a. Water soluble  
 b. Fat soluble  
 c. Direct reaction with Van den Bergh reaction  
 d. Affinity for brain tissue  
 e. Increased in hemolytic anemia
- 87. Pigment stone is composed of?** (Recent Question 2015)  
 a. Ca bilirubinate      b. Ca phosphate  
 c. Ca carbonate      d. Ca gluconate
- 88. Sclerosing cholangitis is associated with-** (Recent Question 2014)  
 a. Ulcerative colitis      b. Celiac sprue  
 c. Wilson's disease      d. Whipple's disease
- 89. Conjugated hyperbilirubinemia is seen in?** (Recent Question 2014)  
 a. Dubin Johnson Syndrome  
 b. Gilbert syndrome  
 c. Crigler Najjar syndrome  
 d. Hemolysis
- 90. Grossly pigmented liver is seen in?** (Recent Question 2014)  
 a. Crigler-Najjar Type I  
 b. Gilberts Syndrome  
 c. Dubin johnson Syndrome  
 d. Rotor's Syndrome
- 91. The following features differentiate Rotor syndrome from Dubin Johnson's syndrome EXCEPT** (APPGMEE 14)  
 a. Liver in patients with Rotor syndrome has no increased pigmentation and appears normal  
 b. In Rotor syndrome, Gall bladder is usually visualized on cholecystography  
 c. Total urinary coproporphyrin is substantially increased in Rotor syndrome  
 d. Fraction of coproporphyrin I in urine is elevated usually more than 80% of the total in Rotor syndrome
- 92. Which of the following statement(s) is/are true about primary sclerosing cholangitis?** (PGI Nov 2017)  
 a. May be complicated by bacterial infection  
 b. Rarely progress to biliary cirrhosis  
 c. Involve only intrahepatic bile duct not extrahepatic bile ducts  
 d. Associated with Inflammatory bowel disease  
 e. Narrowing of bile duct
- 93. Primary biliary cirrhosis is positive for-** (Recent Question 2013)  
 a. ANCA  
 b. Anti-mitochondrial antibody  
 c. Anti nuclear antibody  
 d. Anti-microsomal antibody
- 94. True about cholelithiasis is?** (PGI May 2013)  
 a. Cholesterol stones are most common  
 b. 90% of gallstone are radio-opaque  
 c. Mirrizi syndrome is due to impaction of stone in hartmann's pouch  
 d. Hemolytic anaemia cause black colored stone  
 e. Carcinoma is not a risk associated with gallstone
- 95. All are true about ascending cholangitis except?** (PGI May 2013)  
 a. Most commonly caused by gram positive organisms  
 b. In severe cases collapse can occur  
 c. Urgent removal of stone by ERCP can be done  
 d. Cholecystectomy can be done  
 e. Commonly caused by obstruction of bile duct by stone

- 96. True about obstructive jaundice:** (PGI May 2011)  
 a. Unconjugated bilirubin  
 b. Positive indirect Van den Bergh  
 c. Pruritus  
 d. Pale stools  
 e. Icterus
- 97. In post-hepatic jaundice, the concentration of conjugated bilirubin in the blood is higher than that of unconjugated bilirubin because:** (AIIMS Nov 10)  
 a. There is an increased rate of destruction of red blood cells.  
 b. The unconjugated bilirubin is trapped by the bile stone produced in the bile duct.  
 c. The conjugation process of bilirubin in liver remains operative without any interference.  
 d. The UDP-glucuronoyltransferase activity is increased manifold in obstructive jaundice.

#### METABOLIC LIVER DISEASE

- 98. Biochemical finding used for diagnosis of Wilson disease include(s):** (PGI May 2019)  
 a. Increased serum ceruloplasmin  
 b. Increased urinary copper excretion  
 c. Increased serum copper  
 d. Increased liver copper content  
 e. Decreased serum copper
- 99. A 1-month-old child with conjugated bilirubinemia and intrahepatic cholestasis. On Liver biopsy and staining with PAS red coloured granules were seen inside the hepatocytes. Probable diagnosis is ?** (AIIMS May 2015)  
 a. Alpha1 Antitrypsin deficiency  
 b. Congenital hepatic fibrosis  
 c. Wilson disease  
 d. hereditary hemochromatosis
- 100. PAS-positive, diastase-resistant globules in hepatocytes are seen in?** (Recent Question 2015)  
 a. Hemochromatosis  
 b. Wilsons disease  
 c. Alpha 1 antitrypsin deficiency  
 d. Acute necrotic hepatitis
- 101. Hemochromatosis is a defect in metabolism of:** (Recent Question 2015)  
 a. Iron      b. Copper  
 c. Magnesium C      d. Calcium
- 102. Gene for Wilson's disease is located on chromosome-** (Recent Question 2014)  
 a. 7      b. 10  
 c. 13      d. 17
- 103. Wilson's disease is characterized by-** (Recent Question 2014), (WBPG 2016)  
 a. Increased serum ceruloplasmin  
 b. Decreased copper excretion in urine  
 c. ↑Ceruloplasmin  
 d. Low ↑Ceruloplasmin high urine copper
- 104. Diabetic patient with liver cirrhosis and hyperpigmentation, diagnosis is-** (Recent Question 2014)  
 a. Wilson's disease  
 b. Hemochromatosis  
 c. Primary sclerosing cholangitis  
 d. Hepatitis B



- 105. Type of inheritance in Wilson's disease-**  
(Recent Question 2014)  
a. Autosomal dominant      b. Autosomal recessive  
c. X-linked dominant      d. X-linked recessive
- 106. PAS positive intrahepatic globules are seen in:**  
(Recent Question 2014)  
a. Wilson disease  
b. Hemochromatosis  
c. Primary Sclerosing Cholangitis  
d. Alpha-1-antitrypsin deficiency
- 107. ATP7B gene is present on chromosome:**  
(Recent Question 2014)  
a. 5      b. 13  
c. 18      d. 21
- 108. In Alpha-1 anti trypsin deficiency, hepatocytes are:**  
(Recent Question 2014)  
a. PAS + ve diastase resistant  
b. Diastase positive PAS resistant  
c. PAS -ve      d. Oilred O positive
- 109. Hemochromatosis leads to deposition of?**  
(Recent Question 2014)  
a. Iron      b. Copper  
c. Zinc      d. Lead
- 110. In Wilson's disease, hepatic copper content usually exceeds \_\_\_\_\_ µg per gram dry weight** (MAHA 16)  
a. 150      b. 250  
c. 350      d. 450
- 111. Most common gene responsible for hereditary hemochromatosis is?**  
(Recent Question 2013)  
a. HJV gene      b. HAMP gene  
c. Tfr2 gene      d. HFE gene
- 112. Which of the following leads to chronic liver disease?**  
(DNB Aug 12 Pattern)  
a. Hepatitis A  
b. EBV  
c. Infectious mononucleosis  
d. α-1-antitrypsin deficiency

### LIVER TUMORS

- 113. Which of the infections is/are predisposes to cholangiocarcinoma?**  
(PGI May 18)  
a. Paragonimus westermani  
b. Fasciola hepatica  
c. Schistosoma hematobium  
d. Opisthorchis viverrini  
e. Clonorchis sinensis
- 114. Which of the following is true about Nodular Regenerative Hyperplasia?**  
(JIPMER 18)  
a. Nodule size 0.1 to 1 cm  
b. Fibrosis septa present  
c. Portal hypertension seen in 50% of patients  
d. AST and ALT are markedly elevated
- 115. Vinyl chloride is associated with?**  
(JIPMER 18)  
a. Hemangiosarcoma      b. Hepatoma  
c. Testicular carcinoma      d. Thyroid malignancy
- 116. True about liver haemangioma are all except?**  
(PGI May 18)  
a. Most common benign tumor of liver  
b. Pregnancy is risk factor  
c. Thrombocytopenia may occur  
d. Diagnosis by MRI and CT scan  
e. Need surgical removal in every case because chances of rupture is high
- 117. Which of the following has best prognosis?**  
(JIPMER 2016)  
a. Fibrolamellar Variant of HCC  
b. Hemangiosarcoma  
c. Adenosarcoma  
d. Hepatocellular Ca
- 118. True about fibrolamellar carcinoma of liver:**  
(PGI May 2016)  
a. Better prognosis than typical hepatocellular carcinoma  
b. Associated with cirrhosis  
c. AFP-positive  
d. Occur in younger adults      e. More common in females
- 119. All are true about focal nodular hyperplasia except:**  
(PGI May 2016)  
a. Multiple nodule may present  
b. More common in male  
c. May be associated with contraceptive pills use  
d. Hypovascular on the arterial-phase and hypervascular on the delayed-phase CT images  
e. CT is less sensitive than MRI in depicting the characteristic central scar
- 120. Regarding carcinoma gall bladder following features are true except:**  
(Recent Question 2016-17)  
a. One can have similar presentation with benign biliary disease  
b. Squamous cell carcinoma is 40% of all cases  
c. Most patients present with advanced disease  
d. Prognosis is poor
- 121. Which of the following is not true about alcoholic cirrhosis?**  
(Recent Question 2016-17)  
a. On many occasions alcoholic hepatitis and alcoholic cirrhosis coexist.  
b. Concomitant HIV infection accelerates it.  
c. Starts with macronodular and later on changes to micronodular cirrhosis.  
d. 10 - 40 % remains clinically silent.
- 122. Which of the following do not cause Hepatocellular Ca?**  
(Recent Question 2016)  
a. Hepatitis B  
b. Tyrosenemia  
c. Alcoholism  
d. Non alcoholic fatty liver disease
- 123. Angiosarcoma of the liver can occur due to occupational exposure to:**  
(Recent Question 2015)  
a. Asbestos      b. Benzene  
c. Vinyl chloride      d. Toluene
- 124. Hepatic adenoma is most common in**  
(Recent Question 2015)  
a. Young males      b. Young females  
c. Old males      d. Old females
- 125. Which liver tumor has the best prognosis**  
(Recent Question 2015)  
a. Hepatocellular carcinoma  
b. Hemangiosarcoma  
c. Hemangioblastoma      d. Fibrolamellar carcinoma
- 126. Not true about HNF1-α Inactivated hepatocellular adenomas**  
(Recent Question 2015)  
a. Mostly in women  
b. High risk of malignant transformation  
c. Associated with MODY-3  
d. OCPs are implicated in pathogenesis



- 127. Not true about hepatoblastoma (Recent Question 2015)**  
 a. Most common in children  
 b. Mature hepatocytes present  
 c. Not associated with cirrhosis  
 d. Fatal if untreated
- 128. Which of the following is NOT a risk factor for hepatocellular carcinoma? (Recent Question 2015)**  
 a.  $\alpha$ 1-antitrypsin deficiency b. NASH/NAFLD  
 c. Chronic alcoholism d. Hepatitis D
- 129. Which malignancy is associated with liver cirrhosis? (Recent Question 2015)**  
 a. Hepatocellular Ca b. Cholangiocarcinoma  
 c. FibrolamellarCa d. Pancreatic Ca
- 130. Which is a risk factor for Cholangiocarcinoma? (Recent Question 2015)**  
 a. Persistent hepatitis b. Ulcerative colitis  
 c. Crohn's ds d. Chronic cholecystitis
- 131. Vinyl Chloride is associated with which Carcinoma? (Recent Question 2015)**  
 a. Liver b. Spleen  
 c. Lung d. Prostrate
- 132. True about fibrolamellar variant of HCC? (PGI May 2014)**  
 a. Better prognosis than Primary HCC  
 b. More common in elderly  
 c. Raised AFP seen  
 d. Underlying cirrhosis not a risk factor  
 e. Neurotensin is a biomarker
- 133. Periportal fibrosis is caused by (Recent Question 2013)**  
 a. Methotrexate b. Phenytoin  
 c. Thorotrast d. Halothane
- 134. Thorium dioxide causes - (Recent Question 2013)**  
 a. Lymphoma b. Lymphangiosarcoma  
 c. Angiosarcoma d. Hemangioendothelioma
- 135. Which is risk factor for cholangiocarcinoma? (Recent Question 2013)**  
 a. Obesity  
 b. Primary sclerosing cholangitis  
 c. Salmonella carrier state  
 d. HBV infection
- 136. Klatskin tumor is: (Recent Question 2013)**  
 a. Nodular type of cholangiocarcinoma  
 b. Fibrolamellar hepatocellular carcinoma  
 c. Gall bladder carcinoma  
 d. Hepatocellular carcinoma
- 137. Not raised in liver disorder: (PGI May 2013)**  
 a. Lipase b. Amylase  
 c. ALP d. AST  
 e. ALT
- 138. True about serum AFP level: (PGI May 2013)**  
 a. Raised in testicular tumor  
 b. Raised in 50-70% cases of HCC  
 c. Correlation between tumor recurrence after surgery in HCC  
 d. Correlation with HCC size  
 e. Upper limit of normal in the serum is 200ng/mL
- 139. Not true about FibrolamellarCa of liver (Jipmer 2012)**  
 a. Both sexes equally affected  
 b. Young age group  
 c. Arises from cirrhotic liver  
 d. Good prognosis

- 140. Primary sclerosing cholangitis is likely to be associated with: (JIPMER 11)**  
 a. Adenocarcinoma of pancreas  
 b. Cholangiocarcinoma  
 c. Hepatocellular carcinoma  
 d. Adenocarcinoma of gall bladder
- 141. Von-Meyenburg's complexes are seen in? (PGI Nov 10)**  
 a. Brain b. Liver  
 c. Kidney d. Spleen  
 e. Pancreas
- 142. Which virus causes hepatocellular carcinoma-**  
 a. Arbo virus b. Herpes virus  
 c. Hepatitis-A virus d. Hepatitis-B virus

## GALLBLADDER


- 143. Rokitansky-Aschoff sinuses are a feature of: (Recent Question 2015)**  
 a. Adenomyomatosis of gall bladder  
 b. Chronic Cholecystitis  
 c. Acute Cholecystitis  
 d. Ca gall bladder
- 144. The following condition of GB is precancerous - (Recent Question 2013)**  
 a. Cholesterosis  
 b. Porcelain gall bladder  
 c. Biliary atresia  
 d. Choledochal cyst

## PANCREATITIS

- 145. All of the following are etiological factors of Acute Pancreatitis except? (AIIMS May 2014)**  
 a. Hyperlipidemia  
 b. Trauma  
 c. Mutations in trypsin inhibitor (SPINK1) genes  
 d. Islet cell hypertrophy

## TUMORS OF PANCREAS

- 146. Most common gene associated with pancreatic cancer (Recent Question 2016)**  
 a. KRAS b. SMAD  
 c. P53 d. Rb
- 147. Most common site for Ca Pancreas? (Recent Question 2015)**  
 a. Head b. Body  
 c. Tail d. Uncinate process
- 148. Most commonly involved Oncogene in Pancreatic Carcinoma is? (Recent Question 2013)**  
 a. KRAS b. p16/CDKN2A  
 c. TP53 d. SMAD4
- 149. Maximum progression to pancreatic carcinoma occurs in? (PGI May 06, May 2013)**  
 a. Intraductal papillary mucinous neoplasms  
 b. Pseudopancreatic cyst  
 c. Serous cystic neoplasms  
 d. Mucinous cystic neoplasms



## Answers with Explanations

### 1. Ans. (a) GGT

The activities of three enzymes—alkaline phosphatase, 5'-nucleotidase, and -glutamyl transpeptidase (GGT)—are usually elevated in cholestasis

### 2. Ans. (c) Nutmeg liver: Red areas are pericentral necrotic areas, white areas are viable fibrotic periportal area

(Ref: Robbins 9th/pg 863; 8th/pg 872)

The given picture shows **Chronic passive hepatic congestion**. The centrilobular regions are grossly red-brown and slightly depressed (because of cell death) and are prominently visible against the surrounding zones of uncongested tan liver (**nutmeg liver**).

### 3. Ans. (a) Perihepatic obstruction

(Ref: Robbins 9th/830-840; Harrison's 18th/chapter 42)

Condition	Serum Bilirubin	Urine Urobilinogen	Urine Bilirubin	Fecal Urobilinogen
Normal	Direct 0.1-0.4 mg/dl Indirect 0.2-0.7 mg/dl	0.4 mg/24 h	Absent	40-280 mg/24h
Hemolytic anemia	↑Indirect	Increased	Absent	Increased
Hepatitis	↑Direct and indirect	Decreased if micro-obstruction is present	Present if micro-obstruction occurs	Decreased
Obstructive jaundice	↑Direct	Absent	Present	Trace to absent

### 4. Ans. (d) Acetaminophen (Ref: Robbins 9th/pg 864)

### 5. Ans. (a) Centrilobular necrosis (Ref: R 9th/pg 864)

Centrilobular zonal necrosis	Periportal injury
<ul style="list-style-type: none"> <li>Carbon tetrachloride (CCl<sub>4</sub>)</li> <li>Chloroform</li> <li>Cardiac shock</li> <li>Trichloroethylene</li> </ul>	<ul style="list-style-type: none"> <li>Yellow phosphorus poisoning</li> </ul>
	Midzonal necrosis
	<ul style="list-style-type: none"> <li>Yellow fever</li> </ul>

### 6. Ans. (a) Chronic venous congestion of liver

(Ref: Robbins 9th/pg 863)

### 7. Ans. (d) No Risk of hepatocellular carcinoma

(Ref: Robbins 9th/pg 853; 8th/pg 854)

**Alagille Syndrome (Syndromic Paucity of Bile Ducts; Arteriohepatic Dysplasia)**

- An **autosomal dominant** disorder characterized by **absence of bile ducts in portal tracts**.
- Caused by mutations or deletion of gene encoding **Jagged1**, on chromosome 20p.
- Patients can survive into adulthood but **are at risk for hepatic failure & hepatocellular carcinoma**
- Major clinical features:
  - Chronic cholestasis**
  - Peripheral pulmonary artery stenosis**
  - Butterfly-like vertebral arch defects**
  - An eye defect known as **posterior embryotoxon**,
  - A peculiar hypertelic, **triangular facies**.

### 8. Ans. (a) Acetaminophen (Ref: Robbins 9th/pg 864)

### 9. Ans. (d) Allopurinol (Ref: Robbins 9th/pg 863)

#### Fibrin Ring or "Doughnut" Granulomas

- Small, non-necrotizing granulomas with a very distinctive appearance that are usually found in the liver and bone marrow in patients with<sup>o</sup> fever.
- These granulomas characteristically contain a ring-like structure consisting of fibrinoid material; they may or may not have a centrally located fat vacuole(s).
- Other conditions: CMV, EBV, hepatitis A, infectious mononucleosis, visceral leishmaniasis, Lyme disease, toxoplasmosis, Hodgkin disease, non-Hodgkin lymphomas, and drug reactions like allopurinol.

### 10. Ans. (b) Anabolic steroids (Ref: Robbins 9th/pg 863)

#### Peliosis Hepatis

- Peliosis hepatis** is a condition in which there is **primary hepatic sinusoidal dilation**; unknown pathogenesis
- Sinusoidal dilation** occurs in any condition in which **efflux of hepatic blood is impeded**.
- Liver contains **blood-filled cystic spaces**, either unlined or lined with sinusoidal endothelial cells.
- Peliosis hepatis is associated with **cancer, tuberculosis, AIDS, or post-transplantation immunodeficiency**.
- Also associated with exposure to **anabolic steroids, oral contraceptives and danazol**.
- Bartonella species** have been seen in the sinusoidal endothelial cells in **AIDS-associated peliosis**.
- Clinical signs are generally absent, but **fatal intra-abdominal hemorrhage** or **hepatic failure** may occur.





11. Ans. (a) **Alcohol** (Ref: Robbins 9th/pg 843 Refer pretext)

12. Ans. (b) **Methotrexate** (Ref: Harrison 18th/ch 309)

13. Ans. (d) **Bridging fibrosis** (Ref: Robbins 9th/pg 863)

#### Non-cirrhotic Portal Fibrosis (NCPF) and Idiopathic Portal Hypertension

<b>Characterized by</b>	Portal hypertension and a moderate portal fibrosis without cirrhosis.
<b>Epidemiology</b>	<ul style="list-style-type: none"> <li>NCPF is common in India</li> <li>Idiopathic portal hypertension, described in Japan, has a female predominance</li> </ul>
<b>Clinical presentation</b>	<ul style="list-style-type: none"> <li>Upper gastrointestinal bleeding</li> <li>Isolated splenomegaly</li> </ul>
<b>Pathogenesis</b>	<ul style="list-style-type: none"> <li>Idiopathic</li> <li>Bacterial infection of gut causing septic embolization of portal vein.</li> <li>Fibrosis of portal vein branches associated with increased expression of <b>vascular cell adhesion molecule-1 (VCAM-1)</b>.</li> </ul>
<b>Histology</b>	<ul style="list-style-type: none"> <li>Increased <b>connective tissue deposition and fibrosis of portal tracts</b> with thrombotic obliteration of small branches of portal veins.</li> <li>Infiltrates in portal tract</li> <li>Also called '<b>hepatic sclerosis</b>' or '<b>obliterative portal venopathy</b>'</li> </ul>

14. Ans. (b) **Alcohol** (Ref: Robbins 9th/pg 837)

- Alcoholic hepatitis is often accompanied by prominent activation of sinusoidal stellate cells and portal fibroblasts, giving rise to fibrosis.
- Fibrosis begins with sclerosis of central veins.
- Perisinusoidal scar** then accumulates in the space of Disse of the centrilobular region, spreading outward, encircling individual or small clusters of hepatocytes in a **chicken wire fence pattern**

15. Ans. (b) **Stellate cells** (Ref: Robbins 9th/pg 825; 8th/pg 837)

16. Ans. (c) **CCl<sub>4</sub>** (Ref: Robbins 9th/pg 864; 8th/pg 872)

17. Ans. (b) **Cytokeratin** (Ref: Robbins 9th/pg 843; 8th/pg 858)

**Mallory-Denk bodies** (previously called **Mallory Hyaline bodies**<sup>Q</sup>): Clumped, amorphous, eosinophilic material in ballooned hepatocytes. Intermediate filaments-**keratins 8 and 18 with ubiquitin**.

18. Ans. (b) **Pseudomelanin** (Ref: Sternburg /pg 1667-1670)

Discussing options one by one,

- Lipofuscin**: Lipofuscinosis in centrilobular hepatocytes is induced by prolonged intake of Phenacetin, Aminopyrin, Chlorpromazine, Anticonvulsant therapy.
- Pseudomelanin is not deposited in liver but is seen in intestine in melanosis coli of intestine**
- Bile pigment in obstructive jaundice**
- Iron**: Dark brown or black **malarial pigment** in Kupffer cells & in **hemochromatosis**
- Melanin: deposits in hepatocytes in melanoma**

19. Ans. (b) **Cirrhosis** (Ref: Robbins 9th/pg 842-844)

- Alcoholic Cirrhosis** is the **final & irreversible form of alcoholic liver disease** which usually evolves slowly and insidiously but may develop in 1 or 2 years in some cases of alcoholic liver disease.
- Microvesicular & Macrovesicular fatty change is completely reversible** if there is abstention from further intake of alcohol.

20. Ans. (d) **Bridging fibrosis** (Ref: Robbins 9th/pg 863)

21. Ans. (b, c, d); **b. Dubin Johnson syndrome; c. Extrahepatic cholestasis; d. Intrahepatic cholestasis**

(Ref: Liver Pathology by Suriawinata & Thung, 2011/ pg 67)

Bile infarct is seen in **severe cholestatic disease**

Cholestasis is seen in **Dubin Johnson syndrome, Extrahepatic cholestasis & Intrahepatic cholestasis**

**Bile infarcts:**

- In **severe cholestasis**, extravasation of bile leads to hepatocyte necrosis & pale foamy macrophages, forming **bile infarcts**
- Although small bile infarcts may occur in any intense canalicular cholestasis, **large lesions**, especially around portal tracts, **suggest biliary obstruction**.

22. Ans. (b) **HBsAg**

(Ref: Robbins 9th/pg 831-832; 8th/pg 845-846; Harrison 19th/ Pg 2097)

The history in the question given is suggestive of Hepatitis C patient with concurrent Hepatitis B infection (HBcAb) but it is not enough to reliably tell us if it is acute infection or remote recovery as IgM or IgG is not mentioned in the question. So the next best logical step should be HbsAg followed by HBV DNA

23. Ans. (b) **Anti HBc IgM**

24. Ans. (c) **HBe Ag**

(Ref: Harrison 19th ed. Pg. 2007-2015; R9th 831-833)

25. Ans. (c) **Recovery from remote infection**

(Ref: Harrison 19th ed. Pg. 2007-2015; Robbins 9th 831-833)

26. Ans. (b) **Hepatitis B virus** (Ref: Robbins 9th/pg 831-832)


27. Ans. (d) **Hep E** (Ref: Robbins 9th/pg 835; 8th/pg 849)

A characteristic feature of HEV infection is the high mortality rate among pregnant women, approaching 20%

28. Ans. (d) **Hep E** (Ref: Robbins 9th/pg 830; 8th/pg 840)

**Most common mode of Hep E transmission is Faeco-oral**

Transmission	HAV	HBV	HCV	HDV	HEV
<b>Fecal-oral</b>	+++	—	—	—	+++
<b>Percutaneous</b>	Unusual	+++	+++	+++	—
<b>Perinatal (Fetomaternal)</b>	—	+++	±	+	—
<b>Sexual</b>	±	++	±	++	—



29. Ans. (a) **P** (Ref: Robbins 9th/pg 831-832; 8th/pg 845-846)

30. Ans. (b) **Hepatitis E**

- HEV was included in family **Caliciviridae** previously
- Although resembling caliciviruses, HEV is **now thought to be sufficiently distinct** from any known agent to merit a new classification of its own, as a unique genus, **Hepevirus**, within the family Hepeviridae.
- So the best answer here is HEV

31. Ans. (a) **Ongoing viral replication**

(Ref: R 9th/pg 831-832)

#### HBeAg

- The principal clinical usefulness of HBeAg is as an **indicator of relative infectivity**.
- HBeAg, HBV-DNA & DNA polymerase appear in serum soon after HBsAg, and all signify active viral replication.
- Persistence of HBeAg is an important indicator of continued viral replication, infectivity, and probable progression to chronic hepatitis.
- The appearance of anti-HBe antibodies implies that an acute infection has peaked and is on the decline.

32. Ans. (b) **Faeco-oral** (Ref: Robbins 9th/pg 831-835)

33. Ans. (a) **X gene** (Ref: Robbins 9th/pg 831-832)

Gene responsible for mutation of HBV is X gene

- **X gene** codes for **HBxAg**, that is capable of **transactivating transcription** of both viral & cellular genes
- HBxAg effects **calcium release from mitochondria** → **activates signal-transduction pathways** → stimulation of HBV reverse transcription & HBV DNA replication
- It also **enhances replication of HBV**, leading to severe **chronic hepatitis & hepatocellular carcinoma**.
- Can also enhance the transcription & replication of other viruses like **HIV**
- Cellular processes transactivated by X include human **interferon gene & class I MHC genes**
- These effects contribute to **enhanced susceptibility** of HBV-infected hepatocytes to cytolytic T cells.
- The expression of X can also **induce programmed cell death (apoptosis)**.

34. Ans. (a) **IgM Ab of HBc**

(Ref: Robbins 9th/pg 831-832; 8th/pg 845-846; Harrison 18th/Chapter 304)

35. Ans. (c) **ayw**

(Ref: Ismail et al. Molecular epidemiology & genetic characterization of hepatitis B virus in the Indian subcontinent. Int J Infect Dis. 2014 Mar; 20:1-10)

Hepatitis B has **eight subtypes and eight genotypes (A-H)** This question can only be answered by a **recent publication on Hepatitis B**, mentioned above!

Most common genotype/subtype of Hepatitis B in:

- **North-Eastern India: D/ayw** followed by C / adr
- **Southern India: D/ayw**
- **Eastern India: C/adr**

36. Ans. (a, b); **a. HBsAg; b. IgM anti-HBc Ab**

(Ref: Robbins 9th/pg 831-832; 8th/pg 845-846)

37. Ans. (c) **Hepatitis C** (Ref: Robbins 9th/pg 833-834)

**Progression to chronicity in hepatitis C is upto 85% (maximum among the Hepatitis viruses)**

38. Ans. (d) **Hepatocellular carcinoma** (Ref: R 9th/pg 837)

- **Hepatocellular carcinoma is a complication of chronic hepatitis & not acute hepatitis.**
- Rare complications of acute viral hepatitis include pancreatitis, myocarditis, atypical pneumonia, aplastic anemia, transverse myelitis & peripheral neuropathy

39. Ans. (c) **Precore mutant HBV**

(Ref: Robbins 9th/pg 831-832)

**Pre-core Mutant:** Mutated strains of HBV emerge that **do not produce HBeAg despite the presence of serum HBV DNA**.

40. Ans. (b, c, d) **b. Mononuclear cell infiltration; c. Apoptosis of hepatocyte; d. Portal tracts Inflammation**

41. Ans. (d) **Integration of viral DNA to host DNA**

42. Ans. (d) **Periportal fibrosis and bridging fibrosis**

(Ref: Robbins 9th/pg 837; 8th/pg 850)

43. Ans. (b) **Hepatitis B** (Ref: Robbins 9th/pg 837; 8th/pg 852)

Diagnostic hallmark of **Chronic Hepatitis B** is “**ground-glass**” hepatocytes (cells with **endoplasmic reticulum swollen by HBsAg**)

44. Ans. (c) **Acute viral hepatitis** (Ref: Robbins 9th/pg 837)

- **Councilman bodies:** are intensely eosinophilic apoptotic hepatocytes with pyknotic nucleus, seen in acute viral hepatitis

45. Ans. (d) **Hepatitis D** (Ref: Robbins 9th/pg 835; 8th/pg 849)

Acute **HDV infection** has been associated with **microvesicular change** in hepatocytes; this ‘**spongiocytic change**’ or ‘**morula cell degeneration**’ was attributed to accumulation of **small lipid droplets** in damaged hepatocytes.

46. Ans. (c) **Bridging fibrosis** (Ref: Robbins 9th/pg 837)

- **Hallmark of progressive chronic liver damage** is deposition of **fibrous tissue (scarring)** → bridging fibrosis


47. Ans. (d) **Liver biopsy**

(Ref: Rosai and Ackerman's Surgical Pathology, Volume 1, chapter 13, Box 13.2)

Chronic hepatitis is classified on the basis of:

- Diagnosis of chronic hepatitis
- Etiology
- Grade of disease activity: evaluated by clinical symptoms, aminotransferase levels & liver biopsy
- Stage of disease progression

This classification system was used to distinguish between a **milder form (chronic persistent hepatitis – CPH)** with a



low degree of necro-inflammatory activity & **more severe variants (chronic aggressive or active hepatitis – CAH)** featuring higher degrees of necro-inflammatory lesions.

48. **Ans. (a) Etiology** (Ref: Harrison 18th/Chapter 308)

Single most important indicator of likelihood of progression of hepatitis to liver cirrhosis is etiology

49. **Ans. (c) Characteristic liver histology**

(Ref: Rosai and Ackerman's Surgical Pathology, Volume 1, chapter 13, Box 13.2)

50. **Ans. (a) GGT**

Serum levels of GGT differ from those of ALP during pregnancy, in which GGT remains normal even during cholestasis in pregnancy. GGT is often increased in alcoholics even without liver disease; in some obese people; and in the presence of high concentrations of therapeutic drugs, such as acetaminophen and phenytoin and carbamazepine (increased up to five times the reference limits), even in the absence of any apparent liver injury. Studies suggest that alcohol induces mitochondrial damage, resulting in the release of mitochondrial AST, which, besides being the predominant form of AST in hepatocytes, has a significantly longer half-life than do extramitochondrial AST and ALT. This frequently results in the disproportionate elevation of AST over ALT, yielding an AST/ALT quotient, also called the DeRitis ratio, of 3–4:1 in alcohol-induced liver disease

51. **Ans. (d) Cervical**

52. **Ans. (a, b, d) a. Fat droplet in hepatocyte; b. Mallory body may be seen; d. Perisinusoidal fibrosis may be seen**

(Ref: Robbins 9th ed p 845)

Perisinusoidal scarring leads to a classic micronodular or Laennec cirrhosis first described for end-stage alcoholic liver disease.

53. **Ans. (d) Chronic Hep B** (Ref: Robbins 9th/pg 837)

54. **Ans. (b) Hep B** (Ref: Robbins 9th/pg 837; 8th/pg 850)

55. **Ans. (d) Chronic hep B** (Ref: Robbins 9th/pg 837)

56. **Ans. (c) Alcoholic liver injury**

(Ref: Rosai and Ackerman's Surgical Pathology, Volume 1, chapter 13, Box 13.2)

- **Megamitochondria** appear as **eosinophilic, PAS-diastase-negative, round, oval, or cigar-shaped inclusions**
- Better visualized with **chromotrope-aniline blue (CAB) stain** & immunohistochemistry.
- They are **not specific for alcohol-induced liver disease**, but found more frequently in alcohol-related fibrosis

- **Oxyphilic granular hepatocytes or oncocytes** are a common, though not pathognomonic, cellular component in acute and chronic alcoholic liver disease.
- On Electron microscopy, Oncocytes are characterized by a **large number of mitochondria** (mitochondriosis).

57. **Ans. (c) Lymphocytic infiltration of portal tracts**

(Ref: Robbins 9th/pg 837; 8th/pg 850; Rosai and Ackerman's Surgical Pathology, Volume 1, chapter 13)

**Lipogranuloma** (or fat granuloma) represents a focal response to rupture of lipid-laden hepatocytes. It contains macrophages, occasional lymphocytes, eosinophils, and sometimes giant cells; It may be seen in

**Alcoholic liver disease; Neutrophil rather than lymphocytic infiltration is seen in alcoholic liver disease**

58. **Ans. (c) Cytoplasmic vacuole**

(Ref: Robbins 9th/pg 842-844)

- **Alcoholic liver disease is fully reversible in the steatosis stage.**
- In this stage fatty changes appear in the hepatocytes, represented by **cytoplasmic vacuoles** as the lipid is dissolved during processing
- Loss of cell membrane, nuclear karyolysis and pyknosis represents irreversible cell injury and so not reversible.

59. **Ans. (a) NASH** (Ref: Robbins 9th/pg 845-846)

Obesity and diabetes are two important risk factors for NASH

60. **Ans. (b) Microvesicular steatosis**

(Ref: Robbins 8th/pg 857; Sternberg's 4th ed/ch 36)

The given clinical scenario and findings on CNS imaging are suggestive of Reyes Syndrome.

**Reye syndrome** (also called 'Jamshedpur fever')

- Occurs principally, but not exclusively, in **young children**
- Predisposing factors: **Salicylate** use, inherited **disorder of mitochondrial  $\beta$ -oxidation**
- Presents clinically with an initial acute, mild viral illness, followed by **vomiting, lethargy & coma**
- Severe disease with **poor prognosis**, resulting in death in about one-third of patients.
- Histopathology of liver: **Panlobular microvesicular Steatosis**, with smaller droplets in centrilobular areas & larger fat vacuoles in periportal regions & **necrosis** of periportal hepatocytes
- **Glycogen depletion** is also a feature

61. **Ans. (a) Piecemeal necrosis** (Ref: Robbins 9th/pg 837)

**Piecemeal necrosis is suggestive of chronic active hepatitis & not Alcoholic liver disease**

62. **Ans. (b) Alcoholic hepatitis** (Ref: Robbins 9th/pg 842-844)

In the given scenario, the patient has Hyperbilirubinemia with AST: ALT ratio > 3:1 & GGT=44 IU/l. Most probable diagnosis is **Alcoholic liver disease**.



### Features of Alcoholic liver disease:

- Clinical Features are usually non-specific.
- **AST:ALT Ratio =2:1 or 3:1**<sup>Q</sup> (In contrast to other etiologies where AST:ALT =1:1 or 1:2)
- Increased GGT (Normal 10-40 U/L), Bilirubin, Alkaline phosphatase

### 63. Ans. (b) Valproate

(Ref: Robbins 9th/pg 842-844; 8th/pg 857-858; Rosai and Ackerman's Surgical Pathology, Volume 1, chapter 13)

**Steatosis** refers to hepatocellular fat accumulation.

**Macrovesicular steatosis is more commonly seen than Microvesicular steatosis.**

Features	Macrovesicular steatosis	Microvesicular steatosis
<b>Histology</b>	<ul style="list-style-type: none"> <li>• Single large vacuole distends the hepatocyte &amp; displaces the nucleus to one side</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Fine-droplet fatty change</b> due to inhibition of mitochondrial fatty acid <math>\beta</math>-oxidation &amp; mitochondrial dysfunction</li> </ul>
<b>Etiology</b>	<ul style="list-style-type: none"> <li>• Alcoholic liver disease</li> <li>• Obesity; Diabetes.</li> <li>• Cachexia</li> <li>• PEM or Malnutrition (Periportal)</li> <li>• After total parenteral nutrition</li> <li>• Drugs: Steroid, Methotrexate</li> <li>• AIDS</li> <li>• Poisoning: phosphorus, <math>\text{CCl}_4</math></li> </ul>	<ul style="list-style-type: none"> <li>• Alcoholic liver disease</li> <li>• Acute fatty liver of pregnancy</li> <li>• Severe liver injury</li> <li>• Jamaican vomiting sickness</li> <li>• Drugs: Tetracycline, Amiodarone, Valproate</li> <li>• Reye syndrome</li> </ul>

### 64. Ans. (a) Alcoholic fatty liver

(Ref: Robbins 9th/pg 842-844)

Alcoholic fatty liver shows both microvesicular fatty change, initially, followed by macrovesicular fatty change; But rest of the options give rise to microvesicular steatosis only

### 65. Ans. (a, c, d, e) a. More common in female; c. Oral corticosteroids are given in severe cases; d. Associated with other autoimmune disease; e. Rarely lead to cirrhosis (Ref: Robbins 9th ed p 855-857)

Anti-liver kidney microsome-1 (anti-LKM-1) antibody is found in type II subtype.

### 66. Ans. (b) Anti-CCP antibodies (Ref: R 9th/pg 839-840)

Anti-CCP Ab seen in Rheumatoid Arthritis

### 67. Ans. (a) Anti-LKM1 antibody

(Ref: Robbins 9th/pg 842; 8th/pg 857)

### 68. Ans. (a) Autoimmune hepatitis (Ref: R 9th/pg 839-840)

### 69. Ans. (c) ANA (Ref: Robbins 9th/pg 839-840 8th/pg 855-856)

### 70. Ans. (d) Primary biliary cirrhosis (Ref: R 9th/pg 858-859)

- **Most common autoimmune disease of liver in adults is Primary biliary cirrhosis**
- **Primary Sclerosing Cholangitis is an autoimmune disease that predominantly occurs in males**
- **Primary Sclerosing Cholangitis** is associated with inflammatory bowel disease in 70% cases.
- Prevalence of PSC in persons with ulcerative colitis is about 4%

### 71. Ans. (b) ANCA (Ref: Robbins 9th/pg 839-840)

### 72. Ans. (e) Roter's syndrome

### 73. Ans. (a) Increase of serum bilirubin, c. Increase alkaline phosphatase e. Stool sterocobilinogen absent

(Ref: PJM 20th/20; Harrison 19th/281; CMDT 2016/665)

### 74. Ans. (a) Jaundice becomes severe with time.

### 75. Ans. (d) Primary sclerosing cholangitis

The clinical feature of obstructive jaundice (jaundice, pruritis, raised bilirubin and alkaline phosphatase) along with Inflammatory bowel disease in an adult male is suggestive of Primary sclerosing cholangitis.

### 76. Ans. (a, b, d, e) a. More common in female; b. Periportal fibrosis; d. Jaundice may be present; e. Autoimmune disease are seen

(Ref: Robbins 9th/pg 858; 8th/pg 867)

### 77. Ans. (c) Primary biliary cirrhosis

(Ref: Robbins 9th/pg 858; 8th/pg 867)

**In Primary biliary cirrhosis:** Interlobular bile ducts are actively destroyed by lymphoplasmacytic inflammation with or without granulomas (**the florid duct lesion**)

### 78. Ans. (c) Micronodular cirrhosis of liver

(Ref: Robbins 9th/pg 859; 8th/pg 869)

### 79. Ans. (c) Primary sclerosing cholangitis

(Ref: Robbins 9th/pg 859; 8th/pg 869)

### 80. Ans. (a) ABO incompatibility

(Ref: Robbins 9th/pg 852)

**ABO incompatibility causes jaundice due to hemolysis and not cholestasis**

### 81. Ans. (c) Crigler Najjar type II (Ref: R 9th/pg 853-854)

### 82. Ans. (c) Primary biliary cirrhosis

(Ref: Robbins 9th/pg 859)

### 83. Ans. (c) Rotor syndrome (Ref: Robbins 9th/pg 853)





84. Ans. (a) **Causes cirrhosis** (Ref: Harrison 18th/Chapter 303)

#### Gilbert Syndrome

Feature	Gilbert's Syndrome
<b>Inheritance (all autosomal)</b>	Promoter mutation: recessive Missense mutations: 7 of 8 dominant; 1 reportedly recessive
<b>Total serum bilirubin</b>	< 4 mg/dL in absence of fasting or hemolysis
<b>Routine liver tests</b>	Normal
<b>Hepatic histology</b>	Usually normal; increased lipofuscin pigment in some

85. Ans. (c) **Gilbert syndrome**

(Ref: Robbins 9th/pg 853-854)

86. Ans. (b, d, e); **b. Fat soluble; d. Affinity for brain tissue; e. Increased in hemolytic anemia**

(Ref: Robbins 9th/pg 852)

- Unconjugated bilirubin is **increased in hemolytic anemia**
- It is insoluble in water but is **fat soluble, so it can cross the blood brain barrier**
- It is **not filtered through the glomerulus, so does not appear in urine.**
- **Unconjugated bilirubin bound to albumin** is transported to the liver, where it is **taken up by hepatocytes** via a **carrier-mediated** membrane transport.

87. Ans. (a) **Ca bilirubinate**

(Ref: Robbins 9th/pg 877)

Two main types of gallstones:

- **Cholesterol stones (most common type in West)** containing crystalline cholesterol monohydrate.
- **Pigment stones** composed mainly of calcium bilirubinate; contain <20% cholesterol and are classified into "black" & "brown" types, the latter forming secondary to chronic biliary infection.

88. Ans. (a) **Ulcerative colitis** (Ref: Robbins 9th/pg 859-860)

- **PSC** is commonly seen in **association with inflammatory bowel disease** particularly **chronic ulcerative colitis**, which coexists in approximately **70%** of individuals with primary sclerosing cholangitis.
- Conversely, the **prevalence of PSC in persons with ulcerative colitis is about 4%.**

89. Ans. (a) **Dubin Johnson Syndrome**

(Ref: R 9th/pg 853-854)

90. Ans. (c) **Dubin Johnson Syndrome**

(Ref: R 9th/pg 853-854)

91. Ans. (d) **Fraction of coproporphyrin I in urine is elevated usually more than 80% of the total in Rotor syndrome**

(Ref: Harrison 18th/Chapter 303)

#### Differentiation between Dubin-Johnson syndrome & Rotor syndrome:

Characteristic	Dubin-Johnson syndrome	Rotor syndrome
<b>Pigmentation of Liver</b>	Pigmented cytoplasmic globules	Not seen; normal histology
Total urinary coproporphyrin excretion	Normal	Increased
Fraction of coproporphyrin I/III in urine	<b>&gt; 80% of total<sup>a</sup></b>	<b>&lt;70% of the total<sup>a</sup></b>
BSP clearance	<b>Normal<sup>a</sup>, with reflux</b>	<b>Delayed<sup>a</sup>, No reflux</b>

92. Ans. (a, b, e) **a. May be complicated by bacterial infection; b. Rarely progress to biliary cirrhosis; e. Narrowing of bile duct**

(Ref: Robbins 9th ed p 860)

PSC is characterized by inflammation and obliterative fibrosis of intrahepatic and extrahepatic bile ducts with dilation of preserved segments. Inflammatory bowel disease like ulcerative colitis is seen in 70% of individuals with PSC.

93. Ans. (b) **Anti-mitochondrial antibody**

(Ref: Robbins 9th/pg 859-860; 8th/pg 869)

94. Ans. (a, c, d), **a. Cholesterol stones are most common; c. Mirizzi syndrome is due to impaction of stone in hartmann's pouch; d. Hemolytic anaemia cause black colored stone**

(Ref: Robbins 9th/pg 876-877; 8th/pg 868; Harrison 18th/pg chapter 311)

a.	True	90% of gallstones are <i>cholesterol stones</i>
b.	False	10–15% of cholesterol stones and 50% of pigment stones are radiopaque
c.	True	In Mirizzi syndrome, a gallstone becomes impacted in cystic duct or neck of the gallbladder ( <b>hartmann's pouch</b> ) → compression & obstruction of CBD → jaundice.
d.	True	Pigment stones seen in Hemolytic anemia are black or brown in colour
e.	False	Gallstones are associated with an increased risk of gallbladder carcinoma

95. Ans. (a) **Most commonly caused by gram positive organisms** (Ref: Harrison 18th/Chapter 42)

#### Ascending cholangitis

- It is **most commonly caused by gram negative organisms** like *E. coli*, *Klebsiella* spp
- **Obstruction of bile duct by stone** is an important predisposing factor



- Clinical presentation: **Jaundice** associated with the **sudden onset of severe right upper quadrant pain and shaking chills; circulatory collapse** can occur in severe cases
- **IV antibiotics, Urgent removal of stone by ERCP or Cholecystectomy** can be done

**96. Ans. (c, d, e); c. Pruritus; d. Pale stools; e. Icterus**

(Ref: Harrison 18th/Chapter 42)

**Discussing options about obstructive jaundice one by one:**

a.	False	Conjugated Hyperbilirubinemia is seen
b.	False	Positive direct Van den Bergh for conjugated bilirubin
c.	True	Pruritus occurs as bile salts irritates skin
d.	True	Absence of bilirubin excretion causes pale stools
e.	True	Icterus is a yellowish discoloration of tissue resulting from deposition of bilirubin; Scleral icterus indicates a serum bilirubin of at least 3 mg/dL

**97. Ans. (c) The conjugation process of bilirubin in liver remains operative without any interference**

(Ref: Harrison 18th/Chapter 42)

In post-hepatic jaundice, the concentration of conjugated bilirubin in the blood is higher than that of unconjugated bilirubin because:

- **The conjugation process of bilirubin in liver remains operative without any interference.**
- But there is **decreased excretion** of conjugated bilirubin into the bile ductules & **backward leakage** of the pigment into the circulation

**98. Ans. (b) Increased urinary copper excretion; (d) Increased liver copper content** (Ref: R 9th pg 847)

**99. Ans. (a) Alpha1 Antitrypsin deficiency**

(Ref: Robbins 9th/pg 847-851; 8th/pg 861-864; Refer to pretexts)

**100. Ans. (c) Alpha 1 antitrypsin deficiency**

(Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**101. Ans. (a) Iron** (Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**102. Ans. (c) 13** (Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**103. Ans. (d) Low ↑ Ceruloplasmin high urine copper**

(Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**104. Ans. (b) Hemochromatosis** (Ref: Robbins 9th/pg 847-851)

**105. Ans. (b) Autosomal recessive**

(Ref: Robbins 9th/pg 847-851)

**106. Ans. (d) Alpha-1-Antitrypsin deficiency**

(Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**107. Ans. (b) 13** (Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**108. Ans. (a) PAS + ve diastase resistant**

(Ref: R 9th/pg 847-851)

**109. Ans. (a) Iron** (Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**110. Ans. (b) 250** (Ref: Robbins 9th/pg 847-851; 8th/pg 861-864)

**111. Ans. (d) HFE gene** (Ref: Robbins 9th/pg 847-851)

**112. Ans. (d) α-1-antitrypsin deficiency** (Ref: R 9th/pg 847-851)  
**Cirrhosis can occur due to alpha1 antitrypsin deficiency**

**113. Ans. (d, e) d. Opisthorchis viverrini; e. Clonorchis sinensis**

**114. Ans. (c) Portal hypertension seen in 50% of patients**

**115. Ans. (a) Hemangiosarcoma**

**116. Ans. (e) Need surgical removal in every case because chances of rupture is high**

**117. Ans. (a) Fibrolamellar Variant of HCC**

**118. Ans. (a) Better prognosis than typical hepatocellular carcinoma**

(Ref: Robbins (SEA) 9th/873; Harshmohan 7th/620; Harrison 19th/552)

**119. Ans. (b) More common in male**

(Ref: Robbins (SEA) 9th/867)

**120. Ans. (b) Squamous cell carcinoma is 40% of all cases**

(Ref: Robbins 9th pg 879-880)

**Morphology of Gall bladder Ca is mostly Adenocarcinoma<sup>Q</sup> > Squamous cell Ca (5% cases)**

**121. Ans. (c) Starts with macronodular and later on changes to micronodular cirrhosis.**

(Ref: Robbins 9th pg 842-844)

The three distinctive forms of alcoholic liver injury: (1) hepatocellular steatosis or fatty change, (2) alcoholic (or steato-) hepatitis, and (3) steatofibrosis (patterns of scarring typical for all fatty liver diseases including alcohol) up to and including cirrhosis in the late stages of disease.

Iron overload and infections with HIV, HCV and HBV synergize with alcohol, leading to increased severity of liver disease.



**122. Ans. (b) Tyrosenemia** (Ref: Robbins 9th/pg 870)

Among the given options, Tyrosenemia is the least common cause of HCC, so is the answer.

**123. Ans. (c) Vinyl chloride** (Ref: Robbins 9th/pg 870)

- **Angiosarcoma of liver** is associated with **vinyl chloride, arsenic or Thorotrast exposure** and has poor prognosis

**124. Ans. (b) Young females** (Ref: Robbins 9th/pg 870)

**125. Ans. (d) Fibrolamellar carcinoma**

(Ref: R 9th/pg 870)

Fibrolamellar Ca has a good prognosis while others behave as Malignant tumors.

**126. Ans. (b) High risk of malignant transformation**

(Ref: Robbins 9th/pg 870)

Features of **HNF1- $\alpha$  inactivated adenomas**

- Common in feamles
- Virtually **no risk of malignant transformation**,
- Often associated with **OCP** use
- In individuals with **MODY-3**

**127. Ans. (b) Mature hepatocytes present** (Ref: R 9th/pg 870)

Histology of Hepatoblastoma can be of 2 types:

- **Epithelial type**, composed of small polygonal fetal cells or smaller embryonal cells forming acini, tubules, or papillary structures
- **Mixed epithelial and mesenchymal type**, which contains foci of mesenchymal differentiation that may consist of primitive mesenchyme, osteoid, cartilage, or striated muscle.

*Mature appearing hepatocytes are absent.*

**128. Ans. (d) Hepatitis D** (Ref: Robbins 9th/pg 838; 8th/pg 853)

**129. Ans. (a) Hepatocellular Ca** (Ref: Robbins 9th/pg 870)

**130. Ans. (b) Ulcerative colitis** (Ref: Robbins 9th/pg 874)

**Risk Factors for cholangiocarcinoma:**

- Chronic inflammation eg Ulcerative colitis
- Cholestasis
- Primary sclerosing Cholangitis
- Liver flukes (particularly Opisthorchis and Clonorchis species)
- Hepatitis B and C
- Non-alcoholic fatty liver disease

**131. Ans. (a) Liver** (Ref: Robbins 9th/pg 875; 8th/pg 856)

- Oral contraceptives have been implicated in the development of hepatic adenoma and, rarely, hepatocellular carcinoma and hepatic vein occlusion (Budd-Chiari syndrome).

**132. Ans. (a, d, e) a. Better prognosis than Primary HCC; d. Underlying cirrhosis not a risk factor; e. Neurotensin is a biomarker** (Ref: Robbins 9th/pg 873; 8th/pg 879)

**133. Ans. (a) Methotrexate**

(Ref: Robbins 9th/pg 841; 8th/pg 856)

**134. Ans. (c) Angiosarcoma**

(Ref: Robbins 9th/pg 841; 8th/pg 856)

**135. Ans. (b) Primary sclerosing cholangitis**

(Ref: Robbins 9th/pg 874; 8th/pg 880)

**136. Ans. (a) Nodular type of cholangiocarcinoma**

(Ref: Robbins 9th/pg 874; 8th/pg 880)

**(Klatskin tumor)** is Nodular type of cholangiocarcinoma at the junction of the right and left hepatic ducts.

**Intrahepatic-10%.**

**137. Ans. (a, b) a. Lipase; b. Amylase**

(Ref: Robbins 9th/pg 884; 8th/pg 893)

**Lipase and Amylase are raised in pancreatitis rather than liver disorder.**

**138. Ans. (a, b, c, d); a. Raised in testicular tumor; b. Raised in 50-70% cases of HCC; c. Correlation between tumor recurrence after surgery in HCC; d. Correlation with HCC size** (Ref: Harrison 18th/Chapter 92)

AFP:

- Normal Range of AFP in adults: 0–8.5 ng/mL
- **It is a tumor marker for Liver Cancers**, **non-seminomatous<sup>Q</sup>** germ cell tumors of testis
- **Liver diseases with elevated AFP are:** Tumors like **Hepatocellular Carcinoma,<sup>Q</sup> Hepatoblastoma, Infantile hemangioendothelioma** & non-neoplastic lesions like **Amebic liver abscess<sup>Q</sup> & Hepatitis<sup>Q</sup>**
- Adverse prognostic factors in HCC include ascites, jaundice, vascular invasion, and elevated AFP; AFP correlates with HCC size
- **Postoperative AFP level** is a useful tool for **predicting recurrence after curative hepatectomy.**
- A positive level of AFP after operation might suggest a site of residual viable cancer.

**139. Ans. (c) Arises from cirrhotic liver** (Ref: R 9th/pg 873)

**140. Ans. (b) Cholangiocarcinoma** (Ref: Robbins 9th/pg 874)

**141. Ans. (b) Liver** (Ref: Robbins 9th/pg 862; 8th/pg 869)

**Von Meyenburg Complexes**

- Clusters of **dilated bile ducts** embedded in a fibrous, sometimes hyalinized, stroma located close to or within portal tracts.
- These lesions are often referred to as “**bile duct hamartomas**”
- They are without clinical significance except in the **differential diagnosis of metastases to the liver**



**142. Ans. (d) Hepatitis-B virus**

(Ref: Robbins 9th/pg 871-873)

Hepatitis-B & C virus are important risk factors of HCC

**143. Ans. (a) Adenomyomatosis of gall bladder**

(Ref: Robbins 9th/pg 879; 8th/pg 886)

**Rokitansky-Aschoff sinuses<sup>Q</sup>** are Outpouchings of the mucosal epithelium through the wall seen in adenomyomatosis of gall bladder

**144. Ans. (b) Porcelain gallbladder**

(Ref: Robbins 9th/pg 879)

**Porcelain Gallbladder:**

- **Calcium salt deposition within the wall of a chronically inflamed gallbladder**
- May be detected on the plain abdominal film.
- It is **associated with the development of carcinoma** of the gallbladder
- Cholecystectomy is advised in all patients with porcelain gallbladder

**Limey Bile or Milk of Calcium Bile**

- Calcium precipitation & diffuse, hazy opacification of bile due to calcium salts in the lumen of the gallbladder
- Produce or a layering effect on plain abdominal roentgenography.
- Usually clinically innocuous, but cholecystectomy is recommended, especially when it occurs in a hydropic gallbladder.

**145. Ans. (d) Islet cell hypertrophy**

(Ref: Robbins 9th/pg 884)

**146. Ans. (a) KRAS** (Ref: Robbins 9th/pg 892-894)

Gene	Chr	Percentage	
<b>KRAS</b>	12p	90	<b>Most commonly involved Oncogene<sup>Q</sup></b>
<b>p16/CDKN2A</b>	9p	95	<b>Most commonly involved Tumor suppressor gene<sup>Q</sup></b>
TP53	17p	50–70	Involved in Response to DNA damage
SMAD4	18q	55	TGF $\beta$ pathway
<b>BRCA2</b>	13q	10	<b>Germ-line mutation<sup>Q</sup></b>

**147. Ans. (a) Head** (Ref: Robbins 9th/pg 892-894; 8th/pg 900-903)

**Pancreatic Carcinoma is**

Site of	Pancreatic Head (most common) <sup>Q</sup> > Diffuse > Body > Tail
---------	--

**148. Ans. (a) KRAS** (Ref: Robbins 9th/pg 892-894)

**149. Ans. (a) Intraductal papillary mucinous neoplasms**

(Ref: Robbins 9th/pg 892-894; 8th/pg 900-903)

<b>A. Intraductal papillary mucinous neoplasms</b>	<b>IPMNs progress to an invasive cancer</b>
B. Solid-pseudopapillary neoplasms	Usually benign
C. Serous cystic neoplasms	Always benign
D. Mucinous cystic neoplasms	1/3 <sup>rd</sup> associated with invasive adenocarcinoma



[illegible]This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Renal System and its Disorders

## Key Points

- » **Hematuria** is excretion of **intact RBCs >3/hpf** in urine
- » **Sterile pyuria**: presence of **elevated numbers of pus cells (WBCs)** in sterile urine (seen in Tuberculosis)
- » Most common cause of **isolated glomerular hematuria** is **IgA nephropathy**
- » **Mesangial cells** are **mesenchymal origin**, are **contractile, phagocytic**, and **capable of proliferation**
- » **Crescents** are seen in **RPGN** and suggest **poor prognosis**
- » **Microalbuminuria**: Excretion of **30–300 mg/day** of albumin in urine or **30–300 mg/g** of creatinine in urine
- » **Focal Segmental Glomerulosclerosis (FSGS)** is the **most common cause of nephrotic syndrome in adults**
- » **Minimal change disease** is the **most frequent cause of nephrotic syndrome in children**
- » **Most specific histological lesion in diabetic nephropathy** is **Kimmelsteil-Wilson lesions**
- » **Thyroidization of tubules** is a **feature of Chronic pyelonephritis**
- » Major Causes of Papillary Necrosis is **analgesic nephropathy**
- » **Clear cell Ca** is the **most common histological subtype of Renal cell Carcinoma (RCC)**
- » Polycythemia and Hypertension is the **most common paraneoplastic feature of RCC**
- » **Michaelis-Gutmann bodies** are seen in **Malacoplakia**


## Key Recent Updates

- » MPGN type II is now classified as C3 glomerulopathy
- » Fibronectin glomerulopathy occurs due to mutation of FN<sub>1</sub> gene on chromosome 2q34.



## CLINICAL MANIFESTATIONS OF RENAL DISEASES

- **Hematuria:** Excretion of **intact RBCs** > 3/ hpf in urine<sup>Q</sup>
- **Pyuria:** Presence of > 5 **pus cells/hpf**<sup>Q</sup> in urine, typically from bacterial infection
- **Sterile pyuria**<sup>Q</sup>: Presence of **elevated numbers of pus cells** (WBCs) in urine which is **sterile** using standard culture techniques
- **Sterile pyuria** is seen in Renal tuberculosis
- **Oliguria**<sup>Q</sup>: 24-h urine output <400 mL<sup>Q</sup>, usually due to underlying renal failure.
- **Anuria:** Complete **absence of urine formation** (<100 mL).<sup>Q</sup> It can be caused by **total urinary tract obstruction, total renal artery or vein occlusion, shock, cortical necrosis, ATN and RPGN.**<sup>Q</sup>
- **Polyuria:** 24-h urine output >3 L<sup>Q</sup>
- **Azotemia:** Increased serum levels of **nitrogenous waste products** like **urea and creatinine**<sup>Q</sup>
- **Uremia:** Azotemia along with clinical manifestations<sup>Q</sup> due to deranged renal function.



Normal urine microscopy

### Urinary Casts (Refer Annexures for Images and Types)

- **Formed elements of urine**<sup>Q</sup> that have **kidney as their sole site of origin**<sup>Q</sup>
- **Tamm-Horsfall protein**, secreted from the **thick ascending loop of Henle**<sup>Q</sup> forms the **matrix of all casts**<sup>Q</sup>

- **Width** of the cast depends on the **size of the tubule**<sup>Q</sup> in which it was formed
- Cast formation **increases with lower pH, increased ionic concentration and stasis** in nephrons<sup>Q</sup>




### High Yield Facts

- Eosinophils in the urine suggests **allergic interstitial nephritis** or **atheroembolic renal disease.**<sup>Q</sup>
- **Urinary Dipsticks** detect **only albumin as urinary protein**<sup>Q</sup>
- **Dipstick** gives **false-positive** results for:
  - Albumin: when pH >7.0, urine is very concentrated or contaminated with blood.<sup>Q</sup>
  - Hematuria: when myoglobinuria is present, as in rhabdomyolysis.<sup>Q</sup>
- RBCs of **glomerular origin** are often **dysmorphic**<sup>Q</sup>
- Most common cause of **isolated glomerular hematuria** is **IgA nephropathy**<sup>Q</sup>


## GLOMERULUS AND GLOMERULAR DISEASES

### Structure of Glomerular Filtering Membrane

- **Mesangium**
- **Visceral epithelial cells (podocytes)**<sup>Q</sup>:
  - **20-30-nm**-wide filtration slits → size selective barrier
  - Responsible for synthesis of GBM components
- **Glomerular basement membrane (GBM):**
  - Composed of collagen **type 4** (or COL4  $\alpha 1$  to  $\alpha 6$ )<sup>Q</sup>
  - Each molecule consists of a **7S domain** at the **N terminus**, a triple-helical domain in the middle, and a globular **noncollagenous domain (NC1)** at the C terminus.
  - Other components are: **Laminin, polyanionicproteoglycans (negatively charged), mostly heparan sulfate, fibronectin, entactin** and several other glycoproteins.
- **Fenestrated endothelial cells** (70-100 nm)<sup>Q</sup>.



Structure of glomerulus



Types of glomerular deposits



## PATHOLOGIC RESPONSES OF THE GLOMERULUS TO INJURY


Site of glomerular deposits	Type of Glomerulonephritis	Terminologies used in Kidney Biopsy	
<b>Subepithelial deposits</b>	<ul style="list-style-type: none"> <li>PSGN</li> <li>Membranous GN</li> <li>RPGN</li> <li>Heymann Nephritis</li> </ul>	<b>Terminology</b>	<b>Description</b>
<b>Subendothelial deposits</b>	<ul style="list-style-type: none"> <li>Lupus nephritis</li> <li>MPGN-I</li> </ul>	<b>Diffuse</b>	Involving <b>&gt;50%</b> of the glomeruli in the kidney <sup>Q</sup>
<b>Membranous deposits</b>	<ul style="list-style-type: none"> <li>MPGN II</li> </ul>	<b>Global</b>	Involving the glomerulus <b>completely</b> <sup>Q</sup>
<b>Mesangial deposits</b>	<ul style="list-style-type: none"> <li>IgA nephropathy</li> <li>HSP</li> </ul>	<b>Focal</b>	Involving <b>&lt;50%</b> of the glomeruli in the kidney
		<b>Segmental</b>	Affecting <b>a part of each glomerulus</b> <sup>Q</sup>
		<b>Capillary loop Mesangial</b>	Affecting predominantly <b>capillary or mesangial regions</b> <sup>Q</sup>




### High Yield Facts

- Mesangial cells<sup>Q</sup>- **mesenchymal origin**, are contractile<sup>Q</sup>, phagocytic<sup>Q</sup>, and capable of proliferation<sup>Q</sup>
- Mesangial cells and mesangial matrix support the glomerular tuft
- Glomerular filtering membrane is a size- and charge-dependent barrier
- Neutral substance  $\leq 4A$  can get freely filtered through GBM
- GBM** is **negatively charged** due to **sialo-glycoprotein**
- All proteins (negatively charged) are normally repelled by GBM
- Albumin** (smallest molecular wt, 70 kD) is the **1<sup>st</sup> protein to appear in urine in glomerulonephritis**
- AN**ionic antigens form sub**EN**dothelial deposits
- Cationic** antigens form sub**EP**ithelial deposits
- Neutral** antigens form **mesangial** deposits

## PATHOGENESIS OF GLOMERULAR INJURY










## Electron Microscopy

Subepithelial	Intramembranous	Subendothelial	Mesangial	Combined Subendothelial, Subepithelial and Mesangial
Membranous GN	Dense-deposit disease	MPGN C3GN	IgA nephropathy	Lupus (WHO classes III and IV) C3GN
Lupus (WHO class V)	GN related to endocarditis, deep-seated abscesses	Lupus (WHO class III and IV)	Henoch-Schonlein purpura	MPGN type III
Postinfectious GN	Other infections	Cryoglobulinemic GN (microglobular structure)	Lupus (WHO class II) Clq nephropathy Rare other forms of mesangioproliferative GN	GN related to infections. Proliferative GN with monoclonal IgG deposit



## Immunofluorescence

Deposition Pattern	Granular (Probably Immune Complex)	Linear (classic Antiglomerular Basement Membrane [anti-GBM] Antibodies)
		
Region of deposition	Capillary wall—membranous nephropathy	Mesangium-Berger disease (IgA nephropathy)
Class of Ig and fraction of complement	<b>Full house</b> of Igs-systemic lupus erythematosus (SLE)	IgA-Berger disease (IgA nephropathy)



## NEPHRITIC SYNDROME

- **Definition:** Sudden onset of **gross hematuria, oliguria, nephritic range proteinuria, mild edema, hypertension, and renal insufficiency**<sup>Q</sup>
- Characterized by **inflammation** in the **glomeruli**<sup>Q</sup>

### Difference between nephritic and nephrotic syndrome

Characteristic	Nephritic Syndrome	Nephrotic Syndrome
<b>Hematuria and RBC casts</b>	Present	Absent/few
<b>Proteinuria</b>	Nephritic range (<3.5g/day) <sup>Q</sup>	Nephrotic range (>3.5g/day) <sup>Q</sup> (>40 mg/m <sup>2</sup> /hr)
<b>Hypertension</b>	Present <sup>Q</sup>	Uncommon
<b>Uremia</b>	Present <sup>Q</sup>	Absent
<b>Oliguria</b>	Present <sup>Q</sup>	Absent


## POSTSTREPTOCOCCAL GLOMERULONEPHRITIS (PSGN)

- **Time course:** 1 to 4 weeks after a streptococcal infection (**sore throat or pyoderma**)<sup>Q</sup>
- **Age group:** most frequently, **children 6 -10 years**<sup>Q</sup>
- **Inciting agent:** **group A β-hemolytic streptococci**<sup>Q</sup> – nephritogenic strains (**12, 4, and 1**)<sup>Q</sup>
- **Complement levels** **C3 decrease** **Anti-Streptococcal-O (ASO) increases**, normal levels of C4



### High Yield Facts

- In PSGN, Serum complement levels (C3): **transiently low**<sup>Q</sup>
- **Principal antigen in PSGN:** streptococcal pyogenic exotoxin B (Spe B)<sup>Q</sup>
- **Early treatment** of sore throat/pyoderma with antibiotics **does not**<sup>Q</sup> prevent PSGN
- Deposits seen in PSGN<sup>Q</sup>: **Subepithelial humps**, Subendothelial, mesangial



### Prognosis

#### In children:

- Self-recovery is seen in **95% (Good prognosis)**<sup>Q</sup>
- **1% become severely oliguric, and develop RPGN**<sup>Q</sup>
- **5% undergo slow progression to chronic glomerulonephritis**<sup>Q</sup>

### Morphology

Light Microscopy	Immunofluorescence Microscopy	Electron Microscopy
<ul style="list-style-type: none"> <li>• Hypercellular glomeruli</li> <li>• Infiltration by WBCs,</li> <li>• Proliferation of endothelial and mesangia cells: <b>Endo- and exocapillary proliferation</b><sup>Q</sup></li> <li>• <b>Crescents:</b> Severe cases<sup>Q</sup></li> </ul>	Granular deposits of <b>IgG, and C3 in mesangium</b> <sup>Q</sup> and along GBM	Discrete, amorphous, electron-dense <b>SUBEPITHELIAL deposits ("Humps")</b> <sup>Q</sup>

## RAPIDLY PROGRESSIVE GLOMERULONEPHRITIS (RPGN)

- Characterized by severe glomerular injury (**crescents**) leading to **rapid and progressive loss of renal function** associated with **severe oliguria** and signs of nephritic syndrome.<sup>Q</sup>
- If untreated, **death** from renal failure occurs **within weeks to months**.<sup>Q</sup>



## Types of RPGN

Entity	Type I (20%)	Type II (25%)	Type III (55%)
<b>Mechanism</b>	Anti-GBM Antibody	Immune Complex	<b>Pauci-immune, c-ANCA/p-ANCA mediated</b>
<b>Etiology</b>	Renal limited <b>Good pasture syndrome<sup>Q</sup></b> (Serum antibodies against <b>alpha 3 NC1 domain of collagen – IV</b> )	<ul style="list-style-type: none"> <li>• <b>Postinfectious</b> <ul style="list-style-type: none"> <li>■ Poststreptococcal glomerulonephritis<sup>Q</sup></li> <li>■ Bacterial endocarditis<sup>Q</sup></li> </ul> </li> <li>• <b>Noninfectious</b> <ul style="list-style-type: none"> <li>■ SLE<sup>Q</sup>, HSP<sup>Q</sup></li> <li>■ Mixed cryoglobulinemia<sup>Q</sup></li> </ul> </li> <li>• <b>Primary Renal Disease</b> <ul style="list-style-type: none"> <li>■ MPGN<sup>Q</sup></li> <li>■ IgA nephropathy<sup>Q</sup></li> </ul> </li> </ul>	<b>ANCA-associated</b> <ul style="list-style-type: none"> <li>• Idiopathic</li> <li>• <b>Granulomatosis with polyangiitis (Wegener granulomatosis)<sup>Q</sup></b></li> <li>• Microscopic polyangiitis<sup>Q</sup></li> <li>• Hypersensitivity vasculitis<sup>Q</sup></li> </ul>
<b>Grossly</b>	Kidneys are enlarged and pale, often with <b>petechial hemorrhages</b> on the cortical surfaces. ( <b>FLEA-BITTEN KIDNEY</b> ) <sup>Q</sup>		
<b>Light m/e</b>	<ul style="list-style-type: none"> <li>• <b>Glomeruli: Crescents</b> are Hallmark<sup>Q</sup></li> <li>• Focal and segmental necrosis<sup>Q</sup>, endothelial and mesangial proliferation<sup>Q</sup></li> <li>• <b>Pauci-immune: Segmental glomerular necrosis</b> is characteristic<sup>Q</sup></li> </ul> <div data-bbox="426 761 1404 1117" data-label="Image"> <p><b>Crescents- Formed by</b></p> <ul style="list-style-type: none"> <li>• Proliferation of <b>parietal cells</b><sup>Q</sup></li> <li>• Infiltration by <b>WBCs</b><sup>Q</sup></li> <li>• <b>Fibrin strands</b>.<sup>Q</sup></li> </ul> <p>Crescents obliterate the urinary space and compress the glomerular tuft, hence <b>More the number of crescents → poorer the prognosis</b><sup>Q</sup></p> <p>Crescentic glomerulonephritis</p> </div>		
<b>Immunofluorescence m/e</b>	<b>Linear GBM fluorescence<sup>Q</sup></b> <p><b>Type I RPGN (Linear)</b></p>	<b>Granular immune deposits<sup>Q</sup></b> <p><b>Type II RPGN (Granular)</b></p>	<b>No deposition of immune reactants<sup>Q</sup></b>  <b>No Deposits seen</b>  <b>(Pauci-immune)</b>
<b>Electron m/e</b>	Ruptures in the GBM <sup>Q</sup> may be present, Type II shows immune complex deposits		

## Clinical Course


- **Hematuria with RBC casts<sup>Q</sup>** in the urine, variable proteinuria, hypertension and edema.
- **Progressive over weeks** and ends in **severe oliguria/renal failure**.<sup>Q</sup>
- **Goodpasture syndrome: Recurrent hemoptysis** or even life-threatening pulmonary hemorrhage (**Necrotizing hemorrhagic interstitial pneumonitis**).<sup>Q</sup>

## NEPHROTIC SYNDROME

### The Manifestations of Nephrotic Syndrome Include

- **Massive proteinuria (>3.5 g/day)<sup>Q</sup>** (>40 mg/m<sup>2</sup>/hr)
- **Hypoalbuminemia<sup>Q</sup>** (plasma albumin <3 g/dL)
- **Generalized edema** (due to **loss of oncotic pressure<sup>Q</sup>** > **sodium and water retention**)<sup>Q</sup>





- **Hyperlipidemia:**<sup>Q</sup> Increased synthesis of lipoproteins in the liver, abnormal transport of circulating lipid particles, and decreased lipid catabolism.
- **Lipiduria** (free fat or as oval fat bodies in urine)<sup>Q</sup>

#### Remember: Proteinuria can be

- **Selective proteinuria:** → Initially; consists mostly of **low-molecular-weight proteins** (albumin, 70 kD; transferrin, 76 kD)
- **Nonselective proteinuria:** → Later, in advanced disease, consists of higher molecular-weight globulins and albumin.

#### Causes of Nephrotic Syndrome

Primary Glomerular Disease	Systemic Diseases
<ul style="list-style-type: none"> <li>• Membranous glomerulonephropathy (MGN)<sup>Q</sup></li> <li>• Minimal-change disease (MCD)<sup>Q</sup></li> <li>• Focal segmental glomerulosclerosis (FSGS)<sup>Q</sup></li> <li>• Membranoproliferative glomerulonephritis (MPGN)</li> <li>• IgA nephropathy<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Diabetes mellitus<sup>Q</sup>, Amyloidosis<sup>Q</sup></li> <li>• Systemic lupus erythematosus (SLE)<sup>Q</sup></li> <li>• Drugs (NSAIDs, penicillamine, Lithium, Pamidronate, heroin injection)<sup>Q</sup></li> <li>• Infections (hepatitis B and C, HIV, malaria, toxoplasmosis, syphilis)<sup>Q</sup></li> <li>• Malignant disease (carcinoma, lymphoma)<sup>Q</sup></li> <li>• Hereditary nephritis, Renal vein thrombosis)</li> </ul>



#### High Yield Facts

- **Microalbuminuria:** Excretion of **30–300 mg/day<sup>Q</sup>** of albumin in urine or **30–300 mg/g of creatinine in urine<sup>Q</sup>**
- **Macroalbuminuria:** Excretion of **300–3500 mg/day** of albumin in urine<sup>Q</sup>
- **Nephrotic range proteinuria:** Excretion of **> 3.5 g/day<sup>Q</sup>** of albumin in urine
- **Selective proteinuria:** Selective excretion of **low-molecular-weight proteins** like **albumin and transferrin<sup>Q</sup>** by the kidney.


## MEMBRANOUS NEPHROPATHY

- **Characterized by:** Diffuse thickening of glomerular capillary wall due to accumulation of deposits containing immune complex deposits along **sub-epithelial side** of basement membrane.<sup>Q</sup>
- **Etiology:**
  - **Primary: 75%:** Associated with **HLA-DQA1<sup>Q</sup>**, Autoantigen: **phospholipase A2 receptor<sup>Q</sup>**
  - **Secondary: 25%**
    - Infections: Chronic hepatitis B, hepatitis C, Syphilis, Schistosomiasis, Malaria<sup>Q</sup>
    - Drugs: Penicillamine, Captopril, Gold, NSAIDs<sup>Q</sup>
    - Carcinoma lung, colon, and melanoma


- **Autoimmune diseases:** SLE (10–15%), Rheumatoid Arthritis, Primary biliary cirrhosis, Dermatitis herpetiformis
- **Systemic diseases:** Fanconi's syndrome, sickle cell anemia, diabetes, **Crohn's disease**, Sarcoidosis,

#### Morphology:


- **Light microscopy:** Uniform, diffuse **thickening of the glomerular capillary wall.**<sup>Q</sup>
- **Immunofluorescence microscopy:** **Granular/Lumpy bumpy<sup>Q</sup>** electron dense immune complexes deposits
- **Electron microscopy:** **Granular deposits, (Ig + complement)**
  - Effacement of podocyte foot processes<sup>Q</sup>
- **On Silver methenamine stain-** prominent **“spikes”** and **“domes”<sup>Q</sup>** of silver-staining matrix



Normal capillary wall



Thickened capillary wall



Spikes and domes on silver stain

#### Clinical course:

- **Persistent proteinuria** in **60%** of patients<sup>Q</sup>
- **40%** develop severe CKD or ESRD<sup>Q</sup>
- **40% recurs in patients who undergo transplantation<sup>Q</sup>**






### High Yield Facts

- Membranous nephropathy is the most frequent cause of nephrotic syndrome in elderly<sup>Q</sup>
- Focal segmental glomerulosclerosis (FSGS) is the most common cause of nephrotic syndrome in adults<sup>Q</sup>
- Minimal change disease is the most frequent cause of nephrotic syndrome in children<sup>Q</sup>
- Mutation in NPHS2 is the most common cause of Steroid Resistant Nephrotic syndrome<sup>Q</sup>


## MINIMAL CHANGE DISEASE

- **Characterized by:** Absence of immune deposits but has an immunologic basis.<sup>Q</sup>
- **Epidemiology:** Most common age group involved: 2 to 6 years<sup>Q</sup>
- **Etiology:** Idiopathic
- **Morphology:**
  - **Light microscopy:** Glomeruli appear normal<sup>Q</sup>
  - **Immunofluorescence microscopy:** No Ig/ complement deposits
  - **Electron microscopy:** Diffuse effacement of foot processes of podocytes ("podocytopathy")<sup>Q</sup>
    - No electron-dense deposits<sup>Q</sup>
  - **Proximal tubules** cells get laden with lipid and protein due to tubular reabsorption of lipoproteins: **Lipoid nephrosis**<sup>Q</sup>
- **Clinical course:**
  - **Excellent prognosis:** >90% respond rapidly to steroids<sup>Q</sup>
  - Renal function remains normal<sup>Q</sup>
  - No hypertension or hematuria.<sup>Q</sup>
  - Selective proteinuria is seen



## FOCAL SEGMENTAL GLOMERULOSCLEROSIS (FSGS)

- **Hallmark** Disruption of visceral epithelial cells with effacement of foot processes (podocytopathy)<sup>Q</sup>
- **Genetic basis:** APOL1 gene on chr 22 is strongly associated
- **Etiology**
  - **Primary:** Idiopathic (10% in children and 35% in adults)<sup>Q</sup>
  - **Secondary: Due to underlying etiology**
    - **Reflux nephropathy**<sup>Q</sup>
    - Hypertensive nephropathy
    - **HIV infection (HIV-associated nephropathy)**<sup>Q</sup>
    - **Heroin addiction (heroin nephropathy)**<sup>Q</sup>
    - **Sickle-cell disease**<sup>Q</sup>
    - Massive obesity
    - Secondary event to focal glomerulonephritis (e.g. **IgA nephropathy**)<sup>Q</sup>
    - **Renal ablation/surgery**<sup>Q</sup>



FSGS: Renal biopsy showing sclerosis of a part of glomerulus

- Congenital anomalies (unilateral **renal agenesis or renal dysplasia**)
- **Inherited forms** of nephrotic syndrome : mutation in **podocin**,  $\alpha$ -actinin 4, and TRPC6 (transient receptor potential calcium channel-6)<sup>Q</sup>
- **Morphology**
  - **Light microscopy**
    - Collapse of capillary loops in sclerotic areas
    - Deposition of plasma proteins along capillary wall (**hyalinosis**)
  - **Immunofluorescence microscopy**
    - IgM + C3 deposition in sclerotic areas and/or in mesangium.
  - **Electron microscopy**
    - **Diffuse effacement** of foot processes of podocytes
    - **Focal detachment** of the epithelial cells
    - **Denudation** of the underlying GBM.
- **Clinical course**
  - 20% of patients follow rapid course, with massive proteinuria ending in renal failure within 2 years.
  - Histologic subtype (collapsing variant → unfavorable course; tip variant → good prognosis)
  - 25–50% recurs in patients who undergo transplantation



## Nephrotic Syndrome in Children due to Genetic Disorders of the Podocytes

Gene	Name	Chr	Inheritance	Renal Disease
STEROID-RESISTANT NEPHROTIC SYNDROME				
<i>NPHS1</i> <sup>a</sup>	Nephrin <sup>a</sup>	19q	Recessive <sup>a</sup>	Finnish-type <sup>a</sup> congenital nephrotic syndrome
<i>NPHS2</i> <sup>a</sup>	Podocin <sup>a</sup>	1q	Recessive <sup>a</sup>	FSGS <sup>a</sup>
<i>FSGS1</i>	α-actinin-4 ( <i>αACTN4</i> )	19q	Dominant	FSGS <sup>a</sup>
<i>FSGS2</i>	Unknown	11q	Dominant	FSGS
<i>WT1</i>	Wilms tumor-suppressor gene	11p	Dominant	Denys-Drash syndrome Frasier's syndrome <sup>a</sup>
LMX1BQ	LIM-homeodomain protein	9q	Dominant	Nail-patella syndrome <sup>a</sup>
SMARCAL1	SW1/SNF2-related	2q	Recessive	Schimkeim- muno-osseous dysplasia with FSGS
STEROID-RESPONSIVE NEPHROTIC SYNDROME				
Unknown	Unknown	Unknown	Recessive	MCNS

## HIV-associated Nephropathy (HIVAN)

- Associated with 5–10% of HIV-infected individuals

### Morphology

Light microscopy	Electron microscopy
<p><b>Collapsing variant of FSGS:</b></p> <ul style="list-style-type: none"> <li>Most characteristic lesion<sup>a</sup></li> <li>Poor prognosis<sup>a</sup></li> </ul> <p>Characterized by collapse of entire glomerular tuft along with proliferation and hypertrophy of glomerular visceral epithelial cells<sup>a</sup> (also seen in pamidronate toxicity).</p>	<ul style="list-style-type: none"> <li>Focal cystic dilation of tubule segments filled with proteinaceous material, and inflammation and fibrosis</li> <li>Tubuloreticular inclusions<sup>a</sup> within endothelial cells (also seen in SLE)</li> </ul>

## MEMBRANOPROLIFERATIVE GLOMERULONEPHRITIS (MPGN)

### Etiology

<b>Primary</b>	Idiopathic
<b>Secondary</b>	Invariably type I: More common in adults <sup>a</sup>
	<b>Autoimmune diseases</b> SLE <sup>a</sup>
	<b>Infections</b> Hep C infection, usually with cryoglobulinemia <sup>a</sup> , Hep B, HIV, Schistosomiasis <sup>a</sup>
	<b>Other causes</b> α1-Antitrypsin deficiency <sup>a</sup> , CLL

	MPGN I	MPGN II (Now called C3 glomerulopathy)
<b>Complement activation</b>	Both classical and alternate pathway	Only alternate pathway
<b>Complement levels</b>	All factors reduced	C1, C2 and C4 normal; C3 Nephritic factor seen;
<b>Light Microscopy</b>	Mesangial and GBM proliferation (tram track appearance)	Dense deposits (also called dense deposit disease)
<b>Type of deposits</b>	Subendothelial	Electron dense intramembranous

## ISOLATED GLOMERULAR DISEASES

### IgA Nephropathy (Berger's Disease)

- Most common type of glomerulonephritis in adults, worldwide.
- Most common cause of gross hematuria

**Characterized by:** Recurrent hematuria and presence of IgA deposits in the mesangium.<sup>a</sup>


### Pathogenesis

- Aberrantly glycosylated **IgA-1** deposition in mesangium<sup>a</sup> or Autoimmune response to IgA-1

Mesangial immune deposits **activate mesangial cells<sup>a</sup>** to proliferate, produce increased amounts of **extracellular matrix**, and secrete numerous cytokines and growth factors.


### Etiology

- Primary:** Idiopathic
- Secondary IgA nephropathy:** Due to underlying causes-
  - Gluten enteropathy (celiac disease):** Intestinal mucosal defects<sup>a</sup>
  - Liver disease:** Defective hepatobiliary clearance of IgA<sup>a</sup>



### Morphology

Light microscopy	Immunofluorescence microscopy	Electron microscopy
<ul style="list-style-type: none"> <li>• <b>Mesangioproliferative glomerulonephritis:</b> mesangial widening and endocapillary proliferation<sup>Q</sup></li> <li>• <b>Focal proliferative glomerulonephritis</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Mesangial deposition of IgA+ C3 and properdin, IgG, IgM+/-<sup>Q</sup></b></li> <li>• Early complement components are absent</li> </ul>	<p>Electron-dense deposits in mesangium<sup>Q</sup></p>



### Clinical Course

- Two most common presentations of IgA nephropathy are:
  - Recurrent episodes of **macroscopic hematuria** 1-2 days following an **upper respiratory infection** often accompanied by **proteinuria** or
  - Persistent asymptomatic microscopic hematuria.
- **Nephrotic syndrome, however, is uncommon**
- **15-40% cases progress to ESRD in 20 years.<sup>Q</sup>**
- **15% recurs in patients who undergo transplantation<sup>Q</sup>**



### High Yield Facts

- **Collapsing type** of FSGS has **worst prognosis**
- **Types II and III MPGN** are associated with complement factor H deficiency, presence of C3 nephritic factor, partial lipodystrophy (type II MPGN) or complement receptor deficiency (type III MPGN)
- Kidney changes in **AIDS** patient include **Collapsing variant of FSGS, MPGN, DPGN, IgA nephropathy (Mesangioproliferative glomerulonephritis), MCD and MGN**
- **Most specific histological lesion in diabetic nephropathy** (see chapter 18 Endocrine system) is **Nodular glomerulosclerosis or Kimmelsteil-Wilson lesions**
- **Most common pathological lesion in diabetes nephropathy** is **diffuse GBM thickening > diffuse glomerulosclerosis**

### Hereditary Nephritis

Diseases with **mutations in collagen genes** that manifest primarily with **glomerular injury**.

**Includes:** Alport syndrome and Thin basement membrane lesion (**MC cause of benign familial hematuria**)<sup>Q</sup>


#### Alport Syndrome

- **Triad of:**
  - **Hematuria, sensorineural deafness, Eye disorders:** lens dislocation, posterior cataracts, and corneal dystrophy<sup>Q</sup>
- **Inheritance:**
  - **X-linked dominant<sup>Q</sup>** (85%, **MC** mode of inheritance) > Autosomal
- **Pathogenesis:** Mutations in subunits of collagen → Defective GBM synthesis
  - **COL4A5 (X-linked)<sup>Q</sup>, COL4A3/4 (Autosomal)<sup>Q</sup>**
- **Morphology:**
  - **Light microscopy:** Mesangial proliferation, Capillary wall thickening,
    - **Foam cells<sup>Q</sup>:** Lipid containing tubular or interstitial cells
  - Immunofluorescence microscopy, **Absence of staining with COL4A5<sup>Q</sup>** (Not diagnostic)
  - **Electron microscopy**
    - **GBM: Irregular foci of thickening alternating with thinning<sup>Q</sup>**
    - Pronounced splitting and lamination of the **lamina densa:** distinctive **basket-weave appearance** (diagnostic feature of Alport syndrome)<sup>Q</sup>.

### THIN BASEMENT MEMBRANE LESION (BENIGN FAMILIAL HEMATURIA)

- **Inheritance:** Autosomal inheritance with **defective collagen 4  $\alpha$ 3/4 (COL4A3/4)<sup>Q</sup>**
- **Morphology** on Light microscopy: Diffuse thinning of the GBM to widths between **150 and 225 nm<sup>Q</sup>** (compared with 300 to 400 nm in healthy adults)
- **Prognosis:** Excellent






Alports' Syndrome (GBM showing splitting and lamination) arrow

## TWO SPECIAL CONDITIONS NEED MENTION

### Diabetic Nephropathy

This is nodular glomerulosclerosis (Kimmelstiel-Wilson lesion) of diabetes mellitus.



### Renal Vascular Lesions

**Arteriosclerosis** affecting both **afferent & efferent arteriole**.<sup>Q</sup>

**Pyelonephritis, Including Necrotizing Papillitis**


Starts in interstitial tissue & spreads to tubules; called **necrotizing papillitis (papillary necrosis)**<sup>Q</sup>

### SLE Kidney

Classification of Lupus Nephritis (International Society of Nephrology)

- **Class I : Minimal Mesangial**<sup>Q</sup>
- **Class II : Mesangial Proliferative**
- **Class III : Focal Lupus Nephritis**<sup>Q</sup>
- **Class IV : Diffuse Lupus Nephritis**<sup>Q</sup>
- **Class V : Membranous Lupus Nephritis**
- **Class VI : Advanced Sclerotic Lupus Nephritis**

In lupus there is a full house pattern. They are all IgG, IgA, IgM. Alternative and classical cascade are involved



### Glomerular Lesions

<b>A. Capillary Basement Membrane Thickening</b>	<b>Most characteristic lesion</b> <sup>Q</sup> , GBM thickening & mesangial widening <sup>Q</sup> Along with thickening of <b>tubular basement membranes</b> <sup>Q</sup>
<b>B. Diffuse Mesangial Sclerosis</b>	<b>Most common change</b> <sup>Q</sup> , Diffuse increase in <b>mesangial matrix (PAS-positive)</b> ; <sup>Q</sup> correlates with deteriorating renal function like proteinuria <sup>Q</sup>
<b>C. Nodular or Intercapillary Glomerulosclerosis or Kimmelstiel-Wilson disease</b> <sup>Q</sup>	<ul style="list-style-type: none"> <li>• Spherical, laminated, <b>nodules of matrix</b> at the <b>periphery of glomerulus</b>, which are <b>PAS-positive</b><sup>Q</sup></li> <li>• Capillary microaneurysms</li> </ul> <p><b>Insudative lesions</b></p> <ul style="list-style-type: none"> <li>• <b>Fibrin caps</b><sup>Q</sup>- Accumulation of hyaline material in capillary loops</li> <li>• <b>Capsular drops</b><sup>Q</sup> - hyaline material adherent to Bowman's capsules</li> </ul>

## CHRONIC GLOMERULONEPHRITIS


**It is end-stage glomerular disease** as a result of glomerulonephritis.





Following primary glomerular diseases commonly lead to chronic glomerulonephritis:

- **Rapidly progressive (Crescentic) GN (90%) (most common)<sup>Q</sup>**
- **Focal segmental glomerulosclerosis (50% to 80%)**
- Membranoproliferative GN (50%), Membranous, IgA nephropathy, **Post-streptococcal**




Granular contracted kidney

### Morphology

- **Symmetrically contracted kidneys<sup>Q</sup>** with **diffusely granular cortical surfaces**
- On section, the **cortex is thinned<sup>Q</sup>**, and there is an increase in peripelvic fat.
- Replacement of glomeruli by **blue-staining collagen (Masson trichrome stain)<sup>Q</sup>**
- **Tubular atrophy**, irregular **interstitial fibrosis**, and **mononuclear leukocytic infiltration** of the interstitium.

### Causes of Granular Contracted Kidney



## Mnemonic

**Flea-bitten kidney** is seen in  
"World Health PSM"

- **W**orld - Wegener's granulomatosis
- **H**earth - Henoch Schonlein purpura
- **P** - Post-streptococcal Glomerulonephritis (PSGN). Polyarteritis nodosa
- **S** - Sub acute bacterial endocarditis (SABE), SLE, Good Pasture syndrome
- **M** - Malignant hypertension

## THIN BASEMENT MEMBRANE LESION (BENIGN FAMILIAL HEMATURIA)

**Inheritance:** Autosomal inheritance with **defective collagen 4  $\alpha 3/4$  (COL4A3/4)<sup>Q</sup>**

**Morphology** on Light microscopy: Diffuse thinning of the GBM to widths between **150 and 225 nm<sup>Q</sup>** (compared with 300 to 400 nm in healthy adults)

**Prognosis:** Excellent

## CHRONIC GLOMERULONEPHRITIS

**End-stage glomerular disease** as a result of glomerulonephritis. Following primary glomerular diseases commonly lead to chronic glomerulonephritis:

- **Rapidly progressive (Crescentic) GN (90%) (most common)<sup>Q</sup>**
- **Focal segmental glomerulosclerosis (50% to 80%)**
- Membranoproliferative GN (50%)
- Membranous (30% to 50%)
- IgA nephropathy (30% to 50%)
- **Post-streptococcal (1% to 2%)**

### Morphology

- **Symmetrically contracted kidneys<sup>Q</sup>** with **diffusely granular cortical surfaces**
- On section, the **cortex is thinned<sup>Q</sup>**, and there is an increase in peri-pelvic fat.
- Replacement of glomeruli by **blue-staining collagen (Masson trichrome stain)<sup>Q</sup>**
- **Tubular atrophy**, irregular **interstitial fibrosis**, and **mononuclear leukocytic infiltration** of the interstitium.


### Causes of Granular Contracted Kidney

Symmetric	Asymmetric
Chronic Glomerulonephritis <sup>Q</sup>	Chronic Pyelonephritis <sup>Q</sup>
Diabetes (rare)	
Benign Nephrosclerosis	



## ACUTE KIDNEY INJURY (AKI)/ACUTE RENAL FAILURE (ARF)

### Causes of Acute Kidney Injury



**Definition:** AKI is defined as any of the following:

- Increase in Serum Cr by 0.3 mg/dl within 48 hours; or
- Increase in Serum Cr to 1.5 times baseline within the prior 7 days; or
- Urine volume <0.5 mL/kg/hr for 6 hours

### Acute Tubular Injury/Necrosis (ATN)

- Structural tubular injury<sup>Q</sup> due to renal hypoperfusion.
- Ischemic and toxic ATN account for 90% of cases of acute intrinsic renal failure.

### Laboratory Findings in Acute Renal Failure

Index	Prerenal Azotemia	Oliguric Acute Renal Failure
BUN/P <sub>cr</sub> ratio <sup>Q</sup>	>20:1 <sup>Q</sup>	10-15:1
Urine sodium (U <sub>Na</sub> ), meq/L <sup>Q</sup>	<20 <sup>Q</sup>	>40
Urine osmolality (mosmol/L)	>500	<350
Fractional excretion of sodium	<1% <sup>Q</sup>	>2%
Urine/plasma creatinine (U <sub>cr</sub> /P <sub>cr</sub> )	>40	<20

### New Biomarkers of Acute Kidney Injury

Biomarker	Comments
<i>N-Acetyl--(D) glucosaminidase (NAG)</i> <sup>Q</sup>	Proximal tubule lysosomal enzyme
<i>Retinol-binding protein</i>	Early marker of tubular dysfunction
<i>Cystatin C</i> <sup>Q</sup>	Elevated urinary levels reflect tubular dysfunction; high levels may predict poorer outcome
<i>Kidney injury molecule-1 (KIM-1)</i> <sup>Q</sup>	Elevated urinary levels highly sensitive and specific for AKI
<i>Clusterin</i>	Elevated kidney and urinary levels are very sensitive for AKI
<i>Neutrophil gelatinase associated lipocalin (NGAL)</i> <sup>Q</sup>	Early indicator of AKI following cardiopulmonary bypass
<i>Interleukin-18 (IL-18)</i>	Elevated urinary level is an early marker of AKI and independent predictor of mortality in critically ill patients
<i>Liver fatty acid-binding protein (L-FABP)</i>	A biomarker in CKD and diabetic nephropathy
<i>Sodium/hydrogen exchanger isoform (NHE3)</i>	Urinary levels found to discriminate between prerenal azotemia and AKI in ICU patients
<i>Exosomal fetuin-A</i>	Acute phase protein synthesized in the liver; High Urinary levels in ICU patients with AKI



## TUBULOINTERSTITIAL NEPHRITIS

### Characterized by

- Inflammatory injuries to tubules and interstitium<sup>Q</sup> Insidious in onset
- Principally manifested by azotemia<sup>Q</sup>

### Two Forms

Acute tubulointerstitial nephritis	Chronic interstitial nephritis
<b>Rapid clinical onset<sup>Q</sup></b> <b>Histologically:</b>	<b>Insidious onset<sup>Q</sup></b> <b>Histologically:</b>
<ul style="list-style-type: none"> <li>• Interstitial edema<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Infiltration with mononuclear WBCs</li> </ul>

<ul style="list-style-type: none"> <li>• Neutrophilic infiltration in interstitium and tubules<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Prominent interstitial fibrosis</li> </ul>
<ul style="list-style-type: none"> <li>• Tubular injury</li> </ul>	<ul style="list-style-type: none"> <li>• Tubular atrophy<sup>Q</sup></li> </ul>

### Distinguished Clinically from the Glomerular Diseases by the Following Hallmarks:

- **Absence of nephritic or nephrotic syndrome<sup>Q</sup>**
- **Presence of defects in tubular function<sup>Q</sup>**
  - E.g.: **Impaired** ability to **concentrate urine** (polyuria or nocturia)
  - **Salt wasting**
  - **Diminished** ability to **excrete acids** (metabolic acidosis)
  - Isolated defects in **tubular reabsorption** or secretion

## PYELONEPHRITIS AND URINARY TRACT INFECTION

**Characterized by:** Inflammation of tubules, interstitium, and renal pelvis

**Pathogenesis of Acute Pyelonephritis:** Ascending infection is the **most common** cause.

Characteristics	Acute pyelonephritis	Chronic pyelonephritis
<b>Definition</b>	Acute <b>suppurative inflammation</b> of the kidney	Chronic <b>tubulointerstitial inflammation</b> and <b>scarring</b> involve the calyces and pelvis
<b>Etiology</b>	Ascending urinary tract infection MC: <i>E. coli (MC)</i> <sup>Q</sup> , <i>Proteus</i> , <i>Klebsiella</i> , and <i>Enterobacter</i>	Bacterial infection plays a dominant role; 2 forms: <ul style="list-style-type: none"> <li>• <b>Reflux nephropathy<sup>Q</sup></b>: Urinary infection on congenital <b>vesicoureteral reflux</b></li> <li>• <b>Chronic obstructive pyelonephritis<sup>Q</sup></b></li> </ul>
<b>Morphology: Gross</b>	<ul style="list-style-type: none"> <li>• Discrete focal abscesses</li> <li>• Large wedge like areas</li> </ul>	<ul style="list-style-type: none"> <li>• Coarse, discrete, <b>Corticomedullary scars<sup>Q</sup> (Hallmark)</b> overlying dilated, blunted, or deformed calyces</li> <li>• <b>Flattening of the papillae<sup>Q</sup></b></li> </ul>
<b>Light Microscopy</b>	<ul style="list-style-type: none"> <li>• <b>Patchy interstitial suppurative inflammation (hallmark)</b></li> <li>• Intratubular aggregates of neutrophils</li> <li>• <b>Neutrophilic tubulitis<sup>Q</sup></b></li> <li>• <b>Tubular necrosis<sup>Q</sup></b></li> </ul>	<ul style="list-style-type: none"> <li>• Involves <b>tubules and interstitium</b></li> <li>• <b>Thyroidization of tubules<sup>Q</sup></b>: Dilated tubules filled with casts resembling thyroid colloid</li> <li>• Chronic interstitial inflammation and fibrosis</li> <li>• Vessels demonstrate <b>obliterative intimal sclerosis</b></li> <li>• <b>Hyaline arteriolo sclerosis<sup>Q</sup></b></li> <li>• <b>Periglomerular fibrosis<sup>Q</sup></b></li> <li>• <b>Secondary FSGS<sup>Q</sup></b> may be seen.</li> </ul>
<b>Complications</b>	<ul style="list-style-type: none"> <li>• <b>Papillary Necrosis</b> <ul style="list-style-type: none"> <li>■ Major Causes <ul style="list-style-type: none"> <li>◆ Analgesic nephropathy (Most common)<sup>Q</sup></li> <li>◆ Sickle cell nephropathy</li> <li>◆ Diabetes with urinary tract infection</li> </ul> </li> </ul> </li> <li>• <b>Pyonephrosis<sup>Q</sup></b></li> <li>• <b>Perinephric abscess<sup>Q</sup></b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Xanthogranulomatous pyelonephritis:</b> <ul style="list-style-type: none"> <li>■ A rare form of <b>chronic pyelonephritis<sup>Q</sup></b></li> <li>■ Often associated with <b>Proteus infections<sup>Q</sup></b> and obstruction</li> <li>■ M.C age group: <b>5<sup>th</sup> -6<sup>th</sup> decade<sup>Q</sup></b></li> <li>■ <b>Females</b> are most commonly affected</li> <li>■ Associated features: <b>Large staghorn calculi<sup>Q</sup></b> and <b>hydronephrosis<sup>Q</sup></b></li> <li>■ Gross morphology: <b>large, yellowish orange</b> nodules that may be grossly confused with renal cell carcinoma.</li> <li>■ <b>Microscopy:</b> Accumulation of <b>foamy (lipid laden) macrophages (Xanthoma cells)<sup>Q</sup></b> with <b>plasma cells, lymphocytes</b>.</li> </ul> </li> </ul>




## High Yield Facts

- **Most important factor in pyelonephritis: Vesicoureteral reflux<sup>Q</sup>**
- In the absence of vesicoureteral reflux, infection usually remains localized in the bladder
- An emerging **viral pathogen** causing **pyelonephritis** in **kidney transplantation** is **Polyomavirus** (> 6 months post-transplant)
- **Analgesic nephropathy** is a high risk for development of **transitional cell Carcinoma kidney**

### Pyelonephritis can occur in following steps


- **Colonization<sup>Q</sup>** of the distal urethra and introitus (in the female) by coliform bacteria
- **Retrograde spread<sup>Q</sup>** from the **urethra to the bladder**
- Further spread from **bladder to kidneys** by the following mechanisms:
  - **Urinary tract obstruction** and **stasis** of urine<sup>Q</sup>
  - **Vesicoureteral reflux<sup>Q</sup>**-Incompetence of the vesicoureteral valve that allows bacteria to ascend the ureter into the renal pelvis.
  - Open ducts at the tips of the papillae (**intrarenal reflux**)<sup>Q</sup>




## INHERITED CYSTIC KIDNEY DISEASES

Disease		Gene	Protein	Renal Abnormality	Extrarenal Abnormality
<b>Autosomal dominant polycystic kidney disease<sup>Q</sup></b>	AD	<i>PKD1</i>	Polycystin-1	Cortical and medullary cysts	<b>Cerebral aneurysms;<sup>Q</sup> liver and spleen cysts<sup>Q</sup></b>
	AD	<i>PKD2</i>	Polycystin-2		
<b>Autosomal recessive polycystic kidney disease<sup>Q</sup></b>	AR	<i>PKHD1</i>	Fibrocystin (polyductin)	Distal tubule and collecting duct cysts	<b>Hepatic fibrosis;<sup>Q</sup> Caroli's disease<sup>Q</sup></b>
<b>Nephronophthisis I (juvenile/adolescent)</b>	AR	<i>NPHP1</i>	Nephrocystin	Small fibrotic kidneys; medullary cysts	<b>Retinitis pigmentosa</b>
<b>Nephronophthisis II (infantile)</b>	AR	<i>NPHP2 (INVS)</i>	Inversin	<b>Large kidneys;<sup>Q</sup> widespread cysts</b>	<b>Situs inversus<sup>Q</sup></b>
<b>Nephronophthisis III (juvenile/adolescent)</b>	AR	<i>NPHP3</i>	Nephrocystin-3	Small fibrotic kidneys; medullary cysts	<b>Retinitis pigmentosa; hepatic fibrosis</b>
<b>Medullary cystic kidney disease</b>	AD	<i>MCKD1 /2</i>	Uromodulin	Small fibrotic kidneys; medullary cysts	<b>Hyperuricemia and gout<sup>Q</sup></b>
<b>Tuberous sclerosis</b>	AD	<i>TSC1/2</i>	Hamartin/ Tuberlin	<b>Renal cysts;<sup>Q</sup> angiomyolipomas;<sup>Q</sup> Renal cell carcinoma<sup>Q</sup></b>	<b>Facial angiofibromas;<sup>Q</sup> CNS hamartomas<sup>Q</sup></b>
<b>Von Hippel-Lindau disease</b>	AD	<i>VHL</i>	pVHL	<b>Renal cysts;<sup>Q</sup> Renal cell carcinoma<sup>Q</sup></b>	<b>Retinal angiomas;<sup>Q</sup> CNS hemangioblastomas;<sup>Q</sup> pheochromocytomas<sup>Q</sup></b>





Autosomal dominant polycystic kidney disease showing distorted outline



Autosomal recessive polycystic kidney disease showing smooth outline



### High Yield Facts

- In ADPKD, patients with PKD1 mutation have higher risk of ESRD<sup>Q</sup> than those with PKD2 mutation
- Extra-renal sites of cysts in ADPKD are: Liver (most common)<sup>Q</sup> > CNS (berry aneurysm) > Spleen > Pancreas > Lung
- Familial juvenile Nephronophthisis is the most common variant of Nephronophthisis
- Nephronophthisis complex is the most common genetic cause of ESRD in children<sup>Q</sup>

### Acquired (Dialysis-Associated) Cystic Disease

- Cystic degeneration in ESRD after prolonged dialysis.
- Increased risk of renal cell carcinoma

## RENAL TUMORS

### Benign Neoplasms

- Renal Papillary Adenoma**
  - Small, discrete adenomas
  - Arises from the **renal tubular epithelium**<sup>Q</sup>
- Angiomyolipoma**
  - Benign neoplasm**<sup>Q</sup> consisting of vessels, smooth muscle, and fat
  - Originates from **perivascular epithelioid**<sup>Q</sup> cells.
- Oncocytoma**
  - Epithelial neoplasm composed of **large eosinophilic cells**<sup>Q</sup> having small, round, **benign-appearing nuclei** that have large nucleoli.
  - Arise from the intercalated cells of collecting ducts
  - E/M: **eosinophilic cells** have **numerous mitochondria**.<sup>Q</sup>

### Malignant Tumors

#### Renal Cell Carcinoma (RCC)


- Age group:** most commonly in 6<sup>th</sup> and 7<sup>th</sup> decade<sup>Q</sup>
- Male: Female ratio = 2:1.**<sup>Q</sup>

#### Risk Factors

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li><b>Cigarette smoking (most important)</b><sup>Q</sup></li> <li><b>Obesity (particularly in women)</b><sup>Q</sup></li> <li>Hypertension<sup>Q</sup></li> <li>Unopposed estrogen therapy</li> <li>Exposure to asbestos</li> <li><b>Petroleum products</b><sup>Q</sup></li> <li>Heavy metals.</li> </ul> | <p>There is also an <b>increased risk</b> in patients with:</p> <ul style="list-style-type: none"> <li><b>End-stage renal disease</b><sup>Q</sup></li> <li><b>Chronic kidney disease</b><sup>Q</sup></li> <li><b>Acquired cystic disease</b><sup>Q</sup></li> <li><b>Tuberous sclerosis.</b><sup>Q</sup></li> </ul> |
|---|---|

#### Classification of RCC

- Sporadic:** Most common type
- Hereditary forms:** Autosomal dominant, Young age affected





### Latest Update

#### 4 Types of Familial Variants

- **Von-Hippel-Lindau (VHL) syndrome:** AD
  - Sporadic and familial forms of clear cell carcinoma
- **Hereditary leiomyomatosis and renal cell cancer syndrome:** AD
  - Caused by mutations of the FH gene which expresses fumarate hydratase

- Characterized by cutaneous and uterine leiomyomata and an aggressive type of papillary carcinoma with increased tendency for metastatic spread.
- **Hereditary papillary carcinoma:** AD
  - Mutation in MET proto-oncogene papillary cell Ca
- **Birt-Hogg-Dubé syndrome:** AD
  - Mutations in BHD gene, which expresses folliculin.
  - Constellation of skin (fibrofolliculomas, trichodiscomas, and acrochordons), pulmonary (cysts or blebs), and renal tumors

#### Clinical Features

- Costovertebral pain, palpable mass, hematuria (**Most reliable sign**)<sup>Q</sup>

#### Metastasis

**Most common** locations of metastasis are: **lungs (50%)<sup>Q</sup>** and **bones (33%)<sup>Q</sup>**

### Latest Update

#### Xp11 translocation



- Rare and seen in young Patients
- Tumor cells with clear cytoplasm and papillary structure
- Translocation of TFE3 on Chr Xp11.2

#### Paraneoplastic Feature of RCC

- Hypertension (**most common**)<sup>Q</sup>
- Polycythemia, Hypercalcemia
- **Hepatic dysfunction (Stauffer's syndrome)**<sup>Q</sup>
- **Feminization or masculinization**<sup>Q</sup>
- **Cushing syndrome (Rare)**<sup>Q</sup>, Eosinophilia, Leukemoid reactions, Amyloidosis.

#### Urothelial Carcinoma of the Renal Pelvis

- **Originate from the urothelium of the renal pelvis**<sup>Q</sup>
- Range from **Benign papillomas** to **invasive urothelial (transitional cell) carcinomas**.<sup>Q</sup>
- Multiple lesions involving the pelvis, ureters, and bladder.
- Often associated with bladder urothelial tumor.
- Increased incidence with **Lynch syndrome** and **analgesic nephropathy**.<sup>Q</sup>
- Infiltration of the wall of the pelvis and calyces indicative of **poor prognosis**.<sup>Q</sup>



#### Microscopy

- Infiltration with **large, foamy macrophages**<sup>Q</sup> with a **Michaelis-Guttman bodies**<sup>Q</sup> with multinucleate giant cells and lymphocytes.
- Laminated mineralized concretions resulting from deposition of calcium in enlarged lysosomes, typically present within the macrophages are called Michaelis Guttmann bodies. These histiocytes have pink-cytoplasm called as **von Hanseman histiocytes**

## DISEASES OF URINARY BLADDER

### MALACOPLAKIA

- A vesical **inflammatory reaction**<sup>Q</sup> is characterized macroscopically by soft, yellow, slightly raised **mucosal plaques** 3 to 4 cm in diameter
- It is a granulomatous disease with **defective intracellular lysosomal digestion** of bacteria in histiocytes. It is mostly caused by **E. coli**.<sup>Q</sup>

## BLADDER CARCINOMA

- **Epidemiology:** More common in men than in women, common age group affected: 50–80 years old
- **Types:**
  - Epithelial tumors include **Urothelial or transitional cell type (most common)**<sup>Q</sup>, Squamous<sup>Q</sup> and Glandular<sup>Q</sup>
- **Precursor lesions:**
  - **Noninvasive** papillary tumors, **Flat noninvasive** urothelial carcinoma<sup>Q</sup>




- **Risk factors:**
  - **Cigarette smoking:** Most important<sup>Q</sup>
  - **Industrial exposure to arylamines**<sup>Q</sup> like 2-naphthylamine, Benzidine, Acrolein
  - **Schistosoma haematobium**<sup>Q</sup> infections (**Squamous Carcinoma** > Transitional Cell Ca)
  - Long-term **analgesic use (Phenacetin)**<sup>Q</sup>
  - Long-term exposure to **cyclophosphamide**, Exposure of bladder to **irradiation** Urolithiasis (predisposes to Squamous Cell Ca)
- **Genetic alterations:**
  - Monosomy 9 or del 9p or 9q (most common)
- **Clinical feature:**
  - **Painless hematuria**, Frequency, urgency, and dysuria may be present
- **Prognosis:**
  - Depends on the **histologic grade**<sup>Q</sup> of the papillary tumor and the **stage** at diagnosis.
  - Involvement of **muscularis mucosa (detrusor muscle)**<sup>Q</sup> is associated with **worst prognosis**



### High Yield Facts

- **Noninvasive high grade urothelial carcinoma:** Loss of TP53 and RB genes, **noninvasive low grade papillary urothelial Carcinoma:** Gain of function FGFR3 and HRAS mutations
- **Squamous cell carcinoma and adenocarcinoma**<sup>Q</sup> are associated with a **worse prognosis**<sup>Q</sup>
- **"Polychronotropism"**<sup>Q</sup>: The tendency to recur over time and in new locations in the urothelial tract



Transitional cell carcinoma



### High Yield Facts

#### Assessment of Glomerular Filtration Rate (GFR)

- **Direct measurement: By Inulin clearance**<sup>Q</sup> (Inulin is filtered at the glomerulus but neither reabsorbed nor secreted throughout the tubule)
- **Serum Creatinine:** Used as a surrogate to estimate GFR. It is the most widely used marker for GFR. Commonly used formulae used to calculate GFR from serum creatinine are:
  - **Cockcroft-Gault**<sup>Q</sup>
  - **Schwartz Formula**<sup>Q</sup>
- **Creatinine clearance**<sup>Q</sup>: An approximation of GFR;  $CrCl = \frac{(U_{vol} \times U_{Cr})}{(P_{Cr} \times T_{min})}$ .
- **Cystatin C**<sup>Q</sup>: **More sensitive marker of early GFR decline** than plasma creatinine

R10<sup>th</sup>


### Latest Update

#### C3 Glomerulopathy

Encompasses the following:

1. Dense-deposit disease (formerly known as type II MPGN) where dense glomerular intramembranous deposits are present in the glomerular capillary loops, which stain for C3 only by IF.
2. C3GN, a proliferative GN resembling MPGN type I without Ig deposits.
3. Rare diseases such as familial MPGN type III and complement factor H-related protein 5 abnormality-associated familial GN cases
  - **Immunofluorescence Characteristics:** C3 shows double-linear appearance of the glomerular capillary wall and a ring appearance around the mesangial deposits (mesangial rings)
  - IgG deposition should not be present
  - Type II MPGN is now classified under C3 glomerulopathy






## NEXT Pattern Questions



Q's

1. A 26-year-old patient presented with proteinuria, hematuria. PAS stain of renal biopsy demonstrates characteristic histological findings. Which of the following cannot be a possible etiology?



- a. Good pasture syndrome    b. Wegener's granulomatosis  
c. IgA nephropathy    d. Polyarteritis nodosa


Ans. (d) **Polyarteritis nodosa**

- With the history of nephritic syndrome and the image is suggestive of Crescent's, this is a case of RPGN. Now in the classification of RPGN, PAN is not a cause.



Q's

2. A 60-year-old male with complaints of hematuria, severe hypertension and mild facial puffiness with occasional hemoptysis. The DIF picture of the patient is given below. Which of the following is not true about the given condition?



- a. Antibody to alpha 3 chain of type 4 collagen part of non-collagenous domain is responsible  
b. Strong association with HLA DR B1  
c. Crescents are seen in >50% of the glomeruli  
d. Dense depositions below podocytes are seen in electron microscopy


Ans. (d) **Dense depositions below podocytes are seen in electron microscopy**

- This is a case of nephritic syndrome and the DIF shows linear deposits, so it's a case of Good Pasture syndrome. Since, it involves Anti GBM antibodies, so option D which is indicative of immune complex is false statement.



Q's

3. A patient presented with pus in urine. Urine culture was done which was negative. After a sudden onset renal failure the patient died. On autopsy the following finding was seen in kidney. What is the most likely diagnosis?



- a. TB kidney    b. Infected renal cysts  
c. Renal cell carcinoma    d. Renal stones


Ans. (a) **TB kidney**

- With the history of sterile pyuria, uremia and white cheesy caseous deposits in the kidney, this is a case of TB kidney.



Q's

4. A 2-year-old child presented with purpuric skin lesions, abdominal pain, intestinal bleeding and arthralgia. Immunofluorescence study demonstrated IgA deposition as shown below. What is your diagnosis?



- a. FSGS  
b. Minimal change disease  
c. Diabetic neuropathy  
d. Henoch-Schönlein purpura


Ans. (d) **Henoch-Schönlein purpura**

- The history of this 2-year-old child is suggestive of Henoch schonlein purpura, and with an IgA deposition in the mesangium the biopsy is suggestive of IgA nephropathy.





5. A 50-year-old male presented with blurring of vision. Urine examination showed proteinuria. Histopathology picture of kidney given below. What is not true about the finding in the associated condition?



- Diffuse increase in mesangial matrix
- Sclerotic mesangial nodules
- GBM thinning and permeable
- Hyaline arteriosclerosis


Ans. (c) **GBM thinning and permeable**

- The history of nephrotic syndrome and blurring of vision. Notice the first image shows Kimmelstiel-Wilson nodule suggestive of Diabetic nephropathy. The histology shows GBM thickening and more permeability.




## Image-Based Questions

1. A 5-year-old male presented with mild hematuria, pedal edema and frothy urine. BP 110/70 mm Hg. Subsequently he was treated with steroid but he did not improve. The renal biopsy was carried out. The electron microscopy finding has been shown. What is your diagnosis?




- Minimal Change disease
- FSGS
- MPGN
- PSGN


2. A 55-year-old diabetic male found to have microalbuminuria. Identify the lesion shown in renal biopsy of this patient.



- Amyloidosis
- Kimmelstiel-wilson lesion
- Wire loop lesions
- Crescentic glomerulonephritis




3. The morphological finding of kidney shown in the figure is associated with which of the following conditions?




- Chronic glomerulonephritis
- Renal amyloidosis
- Malignant hypertension
- Acute pyelonephritis

5. Which of the following conditions usually leads to the finding given below:



- Benign hypertension
- Malignant hypertension
- Chronic pyelonephritis
- Acute pyelonephritis

4. Identify the kidney disease?



- Polycystic disease
- Hydronephrosis
- Pyelonephritis
- Chronic pyelonephritis

6. Most common histology of this gross lesion is:



- Clear cell type
- Papillary Cell Ca
- Bellini duct Ca
- Anaplastic Ca



## Answers of Image-Based Questions

1. Ans. (a) **Minimal Change disease**

- History given here is that of nephrotic syndrome and electron microscopy shows flattening of foot processes of podocytes, a feature of minimal change disease.

2. Ans. (b) **Kimmelstiel-Wilson lesion**

- In this case kidney biopsy from a patient suffering from diabetic nephropathy shows Nodular glomerulosclerosis suggested by diffuse increase in mesangial matrix and characteristic acellular PAS-positive nodules.

3. Ans. (c) **Malignant hypertension**

- The given kidney gross shows small, pinpoint **petechial hemorrhages** may appear on the cortical surface from rupture of arterioles or glomerular capillaries, giving the kidney a peculiar "flea-bitten" appearance. The kidney size varies depending on the duration and severity of the hypertensive disease.

4. Ans. (b) **Hydronephrosis**

- Marked dilation of the pelvis and calyces and thinning of the renal parenchyma.

5. Ans. (c) **Chronic pyelonephritis**

- Gross surface is irregularly scarred. Microscopically shows changes involve predominantly tubules and interstitium. The tubules show atrophy in some areas and hypertrophy or dilation in others. Dilated tubules with flattened epithelium may be filled with casts resembling thyroid colloid (thyroidization)

6. Ans. (a) **Clear cell type**


- The given gross shows Renal cell Carcinoma, the most common histology of which is clear cell type.



## Multiple Choice Questions

### STRUCTURE & FUNCTION

- Maltese cross appearance in urinary sediment seen in which of the following disease other than nephrotic syndrome?** (AIIMS Nov 2019)
  - Felty syndrome
  - Fanconi Syndrome
  - Fabry disease
  - Friedrich's ataxia
- Dysmorphic RBC's in urine is/are seen in?** (PGI May 18)
  - Glomerulonephritis
  - Renal vascular injury
  - Renal stone
  - Pyelonephritis
  - Interstitial nephritis
- Which of the following dyads are correctly matched regarding urinary casts and associated condition?**
  - Hyaline casts- may be normally present in healthy person (PGI Nov 2017)
  - Muddy brown casts-acute tubular necrosis
  - WBC cast - pyelonephritis
  - Epithelial cast - acute glomerulonephritis
  - Myoglobin cast - Rhabdomyolysis
- Urinary cast better seen on microscope by?** (PGI May 2017)
  - Centrifuge the urine first
  - Use immunofluorescence light
  - See at edge of cover slip
  - Increasing the light intensity
  - Acidifying it first
- A post renal transplant patient presented with complains of chronic renal failure. As a part of investigative workup, the patient urine microscopy suggested the following finding, identify the structure marked by arrow?** (Recent Question 2016-17)



- Decoy cell
  - Tubular epithelial cell
  - Charcot leyden crystal
  - Hyaline casts
- Which one of the following tests is best for measuring glomerular function?** (Recent Question 2016-17)
    - Blood urea
    - Serum creatinine
    - Creatinine clearance rate
    - Ultrasound of kidney
  - Match list I with list II and select the correct answer using the code given below the lists:** (Recent Question 2016-17)

#### List I (Urine exam)

- Red cell casts
- Microscopic haematuria
- Proteinuria
- Broad cell casts


#### List II (Disease)

- Nephrotic syndrome
- Chronic renal failure
- Polycystic kidney disease
- Glomerulonephritis

#### Code:

A	B	C	D
a. A/4	B/3	C/1	D/2
b. A/3	B/2	C/1	D/4
c. A/2	B/3	C/4	D/1
d. A/4	B/1	C/2	D/3

- Identify the crystal in the urine analysis:** (AIIMS Nov 2015)



- Oxalate
  - Uric acid
  - Phosphates
  - Cysteine
- Identify the arrow marked structure in Urine routine microscopy?** (AP PGME 2015)



- Uric acid
  - Ca oxalate
  - Struvite
  - Cysteine
- Which of the following dyads are correct:**
    - WBC cast- Acute pyelonephritis (PGI May 2015)
    - Broad cast-CRF
    - Eosinophilic cast-interstitial nephritis
    - RBC cast- Glomerulonephritis
    - Broad cast : chronic pyelonephritis
  - Normal level of serum uric acid in males is:** (Recent Question 2015)
    - 3.1-7 mg/dl
    - 2.5-5.6 mg/dl
    - 1.5-3.3 mmol/L
    - 1.8-4.4 mmol/L
  - The protein in glomerular basement membrane responsible for charge dependent filtration is:** (Recent Question 2015, DPG 10)
    - Albumin
    - Collagen type IV
    - Proteoglycan
    - Fibronectin
  - Cast seen in Acute Glomerulonephritis is:** (WB PG 2015)
    - Hyaline cast
    - Granular cast
    - RBC cast
    - WBC cast
  - What is the minimum number of red blood cells per microliter of urine required for diagnosis of hematuria?** (APPGME 14)
    - 3
    - 5
    - 8
    - 10





15. Urine analysis of a patient with hematuria and hypercalciuria is most likely to reveal?  
 a. Isomorphic RBCs (AIIMS Nov 11)  
 b. RBC casts  
 c. Nephrotic range proteinuria  
 d. Eosinophiluria

16. On kidney biopsy, PAS positive structures are:  
 a. Glomerular basement membrane (PGI Nov 10)  
 b. Tubule c. Neutrophils  
 d. Interstitium e. Mesangial matrix

#### PSGN

17. Examine the renal histopathology slide. What is the probable diagnosis? (JIPMER Nov 2019)  
 a. Membranoproliferative GN  
 b. Rapidly proliferative GN  
 c. PSGN  
 d. Diabetic nephropathy

18. Not seen in post streptococcal glomerulonephritis (PSGN)? (PGI May 2016)  
 a. Nephrotic range proteinuria  
 b. Neutrophilic infiltration of tubules  
 c. Subepithelial deposits  
 d. Linear deposits along glomerular basement membrane

19. Post streptococcal glomerulonephritis presents with  
 a. Asymptomatic hematuria (Recent Question 2015)  
 b. Renal failure  
 c. Massive anasarca  
 d. Massive renomegaly

20. Most common renal lesions in HIV  
 a. MPGN (Recent Question 2015)  
 b. RPGN  
 c. FSGS  
 d. Membranous nephropathy

21. Which is seen in Electron microscopy in PSGN? (Recent Question 2015)  
 a. Epithelial humps  
 b. Spike and dome appearance  
 c. Mesangial deposits  
 d. Subendothelial deposits

22. All are true about poststreptococcal glomerulonephritis except - (Recent Question 2014)  
 a. Crescent formation  
 b. Subepithelial deposits  
 c. Granular deposits of IgG  
 d. Deposition of IgA


23. The pathogenesis of acute proliferative glomerulonephritis - (Recent Question 2014)  
 a. Cytotoxic T-cell mediated  
 b. Immune complex mediated  
 c. Antibody mediated  
 d. Cell-mediated (Type IV hypersensitivity)

24. In Poststreptococcal glomerulonephritis:  
 a. C3 decreases, ASO increases (WBPG 2014)  
 b. C3 increases, ASO increases  
 c. C3 decreases, ASO decreases  
 d. C3 increases, ASO decreases

25. Poststreptococcal reactive arthritis is differentiated from Acute Rheumatic Fever by all except:  
 a. Small joint involvement and often symmetric  
 b. Caused by nongroup A hemolytic streptococci  
 c. Non-responsiveness to salicylate  
 d. Shorter incubation period (WB PG 2011)

#### RPGN

26. A 50-year-old male presented with hematuria. Investigations revealed normal glucose levels, proteinuria and creatinine of 9 mg%. Electron microscopic image is shown below. What other investigations could help in the diagnosis? (Recent Pattern Question 2020)



- a. ANA b. HIV serology  
 c. Electrophoresis d. Anti GBM antibodies
27. Examine the gross kidney image. What is the diagnosis? (JIPMER Nov 2019)  
 a. Granular contracted kidney  
 b. Flea bitten kidney  
 c. Spongy kidney  
 d. Hemorrhagic kidney

28. Which of the following is correctly matched in RPGN? (JIPMER 18)  
 a. Type 1 IgA nephropathy  
 b. Type 2 Anti-GBM antibody  
 c. Type 2- Wegener's granulomatosis  
 d. Type 2- SLE nephritis

29. A patient presented with hemoptysis and hematuria. On renal biopsy, it shows crescentic glomerulonephritis. Immunofluorescence microscopy shows linear IgG and C3 deposits. Which of the following is the most appropriate diagnosis? (PGI May 18)

- a. Thin basement disease  
 b. Good pasture syndrome  
 c. Wegener granulomatosis  
 d. PSGN  
 e. Minimal change disease

30. Rapid advanced renal parameters is clinically termed as Rapidly progressive glomerulonephritis. What exactly is the finding on microscopy? (Recent Question 2016-17)

- a. Membrane thickening b. Segmental sclerosis  
 c. Crescents d. Mesangial expansion





- 31. Type I RPGN is seen in** (Recent Question 2015)
- SLE
  - IgA nephropathy
  - Henoch schonlein purpura
  - Good pasture syndrome

- 32. False regarding nephritic syndrome** (Recent Question 2015)
- Generalize edema
  - Proteinuria <3.5 g/day
  - Hypoalbuminemia
  - Hypertension

- 33. True regarding IgA Nephropathy:** (Recent Question 2015)
- Usually in children < 10 years
  - Microscopic hematuria is the most common presentation
  - Recurrent gross hematuria following respiratory infection
  - Decreased serum IgA

- 34. Mesangial deposits of Lambda light chain is seen in** (Recent Question 2015)
- Amyloidosis
  - FSGS
  - MPGN
  - Membranous nephropathy

- 35. Most common cause of primary nephrotic syndrome in adult** (Recent Question 2015)
- Membranous nephropathy
  - Minimal change disease
  - Membranoproliferative glomerulonephritis
  - Focal segmental glomerulosclerosis

- 36. All of the following causes RPGN Except?** (Recent Question 2015)
- Wegeners
  - Polyarteritis nodosa
  - HSP
  - Microscopic polyangitis

- 37. Good pasture's syndrome is characterized by-** (Recent Question 2014)
- Necrotizing hemorrhagic interstitial pneumonitis
  - Emphysema
  - Patchy consolidation
  - Pulmonary edema


- 38. Characteristic feature of Goodpasture's syndrome** (APPGMEE 14)
- Lumpy-bumpy deposits on immunofluorescence
  - Serum antibodies against alpha 3 NC1 domain of collagen - IV
  - Serum antibodies against alpha 1 NC1 domain of collagen III
  - Anti DNase antibodies positive

- 39. Feature of Goodpasture syndrome is/are:** (PGI May 2013)
- Antibody to  $\alpha$ -chain of Type IV collagen (COL-4A)
  - Basement membrane involvement
  - Pulmonary hemorrhage
  - Crescent formation
  - Subendothelial deposits

#### OTHER GLOMERULONEPHRITIS

- 40. Wire loop lesion seen in lupus nephritis is due to:** (JIPMER Nov 2019)
- Capillary wall thickening
  - Basement membrane thickening
  - Subepithelial deposits
  - Sclerosis of mesangium

- 41. Immunofluorescence staining pattern from a kidney biopsy from a 35yr old patient presenting with proteinuria has been shown below. What is the most probable cause?** (AIIMS Nov 18)



- FSGS
  - PSGN
  - Lupus nephritis
  - Good pasteur's syndrome
- 42. True about membranoproliferative glomerulonephritis is?** (AIIMS May 2017)

- Low C4 level
- Double basement membrane appearance on light microscopy
- Dense deposit along basement membrane on electron microscopy
- Mesangial hypocellularity
- GBM thickening


- 43. Which of the following disease has deafness as well as hematuria?** (JIPMER 2016)
- Alport Syndrome
  - Good pasture syndrome
  - IgA nephropathy
  - Cryoglobulinemia

- 44. A 4 years old child presents with rash on lower limbs, arthritis, and abdominal pain. Urine examination reveals microscopic hematuria. The most likely diagnosis is:** (Recent Question 2016-17)
- Thrombaesthesia
  - Idiopathic thrombocytopenic purpura
  - Systemic lupus erythematosus
  - Henoch Schonlein purpura

- 45. In thin basement membrane disease the defect is in** (Recent Question 2015)
- Alpha-1 and alpha-2 chains of collagen type IV
  - Alpha-3 and alpha -4 chains of collagen type IV
  - Alpha-5 chain of collagen type IV
  - Alpha-7 chain of collagen type IV

- 46. Deposits in MPGN are** (Recent Question 2015)
- Subepithelial
  - Subendothelial
  - Intramembranous
  - All of the above

- 47. Tram track appearance of glomerular capillary wall is seen in** (Recent Question 2015)
- Membranoproliferative glomerulosclerosis
  - Focal segmental glomerulosclerosis
  - IgA nephropathy
  - Good pasture syndrome



- 48. Subepithelial humps are characteristic of**  
(Recent Question 2015)
- Rapidly progressive glomerulonephritis
  - Focal segmental glomerulonephritis
  - Acute proliferative glomerulonephritis
  - Membranoproliferative glomerulonephritis
- 49. All the following are true regarding IgA nephropathy except**  
(Recent Question 2015)
- Can present as persistent microscopic hematuria
  - IgA 1 deposition in the mesangium
  - ACE inhibitors can be used
  - Decreased serum IgA level
- 50. Subendothelial deposits in glomerulus are seen in which of the following glomerulonephropathies?**  
(Recent Question 2015)
- MPGN
  - PSGN
  - Minimal change disease
  - FSGS
- 51. In renal biopsy of a 14-year-old boy with nephritic syndrome, glomeruli are showing proliferation of mesangial cells with GBM thickening and mesangial cell interposition. What is the most likely diagnosis in this case?**  
(APPGMEE 2015)
- Membranous nephropathy
  - Diffuse proliferative glomerulonephritis
  - Focal segmental glomerulosclerosis
  - Mesangiocapillary glomerulonephritis
- 52. Dysmorphic RBC with ARF is seen in?**  
(Recent Question 2013)
- Glomerular disease
  - Renal carcinoma
  - Proximal tubule disease
  - Distal tubule disease
- 53. Mutation in COL4A5 chain the diagnosis:**  
(AIIMS May 2013)
- Alport's syndrome
  - Good pasture's syndrome
  - Hereditary Non-polyposis Colon Cancer
  - Xeroderma Pigmentosum
- 54. Mesangial cells of IgA Nephropathy overexpresses:**  
(JIPMER 2012)
- CD51
  - CD61
  - CD71
  - CD81
- 55. Rapidly progressive glomerulonephritis is characterised by**  
(JIPMER 11)
- Crescents
  - Splitting of basement membrane
  - Neutrophil infiltration of the mesangium
  - Glomerulosclerosis
- 56. In which of the following are linear IgA deposits in mesangium noted:**  
(MH 11)
- Henoch Schonlein purpura
  - Malaria
  - Good Pasture's syndrome
  - Wegener's granulomatosis
- 57. Foot process effacement is seen on EM in:**  
(PGI Nov 10)
- Minimal change disease
  - Focal segmental GN
  - IgA nephropathy
  - Mesangial proliferative GN

- 58. Increased levels of  $C_3$ NeF are associated with?**  
(DNB Dec 10)
- Type I MPGN
  - Type II MPGN
  - FSGS
  - Berger Disease
- 59. Membranous Lupus Nephritis is:**  
(Recent Question 2014)
- Class II
  - Class III
  - Class IV
  - Class V

## NEPHROTIC SYNDROME


- 60. True statement about minimal change disease**  
(Recent Question 2015)
- Effacement of foot process on light microscopy
  - Severe hypoalbuminemia
  - Non selective proteinuria
  - Cyclosporine is the first line of treatment
- 61. Most common nephropathy in world is?**  
(Recent Question 2015)
- IgA nephropathy
  - FSGS
  - Minimal Change ds
  - Adult PSGN
- 62. Most common cause of nephrotic syndrome in children-**  
(Recent Question 2015, 2014)
- Membranous GN
  - Minimal change disease
  - PSGN
  - RPGN
- 63. True about light microscopy in minimal change disease is:**  
(Recent Question 2014)
- Loss of foot process seen
  - Anti GBM Abs seen
  - IgA deposits seen
  - No change seen
- 64. A child presented with frothy urine, massive proteinuria and edema. Urine examination revealed RBC nil, WBC nil, no casts, no crystal. No prior episode of similar presentation. What is your diagnosis?** (AIIMS May 2014)
- Minimal change disease
  - IgA nephropathy
  - Membranous glomerulonephritis
  - MPGN
- 65. Patients with minimal change disease are at risk of?**  
(PGI May 2014)
- Spontaneous bacterial peritonitis
  - Sepsis
  - DVT
  - Atherosclerosis
  - Renal Ca
- 66. Mutation in NPHS1 gene causes which disease?**  
(APPGMEE 14)
- Alport syndrome
  - Congenital Finnish type nephrotic syndrome
  - Focal segmental glomerulosclerosis
  - Nail patella syndrome
- 67. In which one of the primary Glomerulonephritides the glomeruli are normal by light microscopy but shows loss of foot processes of the visceral epithelial cells and no deposits by electron microscopy**  
(APPGMEE 14)
- Poststreptococcal glomerulonephritis
  - Membranoproliferative glomerulonephritis type
  - IgA nephropathy
  - Minimal change disease



68. Which of the following statement is true about Congenital nephrotic syndrome caused by Nephtrin protein mutation: (PGI May 2013)
- Cause steroid resistant nephrotic syndrome
  - Nephtrin is a key component of the slit diaphragm
  - Coded by NPHS-1 gene
  - Symptom occur only after 1st month of age
  - Autosomal dominant pattern
69. A 7-year-old girl is brought with complaints of generalized swelling of the body. Urinary examination reveals grade 3 proteinuria and the presence of hyaline and fatty casts. She has no history of hematuria. Which of the following statements about her condition is true? (AIIMS May 11)
- No IgG deposits or C3 deposition on renal biopsy
  - Her C3 level will be low
  - IgA nephropathy is the likely diagnosis
  - Alport's syndrome is the likely diagnosis
70. The most common gene defect in idiopathic steroid resistant nephrotic syndrome - (AIIMS Nov 11, May 07, Nov 06, DNB Dec 09)
- ACE
  - NPHS 2
  - HOX II
  - PAX
71. Hypercoagulation in nephrotic syndrome is caused by- (AI 10)
- Loss of antithrombin III
  - Decreased fibrinogen
  - Decreased metabolism of Vitamin K
  - Increase in protein C
72. Edema in nephrotic syndrome is due to - (AIIMS Nov 10)
- Sodium and water retention
  - Increased venous pressure
  - Hypoalbuminemia
  - Hyperlipidemia
73. True about fibronectin nephropathy are all except: (AIIMS Nov 10)
- Autosomal recessive inheritance
  - Glomerular enlargement with PAS+ trichrome+ mesangial deposit
  - Glomerulus do not consistently stain for Ig and complement
  - Ultrastructural feature is presence of large electron dense mesangial or subendothelial deposit


#### MGN

74. A 60-year-old male with complaints for frothy urine and facial puffiness. The DIF and electron microscopic picture of the patient is given below. Diagnosis is?



- Membranoproliferative nephritis (AIIMS May 2017)
- Membranous glomerulopathy
- Minimal change disease
- FSGS

75. Given below is the immunofluorescence image of kidney biopsy. Identify the condition?



- Good pasteurs syndrome (Recent Question 2016-17)
  - Membranous glomerulonephropathy
  - Berger's disease
  - Ig A nephropathy
76. Of all the types of lupus nephritis, the worst outcome is seen in? (Recent Question 2016-17)
- Minimal mesangial
  - Diffuse nephritis
  - Focal segment nephritis
  - Membranous nephritis
77. A 45-year-old man presents with hematuria. Renal biopsy demonstrates a focal necrotizing glomerulitis with crescent formation. The patient gives history of intermittent hemoptysis and intermittent chest pain of moderate intensity. A previous chest X-ray had demonstrated multiple opacities, some of which were cavitated. The patient also has chronic cold like nasal symptoms. What is the most probable diagnosis? (Recent Question 2016-17)
- Aspergillosis
  - Polyarteritis nodosa
  - Renal carcinoma metastatic to lung
  - Wegner's granulomatosis
78. Most common nephropathy associated with malignancy is? (AIIMS May 2015)
- Membranous
  - MCD
  - IgA
  - FSGS
79. All the following are true regarding minimal change disease except (Recent Question 2015)
- Most common cause of nephrotic syndrome in children
  - Massive proteinuria
  - Hematuria
  - Good response to steroids
80. Subendothelial deposits are seen in (Recent Question 2015)
- Membranous nephropathy
  - MPGN type I
  - MPGN type II
  - IgA nephropathy
81. Which component of HBV causes glomerulonephritis- (AIIMS May 2011)
- HbeAg
  - HBcAg
  - HBsAg
  - Anti HBs Ag antibody

#### FSGS

82. Collapsing variant of FSGS is seen in (Recent Question 2015)
- NSAIDs
  - Heroin abuse
  - HIV
  - CMV






- 83. False statement in collapsing glomerulopathy:** (AP 2012)
- It is a morphologic variant of FSGS (focal segmental glomerulosclerosis)
  - Microscopically retraction and /or collapse of the entire glomerular tuft is seen
  - There is no proliferation or hypertrophy of glomerular visceral epithelial cells
  - It may be idiopathic or HIV associated nephropathy
- 84. HIV infection causes -** (Recent Question 2014)
- Membranous glomerulonephritis
  - IgA glomerulopathy
  - Collapsing glomerulonephritis
  - RPGN
- 85. A person with radiologically confirmed reflux nephropathy develops nephrotic range proteinuria. Which of the following would be the most likely histological finding in the patient?** (AIIMS Nov 11)
- Focal segmental glomerulosclerosis
  - Nodular glomerulosclerosis
  - Membranous glomerulopathy
  - Proliferative glomerulonephritis with crescents
- 86. In glomerulus subendothelial deposits are seen in-** (Recent Question 2014)
- Good pasture syndrome
  - IgA nephropathy
  - MPGN type I
  - MPGN type II
- 87. Renal biopsy of a 14-year-old boy with nephritic syndrome shows proliferation of mesangial cells with Glomerular basement membrane thickening and mesangial cell interposition. What is the most likely diagnosis?** (APPGMEE 14)
- Diffuse proliferative Glomerulonephritis
  - Mesangiocapillary Glomerulonephritis
  - Membranous nephropathy
  - Hemolytic Uremic syndrome
- 88. Type I membrano proliferative Glomerulonephritis is commonly associated with all EXCEPT** (APPGMEE 14)
- SLE
  - Persistent hepatitis C infections
  - Partial lipodystrophy
  - Neoplastic diseases

#### DIABETIC NEPHROPATHY

- 89. A 50-year-old male presented with blurring of vision. Urine examination showed proteinuria. Fundus examination showed dot and blot haemorrhages, microaneurysm and cotton wool spots. Histopathology pic of kidney given below. Your diagnosis?** (AIIMS May 2017)




- Kimmelstiel Wilson nodules
- Crescents
- Amyloid
- Segmental sclerosis

- 90. Nodular glomerulosclerosis is seen in** (Recent Question 2015)
- Diabetes mellitus
  - Malignant hypertension
  - Amyloidosis
  - Multiple myeloma
- 91. Kimmelstiel-Wilson lesion is characteristic of-** (Recent Question 13, 2014, AI 98, MH 2016)
- HIV nephropathy
  - Diabetic nephropathy
  - Amyloidosis
  - Malignant hypertension
- 92. Characteristic type of nephropathy in DM is-** (DNB June 11, COMED 09)
- Focal
  - Diffuse
  - Nodular
  - Crescent


#### PYELONEPHRITIS & PAPILLARY NECROSIS

- 93. A patient presented with pus in urine. Urine culture was done which was negative. After a sudden onset renal failure the patient died. On autopsy the following finding was seen in kidney. What is the most likely diagnosis?** (JIPMER 2017)



- TB kidney
  - Infected renal cysts
  - Renal cell carcinoma
  - Renal stones
- 94. Characteristic feature of benign nephrosclerosis** (Recent Question 2015)
- Leather grain appearance
  - Flea bitten appearance
  - Onion skin appearance
  - Hyperplastic arteriosclerosis
- 95. Papillary necrosis is seen in all the following except** (Recent Question 2015)
- NSAIDs
  - Diabetes mellitus
  - Sickle cell anemia
  - Shock
- 96. The most common infectious agent associated with chronic pyelonephritis is -** (Recent Question 2014, AIIMS May 2003)
- Proteus vulgaris
  - Klebsiella pneumonia
  - Staphylococcus aureus
  - Escherichia coli
- 97. ENaC mutation is seen in:** (Recent Question 2015)
- Liddle syndrome
  - Gordon syndrome
  - Bartter syndrome
  - Gitelman syndrome
- 98. Least likely cause of renal papillary necrosis** (APPGMEE 14)
- Sickle cell disease
  - Analgesic nephropathy
  - Posterior urethral valves
  - Diabetes with UTI





**99. Acquired cystic diseases rather than inherited cause a 12- to 18-fold increased risk of renal cell carcinoma, which develops in 7% of dialyzed patients observed for 10 years. All of the following are seen in adult Polycystic kidney disease except? (PGI Nov 2016)**

- a. Large Kidney
- b. Small kidney
- c. Multilobulated on USG
- d. Cysts in liver and brain
- e. 10% progresses to ESRD

**100. Regarding adult polycystic kidney disease, which one of the following statements is not correct? (Recent Question 2016-17)**

- a. Inherited as autosomal dominant with 100% penetrance
- b. Often associated with hepatic cysts
- c. Associated with increased incidence of subarachnoid haemorrhage
- d. Renal cell carcinoma is a frequent association


**101. Hepatic fibrosis is found in (Recent Question 2016)**

- a. Medullary cystic kidney
- b. ADPKD
- c. ARPKD
- d. Nephrophthosis

**102. Chromosome a/w ADPKD (Recent Question 2016)**

- a. 14 and 16
- b. 14 and 13
- c. 16 & 4
- d. 12 & 16

**103. Gross section of kidney depicts what? (APPGMEE 2015)**



- a. RCC
- b. Medullary sponge kidney
- c. Hydatid cyst
- d. Polycystic kidney disease

**104. Balkan nephropathy is caused by? (APPGMEE 2015)**

- a. Fungal toxins
- b. Lead
- c. Calcineurin inhibitors
- d. Aristocholic acid

**105. True about Adult polycystic kidney disease: (PGI May 12)**

- a. Autosomal dominant pattern of inheritance
- b. 80% develop Nephrolithiasis
- c. Caused by mutation in two genes
- d. Somatic mutation can also occur
- e. Loss of heterozygosity may occur


**106. Which of the following statements about Xanthogranulomatous pyelonephritis is not true - (AI 11, 09)**

- a. Foam cells are seen
- b. Associated with tuberculosis
- c. Yellow nodules are seen
- d. Giant cells may be seen

## VASCULAR DISEASES

**107. What is your diagnosis in the figure given below?**

(Recent Question 2016-17)



- a. Flea bitten kidney in malignant hypertension
- b. Amyloidosis
- c. Ch. Glomerulonephritis
- d. Glomerulosclerosis

**108. A 40-year-old hypertensive male was admitted to the hospital with sudden onset of headache and altered sensorium. On Examination his Blood Pressure was observed to be 220/110mm Hg and the patient died four later. What is likely pathological finding in his kidneys - (AI 12)**

- a. Small kidney with granular surface
- b. Small kidney with petechial hemorrhages
- c. Large kidney with waxy appearance
- d. Large kidney with granular surface

## RENAL TUMORS

**109. Sickle cell anemia leads to which variety of Renal cell ca? (JIPMER 2017)**

- a. Medullary
- b. Papillary
- c. Chromophobic
- d. Belini duct Ca

**110. Loss of y chromosome is seen in in which Renal cell Ca? (JIPMER 2017)**

- a. Papillary
- b. Chromophobe
- c. Chromophilic
- d. Belini duct Ca

**111. MC tumors of kidney in children? (PGI Nov 2016)**

- a. Wilms tumor
- b. Neuroblastoma
- c. PCKD
- d. Angioliposarcoma
- e. Malignant rhabdoid tumor

**112. Chromosomal mutation in low grade transitional cell cancer of bladder is? (JIPMER 2016)**

- a. P53
- b. Rb gene
- c. FGFR3
- d. HRAS

**113. Most common histological type of renal cell cancer? (Recent Question 2016-17)**

- a. Clear cell ca
- b. Clear cell sarcoma
- c. Metaplastic nephroma
- d. Xp translocation ca

**114. Hobnail pattern is seen in which type of renal carcinoma (Recent Question 2015)**

- a. Clear cell carcinoma
- b. Papillary carcinoma
- c. Chromophobe cell cancer
- d. Collecting duct carcinoma



**115. Multiple chromosomal losses are associated with the following type of renal cancer** (Recent Question 2015)

- a. Clear cell carcinoma
- b. Papillary carcinoma
- c. Chromophobe cell cancer
- d. Collecting duct carcinoma

**116. Most common histological variant of renal cell carcinoma is:**

- a. Papillary carcinoma (Recent Question 2015)
- b. Clear cell carcinoma
- c. Chromophobe renal carcinoma
- d. Collecting duct carcinoma

**117. Most common cystic disease of kidney is?** (Recent Question 2015)

- a. X linked recessive polycystic kidney disease
- b. X linked dominant polycystic kidney disease
- c. Autosomal recessive polycystic kidney disease
- d. Autosomal dominant polycystic kidney disease

**118. Most common gene associated with renal cell carcinoma is?** (Recent Question 2015)

- a. WT-1
- b. BRCA-1
- c. VHL
- d. PATCH

**119. Clear cell variety of Renal cell carcinoma is related to gene located on chromosome:** (Recent Question 2014)

- a. 3
- b. X
- c. 22
- d. 20

**120. 1 month post renal transplant a patient developed fever. Which of the following is the most likely organism responsible?** (AIIMS May 2014)

- a. Polyoma virus/ BK virus
- b. Hepatitis C virus
- c. Varicella virus
- d. HHV-6

**121. An emerging viral pathogen causing pyelonephritis in kidney allografts** (Recent Question 2015)

- a. Polyoma virus
- b. Marburg virus
- c. Ebola virus
- d. JC virus

**122. Histopathology showing large cells with plant like appearance with perinuclear halo is seen in which type of renal cell carcinoma?** (PGI Nov 12)

- a. Oncocytoma
- b. Papillary cell carcinoma
- c. Transitional cell Carcinoma
- d. Chromophobe RCC
- e. Clear cell carcinoma

**123. Clear cells are seen in:** (PGI May 2011)

- a. Rhabdomyosarcoma
- b. Adrenal cell carcinoma
- c. Renal cell Carcinoma
- d. Hodgkin's lymphoma
- e. NHL

**124. Chromophobe variant of renal cell carcinoma is associated with -** (AI 10)

- a. VHL gene mutations
- b. Trisomy 7 and 17
- c. 3 p deletions
- d. Monosomy of 1 and Y

**125. An emerging organism responsible for causing Pyelonephritis in Renal Allografts is?** (DNB Dec 10)

- a. Polyoma virus
- b. Herpes virus
- c. Hepatitis B virus
- d. Rota virus

**126. Findings of multiple myeloma in kidney are all except-** (DPG 10)

- a. Tubular casts
- b. Amyloidosis
- c. Wire loop lesions
- d. Renal tubular necrosis

#### DISEASES OF URINARY BLADDER

**127. Most common type of bladder cancer** (Recent Question 2015)

- a. Squamous cell carcinoma
- b. Urothelial cell carcinoma
- c. Adenocarcinoma
- d. Carcinoid

**128. Michaelis Gutmann bodies are seen in -** (Recent Question 2014, 2013/ AIIMS May 07/ DNB Dec 09)

- a. Xanthogranulomatous pyelonephritis
- b. Malakoplakia
- c. Nail patella syndrome
- d. Chronic Pyelonephritis

**129. 90% of bladder cancer arise from-** (Recent Question 2014)

- a. Squamous cells
- b. Glandular cells
- c. Transitional cells
- d. Smooth muscle cells


**130. All are carcinogenic for bladder except?** (Recent Question 2014)

- a. Benzidine
- b. Isopropyl alcohol
- c. Acrolein
- d. Phenacetin



## Answers with Explanations

1. Ans. (c) **Fabry disease**



Round particles producing birefringent Maltese crosses under polarized light are commonly seen in the urinary sediment of patients with a nephrotic syndrome. The appearance of the Maltese crosses is due to the birefringence of lipid droplets, which consist mainly of cholesterol esters. These may also be seen in Fabry disease.

2. Ans. (a) **Glomerulonephritis**

3. Ans. (a, b, c, e) a. **Hyaline casts**- may be normally present in healthy person; b. **Muddy brown casts**- acute tubular necrosis; c. **WBC cast** - pyelonephritis; e. **Myoglobin cast** - Rhabdomyolysis

4. Ans. (a, b) a. **Centrifuge the urine first**; b. **Use immunofluorescence light**

5. Ans. (d) **Hyaline casts**

The given figure shows urine cytology in samples obtained during clinical rejection showing **numerous renal tubular cells, lymphocytes, and macrophages** along with few **Hyaline casts** (as shown in the arrow).

6. Ans. (c) **Creatinine clearance rate**

7. Ans. (a) **A/4 B/3 C/1 D/2**

8. Ans. (a) **Oxalate** (Ref: Henry 22nd/457)

9. Ans. (c) **Struvite** (Ref: Henry 22nd/457)

10. Ans. (a, b, d); a. **WBC cast**-Acute pyelonephritis b. **Broad cast**-CRF d. **RBC cast**-Glomerulonephritis

(Ref: Henry 22nd/457)

11. Ans. (a) **3.1-7 mg/dl** (Ref: Harrison 18th/pg Appendix)

### Normal levels of Uric acid

<b>Females</b>	0.15–0.33 mmol/L	2.5–5.6 mg/dL
<b>Males</b>	0.18–0.41 mmol/L	3.1–7.0 mg/dL

12. Ans. (c) **Proteoglycan** (Ref: Robbins 9th/pg 900; 8th/pg 908)

**Glomerular basement membrane (GBM) contains Poly-anionic proteoglycans (negatively charged) (mostly heparan sulfate)** and hence repels negatively charged proteins.

13. Ans. (c) **RBC cast**

(Ref: Henry's 22nd/ 466-469) Refer to Annexure

Cast seen in Acute Glomerulonephritis is RBC cast, which is pathognomonic

14. Ans. (b) **5** (Ref: Henry's 22nd/ 457-458)

### Normal findings in analysis of urine sediment

- **Red blood cells:** 0–2/high-power field
- **White blood cells:** 0–2/high-power field
- **Bacteria:** None
- **Casts:** None except hyaline casts
- **Epithelial cells:** None

15. Ans. (a) **Isomorphic RBCs** (Ref: Henry's 22nd/ 457-458)

- **Hematuria:** Presence of >3 RBCs/hpf of urine
- Hematuria can be isomorphic (morphologically normal RBCs) or dysmorphic (RBCs with cellular protrusions or fragmentation)
- Hematuria due to nephrolithiasis lead to isomorphic RBCs with glomerulonephritis lead to dysmorphic RBCs.

### Etiology of Hematuria

Isomorphic RBCs		Dysmorphic RBCs
Upper Tract	Lower Tract	Renal disease:
<ul style="list-style-type: none"> <li>• Urolithiasis</li> <li>• Pyelonephritis</li> <li>• Renal cell cancer</li> <li>• Transitional cell Ca</li> <li>• Urinary obstruction</li> <li>• Benign hematuria</li> </ul>	<ul style="list-style-type: none"> <li>• Bacterial cystitis (UTI)</li> <li>• Benign prostatic hyperplasia</li> <li>• Strenuous exercise ("marathon runner's hematuria")</li> <li>• Transitional cell carcinoma</li> <li>• Spurious hematuria (e.g. menses)</li> <li>• Instrumentation</li> <li>• Benign hematuria</li> </ul>	<ul style="list-style-type: none"> <li>• Glomerulonephritis</li> <li>• Lupus nephritis</li> <li>• IgA nephropathy (Berger's disease)</li> <li>• Thin glomerular basement membrane disease</li> <li>• Hereditary nephritis (Alport's syndrome)</li> </ul>



16. Ans. (a, b, e); a. Glomerular basement membrane; b. Tubule; e. Mesangial matrix

(Ref: Rosai 10th/pg 1102-1104)

PAS positive structures are GBM, tubule & mesangial matrix

17. Ans. (c) PSGN (Ref: Robbins 9th/pg 909)

18. Ans. (a) Nephrotic range proteinuria b. Neutrophilic infiltration of tubules, d. Linear deposits along glomerular basement membrane

(Ref: Robbins 9th/ 909-12)

19. Ans. (b) Renal failure

(Ref: Robbins 9th/pg 909; 8th/pg 917)

20. Ans. (c) FSGS (Ref: Robbins 9th/pg 918; 8th/pg 926)

Most common renal involvement in HIV is FSGS.

21. Ans. (a) Epithelial humps (Ref: Robbins 9th/pg 909-912)

Electron microscopy is PSGN shows

Discrete, amorphous, electron-dense SUBEPITHELIAL deposits ("Humps")<sup>Q</sup>

22. Ans. (d) Deposition of IgA (Ref: Robbins 9th/pg 909-912)

23. Ans. (b) Immune complex mediated

(Ref: R 9th/pg 909-912)

24. Ans. (a) C3 decreases & ASO increases

(Ref: Robbins 9th/pg 909-912; 8th/pg 917-919)

25. Ans. (b) Caused by non group A hemolytic streptococci

(Ref: Robbins 9th/pg 909-912; 8th/pg 917-919)

Both Post-streptococcal reactive arthritis and Acute Rheumatic fever are caused by non-group A hemolytic streptococci

26. Ans. (d) Anti GBM antibodies (Ref: R 9<sup>th</sup> ed pg 912)

EM image show rupture of GBM. With a history to proteinuria and hypertension and high creatinine, RPGN can be an important differential.

27. Ans. (b) Flea bitten kidney (Ref: Robbins 9th/pg 912)

28. Ans. (d) Type 2- SLE nephritis

29. Ans. (b) Good Pasture syndrome

30. Ans. (c) Crescents (Ref: Robbins 9th ed. Pg. 912,)

31. Ans. (d) Good pasture syndrome

(Ref: Robbins 9th/pg 912)

32. Ans. (a, c) a. Generalize edema, c. Hypoalbuminemia

(Ref: Robbins 9th/pg 909)

Acute nephritic syndromes classically present with hypertension, hematuria, red blood cell casts, pyuria, and mild to moderate proteinuria.

33. Ans. (c) Recurrent gross hematuria following respiratory infection

(Ref: Robbins 9th/pg 909; 8th/pg 917)

IgA nephropathy is characterized by: Recurrent hematuria and presence of IgA deposits in the mesangium

34. Ans. (a) Amyloidosis

(Ref: Robbins 9th/pg 261; 8th/pg 254; Harrison 18th/pg 947)

Kidney involvement in Amyloidosis

- Deposition of monoclonal Ig or  $\lambda$  (or kappa) light chains in the GBM
- Distinctive nodular glomerular lesions resulting from the deposition of non-fibrillar light chains.
- Glomeruli show PAS-positive mesangial nodules, lobular accentuation & mild mesangial hypercellularity.

35. Ans. (d) Focal segmental glomerulosclerosis

(Ref: Robbins 9th/pg 912; 8th/pg 920)

- Most common cause of nephrotic syndrome in children is Minimal change disease
- Most common cause of nephrotic syndrome in adults: FSGS > MGN

36. Ans. (b) Polyarteritis nodosa

(Ref: Robbins 9th/pg 912)

37. Ans. (a) Necrotizing hemorrhagic interstitial pneumonitis

(Ref: Robbins 9th/pg 912-913; 8th/pg 920-921)

38. Ans. (b) Serum antibodies against alpha 3 NC1 domain of collagen - IV (Ref: Robbins 9th/pg 912-913)

Goodpasture syndrome

- It is type I RPGN & a type II hypersensitivity reaction.
- It is characterized by serum antibodies against alpha 3 NC1 domain of collagen - IV.
- This shows linear GBM fluorescence, while all other Glomerulonephritis, which are type 3 Hypersensitivity reactions show granular immune deposits.

39. Ans. (a, b, c, d); a. Antibody to  $\alpha$ -chain of Type IV collagen (COL-4A); b. Basement membrane involvement; c. Pulmonary hemorrhage; d. Crescent formation

(Ref: Robbins 9th/pg 912-913; 8th/pg 920-921)

40. Ans. (a) Capillary wall thickening (Ref: R 9<sup>th</sup> pg 222)





**41. Ans. (c) Lupus nephritis**

Here, the typical "full-house" pattern with intense (++++) granular staining for IgA, IgG, IgM, kappa & lambda & C1 and C3 in a diffuse mesangiocapillary pattern can be seen which is suggestive of SLE.

Good Pasture will show linear IgG and C3 deposits in basement membrane.

PSGN will show granular deposits of IgG & C3 and sometimes IgM in the mesangium and in sometimes GBM

FSGS will show IgM & C3 in sclerotic area or mesangium

**42. Ans. (a, b, c) a. Low C4 level; b. Double basement membrane appearance on light microscopy; c. Dense deposit along basement membrane on electron microscopy**

**43. Ans. (a) Alport syndrome**

**44. Ans. (d) Henoch Schonlein purpura (Ref: R 9/909-912)**

The clinical history is suggestive of nephritic syndrome with arthritis and purpura which is a triad of Henoch Schonlein purpura

**45. Ans. (b) Alpha-3 and alpha-4 chains of collagen type IV**

(Ref: Robbins 9th/pg 925; 8th/pg 932)

**Thin basement membrane disease (TBMD)**

- It is characterized by persistent or recurrent hematuria, but is not typically associated with proteinuria, hypertension, or loss of renal function or extrarenal disease.
- Genetic defects in *COL(IV) α3/COL(IV) α4* loci of type IV collagen
- Autosomal dominant or recessive inheritance
- GBM shows diffuse thinning compared to normal values for the patient's age in otherwise normal biopsies

**46. Ans. (d) All of the above (Ref: Robbins 9th/pg 920)**

- In MPGN type I, **Subendothelial** electron-dense deposits are most commonly seen
- **Mesangial/ subepithelial** deposits may also be present
- In MPGN type II, intramembranous deposits seen

**47. Ans. (a) Membranoproliferative glomerulosclerosis**

(Ref: Robbins 9th/pg 920; 8th/pg 928)

**48. Ans. (c) Acute proliferative glomerulonephritis**

(Ref: Robbins 9th/pg 910; 8th/pg 920)

**49. Ans. (d) Decreased serum IgA level (Ref: R 9th/pg 923)**

Serum IgA level is increased in IgA nephropathy

**50. Ans. (a) MPGN (Ref: Robbins 9th/pg 920; 8th/pg 928)**

**51. Ans. (d) Mesangiocapillary glomerulonephritis**

(Ref: Robbins 9th/pg 920; 8th/pg 928)

Renal biopsy of a young male with nephritic syndrome showing mesangial cell proliferation with Glomerular

basement membrane thickening and mesangial cell interposition is typical biopsy finding of **Mesangiocapillary Glomerulonephritis, also called MPGN**

**52. Ans. (a) Glomerular disease**

(Ref: Henry's 22nd/ 457-458)

Dysmorphic RBCs with ARF is seen in Glomerular disease, while isomorphic RBCs are seen in non-glomerular bleeds;

**53. Ans. (a) Alport's syndrome**

(Ref: Robbins 9th/pg 924-925; 8th/pg 931-932)

**54. Ans. (c) CD71 (Ref: Robbins 9th/pg 923-924)**

**55. Ans. (a) Crescents (Ref: Robbins 9th/pg 912-913)**

**56. Ans. (a) Henoch Schonlein purpura**

(Ref: Robbins 9th/pg 923-924)

**57. Ans. (a, b, c, d); a. Minimal change disease; b. Focal segmental GN; c. IgA nephropathy; d. Mesangial proliferative GN**

(Ref: Robbins 9th/pg 917-918; 8th/pg 924-926)

**Foot process effacement** is seen on EM in:

Minimal change disease, Focal segmental GN, IgA nephropathy, Mesangial proliferative GN

**58. Ans. (b) Type II MPGN (Ref: Robbins 9th/pg 920-921)**

Increased levels of C<sub>3</sub>NeF are associated with **Type II MPGN**.

**59. Ans. (d) Class V (Ref: Robbins 9th/pg 222-224)**

**Renal involvement in SLE:**

**Class I: Minimal Mesangial Lupus Nephritis<sup>Q</sup>**

**Class II: Mesangial Proliferative Lupus Nephritis**

**Class III: Focal Lupus Nephritis<sup>Q</sup>**

**Class IV: Diffuse Lupus Nephritis<sup>Q</sup>**

**Class V: Membranous Lupus Nephritis<sup>Q</sup>**

**Class VI: Advanced Sclerotic Lupus Nephritis**

- 50% patients of SLE have clinically significant **renal involvement<sup>Q</sup>**

- **Class I Lupus Nephritis Class I Lupus Nephritis is least common<sup>Q</sup> & class IV is the most common pattern<sup>Q</sup>**

**"Wire loop lesions"<sup>Q</sup>** characteristically seen in **Class IV Lupus Nephritis<sup>Q</sup>** is due to **sub-endothelial** immune complex deposits, on **light microscopy**- Also seen in **Class III/V Lupus nephritis<sup>Q</sup>**

**60. Ans. (b) Severe hypoalbuminemia (Ref: R 9th/pg 914)**

a. False	Effacement of foot process is seen on electron microscopy, while light microscopy is normal in minimal change disease
b. True	<b>Severe hypoalbuminemia is a feature of nephrotic syndrome due to minimal change disease</b>
c. False	c. Selective proteinuria
d. False	d. Cyclosporine is the first line of treatment



61. Ans. (a) **IgA nephropathy** (Ref: Robbins 9th/pg 914)  
Most common nephropathy in world is IgA nephropathy

62. Ans. (b) **Minimal change disease** (Ref: R 9th/pg 917-918)

63. Ans. (d) **No change seen** (Ref: Robbins 9th/pg 917-918)

On light microscopy

Glomeruli appear normal<sup>Q</sup> minimal change disease

64. Ans. (a) **Minimal change disease** (Ref: R 9th/pg 917-918)

The child in this question presents with **first episode of massive proteinuria, frothy urine suggestive of lipiduria** and edema. Urine microscopy finding is not significant. This is suggestive of **nephrotic syndrome**, for which, the **most common cause in children is Minimal change disease**.

65. Ans. (a, b, c, d); a. **Spontaneous bacterial peritonitis**; b. **Sepsis**; c. **DVT**; d. **Atherosclerosis**

(Ref: R 9th/pg 917-918)

Patients with minimal change disease are at risk of Spontaneous bacterial peritonitis, Sepsis, DVT and Atherosclerosis

**Complications of Nephrotic syndrome:**

<b>Infection<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>Staphylococcal and Pneumococcal infection. Eg Spontaneous bacterial peritonitis, Sepsis</li> <li>Due to <b>loss of immunoglobulins<sup>Q</sup></b> in the urine</li> </ul>
<b>Thrombotic &amp; thromboembolic complications</b>	<ul style="list-style-type: none"> <li>Due to loss of endogenous anticoagulants (<b>Antithrombin-III, protein C, protein S</b>)<sup>Q</sup> in urine</li> <li><b>Hypercoagulable state</b> → Renal vein thrombosis / Deep vein thrombosis (DVT)</li> <li>Particularly in <b>membranous nephropathy<sup>Q</sup></b></li> </ul>
<b>Atherosclerosis</b>	<ul style="list-style-type: none"> <li>Due to hyperlipidemia</li> </ul>
<b>Loss of other low molecular wt proteins</b>	<ul style="list-style-type: none"> <li><b>Transferrin</b> leads to <b>Iron deficiency anemia</b></li> <li><b>Ceruloplasmin</b> leads to <b>copper deficiency anemia</b></li> <li><b>Cholecalciferol binding protein</b> leads to <b>Vit D deficiency</b></li> <li><b>Thyroglobulin binding protein</b> leads to <b>hypothyroidism</b></li> </ul>

66. Ans. (b) **Congenital Finnish type nephrotic syndrome**

(Ref: Nelson 19 thedpg 1803)

Mutation in NPHS1 gene causes Congenital Finnish type nephrotic syndrome

67. Ans. (d) **Minimal change disease** (Ref: R 9th/pg 917-918)

68. Ans. (a, b, c); a. **Cause steroid resistant nephrotic syndrome**; b. **Nephrin is a key component of the slit diaphragm**; c. **Coded by NPHS-1 gene**

(Ref: Nelson 19th/pg 1802)

- Congenital nephrotic syndrome Finnish type is a type of steroid resistant nephrotic syndrome

- It is caused by NPHS-1 gene mutation, which codes for Nephrin protein, a key component of the slit diaphragm

69. Ans. (a) **No IgG deposits or C3 deposition on renal biopsy**

(Ref: Robbins 9th/pg 917-918)

In the given scenario, a 7 year old girl presents with generalized swelling, massive proteinuria, but no hematuria and presence of hyaline and fatty casts. Therefore, she is suffering from Nephrotic Syndrome, the commonest cause of which is **Minimal change disease, in which there is no IgG or C3 deposition** on renal biopsy.

70. Ans. (b) **NPHS 2** (Ref: Nelson 19th ed/pg 1802)

**Most common gene defect in idiopathic steroid resistant nephrotic syndrome NPHS 2 which codes for podocin**;

71. Ans. (a) **Loss of antithrombin III**

(Ref: Robbins 9th/pg 914; 8th/pg 922)


72. Ans. (c) **Hypoalbuminemia** (Ref: Harrison 19th ed/pg 253)

73. Ans. (a) **Autosomal recessive inheritance**

(Ref: Rosai 10th/pg 1153)

**Fibronectin glomerulopathy**

<b>Definition</b>	Characterized by massive fibronectin deposition in the glomeruli.
<b>Genetic basis</b>	Caused by <b>mutations of FN1 gene on chromosome 2q34</b>
<b>Inheritance</b>	Hereditary <b>autosomal dominant (option A)</b> disease; <b>Both sexes</b> are equally affected
<b>Clinical features</b>	<b>Proteinuria</b> (nephrotic range), <b>microscopic hematuria</b> and slow <b>deterioration of renal function</b> over a period of several years.
Light microscopy	<ul style="list-style-type: none"> <li><b>Glomeruli appear enlarged &amp; lobulated</b> with minimal degree of hypercellularity</li> <li>Most characteristic feature: <b>marked enlargement of mesangium &amp; subendothelial (option D) space</b> due to <b>deposition of a homogeneous PAS +ve, Congo red -ve (option B) material</b></li> </ul>
<b>Electron microscopy</b>	<b>Dense granular appearance of the deposit in mesangium or subendothelium</b>
Immunohistochemical studies	<ul style="list-style-type: none"> <li><b>Strong positivity for fibronectin</b> in the areas corresponding to the deposits.</li> <li>Scanty immunoglobulins and complement factors may occasionally be seen, inconsistently. (option C)</li> </ul>
<b>Prognosis</b>	<b>Poor prognosis</b>
Recurrence after renal transplant	The disease has been reported to <b>recur after renal transplantation</b> .



**74. Ans. (b) Membranous glomerulopathy**

(Ref: Robbins 9th ed p 915, 920)

**The given image shows Membranous glomerulopathy with Diffuse capillary wall thickening in EM image and granular (IgG and C3 deposits) in immunofluorescence.**

**75. Ans (b) Membranous glomerulonephropathy**

(Ref: Robbins 9th/pg 231-234; 8th/pg 221-229)

- Whenever you get an immunofluorescence image in a kidney biopsy, you must try to see if the deposits look granular (lumpy bumpy) or linear.
- If its linear the answer will be Good Pasteur's syndrome (Anti GBM antibody ds)
- If it is granular, it can be any other immune complex disease.

Among the given options, the only immune complex deposition disease is Membranous nephropathy.

**76. Ans. (d) Membranous nephritis**

**77. Ans. (d) Wegner's granulomatosis**

**78. Ans. (a) Membranous**

(Ref: Robbins 9th/pg 915; 8th/pg 922)

Most common nephropathy associated with malignancy is **Membranous in adults & minimal change in children**

**79. Ans. (c) Hematuria (Ref: Robbins 9th/pg 917; 8th/pg 924)**

**80. Ans. (b) MPGN type I**

(Ref: Robbins 9th/pg 920; 8th/pg 928)

**81. Ans. (c) HBsAg (Ref: Robbins 9th/pg 914; 8th/pg 922)**

- HBsAg causes Glomerulonephritis with Nephrotic syndrome
- HBsAg, immunoglobulin, and C3 deposition has been found in the glomerular basement membrane.

**82. Ans. (c) HIV (Ref: Robbins 9th/pg 920; 8th/pg 928;**

Collapsing variant of FSGS is seen in HIV

**83. Ans. (c) There is noproliferation or hypertrophy of glomerular visceral epithelial cells**

(Ref: Robbins 9th/pg 920; 8th/pg 928)

**HIV-ASSOCIATED NEPHROPATHY (HIVAN)** is characterized by collapse of entire glomerular tuft along with proliferation and hypertrophy of glomerular visceral epithelial cells<sup>o</sup>

**84. Ans. (c) Collapsing glomerulonephritis**

(Ref: Robbins 9th/pg 920)

**85. Ans. (a) Focal segmental glomerulosclerosis**

(Ref: Robbins 9th/pg 918-919; 8th/pg 926-927)

**Nephrotic syndrome** in a patient with **reflux nephropathy** is caused by **Focal segmental glomerulosclerosis**, as a component of the adaptive response to loss of renal tissue

**86. Ans. (c) MPGN type I (Ref: Robbins 9th/pg 920-921)**

**87. Ans. (b) Mesangiocapillary Glomerulonephritis**

(Ref: Robbins 9th/pg 920-921; 8th/pg 928-929)

**88. Ans. (c) Partial lipodystrophy**

(Ref: Robbins 9th/pg 920-921; 8th/pg 928-929)

Types II & III MPGN are associated with complement factor H deficiency, presence of C<sub>3</sub> nephritic factor, partial lipodystrophy (type II MPGN) or complement receptor deficiency (type III MPGN)

**89. Ans. (a) Kimmelstiel Wilson nodules**

(Ref: Robbins 9th ed p 1118)

**Diabetic Nephropathy:**

The renal lesions include mainly: Glomerular lesions:

- Basement membrane thickening and increased mesangial matrix in patients
- Diffuse glomerulosclerosis
  - Increase in mesangial matrix associated with PAS+ basement membrane thickening, eventually obliterates mesangial cells, found in most individuals with disease of **more than 10 years duration**
- **Nodular glomerulosclerosis**
  - Inter-capillary glomerulosclerosis or Kimmelstiel-Wilson disease

**90. Ans. (a) Diabetes mellitus (Ref: Robbins 9th/pg 1118)**

**Most specific histological lesion in diabetic nephropathy** (see chapter 18 Endocrine system) is Nodular glomerulosclerosis or **Kimmelsteil -Wilson lesions**

**91. Ans (b) Diabetic nephropathy (Ref: R 9th/pg 1118-1119)**

**92. Ans. (c) Nodular (Ref: Robbins 9th/pg 1118-1119)**

**93. Ans. (a) TB kidney**

The clinical history is suggestive of sterile pyuria. The gross morphology shows greyish white are caseating necrotic material which is formed in patches, predominantly in the cortical areas involving the whole circumference of the kidney. Hence, the first possibility is renal TB.

**94. Ans. (a) Leather grain appearance (Ref: R 9th/pg 938)**

**95. Ans. (d) Shock (Ref: Robbins 9th/pg 939; 8th/pg 930)**

**Major Causes of Papillary Necrosis**

- Analgesic nephropathy (Most common)<sup>o</sup>
- Sickle cell nephropathy
- Diabetes with urinary tract infection

**96. Ans. (d) Escherichia coli (Ref: Robbins 9th/pg 933-934)**

**The most common infectious agent associated with chronic pyelonephritis is Escherichia coli**



97. Ans. (a) **Liddle syndrome** (Ref: Harrison 18th/Ch 285)

98. Ans. (c) **Posterior urethral valves**

(Ref: Robbins 9th/pg 932)

99. Ans. (b) **Small kidney** (Ref: Robbins 9/945)

100. Ans. (d) **Renal cell carcinoma is a frequent association**

(Ref: Robbins 9/949-952)

101. Ans. (c, d); c. **ARPKD**, d. **Nephrophthosis**

(Ref: Robbins 9th/pg 945; 8th/pg 956; Refer to pretext)

102. Ans. (c) **16 & 4** (Ref: Robbins 9th/pg 945; 8th/pg 956)

*PKD1* gene is located on chromosome 16p13.3, that encodes a large integral membrane protein named *polycystin-1*. *PKD2* gene, located on chromosome 4q21, encodes *polycystin-2* that accounts for most of the remaining cases of polycystic disease.

103. Ans. (d) **Polycystic kidney disease** (Ref: R 9th/pg 945)

104. Ans. (d) **Aristolochic acid** (Ref: Robbins 9th/pg 929)

**Aristolochic Nephropathy or Balkans Nephropathy:**

- **Chronic tubulointerstitial nephritis** caused by **aristolochic acid**, a supplement found in some herbal remedies in Europe in **Balkan region**
- Associated with **urothelial atypia**, occasionally culminating in tumors of renal pelvis & urethra.
- **Drug forms covalent adducts** with DNA and causes renal failure and interstitial fibrosis associated with a relative paucity of infiltrating leukocytes.
- **Increased incidence of carcinoma** in the kidney and urinary tract.

105. Ans. (a, c, d, e); a. **Autosomal dominant pattern of inheritance**; c. **Caused by mutation in two genes**; d. **Somatic mutation can also occur**; e. **Loss of heterozygosity may occur**

(Ref: Robbins 9th/pg 945-947 Harrison 18th/pg Chapter 284)

- Over **90%** of cases are inherited as an **autosomal dominant** trait, remainder due to spontaneous mutations. (**option A**)
- Mutations in the ***PKD-1*** gene on chromosome 16 (ADPKD-1) account for **85% of cases**, and mutations in the ***PKD-2*** gene on chromosome 4 (ADPKD-2) account for the remainder. (**option C**)
- Direct mutation analysis of isolated cysts suggests that there is **loss of heterozygosity (option E)**, whereby a **somatic mutation in the normal allele (option D)** of a small number of tubular epithelial cells leads to **unregulated clonal proliferation of cells** that ultimately form the cyst lining.
- **Nephrolithiasis** is seen in **20%** of patients with ADPKD (**option B, false**)

106. Ans. (b) **Associated with tuberculosis** (Ref: R 9th/pg 934)

Xanthogranulomatous pyelonephritis is associated with proteus infections and not TB

107. Ans. (a) **Flea bitten kidney in malignant hypertension**

(Ref: Robbins 9th/pg 939; 8th/pg 950-951)

Small, pinpoint petechial hemorrhages appear on the cortical surface from rupture of arterioles or glomerular capillaries, giving the kidney a peculiar "**flea-bitten**" appearance.

108. Ans. (b) **Small kidney with petechial hemorrhages**

(Ref: Robbins 9th/pg 939; 8th/pg 950-951)

This is a case of a middle aged man who died of malignant hypertension

Kidneys in such patients have a characteristic flea-bitten appearance i.e. **Small kidney with petechial hemorrhages**.

109. Ans. (a) **Medullary**

110. Ans. (b) **Chromophobe**

111. Ans. (a) **Wilms' tumor**

(Ref: Nelson 20th ed/ pg 2466-68)

**Wilms' tumor (WT)**, also known as **nephroblastoma**, is the most common primary malignant renal tumor of childhood; It is the second most common malignant abdominal tumor in childhood. The most common sites of metastases are the lungs, regional lymph nodes, and liver.

112. Ans. (c) **FGFR3** (Ref: Robbins 9th/968)

- **Noninvasive high-grade urothelial carcinoma** is associated with loss of the TP53 and RB tumor suppressor genes while **Noninvasive low-grade papillary urothelial carcinoma** is associated with gain of function FGFR3 and HRAS mutations.

113. Ans (a) **Clear cell ca**

114. Ans. (d) **Collecting duct carcinoma** (Ref: R 9th/pg 953)

115. Ans. (c) **Chromophobe cell cancer** (Ref: R 9th/pg 953)

116. Ans. (b) **Clear cell carcinoma**

(Ref: Robbins 9th/pg 953)

117. Ans. (d) **Autosomal dominant polycystic kidney**

(Ref: Robbins 9th/pg 953; 8th/pg 964)

118. Ans. (c) **VHL** (Ref: Robbins 9th/pg 953; 8th/pg 964)

119. Ans. (a) **3**

(Ref: Robbins 9th/pg 953-955; 8th/pg 964-966)

Clear cell variety of Renal cell carcinoma is related to gene located on chromosome 3





120. Ans. (b) **Hepatitis C virus** (Ref: Harrison 18th/Chapter 282)

**Most Common Opportunistic Infections in Renal Transplant Recipients**

Peri-transplant (<1 month)	Early (1–6 months)	Late (>6 months)
<ul style="list-style-type: none"> <li>Wound infections</li> <li>Herpesvirus</li> <li>Oral candidiasis</li> <li>Urinary tract infection</li> </ul>	<ul style="list-style-type: none"> <li><i>Pneumocystis carinii</i></li> <li>Cytomegalovirus</li> <li><i>Legionella</i></li> <li><i>Listeria</i></li> <li>Hepatitis B</li> <li>Hepatitis C</li> </ul>	<ul style="list-style-type: none"> <li><i>Aspergillus</i></li> <li><i>Nocardia</i></li> <li>BK virus (polyoma)</li> <li>Herpes zoster</li> <li>Hepatitis B</li> <li>Hepatitis C</li> </ul>

121. Ans. (a) **Polyoma virus**

(Ref: Robbins 9th/pg 953; 8th/pg 964)

122. Ans. (a, d); **a. Oncocytoma; d. Chromophobe RCC**

(Ref: Rosai 10th/pg 1183-1193)

**Histological distinction of Chromophobe RCC from Oncocytoma can be difficult;**

123. Ans. (c) **Renal cell Carcinoma** (Ref: R 9th/pg 953-955)

124. Ans. (d) **Monosomy of I and Y** (Ref: R 9th/pg 953-955)

125. Ans. (a) **Polyoma virus** (Ref: Harrison 18th/Chapter 282)

**An emerging organism responsible for causing Pyelonephritis in Renal Allografts is Polyoma virus**

126. Ans. (c) **Wire loop lesions** (Ref: Robbins 9th/pg 222-224)

- Wire loop lesions are seen in Lupus nephritis
- Features of kidney involvement in Multiple myeloma are: Tubular casts, Amyloidosis & Renal tubular necrosis

**Kidney involvement in Multiple myeloma:**

- Amyloidosis**, monoclonal light chain
- Nodular glomerular lesions** due to deposition of *non-fibrillar* light chains (PAS +ve).

127. Ans. (b) **Urothelial cell carcinoma**

(Ref: Robbins 9th/pg 964; 8th/pg 974)

128. Ans. (b) **Malakoplakia**

(Ref: Robbins 9th/pg 963; 8th/pg 975)

- Michaelis-Gutmann bodies**<sup>Q</sup> are Laminated mineralized concretions seen in malakoplakia

129. Ans. (c) **Transitional cells** (Ref: Robbins 9th/pg 964-966)

130. Ans. (b) **Isopropyl alcohol** (Ref: Robbins 9th/pg 964-966)

# Male and Female Genitourinary Tract

## Key Points

- » Most common testicular neoplasms in men over the age of 60 years-testicular lymphoma
- » Most common malignant paratesticular tumors are rhabdomyosarcomas in children
- » Yolk sac tumor most common testicular tumor in infants and children up to 3 years of age
- » HPVs are involved in the pathogenesis of cervical, vaginal, and vulvar cancers.
- » Most common tumor to develop in cryptorchid testis is seminoma
- » Primary adenocarcinoma of the fallopian tubes is rare and is usually serous papillary type
- » Superficial inguinal pouch is the most common site of ectopic testis

## Key Recent Updates

- » ITGCN is now renamed as GCNIS i.e. germ cell neoplasm in situ
- » SOX, 2 is associated with embryonal carcinoma.



## MALE GENITAL TRACT

### PENIS


#### Inflammation

- **Balanoposthitis:**<sup>Q</sup> Nonspecific infection of glans & prepuce, mostly caused by **Candida, anaerobes & Gardenella.**<sup>Q</sup>
- Persistence of such infections leads to inflammatory scarring is a **common cause of phimosis.**<sup>Q</sup>

#### Tumors of Penis

##### Benign Tumors

Peyronie Disease	Condyloma Acuminatum
<ul style="list-style-type: none"> <li>• Characterized by fibrous bands involving corpus cavernosum</li> <li>• Results in penile curvature and pain during intercourse.<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Caused by HPV types 6 &amp; 11<sup>Q</sup></li> <li>• Koilocytosis<sup>Q</sup> is characteristic of HPV infection</li> <li>• Recurs but rarely progresses to invasive cancer</li> </ul>



Koilocytes (Perinuclear Halo & nuclear enlargement)

Image shown koilocytosis S/o HPV lesion

#### Carcinoma in situ: Precancerous Lesions

- Cytological changes of malignancy **confined to epithelium**
- 2 distinct lesions of CIS: **Bowen disease<sup>Q</sup> & bowenoid papulosis**

	Bowen's disease	Bowenoid papulosis
Age	Older than age 35 years <sup>Q</sup>	Younger age <sup>Q</sup>
No. of lesions	Solitary lesion <sup>Q</sup>	Multiple lesions <sup>Q</sup>
Prognosis	Transforms into infiltrating squamous cell carcinoma <sup>Q</sup> in 10%	Never <sup>Q</sup> develops into invasive carcinoma & in many cases regresses spontaneously.

#### Malignant Tumors: Squamous Cell Carcinoma

- **Risk factors:** HPV16 (Most common), HPV 18, Cigarette smoking
- Risk is **reduced by circumcision**; therefore it is *rare in Jews and Muslims*.

- Occurs on the glans or shaft of the penis as an **ulcerated infiltrative lesion**<sup>Q</sup>
- Spread to **inguinal nodes**<sup>Q</sup> and infrequently to distant sites.<sup>Q</sup>

### POINTS TO REMEMBER

- Most frequent penile neoplasm is condyloma acuminatum > carcinoma
- Both bowen disease and bowenoid papulosis are associated with HPV
- Verrucous carcinoma (also known as Giant condyloma or Buschke-Lowenstein tumor) refers to exophytic well-differentiated variant of squamous cell carcinoma, which is locally invasive, but rarely metastasize.

## TESTIS AND EPIDIDYMIS

### Cryptorchidism

- **Complete or partial failure of the intra-abdominal testes to descend into the scrotal sac**<sup>Q</sup>
- Associated with **testicular dysfunction and an increased risk of testicular cancer.**<sup>Q</sup>
- Grossly, testis is **small, brown and atrophic.**<sup>Q</sup>
- Microscopically: Atrophic tubules with thickened basement membrane, leydig cells are spared and appear predominant
- Higher **risk for testicular cancer:** from foci of **intratubular germ cell neoplasia within atrophic tubules.**
- Orchiopexy **reduces the risk of sterility and cancer.**<sup>Q</sup>




### High Yield Facts

- Testicular atrophy occasionally occurs as a primary failure of **genetic origin**, such as in **Klinefelter's syndrome**<sup>Q</sup>
- **Ectopic testes** are the deviation of **testes** from normal path of descent.
- Ectopic testis is **fully developed** with **normal spermatogenesis**<sup>Q</sup> whereas undescended testis lacks spermatogenesis
- **Superficial inguinal pouch is the most common site of ectopic testis**<sup>Q</sup>
- **Most common tumor** to develop in cryptorchid testis is **seminoma**<sup>Q</sup>
- **Most common cause of non-specific inflammation of testis in:**
  - A sexually active young patient (< 35 years) - **Chlamydia trachomatis** and **Neisseria gonorrhea**<sup>Q</sup>
  - Men **older than 35 years** - **E. coli and pseudomonas.**<sup>Q</sup>


### Inflammation

- More common in the **epididymis** than in the testis.
- **Gonorrhea & tuberculosis almost invariably arise in epididymis, whereas syphilis affects the testis**<sup>Q</sup> first.





## Testicular Tumors



### Mnemonic

#### Nonseminomatous Germ Cell Tumors

C – Choriocarcinoma  
E – Embryonal Carcinoma  
T – Teratoma  
Y – Yolk Sac Tumors



### Testicular Germ Cell Tumors

Most originate from a precursor lesion called **intratubular germ cell neoplasia (ITGCN)** except **spermatocyte seminoma; teratoma and Yolk Sac tumor in child.**

#### Predisposing Factors for Germ Cell Tumors are

- **Environmental Factors:** Increased risk by in utero exposures to **pesticides and nonsteroidal estrogens.**
- **Testicular dysgenesis syndrome (TDS):** cryptorchidism, hypospadias, and poor sperm quality.
- **Cryptorchidism**
- **Klinefelter syndrome:** increased risk (50 times) of **mediastinal (not testicular) germ cell tumors**
- **Genetic Factors:**
  - **Strong familial predisposition** associated with the development of testicular germ cell tumors.
  - Isochromosome of short arm of chromosome **12, i(12p)**
  - Familial germ cell tumor risk: ligand for the receptor tyrosine kinase **KIT and BAK**

- **Septa show lymphocytic infiltration (T lymphocytes)**
- Classic **seminoma cell** - large and round to polyhedral and has a distinct cell membrane;
- **Clear or watery-appearing cytoplasm;** Cytoplasm contains glycogen.



### POINTS TO REMEMBER

#### Special Features of Testicular Germ Cell Tumors

- **Lymphatic spread** is common to all forms of testicular tumors
- Retroperitoneal para-aortic nodes are **the first to be involved.**
- Hematogenous spread Lung > Liver > Brain > Bones
- **Extragenital site of germ cell tumors** include **mediastinum, retroperitoneum and pineal gland**
- Biopsy is associated with risk of tumor spillage, so contra-indicated
- The histology of metastases may differ from that of the testicular lesion.

### Seminomatous Germ Cell Tumors

#### Seminoma

**Clinical features:** Peak incidence in **third decade**, almost never occur in infants.

#### Morphology:

- Sheets of **Monomorphic cells**
- **Sheets divided into poorly demarcated lobules by septa of fibrous tissue**

**Tumor markers:** **PLAP** (Placental alkaline phosphatase), **GGT** (Gamma glutamyl transpeptidase), **hCG** (15% cases)

**Genetics:** Seminomas contain isochromosome **12p** and express **OCT3/4 and NANOG.**

#### Spermatocytic Tumor

- It is **distinctive tumor both clinically and radiologically** as compared to seminoma





- Do not arise from an intratubular germ cell neoplasia<sup>Q</sup>
- Age of involvement >65 years<sup>Q</sup>
- Slow growing tumor that rarely produces metastases<sup>Q</sup>
- There is no ovarian counterpart<sup>Q</sup>
- Histologically, it is characterized by three cell populations<sup>Q</sup>- small cells, medium cells, giant cells (Most common).
- Prognosis is excellent<sup>Q</sup>



### High Yield Facts

- Germ cell tumors constitute the most common testicular tumor of men<sup>Q</sup>
- Seminoma: most common type germ cell tumors (50%).<sup>Q</sup>
- Dysgerminoma –Female counter part of seminoma
- Spireme chromatin is seen in Spermatocytic seminoma<sup>Q</sup>
- Sheets of monomorphic cells, lymphocytic infiltration (T lymphocytes) is characteristic feature of seminoma

### Nonseminomatous Germ Cell Tumors

#### Choriocarcinomas


- Most aggressive testicular tumor;<sup>Q</sup>.
- Often cause no testicular enlargement, but only a small palpable nodule<sup>Q</sup>
- A mixture of malignant cytotrophoblasts and syncytiotrophoblasts seen
- Tumor marker is hCG

#### Embryonal cell carcinomas


- Present as sheets of undifferentiated cells; focal glandular differentiation may be present
- Elevated AFP and hCG is seen in this tumor<sup>Q</sup>

#### Teratomas


- Occur at any age from infancy to adult life.
- Second most common testicular tumor in infants & children<sup>Q</sup>
- Derived from all three germ layers<sup>Q</sup>
- In children, differentiated mature teratomas are considered benign<sup>Q</sup>
- In post-pubertal males, all teratomas are regarded as malignant.<sup>Q</sup>




Choriocarcinoma: syncytiotrophoblasts (s) are multinucleated and have a dark staining cytoplasm. The cytotrophoblasts (c) are mononuclear and have a pale staining cytoplasm



Embryonal cell ca: Characterized by large highly pleomorphic tumor cells



Teratoma: with >1 germ layers



Mic: Shows Schiller-Duval body S/o YST

#### Yolk sac tumor (YST) or endodermal sinus tumor


- Most common testicular tumor in infants and children up to 3 years of age.<sup>Q</sup>
- Schiller-Duval bodies or glomeruloid structures (resembling endodermal sinuses).<sup>Q</sup>
- These have excellent prognosis.<sup>Q</sup>
- Tumor marker-AFP<sup>Q</sup>



## High Yield Facts

- **YST**- Most common testicular tumor in infants and children up to 3 years of age.
- **Mixed Tumors**: About 60% of testicular tumors are composed of more than one of the “pure” patterns
- **Common mixtures include**: Teratoma, embryonal carcinoma, and yolk sac tumor
- **Teratocarcinoma**- embryonal carcinoma with teratoma.
- **Teratoma with malignant transformation**- malignant non-germ cell tumors arising in teratomas, like Squamous/cell carcinoma
- Secondary tumors are **chemoresistant**<sup>Q</sup>

## Tumors of Sex Cord-Gonadal Stroma

Sertoli cell tumors: Androblastoma <sup>Q</sup>	Leydig cell tumors	Leydig cell tumor
<ul style="list-style-type: none"> <li>• <b>Hormonally silent</b><sup>Q</sup></li> <li>• Present as a <b>testicular mass most commonly</b>.<sup>Q</sup></li> <li>• Most are <b>benign</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Elaborate hormones</b>: androgens or androgens and estrogens</li> <li>• Most common presentation is <b>testicular swelling</b><sup>Q</sup>.</li> <li>• Gynecomastia may be the <b>first symptom</b><sup>Q</sup></li> <li>• Cytoplasm: <b>Lipid droplets, vacuoles, or lipofuscin pigment</b></li> <li>• Characteristic: <b>rod-shaped crystalloids of Reinke</b><sup>Q</sup></li> <li>• Most are <b>benign</b><sup>Q</sup></li> </ul>	 <p>Image showing crystals of Reinke</p>

## Gonadoblastoma

- Contains a **mixture of germ cells and gonadal stromal elements**.
- Almost always arise in gonads with **some form of testicular dysgenesis**
- **Hallmark oncogene**: TSPY gene

## Testicular Lymphoma


- **Most common testicular neoplasms in men over the age of 60 years**.<sup>Q</sup>
- **Diffuse large B-cell lymphoma > Burkitt lymphoma > EBV-positive extranodal NK/T cell lymphoma**<sup>Q</sup>
- Prognosis is **extremely poor**; have a higher propensity for **central nervous system involvement**<sup>Q</sup>

## High Yield Facts

- **Varicocele** results from **dilatation of testicular veins in pampiniform plexus**.<sup>Q</sup> (left > right side)
- **Varicocele** is associated with oligospermia (<20 million spermatozoa/mL of semen) and **is cause of infertility**.<sup>Q</sup>
- **Spermatocele** is cystic enlargement of **efferent ducts or rete testis**<sup>Q</sup> with numerous spermatocytes present
- Most common **benign** paratesticular tumor is **adenomatoid tumor**.<sup>Q</sup>
- Most common **malignant** paratesticular tumors are **rhabdomyosarcomas in children**<sup>Q</sup>
- Most common **malignant** paratesticular tumors are **liposarcomas in adults**.<sup>Q</sup>

## PROSTATE

- Prostate is a **combined tubuloalveolar organ**,<sup>Q</sup> weighing about 20 g in a normal adult;
- **Benign glands are two layered (basal cells and columnar cells), while in cancer single layered cells seen**<sup>Q</sup>








## Nodular Hyperplasia—Benign Prostatic Hyperplasia (BPH)

- The most common benign prostatic disease in men older than age 50 years.
- It mostly originates from **transitional zone of prostate (carcinoma mostly arises from peripheral zone)**.<sup>Q</sup>
- Nodular hyperplasia is not considered to be a premalignant lesion.<sup>Q</sup>
- Median lobe hypertrophy—nodular enlargement may project up into floor of urethra as a hemispherical mass.<sup>Q</sup>


- The main androgen in the prostate, is **dihydrotestosterone (DHT)**.
- **5 $\alpha$ -reductase** converts testosterone into the more potent **dihydrotestosterone**
- Both glands and stroma can become hyperplastic. Stromal or epithelial predominant hyperplasia may occur.
- Medium or large glands with 2 benign cell layers (secretory and basal) showing some architectural complexity including papillary infoldings.



Gross showing nodular hyperplasia



Microscopy showing normal gland stroma



Microscopy showing glands (All are lined by two layer of cells) and stroma S/O BPH



### High Yield Facts

- Chronic abacterial prostatitis is the **most common form** of prostatitis. MC association is **Chlamydia** or associated with **Ureaplasma**.
- MC cause of Granulomatous prostatitis is intravesical administration of BCG (used for treatment of superficial bladder carcinoma).


- **Primary grade is assigned to dominant pattern** and secondary grade to sub-dominant pattern.
- Combined Gleason score is derived by addition of these two grades
- **Staging of prostatic cancer is also important in the selection of the appropriate form of therapy**

## Tumors of Prostate


- **Adenocarcinoma of prostate** is most common form of cancer in men.<sup>Q</sup>
- Most of the prostatic carcinomas are **acinar adenocarcinoma**.<sup>Q</sup>
- Characterized by small glands that appear “back to back” without intervening stroma
- Most arise in peripheral zone, classically in a posterior location.<sup>Q</sup>

### Grade and Stage are the Best Prognostic Predictors

- Prostate cancer is **graded** using the **Gleason system**
- **Five grades** on the basis of differentiation.<sup>Q</sup>
- Grade I is well differentiated and Grade 5 shows no glandular differentiation. Grade 2, 3 and 4 are in-between.



Characteristic morphologic features of prostatic adenocarcinoma (10x)



### High Yield Facts

#### Prostatic Cancer

- Local extension **most commonly** involves periprostatic tissue, seminal vesicles, and the base of the urinary bladder
- Fascia of Denonvilliers** prevents the backward extension of the tumor.<sup>Q</sup>
- Metastases spread via **lymphatics** to the obturator nodes<sup>Q</sup> and eventually to the para-aortic nodes.
- Hematogenous spread** occurs chiefly to **bones (osteoblastic secondaries)** most commonly to lumbar spine
- Most common epigenetic alteration** in prostate cancer is hypermethylation of the **glutathione S-transferase (GSTP1) gene**, which down-regulates GSTP1 expression<sup>Q</sup>

#### Genes that increase risk in prostatic carcinoma

- Germline mutations BRCA2: 20-fold increased risk
- Germline mutation in **HOXB13**: Homeobox gene that regulates prostatic development

## FEMALE GENITAL TRACT

### INFECTIONS OF FEMALE GENITAL TRACT

- Neisseria gonorrhoeae* and *Chlamydia* infections, are major causes of female infertility<sup>Q</sup>
- Trichomonas vaginalis*- **strawberry cervix**.<sup>Q</sup>
- Gardnerella vaginalis*-**bacterial vaginosis**<sup>Q</sup>: Causes **green-gray, malodorous (fishy) vaginal discharge**<sup>Q</sup>, Premature Labor

### VULVA

#### Non-neoplastic Epithelial Disorders: Leukoplakia


**Definition:** White plaques on the vulva are **clinically** referred to as leukoplakia. It can be caused by:

- Inflammatory dermatoses (e.g., psoriasis, chronic dermatitis)
- Lichen sclerosus and squamous cell hyperplasia
- Neoplasias, such as vulvar intraepithelial neoplasia (VIN), Paget disease, and invasive carcinoma

#### Cysts of Vulva

##### Benign Exophytic Lesions


- Fibroepithelial polyps**, or skin tags<sup>Q</sup>
- Squamous papillomas**<sup>Q</sup>



### Squamous Neoplastic Lesions

#### Vulvar Squamous Cell Carcinomas are Divided into Two Groups

	Basaloid & Warty Carcinomas	Keratinizing Squamous Cell Carcinomas
<b>Epidemiology</b>	Less common <sup>Q</sup>	More common <sup>Q</sup>
<b>Age affected</b>	Younger ages <sup>Q</sup>	Older women <sup>Q</sup>
<b>HPV association</b>	High risk HPVs (type 16) <sup>Q</sup>	Unrelated to HPV (70% of cases) <sup>Q</sup>
<b>Precursor lesions</b>	Classic VIN (vulvar intraepithelial Neoplasia)	Long-standing lichen Sclerosis/ squamous cell hyperplasia <sup>Q</sup> Differentiated VIN or VIN simplex. <sup>Q</sup>



MIC: Keratinizing squamous cell carcinoma





## Glandular Neoplastic Lesions

### Papillary Hidradenoma

- Tubular ducts lined by a **layer of nonciliated columnar cells**<sup>Q</sup> with a layer of flattened “**myoepithelial cells**” underlying the epithelium

### Extramammary Paget's Disease

- Involved:** Epidermal apocrine gland-bearing areas, esp. **vulva, scrotum & perianal areas**
- Gross:** Pruritic, erythematous, crusted, dry raised lesions
- Microscopy:** Single or small clusters of large cells; Faintly basophilic or **vacuolated cells (Paget's cells)**, Large nuclei and prominent nucleoli, Amorphous, granular cytoplasm
- Stains used:** Stain positively with **PAS or mucicarmine stains**.<sup>Q</sup>
- Immunohistochemistry:** CEA, S100, Melan-A, CAM 5.2, EMA, CK 7, GCDFP-15<sup>Q</sup>
- D/D:** Malignant melanoma (malignant cells **melanin stain** or S100 immunoperoxidase stain)





Image shows Paget's cells


### POINTS TO REMEMBER

- Extramammary Paget's disease** is usually seen in **isolation** & associated with an underlying malignancy in **12%**<sup>Q</sup>
- Mammary Paget's disease** is **almost always** associated with an underlying malignancy of the breast


## VAGINA



Gross: Grape like clusters S/O ERMS



Mic: shows cambium layer S/O ERMS



## High Yield Facts

- **Gartner duct cysts** derived from **Wolffian (mesonephric) duct nests**, are found along **lateral walls of vagina**<sup>Q</sup>
- HSVs cause painful genital ulcerations<sup>Q</sup>
- HPV are involved in the pathogenesis of cervical, vaginal, and vulvar cancers.<sup>Q</sup>
- Squamous cell carcinoma is the most common histologic type of vulvar cancer<sup>Q</sup>
- Most common malignant tumor to involve the vagina is carcinoma spreading from the cervix<sup>Q</sup>
- **Most common primary malignant tumor of vagina-squamous cell carcinoma**
- **Most common vaginal malignancy in infants: embryonal rhabdomyosarcoma (sarcoma botryoides).**<sup>Q</sup>
- Lesions in the lower **2/3<sup>rd</sup> of the vagina metastasize** to the **inguinal nodes**<sup>Q</sup>
- Lesions in the **upper vagina** tend to spread to **regional iliac nodes**<sup>Q</sup>
- **Vaginal adenosis** is presence of metaplastic cervical or endometrial epithelium within the vaginal wall, derived from **persistent Müllerian epithelium islets in postembryonic life.**<sup>Q</sup>
- **Benign tumors** of the vagina include **stromal tumors (stromal polyps), leiomyomas, and hemangiomas**

## CERVIX

### Intraepithelial and Invasive Squamous Neoplasia

- CIN can be divided into three grades; CIN I, CIN II and CIN III

Cervical Dysplasia	CIN	Besthesda system	HPV type	Morphology
Mild	CIN I	Low grade SIL (L-SIL)	6,11	Koilocytic atypia, flat condyloma
Moderate, severe, carcinoma in situ	CIN II & CIN III	High grade SIL (H-SIL)	16,18	Progressive cellular atypia, loss of maturation

- Most common cancer subtype in cervix is squamous cell carcinoma (80% cases) > Adenocarcinoma (15%)

### POINTS TO REMEMBER

#### Natural history of squamous intraepithelial lesions with approximate 2-year follow-up

Lesion	Regress	Persist	Progress
LSIL	60 <sup>Q</sup>	30	10 % to HSIL
HSIL	30	60 <sup>Q</sup>	10% TO CARCINOMA



## High Yield Facts

- **Koilocytic atypia** -The nuclear changes of LSIL, often accompanied by **cytoplasmic “halos.”** These “halos” consist of perinuclear vacuoles, a cytopathic change created by **HPV-encoded protein called E5<sup>Q</sup>** that localizes to the membranes of the **endoplasmic reticulum.**<sup>Q</sup>
- Diagnosis of SIL is based on identification of **nuclear atypia.**
- **HPVs infect immature basal cells of the squamous epithelium** in areas of epithelial breaks, or **immature metaplastic squamous cells** present at the squamocolumnar junction<sup>Q</sup>
- **The ability of HPV** to act as a carcinogen depends on the viral proteins **E6 and E7**
- **High risk HPV types for cervical carcinoma:** 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68 and others
- **Low risk HPV types for cervical carcinoma:** 6, 11, 42, 44 (associated with condyloma)
- **Nabothian cysts:** Obstruction of *mucous gland ducts in endocervix* results in small mucous (Nabothian) cysts.

## UTERUS

- Endometrium and myometrium are relatively resistant to infections.

### Endometrial Hyperplasia

- Increase in the number of glands relative to the stroma.
- It is divided into non-atypical and atypical hyperplasia based on nuclear atypia.<sup>Q</sup>
- Atypical hyperplasia is associated with an increased risk of endometrial carcinoma.<sup>Q</sup>

### Malignant Tumors of Endometrium & Myometrium

#### Tumors of Endometrium

- Endometrial carcinoma is the most common invasive cancer of the female genital tract.
- Most of the endometrial carcinomas are **adenocarcinomas.**
- If there are areas of **benign** squamous differentiation within Endometrial Ca, they are called **adenoacanthomas.**

- If there are areas of **malignant** squamous differentiation, they are called **adenosquamous Endometrial Ca**
- **Endometrial carcinoma are divided into type 1 and type 2 tumors** <sup>Q<sup>Q</sup></sup>
- All tumors in type II endometrial tumors category are classified as **grade 3 irrespective of histologic pattern**

	Type 1	Type 2
<b>Morphology</b>	Endometrioid	Serous, Mixed Mullerian Tumor, Clear Cell
<b>Genes</b>	PTEN & (Most common), KRAS, FGF2	P 53, Aneuploidy
<b>Prognosis</b>	Indolent	Aggressive

#### Tumors of the Endometrium with Stromal Differentiation

##### Malignant Mixed Müllerian Tumors (Carcinosarcomas)

- Endometrial adenocarcinomas with a malignant mesenchymal component






- Stroma differentiates into malignant mesodermal components, including muscle, cartilage
- Resemble endometrial Ca genetically & have **poor outcomes** with current therapies.

## Adenosarcomas

Malignant appearing stroma, with **benign but abnormally shaped endometrial glands**.

## Stromal Tumors

May be benign stromal nodules or endometrial stromal sarcomas.



Multiple uterine leiomyomas; submucous, subserous, intramural


**Tumors of Myometrium**

**Fibroids (Leiomyoma)**

- Benign smooth-muscle tumors on cut section—sharply circumscribed and whorled.
- Most common tumor in women**
- Associated with MED12 mutations.**


**Leiomyosarcoma**

- A rare, malignant smooth muscle tumor of the uterus. **Mitoses are the most important criteria** in assessing malignancy in smooth-muscle tumors of the uterus.



Mic: Leiomyosarcoma

Highly pleomorphic. Hyperchromatic cells



Leiomyoma: interlacing bundles of smooth muscle

## FALLOPIAN TUBES

- Most common disorders** infections followed by ectopic (tubal) pregnancy and endometriosis.
- Suppurative salpingitis** *Gonococcus*—Most common (>most common primary lesion spartatubal cysts<sup>Q</sup>)
- Hydatids of Morgagni paratubal cysts**<sup>QQ</sup>
- Adenomatoid tumors**<sup>Q</sup>
- Primary adenocarcinoma of the fallopian tubes is rare and is usually **serous papillary type**<sup>Q</sup>

## OVARIES


### Cysts

- Follicular cysts:** These are benign cysts of the ovary.
- Chocolate cysts:** These refer to cystic areas of endometriosis that include hemorrhages and blood clots.
- Polycystic ovarian disease (PCOD)

### Polycystic Ovarian Disease (Stein-Leventhal Syndrome)

Characterized by	Endocrine abnormalities	Abnormalities in ovary
<ul style="list-style-type: none"> <li>Hyperandrogenism</li> <li>Menstrual abnormalities</li> <li>Polycystic ovaries</li> <li>Chronic anovulation</li> <li>Decreased fertility<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li><b>Excess androgens</b> (androstenedione)Q</li> <li><b>Increased estrogen levels</b><sup>Q</sup></li> <li>Increased LH &amp; decreased FSH levels</li> <li><b>High LH/FSH ratio</b></li> <li>Increased GnRH levels</li> </ul>	<ul style="list-style-type: none"> <li>Enlarged with thick capsules</li> <li>Hyperplastic ovarian stroma</li> <li>Numerous follicular cysts lined by a hyperplastic theca interna.</li> </ul>

## Ovarian Tumors



### Surface Epithelial Tumors

- Derived from the surface coelomic epithelium, which embryonically gives rise to the Mullerian epithelium.
- Epithelial ovarian tumors are classified into benign, borderline or malignant
- Benign tumors are composed of **well-differentiated epithelial cells with minimal proliferation**.<sup>Q</sup>
- Borderline tumors show increased cell proliferation, but **lack stromal invasion**.<sup>Q</sup>
- Malignant tumors show increased epithelial atypia and are defined by the **presence of stromal invasion**.<sup>Q</sup>
- About 80% of all ovarian epithelial tumors are benign and occur in young women<sup>Q</sup>


### Serous Ovarian Tumors

- Associated Mutation BRCA1 and BRCA2, p53—High grade, KRAS, BRAF—low grade**
  - Most common malignant ovarian tumors; account for approximately 40% of all cancers of ovary



- Composed of ciliated columnar serous epithelial cells, similar to **the lining cells of the fallopian tubes**.<sup>Q</sup>

- They commonly involve the **surface of ovary, bilaterally**
- **Concentric calcifications (psammoma bodies)** seen





## Mucinous Ovarian Tumors

- **KRAS Consistent** → mutation associated
  - Account for 20% to 25% of all ovarian neoplasms; **Most common- gastric or intestinal type differentiation**.
  - Grossly unilateral, **more cysts and no surface involvement**.
  - Tend to **produce larger cystic masses**
  - **Mutation of the KRAS proto-oncogene is a consistently seen in mucinous tumors of ovary** <sup>RO</sup>

## Brenner Tumor (Transitional Cell)

- 10% of ovarian tumors
- Unilateral
- Benign
- Contain Neoplastic epithelial cells resembling urothelium








## Endometrioid Ovarian Tumors

- **20% endometriosis**
- 10% to 15% of all ovarian cancers;<sup>Q</sup> 15% to 20% of cases coexist with endometriosis<sup>Q</sup>
- Distinguished by the presence of tubular glands resembling benign or malignant endometrium.<sup>Q</sup>
- Molecular studies-striking similarities to endometrial endometrioid carcinoma; PI3K/AKT pathway signaling and mutations in mismatch **DNA repair genes** and **CTNNB1** ( $\beta$ -catenin).<sup>Q</sup>


## Germ Cell Tumors



## Teratomas

Divided into three categories: mature (benign), immature (malignant), monodermal or highly specialized

Mature (Benign) Teratomas	Immature Malignant Teratomas
<ul style="list-style-type: none"> <li>• Most benign teratomas are <b>cystic (dermoid cysts)</b></li> <li>• Arise from ectodermal differentiation of totipotential cells.<sup>Q</sup></li> <li>• Arise from an ovum <b>after the first meiotic division</b><sup>Q</sup></li> </ul> <p><b>Morphology:</b></p> <ul style="list-style-type: none"> <li>• Unilocular cysts<sup>Q</sup> containing hair &amp; cheesy sebaceous material with <b>tooth structures &amp; areas of calcification</b>.<sup>Q</sup></li> <li>• <b>Rokitansky protuberance</b><sup>Q</sup> -The inner lining of cyst contains white shiny masses projecting from the wall toward the center of the cysts. Hair, other dermal appendages, bone and teeth are present, they usually arise from this protuberance</li> </ul>	<ul style="list-style-type: none"> <li>• Resembles <b>fetus or embryo tissues</b> rather than adult.</li> <li>• Seen chiefly in <b>prepubertal adolescents &amp; young women</b>, the mean age being 18 years.</li> <li>• Microscopy: Immature neuroepithelium &amp; cartilage;</li> <li>• Risk of extraovarian spread is dependent on <b>histologic grade of tumor</b></li> <li>• Most recurrences <b>develop in the first 2 years</b></li> </ul>



## Dysgerminoma


**Characteristics:** Ovarian counterpart of the **seminoma** of the testis; usually **unilateral** (80% to 90%) and **solid**

**Genetics:** Express **OCT-3, OCT4, and NANOG**. Onethird have activating mutations in the **KIT gene**

**Morphology:** Sheets or cords of cells separated by scant fibrous stroma, infiltrated with mature **lymphocytes**<sup>Q</sup> & occasional granulomas

**Hormones elaborated:** Most of these tumors have **no endocrine function (few produced hCG)**


- Tumor marker- **LDH, PLAP**
- All dysgerminomas are **malignant**



Dysgerminoma

## Endodermal Sinus (Yolk Sac) Tumor

- **Second most common malignant tumor** of germ cell origin; usually **unilateral**
- Derived from differentiation of malignant germ cells **toward extraembryonic yolk sac structure**.
- Histology - Glomerulus-like structure composed of a central blood vessel enveloped by germ cells within a space lined by germ cells (**Schiller-Duval body**).




Mic: Shows Schiller-Duval body S/o YST

- Intracellular and extracellular **hyaline droplets** - composed of **α-fetoprotein** and **α1-antitrypsin**.
- Tumor Marker- **AFP**

### Choriocarcinoma

- Most common of **placental origin**, is e.g. of **extra-embryonic differentiation of malignant germ cells**.
- Pure **ovarian** choriocarcinomas are rare – exist in combination with other germ cell tumors
- Elaborate high levels of **chorionic gonadotropins**<sup>Q</sup>
- Most common site of metastasis - **lung**.<sup>Q</sup>




Mic: Admixed syncytiotrophoblasts & cytotrophoblasts S/O Choriocarcinoma

### Sex-Cord Stromal Tumors

#### Granulosa Cell Tumor

- **Most common** malignant tumor in this category;<sup>Q</sup> usually **unilateral** (80% to 90%) and solid cystic

- Hormonally active tumors have a **yellow coloration** due to intracellular lipids.
- Composed of cells that **stain positively with inhibin**.<sup>Q</sup>



Mic: Shows cell exner bodies S/o granulosa cell Tumor

- Cells have coffee bean nuclei with grooves
- **Call Exner bodies**<sup>Q</sup> (small follicle like structure with eosinophilic material) are characteristic features
- **Clinical features depend upon the estrogenic activity of the tumor**<sup>Q</sup>
- Behavior of the **endometrium closely resembles that of metropathia hemorrhagica**<sup>Q</sup>
- Tumor cells may secrete estrogens → **precocious sexual development in girls/ ↑endometrial hyperplasia & Ca**
- Less commonly granulosa cell tumors **can secrete androgens** and produce masculinization.
- **Metastases 1<sup>st</sup> involve opposite ovary**<sup>Q</sup> followed by lumbar region, mesentery, liver and mediastinum.
- Tumor marker-Inhibin
- **FOXL2 gene** mutation (97% cases of Adult granulosa cell tumor)

### High Yield Facts

Findings	Tumors
<b>Call exner bodies</b>	Granulosa cell tumor <sup>Q</sup>
<b>Reinke's crystal</b> <sup>Q</sup>	Hilus cell tumor/Leyding cell tumor
<b>Signet ring cell</b> <sup>Q</sup>	Krukenberg tumor
<b>Hobnail cells</b>	Clear cell carcinoma
<b>Psammoma bodies</b>	Papillary serous cystadenoma of ovary
<b>Schiller duval body</b>	Endodermal sinus (Yolk sac) tumour

### Thecomas

- Are composed of **spindle-shaped cells with vacuolated cytoplasm**.
- Vacuolated because of **steroid hormone (estrogen) production**, which can be stained with an **Oil Red O** stain.





### Sertoli-Leydig Tumors (Androblastomas)

- **Functional**<sup>Q</sup> tumors, which commonly produce **masculinization** due to secretion of androgens
- Peak incidence is in the **second and third decades**<sup>Q</sup> are usually **unilateral**<sup>Q</sup>



- Patients with PCOD have risk of endometrial hyperplasia carcinoma due to *excess estrogen production*
- **Pseudomyxoma peritonei** results from spread of **mucinous tumor in appendix** & is marked by **mucinous ascites, cystic epithelial implants on peritoneal surfaces, adhesions & frequent involvement of ovaries**<sup>Q</sup>
- In contrast to choriocarcinomas arising in placental tissue, those arising in the ovary are generally unresponsive to chemotherapy and are often fatal.<sup>Q</sup>
- **Most common germ cell tumor in females is benign cystic teratoma**<sup>Q</sup>
- **Benign cystic teratoma** can be associated with paraneoplastic syndromes like **inflammatory limbic encephalitis**. <sup>Q</sup>

- Cut surface is usually solid and varies from gray to **golden brown**<sup>Q</sup> in appearance
- **Heterologous elements, such as mucinous glands, bone, and cartilage, may be present in some tumors**<sup>Q</sup>

### High Yield Facts


- About 1% of the dermoids undergo malignant transformation, **most commonly to squamous cell carcinoma**<sup>Q</sup>
- **Most common specialized/ monodermal teratoma** are struma ovarii and carcinoid
- Metastatic intestinal carcinoid **involving the ovaries is always bilateral**.<sup>Q</sup>
- **Stromal carcinoid** is a combination of struma ovarii and carcinoid in the same ovary.
- Gonadoblastoma have coexistent dysgerminoma in 50% cases
- Sertoli leydig cell tumors have mutations in **DICER, gene**
- MC mutation in Granulosa cell tumor is in **FOXL2 gene**

### Metastatic Tumors of Ovary


- **Most common** occur from **tumors of Mullerian origin**: Uterus, fallopian tube, contralateral ovary, or pelvic peritoneum.
- **Most common extramullerian primaries** are **breast & GIT** including colon, stomach, biliary tract, and pancreas.<sup>Q</sup>

### Krukenberg Tumor

- Refers to a **metastatic bilateral ovarian** malignancy whose primary site is **GIT or breast**.<sup>Q</sup>
- Composed of **mucin-producing, signet-ring cancer cells, most often of gastric origin**.<sup>Q</sup>
- Ca colon, appendix, breast (**especially invasive lobular carcinoma**), pancreas and gall bladder are other **primary sites**.



Gross B/L enlarged ovaries



Mic: Shows cells with nucleus pushed to periphery S/O signet ring cell

## IMPORTANT ADDITIONAL INFORMATION TO REMEMBER IN GENITAL SYSTEM

### SEMEN ANALYSIS

In 2010 the World Health Organization (WHO) updated its reference values for the Semen Analysis


Cut-off reference values for semen characteristics as published in consecutive WHO manuals

The most common reasons for laboratory **semen analysis** in humans are

- As part of a couple's infertility investigation

- After a vasectomy to verify that the procedure was successful.
- Testing human donors for sperm donation,

Semen characteristics	WHO 2010
Volume (mL)	≥ 1.5
Sperm count (10 <sup>6</sup> /mL)	≥ 15
Total sperm count (10 <sup>6</sup> )	≥ 39
Total motility (%)	≥ 40
Progressive motility	≥ 32%
Vitality (%)	≥ 58
Morphology (%)	≥ 4
Leukocyte count (10 <sup>4</sup> /mL)	< 1.0



## Prerequisites for Analysis

**Liquefaction:** Process when the gel formed by proteins from the seminal vesicles is broken up and the semen becomes more liquid. In the NICE guidelines, a liquefaction time < 60 minutes -normal ranges. Semen analysis should be done **after liquefaction with through mixing**

## Fructose Level

WHO specifies a normal level of 13  $\mu\text{mol}$  per sample. Absence of fructose may indicate a problem with the seminal vesicles

## Normal Values

### PH

- Acidic ejaculate-one or both of the seminal vesicles are blocked.
- Basic ejaculate-infection

### Motility

- Grade a:** Sperm with progressive motility-swim fast in a straight line.
- Grade b:** (non-linear motility): These also move forward but in a curved motion
- Grade c:** These have **non-progressive motility**-they move their tails only.
- Grade d:** These are immotile

### Morphology

- A *motile sperm organelle morphology examination* (MSOME) is a particular morphologic investigation which increases light microscope's resolution 6000x times to assess sperm morphology

### Vitality

- Number of live sperms is called viable
- A viable sperm will have intact cell membrane and won't take up **eosin Y dye**

## Nomenclature Related to Sperm Quality

- Aspermia:**
  - No semen (no or retrograde ejaculation)
- Asthenozoospermia:**
  - Percentage of progressively motile (PR) spermatozoa **below** the lower reference limit
- Azoospermia:**
  - No spermatozoa** in the ejaculate
- Cryptozoospermia:**
  - Spermatozoa absent from fresh preparation but **observed** in a **centrifuged** pellet
- Necrozoospermia:**
  - Low percentage of live, and high percentage of immotile, spermatozoa in the ejaculate
- Oligozoospermia:**
  - Total **number** (or concentration, depending on outcome reported) of spermatozoa below the lower reference limits

## Teratozoospermia:

- Percentage of **morphologically normal** spermatozoa **below** the lower reference limit.

## PAP SMEAR


### Specimen Adequacy

#### Minimum Squamous Cellularity Criteria

- Conventional smear-8000 to 12000 well-preserved, well-visualised squamous cells.
- Liquid based prep-min 5000.

#### Endocervical Zone Component


- Atleast 10 well-preserved endocervical or squamous metaplastic cells, singly or in clusters.





### Morphology

- Superficial cells display the most maturity, having been affected by estrogen
- Intermediate cells display mild maturation, having been affected by progesterone
- Parabasal cells are the least mature cells having not been affected by estrogen or progesterone
- MATURATION INDEX (MI)** is a ratio obtained through performing a random count of three major cell types (parabasal cells, intermediate cells and superficial cells) that are shed from the squamous epithelium
- The cell count is expressed as a percentage that reads as follows:
- MI = % parabasal cells/% intermediate cells/% superficial cells.







Schematic diagram of cervical epithelial layers




Endocervical cells




## Few Images of Pap Smear




### Bacterial Vaginosis



### Squamous Cell Carcinoma, Pap




### Herpes Simplex




- Nuclear moulding, Multinucleation.
- Ground glass chromatin with prominent nuclear membrane

### Human Papilloma Virus




- Koilocytosis-superficial and intermediate cells
- Koilocyte: Perinuclear halo with raisinoid nuclei

### Trichomonas



- Pear shaped, oval, cyanophilic organisms, 15-30μ
- Pale vesicular eccentrically located nucleus


### Candida



- Double contoured pale pink hyphae and pseudohyphae
- Pseudohyphae appear septate
- Spores are eosinophilic




## Menopause



Parabasal cells

## Actinomyces




Usually seen with intrauterine device

R10<sup>th</sup>


### Latest Update

- ITGCN is now renamed as GCNIS i.e. germ cell neoplasm in situ
- SOX, 2 is associated with embryonal carcinoma




## Image-Based Questions

1. A 32-year-old female complained of menorrhagia. On pap smear, identify the cells marked in circle



- a. Koilocytes  
b. Normal squamous cells  
c. Candida  
d. CMV

2. A 55-year-old female presented with Pruritic, erythematous, crusted, dry raised lesions over vulva. Diagnosis?




- a. Paget's disease  
b. Carcinoma vulva  
c. CIS  
d. Leukoplakia

3. A 25-year-old pregnant female has hcg levels which are markedly high. On USG, size of uterus exceeds the gestational age. Diagnosis:



- a. Choriocarcinoma  
b. Hydatidiform mole  
c. Normal pregnancy  
d. PSTT

4. Identify the molecular subtype of breast cancer:



- a. Luminal A  
b. Luminal B  
c. Her2 Neu positive  
d. Triple negative



## Answers of Image-Based Questions

### 1. Ans. (a) Koilocytes

- A **Koilocyte** is a squamous epithelial cell that has undergone a number of structural changes, which occur as a result of infection of the cell by human papillomavirus
- Koilocytes may have the following cellular changes:
  - Nuclear enlargement (two to three times normal size)
  - Irregularity of the nuclear membrane contour
  - A darker than normal staining pattern in the nucleus, known as Hyperchromasia
  - A clear area around the nucleus, known as a perinuclear halo.

### 2. Ans. (a) Paget's disease

- Here we see Single and small clusters of large cells; **vacuolated cells (Paget's cells)**, Large nuclei and prominent nucleoli, Amorphous, granular cytoplasm

### 3. Ans. (b) Hydatidiform mole

- Complete hydatidiform mole, consisting of numerous swollen (hydropic) villi.

### 4. Ans. (c) Her2neu +

- ER, PR are nuclear stains whereas Her2neu is cytoplasmic and membranous stain. So here we see cytoplasmic and membranous positivity, hence Her2neu + cancer.





## Multiple Choice Questions

### MALE GENITAL SYSTEM

#### PENIS

1. Which of the following does not progress to carcinoma?  
(Recent Question 2016)
  - a. Bowen's disease
  - b. Bowenoid papulosis
  - c. Leukoplakia
  - d. Erythroplakia
2. Corbus disease is:  
(AP 2013)
  - a. Dense fibrosis of dermis and buck fascia of penile corpus
  - b. Balanitis circumscrip taplasma cellularis
  - c. Gangrenous balanitis
  - d. Erythroplasia of queyrat
3. Verrucous carcinoma is-  
(Recent Question 2014, Kerala 2K)
  - a. Extremely well differentiated squamous cell carcinoma
  - b. Poorly differentiated squamous cell Ca
  - c. Example of condyloma
  - d. An example of adenocarcinoma

#### TESTIS

4. Post surgery image of a tumor in scrotum is shown below. What could be your possible diagnosis?  
(Recent exam 2018)



- a. Teratoma
  - b. Seminoma
  - c. Yolk sac tumor
  - d. Lymphoma
5. A surgeon suspecting testicular carcinoma in a patient asks the intern to send the sample for histopathology, what is the fluid in which the intern should send the sample to the pathologist?  
(AIIMS May 16)
  - a. Bouvin solution
  - b. 10% formalin
  - c. 95 % ethanol
  - d. Alcohol
6. All are true about seminomas except:  
(Recent Question 2015)
  - a. Most common type of germ cell tumor
  - b. Anaplastic seminomas is associated with a worse prognosis
  - c. Almost never occur in infants
  - d. Spermatocytic seminoma is slow growing with good prognosis

7. Intratubular germ cell neoplasia is implicated as a cause for the following testicular tumor?  
(Recent Question 2015)
  - a. Pediatric yolk sac tumors
  - b. Pediatric teratomas
  - c. Seminomas
  - d. Adult spermatocytic seminomas
8. All the following are testicular dysgenesis syndromes except:  
(Recent Question 2015)
  - a. Cryptorchidism
  - b. Epispadias
  - c. Poor sperm quality
  - d. Hypospadias
9. Not seen in children  
(Recent Question 2014-15)
  - a. Neuroblastoma
  - b. Retinoblastoma
  - c. Hepatoblastoma
  - d. Seminoma
10. Schiller-Duval bodies is seen in:  
(Recent Question 2014, DNB 11)
  - a. Choriocarcinoma
  - b. Embryonal cell Ca
  - c. Endodermal sinus tumour
  - d. Immature teratoma
11. Alkaline phosphatase is a tumor marker of which tumor-  
(Recent Question 2014)
  - a. Seminoma
  - b. Embryonal carcinoma
  - c. Yolk sac tumor
  - d. Embryonal sinus tumor
12. All are germ cell tumors except- (Recent Question 2014)
  - a. Seminoma
  - b. Leydig cell tumor
  - c. Embryonal carcinoma
  - d. Endodermal sinus tumor
13. Microscopic feature of seminoma include all of the following except (Recent Question 2013, AP PGME 14)
  - a. Gland formation
  - b. Lymphocytic infiltration
  - c. Monomorphic cells
  - d. Destruction of seminiferous tubules
14. True about serum AFP level:  
(PGI May 2013)
  - a. Raised in testicular tumor
  - b. Raised in 70% cases of HCC
  - c. Correlation between tumor recurrence after surgery in HCC
  - d. Correlation with HCC size
  - e. Upper limit of normal in the serum is 200 ng/ml
15. Commonest histological type of carcinoma testis is -  
(DNB 2012)
  - a. Teratoma
  - b. Yolk sac tumour
  - c. Seminoma
  - d. Chorio carcinoma
16. AFP is elevated in:  
(PGI May 2011)
  - a. HCC
  - b. Hepatoblastoma
  - c. Infant hemangioendothelioma
  - d. Amebic liver abscess
  - e. Embryonic sarcoma
17. Tumour marker for Endodermal Sinus Tumour
  - a. PLAP
  - b. hCG
  - c. Alfa feto protein
  - d. Cytokeratin



18. A glomerulus-like structure composed of central blood vessel enveloped by germ cells within a space lined by germ cells, is seen in - (Karnataka 11)
- Sertoli-Leydig cell tumor
  - Granulosa cell tumor
  - Endodermal sinus tumor
  - Sex cord tumor with annular tubules

### PROSTATE

19. NKX3-1 immunohistochemical used for diagnosis of: (JIPMER Nov 2019)
- Colorectal carcinoma
  - Pancreatic carcinoma
  - Prostate
  - Renal cell carcinoma
20. Gleason's grading system is for - (Recent Question 2014-15)
- Carcinoma testis
  - Carcinoma colon
  - Carcinoma thyroid
  - Carcinoma prostate
21. Which of the following is not a variant of Prostate specific antigen (PSA)? (Recent Question 2014)
- PSA density'
  - PSA velocity
  - PSA Nodularity
  - Ratio of free and bound PSA in the serum

### SEMEN ANALYSIS

22. According to the 2010 WHO criteria what are the characteristics of normal semen analysis (AIIMS May 2015)
- Volume 1.5 ml, count 15 million, morphology 4% progressive motility 32%
  - Volume 2.0 ml, count 20 million, morphology 4% progressive motility 32%
  - Volume 1.5 ml, count 20 million, morphology 4% progressive motility 32%
  - Volume 2.0 ml, count 15 million, morphology 40% progressive motility 32%
23. Semen analysis is to be done? (Recent Question 2014)
- As early as possible in semisolid state
  - After 15-30 minutes irrespective of liquefaction
  - After 30-60 minutes irrespective of liquefaction
  - After liquefaction with thorough mixing
24. Teratozoospermia refers to? (Recent Question 2013)
- Absence of semen
  - Absence of sperm
  - All dead sperms in ejaculate
  - Morphologically defective sperms

### FEMALE GENITAL SYSTEM

#### VULVA


25. Extramammary paget's is seen in? (Recent exam 2018)
- Uterus
  - Vulva
  - Vagina
  - Ovary
26. Predisposing factor for Carcinoma Vulva is all except: (Recent Question 2015)
- Smoking
  - Human papilloma virus (HPV) infection
  - Fibroepithelial polyps
  - Leukoplakia

### VAGINA


27. A grape like, polypoid, bulky mass protruding through vagina in a 4 yr old girl is characteristic of (MHPG 2014)
- Fibrosarcoma
  - Sarcoma botryoides
  - Leiomyosarcoma
  - Inflammatory polyp
28. Sarcoma botryoides is a type of? (Recent Question 2015)
- Rhabdomyosarcoma
  - Lymphangioma
  - Leiomyoma
  - Rhabdomyoma

### CERVIX

29. Identify the PAP Smear given below? (AIIMS May 16)



- Trichomonas
  - Chlamydia
  - Actinomycetes
  - Herpes simplex type 2
30. Findings of Cervical PAP smear taken during the late menstrual peri- od from a 45 year old lady suffering with a ovarian tumor is shown below. Which ovarian tumor is she most likely suffering from? (AIIMS May 16)



- Dysgerminoma
  - Granulosa cell tumor
  - Mucinous cyst adeno Ca
  - Serous cyst adeo ca
31. Which of the following marker favours diagnosis of preinvasive & invasive cervical cancer: (PGI May 16)
- Ki67
  - Oncoprotein E6
  - p16INK4, cyclin E, and Ki-67
  - Oncoprotein E8

### CERVIX

32. Most common cause of Cervical neoplasia is? (Recent Question 2015)
- HPV-6
  - HPV-11
  - HPV-16
  - HHV




- 33. Risk factor of CA cervix** (Recent Question 2014-15)
- Smoking
  - Sex at 25 yrs
  - Decreased parity
  - Single sexual partner

- 34. 100/0/0 maturation index denotes** (Recent Question 2015)
- Atrophic smear
  - Pregnancy
  - Reproductive age female
  - None

### UTERUS


- 35. Hysterectomy from a 35-year-old female showed the following gross and histological features. What could be your possible diagnosis?** (Recent exam 2018)




- Carcinoma endometrium
  - Leiomyoma
  - Malignant mixed Müllerian tumor
  - Leiomyosarcoma
- 36. Endometriotic lesion histology represents it's?** (AIIMS May 2017)
- High estrogen
  - High progesterone
  - High cholesterol
  - High levels of prolactin
- 37. Gene most commonly involved in Endometrial carcinoma** (JIPMER 2016)
- PTEN
  - Braf mutation
  - RAS
  - REL
- 38. Swiss cheese pattern endometrium is seen in** (Recent Question 2015)
- Carcinoma endometrium
  - Metropathia hemorrhagica
  - Hydatidiform mole
  - Halban's disease
- 39. Endometrial Carcinoma risk in?** (Recent Question 2014)
- Sertoli Leydig cell
  - Immature teratoma
  - Gonadoblastoma
  - Granulosa theca cell tumor
- 40. Complete mole can be differentiated from partial mole by:** (Recent Question 2014, PGI May 12)
- P57
  - P53
  - P16inkga
  - P63
  - PED
- 41. True about Complete (classic) moles A/E** (Recent Question 2014)
- 90% have a 46, XX karyotype
  - Androgenesis
  - Duplication of the genetic material of one sperm
  - Arise from the fertilization of a single egg by two sperm

### OVARY


- 42. A 35-year-old female with adnexal mass, CA 125 normal, LDH raised and CA 19.9 was normal. Gross and histopathology has been given below. What is your diagnosis?** (AIIMS Nov 16)



- Choriocarcinoma
  - Dysgerminoma
  - Teratoma
  - Papillary serous cystadenoma
- 43. A 25 female patient presented with complains of feeling of mass per abdomen. On examination, a mass was found on left adnexal lesion with multiple solid cystic areas. The patient was operated and the mass was removed, which has been shown below. What is the true statement regarding the condition?** (AIIMS Nov 2016)



- Specimen shows multiple solid cystic areas suggestive of dermoid cyst.
  - Specimen shows cystic areas suggestive of serous cyst adenoma
  - Specimen shows Choriocarcinoma
  - Specimen shows Dysgerminoma
- 44. Call exner bodies are seen in:** (Recent Question 2016-17)
- Dysgerminoma
  - Granulosa cell tumor
  - Thecoma
  - Arrhenoblastoma
- 45. BA lady with abdominal mass was investigated. On surgery, she was found to have b/l ovarian masses with smooth surface. On microscopy they revealed mucin secreting cells with signet ring shapes. What is your diagnosis?** (AIIMS May 2015)
- Dysgerminoma
  - Krukenberg tumour
  - Primary Adenocarcinoma of the ovaries
  - Dermoid cyst



- 46. One of the following is a germ cell tumor of ovary:**  
(MH PGMEET 2016, AP 2015)
- Granulosa cell tumor
  - Mucinous cystadenoma
  - Brenner tumor
  - Benign cystic teratoma
- 47. Fibroma belongs to** (Recent Question 2014-15)
- Germ cell tumor
  - Sex cord stromal tumor
  - Surface epithelial stromal tumors
  - Metastatic tumors from non ovarian primary
- 48. Which hormone is increased in PCOS?** (Recent Question 2015)
- LH
  - FSH
  - Inhibin
  - Estrogen
- 49. Which of these tumors is unique to pregnancy?** (Recent Question 2015)
- Luteoma
  - Serous cystadenoma
  - Mucinous cystadenoma
  - Teratoma
- 50. Most common ovarian tumor** (Recent Question 2013)
- Serous cystadenoma
  - Choriocarcinoma
  - Teratoma
  - Fibroma
- 51. Rokitanski protruberences are seen in -** (Recent Question 2013)
- Mucinous carcinoma
  - Teratoma
  - Epidermal cystoids adenoma
  - Papillary carcinoma
- 52. Marker for ovarian carcinoma in serum is -** (Recent Question 2013)
- CA-125
  - Fibronectin
  - Acid Phosphatase
  - PSA
- 53. True about CA-125:** (PGI May 2013)
- Glycoprotein
  - It is a specific marker
  - Increased in colon carcinoma
  - Normal range in pre menopausal females is 200 U/ml
  - May be elevated in Pelvic inflammatory disease
- 54. CA 125 is used for?** (DNB Aug 12)
- Follow up of ovarian cancer
  - Diagnosis of pancreatic cancer
  - Diagnosis of stomach cancer
  - Diagnosis of ovarian cancer
- 55. A 20-year-old female is diagnosed with granulosa cell tumor of the ovary. Which of the following biomarkers would be most useful for follow-up of patient?** (AIIMS 10, 11)
- CA 19-9
  - CA 50
  - Inhibin
  - Neuron-specific-enolase
- 56. Which one of the following is not true regarding chorio-carcinoma?** (DNB 11)
- Aggressive malignancy
  - Raised HCG levels
  - Common below 20 years of age
  - Gonadal type is chemosensitive





## Answers with Explanations

### 1. Ans. (b) **Bowenoid papulosis** (Ref: Robbins 9th/pg 970-71)

Bowen disease	Bowenoid papulosis
Transforms into <b>infiltrating squamous cell carcinoma</b> <sup>Q</sup> in 10%	<b>Never<sup>Q</sup></b> develops into <b>invasive carcinoma</b> & in many cases regresses spontaneously.

### 2. Ans. (c) **Gangrenous balanitis**

(Ref: [www.pathologyoutlines.com/topic/penscrotumgangrenousbalanitis.html](http://www.pathologyoutlines.com/topic/penscrotumgangrenousbalanitis.html))

- Corbus disease is rapidly progressing necrotizing inflammatory disease due to anaerobes in glans penis (Gangrenous balanitis)

### 3. Ans. (a) **Extremely well differentiated squamous cell carcinoma** (Ref: Robbins 9th/pg 971; 8th/pg 984)

**Verrucous carcinoma** [also known as **Giant condyloma** or Buschke-Lowenstein tumor]

- Exophytic well-differentiated variant of squamous cell carcinoma<sup>Q</sup>**
- Locally invasive, but rarely metastasize.<sup>Q</sup>**
- It invades the underlying tissue along a broad fronts**

### 4. Ans. (b) **Seminoma**

(Ref: Robbins 9th ed p 976)

Seminomas produce bulky masses, sometimes ten times the size of the normal testis. The typical seminoma has a homogeneous, gray-white, lobulated cut surface, usually devoid of hemorrhage or necrosis. Generally the tunica albuginea is not penetrated, but occasionally extension to the epididymis, spermatic cord, or scrotal sac occurs. Uniform cells divided into poorly demarcated lobules by delicate fibrous septa containing a lymphocytic infiltrate. The classic seminoma cell is large and round to polyhedral and has a distinct cell membrane; clear or watery-appearing cytoplasm; and a large, central nucleus with one or two prominent nucleoli.

### 5. Ans. (b) **10% formalin**

(Ref: Complete review of pathology 2<sup>nd</sup> ed / Annexure 4. <http://www.adasp.org/Surveys/Question-about-Bouins-fixative-for-testis-biopsy.html>)

**This is a tricky Question !**

When you first see the term analysis for sperm, the answer is bouin's fluid. But what the examiner wants to know is whether you know that for histological diagnosis, the fixative is 10% formalin

**Remember:** Bouins fluid should be used for testicular analysis in cases of infertility or even CIS( carcinoma in situ) as Formalin induces marked shrinkage → difficulty in germ cell and CIS recognition.

But once you have known case of testicular carcinoma, then you need to make correct diagnosis on the basis of morphology, also preserve antigens for IHC and do molecular tests, which comes best when you fix the tissue in 10% buffered neutral formalin

### 6. Ans. (b) **Anaplastic seminomas is associated with a worse prognosis**

(Ref: Genitourinary Pathology: A Volume in the Series: pg 610)

The term anaplastic seminoma is used when seminoma shows pleomorphism, robust mitotic activity and scant lymphocytic infiltrate. Anaplastic semioma has **same** prognosis as classical seminoma.

### 7. Ans. (c) **Seminomas** (Ref: Robbins 9th/pg 977; 8th/pg 989)

Testicular germ cell tumour originate from a precursor lesion called **intratubular germ cell neoplasia (ITGCN)** except **spermatocyte seminoma**; pediatric **teratoma** and **Yolk Sac tumor**

### 8. Ans. (b) **Epispadias** (Ref: Robbins 9th/pg 975; 8th/pg 988)

**Testicular dysgenesis syndrome (TDS):** cryptorchidism, hypospadias, and poor sperm quality

### 9. Ans. (d) **Seminoma** (Ref: Robbins 9th/pg 977; 8th/pg 989)

### 10. Ans. (c) **Endodermal sinus tumour** (Ref: R 9th/pg 977)

Yolk sac tumor or endodermal sinus tumor is the most common **testicular tumor in infants and children <3 yrs of age**. It has good prognosis

### 11. Ans. (a) **Seminoma** (Ref: R 9th/pg 975-76; 8th/pg 988-89)

#### **Seminoma**

**Most common type** of germ cell tumors (50%).<sup>Q</sup>

**Tumor markers:**

**PLAP** (Placental alkaline phosphatase).<sup>Q</sup> **GGT** (Gamma glutamyl transpeptidase)<sup>Q</sup>, **hCG** (15%)

### 12. Ans. (b) **Leydig cell tumor** (Ref: Robbins 9th/pg 975)

### 13. Ans. (a) **Gland formation**

(Ref: Robbins 9th/pg 976; 8th/pg 988-89)



14. Ans. (a, b, c, d); a. **Raised in testicular tumor**; b. **Raised in 70% cases of HCC**; c. **Correlation between tumor recurrence after surgery in HCC**; d. **Correlation with HCC size**

(Ref: Walker's Pediatric Gastrointestinal Disease 5th ed; volume 2 :914, Pediatric Hematology and Oncology, 18:11-26, 2001)

**Increased AFP levels are seen in**

- Omphalocele
- Hepatocellular carcinoma/hepatoma
- Hepatoblastoma
- Neural tube defects: ↑  $\alpha$ -fetoprotein in amniotic fluid and maternal serum
- **Non-seminomatous germ cell tumors (option A)**
  - Yolk sac tumor
  - Immature teratoma (rarely)
- Ataxia telangiectasia:
- Serum AFP has significant correlation with the size of tumour.
- Its concentration in adult serum is less than 20 ng/ml.- **option E is false**

15. Ans. (c) **Seminoma**

(Ref: Robbins 9th/pg 975; 8th/pg 988-89)

16. Ans. (a, b, c, d); a. **HCC**; b. **Hepatoblastoma**; c. **Infant hemangioendothelioma**; d. **Amebic liver abscess**

(Ref: Walker's Pediatric Gastrointestinal Disease 5th ed; volume 2 :914)

Infantile hemangioendothelioma - most common hepatic vascular tumor in infants < 6 months of age; same cases can be associated with raised AFP

Elevated AFP levels are commonly found in acute liver disorders like Fulminant acute hepatitis, Liver cirrhosis and few cases of Liver abscess (pyogenic, amebic)

Embryonal sarcoma of liver is not associated with raised AFP

17. Ans. (c) **Alfa fetoprotein** (Ref: Robbins 9th/pg 979-80)

#### Tumour Markers For Testicular Tumors

Oncofetal substances	Cellular Enzymes
<ul style="list-style-type: none"> <li>• <math>\alpha</math>-FP: (Increases in YST)<sup>a</sup> <ul style="list-style-type: none"> <li>■ Produced by <b>trophoblastic cells</b><sup>a</sup></li> <li>■ <b>Increases in Yolk sac tumour, embryonal carcinoma and terato carcinoma. (YET)</b><sup>a</sup></li> <li>■ <b>Doesn't increase in pure choriocarcinoma</b><sup>a</sup></li> <li>■ Metabolic half life: 5-7 days<sup>a</sup></li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• LDH: <ul style="list-style-type: none"> <li>■ Not a specific tumour marker</li> <li>■ Most useful as a marker for "bulk" disease</li> <li>■ Raised serum LDH has poor prognosis</li> </ul> </li> <li>• PLAP (Placental alkaline phosphatase): <ul style="list-style-type: none"> <li>■ Elevated in <b>seminoma</b><sup>a</sup></li> </ul> </li> <li>• GGT (Gamma glutamyl transpeptidase): <ul style="list-style-type: none"> <li>■ Marker of <b>seminoma testis</b><sup>a</sup></li> <li>■ Marker for "bulk" disease</li> </ul> </li> </ul>

18. Ans. (c) **Endodermal sinus tumor**

(Ref: Robbins 9th/pg 977)

**Yolk sac tumor or endodermal sinus tumor show Schiller-Duval bodies or glomeruloid structures**

19. Ans. (c) **Prostate**

20. Ans. (d) **Carcinoma prostate** (Ref: Robbins 9th/pg 982)

21. Ans. (c) **PSA Nodularity** (Ref: Robbins 9th/pg 982)

Six variants in PSA Value

● <b>Age specific reference range</b>		● <b>PSA density:</b>
Age	Cut off	■ It is the ratio of serum PSA value and volume of prostate (PSA/volume prostate)
40-49 years	0 to 2.5 ng/ml	■ Upper normal limit is 0.15.
50-59 years	0 to 3.5 ng/ml	■ Butter discriminator between benign and malignant cancer than PSA
60-69 years	0 to 4.5 ng/ml	
70-79 years	0 to 6.5 ng/ml	
Raising the PSA threshold in older men improves specificity		
● <b>PSA velocity:</b>		● <b>Ratio of free and bound PSA in the serum:</b>
■ Rate of change of PSA with time.		■ It is calculated as $\text{Free PSA} / \text{Total PSA} \times 100$
■ <b>According to BLSA study PSA Velocity Of more than 0.75 ng/ml/Year shows increased of prostate cancer.</b>		■ <b>Free PSA less than 10% indicates high risk of carcinoma.</b>
● <b>Free versus total PSA</b>		● <b>Pro-PSA.</b>
■ Free PSA (not bound to other proteins)/ total amount of PSA (free plus bound)		■ Pro-PSA refers to several different inactive precursors of PSA.
■ Lower proportion of free PSA-more aggressive cancer.		■ Pro-PSA is more strongly associated with prostate cancer

22. Ans. (a) **Volume 1.5 ml, count 15 million, morphology 4% progressive motility 32%**

(Ref: Who 2010 Manual Sperm Analysis)


In 2010 the World Health Organization (WHO) updated its reference values for the Semen Analysis

Semen characteristics	WHO 2010
Volume (mL)	≥ 1.5
Sperm count (10 <sup>4</sup> /mL)	≥ 15
Total sperm count (10 <sup>6</sup> )	≥ 39
Total motility (%)	≥ 40
Progressive motility	≥ 32%

23. Ans. (d) **After liquefaction with thorough mixing**

(Ref: [www.who.int/reproductivehealth/publications/infertility](http://www.who.int/reproductivehealth/publications/infertility))

<b>Liquefaction</b>	Process when the gel formed by proteins from the seminal vesicles is broken up and the semen becomes more liquid In the NICE guidelines, a liquefaction time <b>&lt;60 minutes</b> -normal ranges Semen analysis should be done <b>after liquefaction with through mixing</b>
---------------------	---



**24. Ans. (d) Morphologically defective sperms**

(Ref: [www.who.int/reproductivehealth/publications/infertility](http://www.who.int/reproductivehealth/publications/infertility))

**25. Ans. (b) Vulva** (Ref: Robbins 9th ed p 999)

Extramammary Paget's is rare lesion of the vulva is similar in its manifestations to Paget disease of the breast. In the vulva, it presents as a pruritic, red, crusted, maplike area, usually on the labia majora. aget disease is a distinctive intraepithelial proliferation of malignant cells. Paget cells are larger than surrounding keratinocyte and are seen singly or in small clusters within the epidermis. The cells have pale cytoplasm containing mucopolysaccharide that stains with periodic acid-Schiff (PAS), Alcian blue, or mucicarmine stains. In addition, the cells express cytokeratin 7.

**26. Ans. (c) Fibroepithelial polyps** (Ref: Robbins 9th/pg 997)

**Predisposing factors for ca vulva**

- Cigarette use
- Human papillomavirus (HPV) infection
- Lichen sclerosus
- Premalignant lesions like leucoplakia
- Cervical cancer.
- Patients that are infected with HIV tend to be more susceptible to vulvar cancer as well.

Option C- **Fibroepithelial polyps are also known as skin tags- they are benign exophytic lesions of vulva**

**27. Ans (b) Sarcoma botryoides** (Ref: Robbins 9th/pg 1001)

**28. Ans. (a) Rhabdomyosarcoma** (Ref: Robbins 9th/pg 1001)

**29. Ans. (c) Actinomycetes**

(Ref: Textbook of Interpretation of PAP smear, refer to pretext for details)

The image given shows a wooly appearance; periphery containing swollen filaments with clubs of filament organism. This is suggestive of **Actinomycetes**.

**30. Ans. (b) Granulosa cell tumor**

(Ref: Textbook of Interpretation of PAP smear, refer to pretext for details)

In the given PAP smear, predominance of Mature cells (orange coloured cells compared to parabasal cells: blue coloured), might be due to estrogen secretion from Granulosa cell tumor.

**31. Ans. (a, b, c) a. Ki67 b. Oncoprotein E6 c. p16INK4, cyclin E, and Ki-67** (Ref: <http://>)

**32. Ans. (c) HPV-16**

(Ref: <https://aidsinfo.nih.gov/guidelines/html/>)

At least 12 HPV types are considered oncogenic, including HPV16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, and 59

- HPV16 alone, though, accounts for approximately 50% of cervical cancers in the general population and HPV18 for another 10% to 15%. The other oncogenic HPV types each individually account for fewer than 5% of tumors.
- HPV types 6 and 11 cause 90% of genital warts, but are not considered oncogenic

**33. Ans. (a) Smoking**

<b>Risk factors for ca cervix are</b>	<ul style="list-style-type: none"> <li>• Early age at first intercourse<sup>a</sup></li> <li>• Multiple sexual partners<sup>a</sup></li> <li>• Increased parity<sup>a</sup></li> <li>• High-risk HPVs (most important factor)</li> </ul>	<ul style="list-style-type: none"> <li>• Cigarette smoking, use of OCPs</li> <li>• <b>Genital infections<sup>a</sup></b></li> <li>• History of HSIL</li> <li>• <b>Certain HLA &amp; viral subtypes<sup>a</sup></b></li> </ul>
---------------------------------------	--	---

**34. Ans (a) Atrophic smear** (Ref: Dutta Gyne 4th ed:105)

MI is a ratio obtained through performing a random count of three major cell types (parabasal cells, intermediate cells and superficial cells) that are shed from the squamous epithelium.

The cell count is expressed as a percentage that reads as follows: **MI = % parabasal cells/ % intermediate cells/ % superficial cells.**

- **Atrophic smear** -100/0/0 means majorly parabasal cells i.e. no progesterone or estrogen.

**35. Ans. (b) Leiomyoma** (Ref: Robbins 9th ed p 1020)

Leiomyomas are sharply circumscribed, discrete, round, firm, gray-white tumors varying in size from small, barely visible nodules to massive tumors that fill the pelvis. Except in rare instances, they are found within the myometrium of the corpus. Leiomyomas are typically composed of bundles of smooth muscle cells that resemble the uninvolved myometrium. Usually, the individual muscle cells are uniform in size and shape and have the characteristic oval nucleus and long, slender bipolar cytoplasmic processes. Mitotic figures are scarce.

**36. Ans. (a) High estrogen** (Ref: R 9th/p 721)

**Increased estrogen production by endometriotic stromal cells**, due in large part to high levels of the key steroidogenic enzyme aromatase, which is absent in normal endometrial stroma. Mutations in specific genes (PTEN and ARID1A) in endometriotic cysts.

**37. Ans. (a) PTEN** (Ref: Robbins 9th/pg 1015)

Type 1 endometrial ca is MC and PTEN is MC involved in it.



**38. Ans. (b) Metropathia hemorrhagica**

(Ref: Textbook of gynecology by Rao:65)

**Metropathia hemorrhagica**

Anovulatory DUB associated with endometrial hyperplasia with acyclical bleeding  
Overgrowth of endometrial stroma and glands producing swiss cheese appearance

**39. Ans. (d) Granulosa theca cell tumor** (Ref: R 9th/pg 1032)

Granulosa cell tumor

- The **most common** malignant **Sex-Cord Stromal Tumors of ovary**
- The main **clinical features depend upon the oestrogenic activity of the tumour**<sup>Q</sup>
- The tumor cells may secrete estrogens and **cause precocious sexual development in girls** or increase the risk for **endometrial hyperplasia and carcinoma in women**.

**40. Ans. (a) P57** (Ref: Robbins 9th/pg 1039)

Gestational trophoblastic diseases include: benign hydatidiform mole (partial and complete), invasive mole, placental site trophoblastic tumor and choriocarcinoma


**Hydatidiform mole:**

There are two types of benign, noninvasive moles—complete and partial Both partial and complete, are composed of *avascular, grape-like structures* that **do not invade the myometrium**

Complete (classic) moles	Partial moles
All the chorionic villi are abnormal and fetal parts are not found	Only some of the villi are abnormal and fetal parts may be seen.
<b>46, XX diploid pattern</b>	Triploid or a tetraploid karyotype
MORPHOLOGY	
The chorionic villi are enlarged, scalloped in shape with central cavitation ( <b>cisterns</b> ) <b>Extensive</b> <sup>Q</sup> trophoblast proliferation	Focal villi enlargement Focal trophoblast proliferation
Risk of choriocarcinoma- <b>2.5%</b> Risk of persistent or invasive mole- <b>15%</b>	Have an increased risk of persistent molar disease, but are <b>not associated with choriocarcinoma</b> . <sup>Q</sup>

Immunostaining for p57, which is a gene that is paternally imprinted (inactivated) helps in differentiating the two moles. Because the complete mole arises only from paternal chromosomes, **immunostaining for p57 will be negative**.

**41. Ans. (d) Arise from the fertilization of a single egg by two sperm** (Ref: Robbins 9th/pg 1039)



**Figure:** Origin of complete and partial hydatidiform moles.

A. Complete moles most commonly (90%) arise from fertilization of an empty ovum by a single sperm that undergoes duplication of its chromosomes—karyotype 46XX. (**a phenomenon called androgenesis**<sup>Q</sup> as it is derived from paternal chromosome).

B. Less commonly, (10%) complete moles arise from dispermy in which two sperm fertilize an **empty ovum**—**karyotype**<sup>Q</sup> 46XX or 46XY karyotype.

C. Partial moles arise from **two sperm fertilizing a single ovum**.—**OPTION D IS FALSE**

**42. Ans. (b) Dysgerminoma**

Here, you see, solid ovarian neoplasm with sheets of large polygonal cells separated by septa prominent nucleon. The septa is infiltrated by lymphocytes.

**43. Ans. (a) Specimen shows multiple solid cystic areas suggestive of dermoid cyst.**

(Ref: Robbins 9th/pg 1029)

- Dermoid cysts are the **commonest solid ovarian neoplasm found in young women**.





44. Ans. (b) **Granulosa cell tumor**

(Ref: Robbins's 9<sup>th</sup>/pg 1029)

45. Ans. (b) **Krukenberg tumour**

(Ref: Robbins's Pathology 9th Ed/Pg 1032; Robbins's Pathology 8th ED/PG 1050; Shaw's Textbook of Gynaecology 15/e pg 425)

**Krukenberg Tumor**

- Refers to a **metastatic bilateral ovarian** malignancy whose primary site is **GIT or breast**.<sup>Q</sup>
- Composed of cellular or myxomatous stroma with scattered **mucin-producing, signet-ring cancer cells, most often of gastric origin (70%)**<sup>Q</sup>
- Ca colon, appendix, breast (**sp invasive lobular carcinoma**), pancreas and gall bladder are other **primary sites**.

46. Ans. (d) **Benign cystic teratoma** (Ref: Robbins 9th/pg 1030)

47. Ans. (b) **Sex cord stromal tumor** (Ref: Robbins 9th/pg 1032)

48. Ans. (a) **LH** (Ref: Robbins 9th/pg 1022)

**Polycystic ovarian disease (Stein-Leventhal syndrome)**

Characterized by	Endocrine abnormalities	Abnormalities in ovary
<ul style="list-style-type: none"> <li>• Hyperandrogenism</li> <li>• Menstrual abnormalities</li> <li>• Polycystic ovaries</li> <li>• Chronic anovulation</li> <li>• Decreased fertility<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Excess androgens</b> (androstenedione)<sup>Q</sup></li> <li>• <b>Increased estrogen levels</b><sup>Q</sup></li> <li>• Increased LH &amp; decreased FSH levels</li> <li>• <b>High LH/FSH ratio</b></li> <li>• Increased GnRH levels</li> </ul>	<ul style="list-style-type: none"> <li>• Enlarged with thick capsules</li> <li>• Hyperplastic ovarian stroma</li> <li>• Numerous follicular cysts lined by a hyperplastic theca interna.</li> </ul>

49. Ans. (a) **Luteoma**

(Ref: Best Practice and Research Clinical Endocrinology and Metabolism (Elsevier Ltd.) 25:985-992)

A **luteoma** is a tumor that occurs in the ovaries during pregnancy.

It is associated with an increases of sex hormones, primarily progesterone and testosterone

Benign

Resolve themselves after delivery

50. Ans. (a) **Serous cystadenoma** (Ref: Robbins 9th/pg 1023)

- Most ovarian tumors are surface epithelial (65-70%)
- Serous cystadenoma is the most common surface epithelial tumor

51. Ans. (b) **Teratoma** (Ref: R 9th/pg 1029; 8th/pg 1057-48)

- **Mature (Benign) Teratomas**
- **Rokitansky protuberance**<sup>Q</sup>-The inner lining of cyst contains single or multiple white shiny masses projecting from the wall toward the center of the cysts. Hair, other dermal appendages, bone and teeth are present, they usually arise from this protuberance

52. Ans. (a) **CA-125**

(Ref: Robbins 9th/pg 1033; Am J Prev Med 13 (6): 444-6)

- CA125- is a glycoprotein
- Elevated levels in **epithelial ovarian cancers**
- CA-125 is the most frequently used biomarker for ovarian cancer detection

**Elevated CA125- not a specific marker**

- In other cancers, including **endometrial cancer, fallopian tube cancer, lung cancer, breast cancer and gastrointestinal cancer**.
- **Benign conditions**, such as endometriosis, PID, cirrhosis and diabetes mellitus

53. Ans. (a, c, e); a. **Glycoprotein**; c. **Increased in colon carcinoma**; e. **May be elevated in Pelvic inflammatory disease**

(Ref: Robbins 9th/pg 1033; 8th/pg 1052, Am J Prev Med 13 (6): 444-6)

54. Ans. (a) **Follow up of ovarian cancer**

(Ref: Robbins 9th/pg 1033; 8th/pg 1052, Am J Prev Med 13 (6): 444-6)

55. Ans. (c) **Inhibin** (Ref: Robbins 9th/pg 1032)

Elevated tissue and serum levels of **inhibin**,<sup>Q</sup> a product of granulosa cells, are associated with granulosa cell tumors

Option a- CA19-9- pancreatic cancer (Ref: Eur J Surg Oncol. Apr 2007;33(3):266-70)

Option D- NSE- Neuroblastoma

56. Ans. (d) **Gonadal type is chemosensitive**

(Ref: Robbins 9th/pg 1031)

- **In contrast to choriocarcinomas arising in placental tissue, those arising in the ovary are generally unresponsive to chemotherapy and are often fatal.**<sup>Q</sup>


# Breast

## Key Points

- » Most common benign tumor of the female breast—fibroadenoma
- » Most common malignant tumor of the female breast—ductal carcinoma
- » Most common tumor of the female breast to be seen bilaterally—lobular carcinoma
- » Extramammary Paget's disease is usually seen in isolation & associated with an underlying malignancy in 12%

## Key Recent Updates

- » ITGCN is now renamed as GCNIS i.e. germ cell neoplasm in situ
- » SOX, 2 is associated with embryonal carcinoma.



## DISORDERS OF DEVELOPMENT

- **Congenital nipple inversion**
  - Congenitally inverted-little significance
  - Correct spontaneously<sup>Q</sup> during pregnancy, or can be everted by simple traction
- **Milk-line remnants**
  - Result from the persistence of epidermal thickenings along the milk line (axilla to the perineum)
  - Most commonly come to attention as a result of painful premenstrual enlargements.
- **Accessory Axillary Breast Tissue**

## Clinical Presentations of Breast Disease

- **Pain** (mastalgia or mastodynia) is **the most common<sup>Q</sup>** breast symptom.
- Discrete palpable masses are the **second most common** breast symptom.
- **Most common palpable lesions** are cysts, fibroadenomas, and invasive carcinomas
- Benign palpable masses are **most common in premenopausal women<sup>Q</sup>**
- Mass becomes palpable when size >2 cm
- Most common site of carcinomas-upper outer quadrant (50%)

**Nipple discharge** is a less common presenting symptom, but is of concern when it is spontaneous and unilateral.

Type of discharge	Cause
Milky discharges ( <i>galactorrhea</i> )	<ul style="list-style-type: none"> <li>• Elevated prolactin levels (e.g., by a pituitary adenoma), hypothyroidism, endocrine an ovulatory syndromes, OCP, tricyclic antidepressants, methyl dopa, or phenothiazines</li> <li>• <b>Galactorrhea is not associated with malignancy.<sup>Q</sup></b></li> </ul>
Bloody or serous discharges	<ul style="list-style-type: none"> <li>• Solitary large duct papilloma, cysts, or carcinoma</li> </ul>

## Mammographic Signs of Breast Carcinoma are Densities and Calcifications

Densities	Calcifications
<ul style="list-style-type: none"> <li>• Most neoplasms are radiologically denser than normal breast.</li> <li>• Rounded densities are <b>most commonly benign lesions</b> such as fibroadenomas or cysts</li> </ul>	<ul style="list-style-type: none"> <li>• <i>Often associated with benign lesions eg clusters of apocrine cysts, hyalinized fibroadenomas &amp; sclerosing adenosis</i></li> <li>• <i>Calcifications associated with malignancy are usually small, irregular, numerous, and clustered.<sup>Q</sup></i></li> </ul>

## Inflammatory Disorders

Account for less than 1% of breast symptoms.

<b>Acute mastitis</b>	<ul style="list-style-type: none"> <li>• Occurs during the first month<sup>Q</sup> of breastfeeding; Most common cause is <b><i>Staphylococcus aureus</i><sup>Q</sup></b></li> </ul>
<b>Squamous metaplasia of lactiferous ducts</b>	<ul style="list-style-type: none"> <li>• Also known as <b>recurrent subareolar abscess, periductal mastitis, and Zuska disease.<sup>Q</sup></b></li> <li>• Present with a painful erythematous subareolar mass; &gt;90% patients are <b>smokers.<sup>Q</sup></b></li> <li>• Relative <b>deficiency of vitamin A<sup>Q</sup></b> associated with smoking or toxic substances in tobacco smoke alter the differentiation of the lactiferous ducts</li> <li>• <b>Keratinizing squamous epithelium<sup>Q</sup></b> extending to an abnormal depth into orifices of nipple ducts.</li> </ul>
<b>Duct ectasia</b>	<ul style="list-style-type: none"> <li>• Occur in the fifth or sixth decade of life<sup>Q</sup>, usually in multiparous women</li> <li>• Periareolar mass<sup>Q</sup> associated with thick, white nipple secretions &amp; occasionally skin retraction</li> <li>• Not associated with cigarette smoking.<sup>Q</sup></li> </ul>
<b>Fat necrosis</b>	<ul style="list-style-type: none"> <li>• Painless palpable mass, skin thickening or retraction, a mammographic density or calcifications.</li> <li>• About half of affected women have a history of breast trauma or prior surgery<sup>Q</sup></li> <li>• Grossly-ill-defined, firm, gray-white nodules containing small chalky-white foci<sup>Q</sup></li> <li>• Microscopically- central areas of liquefactive fat necrosis with neutrophils and macrophages<sup>Q</sup></li> </ul>
<b>Granulomatous mastitis</b>	<p>Can be a manifestation of systemic granulomatous diseases (e.g., granulomatosis with polyangiitis, sarcoidosis, TB, fungi) or of disorders that are localized to the breast (granulomatous lobular mastitis)<sup>Q</sup></p>





## Latest Update

### Lymphocytic Mastopathy (Sclerosing Lymphocytic Lobulitis)

- Presents with single or multiple hard palpable masses or mammographic densities<sup>Q</sup>
- Most common in women with **type 1 (insulin-dependent) diabetes** or **autoimmune thyroid disease**
- Granulomatous lobular mastitis is an uncommon disease that only occurs in parous women. <sup>Q</sup>-Caused by a hypersensitivity reaction to antigens expressed during lactation<sup>Q</sup>
- Cystic neutrophilic granulomatous mastitis is caused by *Corynebacteria*<sup>Q</sup>

## Benign Epithelial Lesions

Non-proliferative breast changes (fibrocystic changes):

Cysts	Fibrosis	Adenosis
<ul style="list-style-type: none"> <li>• Form by the dilation of lobules</li> <li>• <b>Bluedome cysts</b>-turbid, semi-translucent fluid of a brown or blue color in unopened cysts</li> <li>• Calcifications are common and may be detected by mammography</li> </ul>	<p>Cysts frequently rupture ↓ Release of secretory material into adjacent stroma ↓ Chronic inflammation &amp; fibrous scarring lead to palpable firmness of the breast.</p>	<ul style="list-style-type: none"> <li>• Increase in number of acini per lobule</li> <li>• Acini - lined by columnar cells,</li> <li>• Benign or <b>show nuclear atypia ("flat epithelial atypia")</b>.</li> </ul>



## High Yield Facts

- Flat epithelial atypia is a **clonal proliferation associated with deletions of chromosome 16q**.<sup>Q</sup>
- **Earliest recognizable precursor of low-grade breast cancers**, but does not convey an increased cancer risk



## Proliferative Breast Disease without Atypia

Associated with a small increase in the risk of subsequent carcinoma in either breast<sup>Q</sup>

<b>Epithelial Hyperplasia</b>	<ul style="list-style-type: none"> <li>• Defined by the <b>presence of more than two cell layers</b><sup>Q</sup></li> <li>• Normally, only myoepithelial cells &amp; <b>single layer of luminal cells</b> are present</li> </ul>
<b>Sclerosing Adenosis</b>	<ul style="list-style-type: none"> <li>• Increased number of acini <b>compressed &amp; distorted in central portion</b> of lesion but <b>dilated at the periphery</b>; Myoepithelial cells are usually prominent</li> </ul>
<b>Complex Sclerosing Lesion</b>	<ul style="list-style-type: none"> <li>• Have components of sclerosing adenosis, papillomas &amp; epithelial hyperplasia.</li> <li>• One member of this group, the radial sclerosing lesion ("radial scar")</li> <li>• Radial scars are stellate lesions with a <b>central nidus of entrapped glands in a hyalinized stroma</b></li> </ul>
<b>Papillomas</b>	<ul style="list-style-type: none"> <li>• Papillomas grows within a <b>dilated duct</b></li> <li>• <b>Large duct papillomas</b> are situated in <b>lactiferous sinuses of the nipple</b> (<b>solitary</b> with nipple discharge)</li> <li>• <b>Small duct papillomas</b> are <b>multiple &amp; located deeper within the ductal system</b> &amp; usually present as <b>small palpable masses, or densities/calcifications on mammogram</b></li> </ul>

## Proliferative Breast Disease with Atypia

- Proliferative disease with atypia includes (ADH) and (ALH).
- Atypical hyperplasia is a clonal proliferation having some, but not all, of the histologic features that are required for the diagnosis of carcinoma in situ.

	Atypical ductal hyperplasia (ADH)	Atypical lobular hyperplasia (ALH)
Morphology	Histologic resemblance to ductal carcinoma in situ Monomorphic cells only partially fill involved ducts. <sup>Q</sup>	Resembles lobular carcinoma in situ Cells do not fill or distend <b>more than 50%</b> <sup>Q</sup> of the acini within a lobule
Mutation	Loss of 16q or gain of 17p <sup>Q</sup>	Loss of E-cadherin <sup>Q</sup>
		





## Epithelial Breast Lesions and the Risk of Developing Invasive Carcinoma

Pathological lesion	Relative Risk (Absolute Lifetime Risk)*
Nonproliferative Breast Changes (Fibrocystic changes)	1 (3%) <sup>Q</sup>
Proliferative Disease Without Atypia	1.5 to 2 (5%-7%) <sup>Q</sup>
Proliferative Disease with Atypia	4 to 5 (13%-17%) <sup>Q</sup>
Carcinoma in Situ	8 to 10 (25%-30%) <sup>Q</sup>

## CARCINOMA OF THE BREAST

Based on the expression of estrogen receptor and HER2:

### Hereditary Breast Cancer

Gene (Location)	% of "Single Gene"	Risk by Age 70	Other Associated Cancers	Comments
BRCA1 (17q) <sup>Q</sup>	52% <sup>Q</sup>	40%- 90%	Ovarian, male breast cancer (but lower than BRCA2), prostate, pancreas, fallopian tube	Poorly differentiated and triple negative <sup>Q</sup> (basal-like)
BRCA2 (13q) <sup>Q</sup>	32%	30%-90%	Ovarian, male breast cancer, prostate, pancreas, stomach, melanoma, gallbladder, bile duct, pharynx	Bi-allelic germline Mutations; Fanconi anemia <sup>Q</sup>
TP53 (17p) Li-Fraumeni	3%	>90% <sup>Q</sup>	Sarcoma, leukemia, brain tumors, adrenocortical carcinoma	MC sporadic breast Ca <sup>Q</sup>
CHEK2 (22q)	5%	10%-20%	Prostate, thyroid, kidney, colon	Increase risk for breast Ca after radiation exposure <sup>Q</sup>



### High Yield Facts

- **BRCA1 and BRCA2** 80- 90% of "single gene" familial breast cancers and 3% of all breast cancers
- **BRCA1-associated breast cancers** are poorly differentiated, have "medullary features" (a syncytial growth pattern with pushing margins and a lymphocytic infiltrate) and are **triple negative** (ER, PR, Her2neu negative)

- **BRCA2-associated breast carcinomas** also poorly differentiated, but are **ER-positive**
- **Three other tumor suppressor genes**—PTEN (Cowden syndrome), STK11 (Peutz-Jeghers syndrome), and ATM (ataxia telangiectasia)→<1% of all familial breast cancers.

## CLASSIFICATION OF BREAST CARCINOMA

### Noninvasive Carcinomas (Carcinoma in situ)

Noninvasive carcinomas (carcinoma in situ) may be located within the ducts (intraductal carcinoma) or within the lobules (lobular carcinoma in situ).

### DCIS (Ductal Carcinoma in situ)

- Among the mammographically detected cancer, almost half are DCIS.<sup>Q</sup>

- Estrogen receptor (ER)-positive, HER2-negative (50% to 65% of tumors);<sup>Q</sup>
- HER2-positive (10% to 20% of tumors, which may either be ER-positive or ER-negative);<sup>Q</sup>
- ER-negative, HER2-negative (10% to 20% of tumors).<sup>Q</sup>

### Etiology and Pathogenesis



- **Major risk factors** for the development of breast cancer are **genetic and hormonal**<sup>Q</sup>
- 12% of breast cancers occur due to inheritance of an identifiable susceptibility gene or genes.<sup>Q</sup>
- **Major risk factors for sporadic breast cancer are related to hormone exposure:** gender, age at menarche and menopause, reproductive history, breastfeeding, and exogenous estrogens.<sup>Q</sup>


- DCIS most frequently presents as **mammographic calcifications**.<sup>Q</sup>

### Histological types of DCIS (five types)

- Comedo carcinoma<sup>Q</sup>
- Cribriform<sup>Q</sup>
- Micropapillary<sup>Q</sup>
- Solid
- Papillary<sup>Q</sup>



Ductal carcinoma in situ (DCIS)	Lobular carcinoma in situ (LCIS)
<ul style="list-style-type: none"> <li>• Malignant clonal proliferation of epithelial cells limited to <b>ducts</b> by the <b>basement membrane</b>.</li> <li>• <b>Myoepithelial cells are preserved<sup>Q</sup></b> in involved ducts/lobules</li> <li>• <b>Detected</b> by mammography as <b>calcifications<sup>Q</sup></b></li> <li>• Divided into comedo and non-comedo- cribriform carcinoma, and intraductal papillary carcinoma</li> <li>• <b>Comedocarcinoma-solid intraductal sheet of cells</b> with a central area of <b>necrosis</b>-undergoes calcification, can rarely present as <b>palpable mass<sup>Q</sup></b></li> <li>• Associated with <b>erb B2/neu oncogene</b> and <b>poor prognosis</b>.</li> <li>• Cribriform carcinoma: <b>round, duct like structures<sup>Q</sup></b> within the solid intraductal sheet of epithelial cells</li> <li>• Intraductal papillary carcinoma has a <b>papillary pattern<sup>Q</sup></b>.</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Incidental biopsy finding<sup>Q</sup></b></li> <li>• <b>Not associated with calcifications</b> or stromal reactions that produce mammographic densities</li> <li>• <b>Bilateral in 20% to 40% of cases</b></li> <li>• LCIS is more common in <b>young women</b>,</li> <li>• 80 - 90% of cases occurring before menopause.</li> <li>• Cells <b>lack the cell adhesion protein E-cadherin</b></li> <li>• LCIS always expresses <b>ER and PR</b></li> </ul>
	


**Latest Update**

**DCIS with microinvasion**

- When there is an area of invasion through the basement membrane into stroma measuring no more than 0.1 cm. <sup>Q</sup>

## INVASIVE (INFILTRATING) CARCINOMA

- Divided on the basis of molecular and morphologic characteristics

**Molecular subtypes:**

**Molecular profiling** of breast cancer is done on gene **profiling<sup>Q</sup>**

**A. ER-positive, HER- negative (also termed "luminal" 60-70%):**

- **Most common form<sup>Q</sup> of invasive breast cancer.**
- Essentially all are well differentiated carcinomas.
- Mucinous, papillary, cribriform, and lobular patterns may be present in this group<sup>Q</sup>

Based on proliferation rates, it is further divided into two subgroups.

<b>ER-positive, HER2-negative, low proliferation (40% -55%)</b> <b>Luminal A</b>	<ul style="list-style-type: none"> <li>• <b>Most common form of invasive breast cancer.</b> <sup>Q</sup></li> <li>• Majority of cancers in <b>older women and in men.</b> <sup>Q</sup></li> <li>• MC type detected by mammographic screening and in women treated with menopausal hormone therapy <sup>Q</sup></li> <li>• <b>Lowest incidence</b> of local <b>recurrence</b> &amp; often <b>cured by surgery</b> <sup>Q</sup></li> <li>• Respond well to hormonal treatment with long survival <sup>Q</sup></li> <li>• <b>Metastasise late</b> but usually to bone.<sup>Q</sup></li> </ul>
<b>ER-positive, HER2-negative, high proliferation (10%):</b> <b>Luminal B</b>	<ul style="list-style-type: none"> <li>• Most common type of carcinoma associated with <b>BRCA2 germline mutations<sup>Q</sup></b></li> </ul>



**B. HER2-positive (20% of cancers): 2<sup>nd</sup> MC subtype of invasive breast cancer**

- **More common in young women and in non-white women.** <sup>Q</sup>
- Majority of these carcinomas are poorly differentiated tumors <sup>Q</sup>
- **No specific morphologic pattern associated with this cell type**
- **50% of apocrine<sup>Q</sup> Carcinomas and 40% of micropapillary carcinomas belong to this category.**
- **The associated DCIS is often extensive** <sup>Q</sup>
- **>50% tumors with germlineTP53 mutations** (Li-Fraumeni syndrome) develop carcinomas of this subtype
- Responds to trastuzumab (Herceptin),- humanized monoclonal antibody that specifically binds and inhibits HER2






### C. ER-negative, HER2-negative tumors ("basal-like" triple negative Ca; 15%)-young premenopausal women as well as African American and Hispanic women




- The majority of carcinomas arising in women with BRCA1 mutations are of this type <sup>a</sup>
- Almost all of these tumors are poorly differentiated.
- Spindle cell, squamous, and matrix producing patterns can also be seen. <sup>a</sup>
- DCIS is generally very limited or not present. <sup>a</sup>
- Cancers can metastasize when small in size, frequently to viscera and to the brain. <sup>a</sup>
- Characterised by rapid growth, high proliferation-present as a palpable mass in the interval between mammographic screenings <sup>a</sup>
- 30% completely respond to chemotherapy <sup>a</sup>
- Local recurrence is common <sup>a</sup>
- Express markers of typical myoepithelial cells (e.g. basal keratins, P-cadherin, p63 or laminin), progenitor cells or putative stem cells (cytokeratin 5 and 6) <sup>a</sup>



LUMINAL TYPE



HER2NEU TYPE




TRIPLE  
NEGATIVE

### On the basis of morphology:

#### Invasive breast carcinoma 2 types: No- special type carcinoma (Intraductal) & Special carcinoma


#### Invasive Carcinoma, No Special Type (NST; Invasive Ductal Carcinoma) <sup>a</sup>

- Majority of carcinomas (70 to 80%) belong to this subtype.
- They MC present as a hard, irregular radiodense mass associated with a desmoplastic stromal reaction <sup>a</sup>
- On gross examination, most carcinomas are firm to" hard and have an irregular border.
- All types of invasive carcinoma are graded using **Modified Bloom Richardson Score**. <sup>a</sup>
- Carcinomas are scored for **tubule formation, nuclear pleomorphism, and mitotic rate** <sup>a</sup>
- The points added to divide carcinomas into grade I (well differentiated), grade II (moderately differentiated), and grade III (poorly differentiated) types.



Invasive ductal Ca: Malignant glands (M) invading stroma (S)





## Special -subtypes

- |   |   |  |
|---|---|--|
| <ul style="list-style-type: none"> <li>Lobular</li> <li>Medullary</li> <li>Metaplastic</li> <li>Apocrine carcinoma</li> </ul> | <ul style="list-style-type: none"> <li>Cribriform</li> <li>Tubular</li> <li>Inflammatory carcinoma</li> </ul> | <ul style="list-style-type: none"> <li>Colloid</li> <li>Papillary</li> <li>Micropapillary carcinoma</li> </ul> |
|---|---|--|

Special histologic types of breast cancer often harbor unique genetic aberrations, sometimes have **distinct gene signatures**

### 1. Lobular carcinoma (invasive) of breast

- Most cases show biallelic loss of expression of CDH1<sup>Q</sup>, the gene that encodes E-cadherin.<sup>Q</sup>
- Loss of E-cadherin, lobular carcinomas are discohesive<sup>Q</sup> and fail to incite a desmoplastic response<sup>Q</sup>
- Seen bilaterally
- Most common type of breast carcinoma to present as an occult primary<sup>Q</sup>
- Characteristic patterns of metastatic spread:** involving the peritoneum and retroperitoneum, the leptomeninges (carcinomatous meningitis), the gastrointestinal tract, and the ovaries and uterus.<sup>Q</sup>
- Histologic hallmark:** presence of discohesive infiltrating tumor cells as single cells in **single file pattern**<sup>Q</sup>, including **signet-ring cells**<sup>Q</sup> containing intracytoplasmic mucin droplets<sup>Q</sup>
- Tubule formation is absent.
- Males and females with heterozygous germline mutations in CDH1 also have a greatly increased risk of **gastric signet ring cell carcinoma**<sup>Q</sup>



Indian file pattern S/o Lobular Ca Breast

### 2. Medullary Carcinoma

- Characterised by features that are characteristic of BRCA1-associated carcinomas.
- Comes under the category of **ER-negative, HER2-negative tumors**<sup>Q</sup>
- DCIS is **minimal or absent**.<sup>Q</sup>
- The tumor has a **soft, fleshy consistency** and is **well-circumscribed**
- Medullary carcinomas have **better prognosis** than do NST carcinomas<sup>Q</sup>

### Characterized by

- Solid, syncytium-like sheets (occupying more than 75% of the tumor) of large cells with vesicular, pleomorphic nuclei, containing prominent nucleoli
- Frequent mitotic figures
- A moderate to marked lymphoplasmacytic infiltrate surrounding and within the tumor
- Pushing (non-infiltrative) border



### Latest Update

Current WHO classification system recommends grouping medullary carcinomas with similar carcinomas into one group termed "carcinomas with medullary features."<sup>Q</sup>

#### ER-negative, HER2-negative tumors

- Secretory carcinoma
- Spindle cell carcinoma
- Low-grade adenosquamous carcinoma
- Adenoid cystic carcinoma
- Medullary carcinoma



### Latest Update

Two special histologic types frequently overexpress HER2

Apocrine carcinoma	Micropapillary carcinoma
Resemble the cells that line sweat glands.	Shows a characteristic pattern of anchorage-independent growth. <sup>Q</sup> Although the cells are adherent to each other and express E-cadherin, they lack adhesion to the stroma <sup>Q</sup>



### High Yield Facts

- Special type carcinomas carry good prognosis but inflammatory carcinoma has poor prognosis
- Both lobular carcinoma of breast and signet ring carcinoma of GIT are characterized by the loss of E-cadherin<sup>Q</sup>

## PROGNOSTIC FACTORS

### Major

- Invasive carcinoma** has worse prognosis than in-situ carcinoma
- Distant **metastasis** indicates bad prognosis.

#### Lymph node status:

- Axillary lymph node status:** most important prognostic factor for invasive ca in the absence of distant metastases.<sup>Q</sup>
- Sentinel lymph node status:** If negative for metastasis, it is unlikely that other more distant nodes will be involved and the patient can be spared the morbidity of a complete axillary dissection<sup>Q</sup>





**Lymphovascular invasion: strongly associated with the presence of lymph node metastases with poor prognostic factor**

- **Tumor Size** <1 cm good prognosis, > 2 cm bad prognosis.
- **Local invasion** into skeletal muscle carries poor prognosis.
- **Inflammatory carcinoma** has poor prognosis

## Minor

- **Molecular subtype-** explained above
- **Histological type:** Invasive ductal carcinoma (no special type; NST) carries poor prognosis.
- **Tubular, mucinous, lobular, papillary, adenoid cystic** has better prognosis
- **Metaplastic carcinoma or micro-papillary carcinoma** have a poorer prognosis<sup>Q</sup>
- Nottingham histological score (Scarff-Bloom-Richardson grade): Grade 1 good prognosis, grade 3 poor
- **Estrogen and Progesterone receptor** positivity indicates good response to anti-estrogen therapy.
- **Strongly ER-positive cancers** are less likely to respond to chemotherapy<sup>Q</sup>
- **HER2/neu** overexpression: Poor prognosis
- High **proliferative rate** indicates worse prognosis
- **Aneuploidy** indicates bad prognosis

## Stromal Tumors

Origin from intralobular carcinoma		Origin from interlobular carcinoma	
<b>Fibroadenoma:</b> <ul style="list-style-type: none"> <li>• Most common benign tumor of the female breast.<sup>Q</sup></li> <li>• "Proliferative changes without atypia"</li> </ul>		<b>Benign- Myofibroblastoma</b> only breast tumor that is equally common in males <sup>Q</sup>	
<b>Phyllodes Tumor/Cystosarcoma phyllodes:</b> Gains in <b>chromosome 1q</b> being the most frequent mutation <sup>Q</sup> <b>Gross:</b> "leaf like due to stromal proliferations" <b>Grade:</b> benign, intermediate and malignant		<b>Malignant:</b> <ul style="list-style-type: none"> <li>• <b>Angiosarcoma:</b> most common stromal malignancy</li> <li>• Sporadic or associated with <b>radiation exposure</b> or <b>lymphedema</b>.<sup>Q</sup></li> </ul>	

## Male Breast Cancer


The incidence in breast cancer in men is only 1% of that in women

## Risk Factors

- Increasing age<sup>Q</sup>
- First-degree relatives with breastcancer<sup>Q</sup>
- Exposure to exogenous estrogens or ionizing radiation<sup>Q</sup>
- Infertility<sup>Q</sup>
- Obesity<sup>Q</sup>
- Prior benign breast disease<sup>Q</sup>
- From 3% to 8% of cases are associated with Klinefelter syndrome and decreased testicular function.<sup>Q</sup>
- From 4% to 14% of cases in males are attributed to germline BRCA2 mutations<sup>Q</sup>
- Male breast cancer is also observed in BRCA1 families, although not as frequently<sup>Q</sup>
- Most common histopathology- breast Carcinoma NSTQ
- ER positivity is more common<sup>Q</sup>
- Distant metastases to the lungs, brain, bone, and liver are common.
- Most cancers are treated locally with mastectomy and axillary node dissection




**Complex fibroadenomas**—fibroadenomas associated with cysts larger than 0.3 cm<sup>Q</sup>, sclerosing adenosis<sup>Q</sup>, epithelial calcifications<sup>Q</sup>, or papillary apocrine change<sup>Q</sup>



## NEXT Pattern Question




1. 24/f presented with painless left sided breast lump for last 2 months. Local examination revealed soft mobile, non tender mass. Excision biopsy was done and histopathological image is as shown below. What is your diagnosis?



- a. Phyllodes tumor      b. Intraductal ca breast      c. Fibroadenoma      d. Ductal ca in situ

**Ans. (c) Fibroadenoma**

- Painless left sided breast lump which is soft mobile, non-tender mass looks like a benign lesion. The histopathology shows proliferation of pinkish stroma and ducts which have become distorted. This is suggestive of fibroadenoma.



## Multiple Choice Questions

### BREAST

#### INFLAMMATORY DISORDERS


1. **Granulomatous mastitis is caused by all except-** *(Recent Question 2014)*
- TB
  - Fungus
  - Staphylococcus
  - Antibodies to milk antigens

#### BENIGN EPITHELIAL LESIONS

2. **All are benign conditions except** *(Recent Question 2014-15)*
- Fibroadenoma
  - Cystosarcoma phyllodes
  - Pagets disease of nipple
  - Galactocele
3. **A 17/F underwent FNAC for a lump in the breast which was non-tender, firm and mobile. Which of the following features would suggest finding of a benign breast disease?** *(AIIMS Nov 14)*
- Dyscohesive ductal epithelial cells without cellular fragments
  - Tightly arranged ductal epithelial cells with bare nuclei
  - Stromal predominance with spindle cells
  - Polymorphism with single or arranged ductal epithelial cells
4. **Lesions affecting the terminal duct lobular unit (TDLU) in breast are all except -** *(DPG 11)*
- Nipple adenoma
  - Blunt duct adenosis
  - Intraductal papilloma
  - Fibroadenoma

#### CARCINOMA OF THE BREAST

5. **A 30-year-old female presented with 4 cm mass in the right breast. Biopsy showed densely packed cells with bland nuclei and mucin infiltrating the stroma. What is your diagnosis?** *(Recent Pattern Question 2020)*




- Invasive papillary carcinoma
- Medullary carcinoma
- Apocrine carcinoma
- Colloid carcinoma

6. **Van Nuys prognostic indicator for DCIS does not include which of the following parameter?** *(Recent exam 2018)*

- DCIS size
- Age of the patient
- Type of DCIS
- Excision margin

7. **The pictures below show breast biopsy specimen on which immuno histochemical for ER, PR and Her2 neu have been done. Which of the following is true about the prognosis of the patient?** *(AIIMS May 16)*



- Good prognosis
- Poor prognosis
- Good prognosis with trastuzumab therapy can be given
- Good prognosis without trastuzumab therapy should not be given

8. **Most important prognostic factor in Ca. Breast.**

- Lymph Node status
- Tumor Size
- Progesterone receptor status
- Stage

9. **Somatic mutation E17K in the PH domain of AKT-1 gene mutation is associated with?**

*(Recent Question 2016)*

- Stomach
- Breast
- Ovary
- Pancreas

10. **Cancer detected in one breast which to be screened in contralateral breast** *(Recent Question 2016)*

- Lobular
- Ductal
- medullary
- colloid

11. **Male breast cancer wrong statement-**

- Brca2 seen in 6% cases
- Lobular carcinoma is common
- DUCTAL carcinoma is most common subtype
- Colloid carcinoma can be seen

12. **Cystosarcoma phylloides, true is?** *(PGI Nov 2015)*

- It has a spectrum of benign to malignant
- Can be rarely seen in children
- Always malignant
- Always benign
- Seen in elderly females

13. **Following is not true about the gene mutations leading to breast carcinoma** *(Recent Question 2014-15)*

- Most common mutation in inherited breast carcinoma is BRCA1
- BRCA 1 mutation is present in most of the cases of breast carcinoma
- Inherited breast carcinomas make about 3 % of the total cases
- p53 mutation also increases chances of colon and brain cancer





- 14. Breast CA with best prognosis (Recent Question 2015)**  
a. Mucinous                      b. Medullary  
c. Invasive ductal              d. Lobular Ca
- 15. All of the following are invasive carcinoma breast except- (Recent Question 2015)**  
a. Comedo carcinoma  
b. Colloid carcinoma  
c. Lobular carcinoma  
c. Medullary carcinoma
- 16. The type of mammary ductal carcinoma in situ (DCIS) most likely to result in a palpable abnormality in the breast is: (Recent Question 2014)**  
a. Apocrine DCIS  
b. Neuroendocrine DCIS  
c. Well differentiated DCIS  
d. Comedo DCIS
- 17. Molecular classification of breast cancer is based on? (AIIMS Nov 14)**  
a. Gene profiling              b. ER, PR, and HER-2 neu  
c. histology                      d. Mutations
- 18. The most common site for lymphagiosarcoma is: (AP PGME 14)**  
a. Liver  
b. Spleen  
c. Post mastectomy arm  
d. Retroperitoneum
- 19. True about histology in infiltrating lobular breast carcinoma: (JIPMER 11)**  
a. Single file pattern              b. Pleomorphic cells in sheets  
c. Cribiform pattern              d. Pin wheel pattern
- 20. Modified Bloom Richardson criteria for CA breast includes: (PGI May 2010)**  
a. Desmoplasia                      b. Lymphovenous embolism  
c. Mitotic rate                      d. Tubule formation  
e. Nuclear polymorphism
- 21. Breast Ca is not a/w: (PGI May 10)**  
a. BRCA I & BRCA 2  
b. Apocrine metaplasia  
c. Atypical ductal hyperplasia  
d. Fibroadenoma  
e. Moderate hyperplasia
- 22. Most common carcinoma of breast is: (Recent Question 2016, MH PGME 2016, TN 97)**  
a. Ductal carcinoma  
b. Colloid carcinoma  
c. Lobular carcinoma  
d. Sarcoma phylloides
- 23. Malignancy of the Breast is likely to be associated with (MH PGME 16, AI 94)**  
a. Sclerosing adenosis  
b. Atypical epithelial hyperplasia  
c. Cystic change  
d. Apocrine metaplasia



## Answers with Explanations

### 1. Ans. (c) **Staphylococcus** (Ref: Robbins 9th/pg 1047)

- **Granulomatous mastitis:** Can be a manifestation of systemic granulomatous diseases (e.g., granulomatosis with polyangiitis, sarcoidosis, TB, fungi) or of disorders that are localized to the breast (granulomatous lobular mastitis)<sup>Q</sup>

**Granulomatous lobular mastitis-** only occurs in parous women, caused by a **hypersensitivity reaction to antigens expressed during lactation**<sup>Q</sup>

### 2. Ans. (c) **Pagets disease of nipple**

(Ref: Robbins 9th/pg 1057)

Option a and d are clearly benign

We get confused in option b and c

Now cystosarcoma phyllodes are usually benign but can be borderline or rarely malignant also

Mammary Pagets disease is always associated with an underlying carcinoma of the breast. Mammary Paget cells are malignant epithelial cells derived from underlying ductal adenocarcinoma of the breast that invade into the skin of nipple and areolar areas.

### 3. Ans. (b) **Tightly arranged ductal epithelial cells with bare nuclei**

(Ref: Gray Diagnostic Cytopathology 2nd ed: 279-80)

- Option A- Dyscohesive ductal epithelial cells without cellular fragments- feature of ductal carcinoma
- Option B- **Tightly arranged ductal epithelial cells with bare nuclei- fibroadenoma (benign disease)**
- Option C- Stromal predominance with spindle cells- phyllodes tumor
- Option D- Polymorphism with single or arranged ductal epithelial cells- tubular carcinoma

### 4. Ans. (a) **Nipple adenoma** (Ref: Robbins 9th/pg 1044)

**Lesions arising from various parts of breast:**

Normal	Lesions
<b>Lobules and terminal ducts (Terminal Duct Lobular Unit or TDLU)</b>	<ul style="list-style-type: none"> <li>• Cyst, Adenosis, Multiple papillomatosis</li> <li>• Sclerosing adenomas</li> <li>• Hyperplasia, Atypical hyperplasia</li> <li>• Carcinoma</li> </ul>
<b>Large ducts</b>	<ul style="list-style-type: none"> <li>• Duct ectasia, Single intraductal papilloma</li> <li>• <b>Nipple adenoma</b><sup>Q</sup>, large duct adenoma</li> <li>• Squamous metaplasia of lactiferous ducts</li> <li>• Paget disease<sup>Q</sup></li> </ul>
<b>Intralobular stroma</b>	<ul style="list-style-type: none"> <li>• Fibroadenoma (focal hyperplasia of stroma &amp; epithelial component of TDLU)</li> <li>• Phyllodes tumor</li> </ul>
<b>Interlobular stroma</b>	<ul style="list-style-type: none"> <li>• Fat necrosis</li> <li>• Lipoma</li> <li>• Sarcoma</li> </ul>

### 5. Ans. (d) **Colloid carcinoma** (Ref: Robbins 9th/pg 1057)

### 6. Ans. (c) **Type of DCIS**

(Ref: Sterenberg diagnostic surgical pathology 5th ed p 312)

The Van Nuys Prognostic Index (VNPI) classifies patients with DCIS to guide decisions on the best treatment option. The index uses patient age, tumour size, tumour growth patterns (histological grade) and the amount of healthy tissue surrounding the tumour after removal (resection margin width) to predict the risk of cancer returning.

### 7. Ans. (b) **Poor prognosis**

(Ref: Robbins 9th/pg 1064 Complete review of pathology 2nd ed/ pg 656)

- The given image shows the breast tissue negatively stained for all three i.e ER, PR and Her2 neu.(called Triple negative); it signifies poor prognosis

### 8. Ans. (a) **Lymph Node status**

(Ref: Robbins 9th/pg 1064 Complete review of pathology 2nd ed/ pg 656)

**Axillary LN status is the most important prognostic indicator in breast ca**

### 9. Ans. (b) **Breast**

(Ref: Atlasgeneticsoncology.org/Genes/AKT1)

Hyper-activation of AKT1 has been found associated to several human cancers:

- Thyroid carcinoma
- Breast carcinoma
- Non-small cell lung carcinoma
- Gastric carcinoma

Now coming to question: Somatic mutation E17K occurs in the PH domain of AKT1 in 8% of human breast cancers. This mutation also occurs in 6% of colorectal cancers.

### 10. Ans. (a) **Lobular** (Ref: Robbins 9th/pg 1065; 8th/pg 1085)

Lobular carcinomas are usually bilateral

### 11. Ans. (b) **Lobular carcinoma is common**


(Ref: Robbins 9th/pg 1054)

From 4% to 14% of cases in males are attributed to germline BRCA2 mutations –OPTION A IS TRUE

Structure of the male breast does not have lobules and acini, lobular carcinoma cases are seen infrequently- OPTION B IS FALSE

Most common breast cancer in male is ductal carcinoma NOS

Colloid carcinoma can be seen.



12. Ans. (a, b, e); a. **it has a spectrum of benign to malignant**; b. **Can be rarely seen in children**; e. **Seen in elderly females** (Ref: Robbins 9th/pg 1051)

**It has a spectrum of benign to malignant- true**

Phyllodes tumors occurs in a median age of fifth decade of life

Rare reports of phyllodes tumor in children have been described.

13. Ans. (b) **BRCA 1 mutation is present in most of the cases of breast carcinoma**

(Ref: Robbins 9th/pg 1054; 8th/pg 1077)

- a. Most common gene involved in familial breast cancer-BRCA1(52%) > BRCA1(32%)
- b. **p53 mutation is present in most of the cases of breast carcinoma**
- c. Inherited breast carcinomas make about 3 % of the total cases
- d. p53 mutation increases risk of all cancers

14. Ans. (a) **Mucinous** (Ref: Robbins 9th/pg 1064-65)

- **Histological type:** Invasive ductal carcinoma (no special type; NST) carries poor prognosis.
  - **Tubular, mucinous, lobular, papillary, adenoid cystic has better prognosis**
    - **Metaplastic carcinoma or micro-papillary carcinoma have a poorer prognosis<sup>Q</sup>**
- Remember mucinous > lobular

15. Ans. (a) **Comedo carcinoma** (Ref: Robbins 9th/pg 1057)

16. Ans. (d) **Comedo DCIS** (Ref: 9th/pg 1057; 8th/pg 1080)

**Comedocarcinoma-solid intraductal sheet of cells** with a central area of **necrosis**- undergoes calcification, can rarely present as **palpable mass<sup>Q</sup>**

17. Ans. (a) **Gene profiling**

(Ref: R 9th/pg 1061; 8th/pg 1080)

**Molecular profiling of breast cancer is done on gene profiling<sup>Q</sup>**

**Gene profiling, measures relative quantities of mRNA for every gene, has 5 major patterns of gene expression.**

18. Ans. (c) **Post mastectomy arm** (Ref: Cancer 1948;1:64-81)

- Lymphangiosarcoma is a misnomer because this malignancy seems to arise from blood vessels instead of lymphatic vessels.
- Most commonly, this tumor is a result of **lymphedema induced by radical mastectomy**
- **Stewart-Treves syndrome** is a rare, cutaneous angiosarcoma that develops in long-standing chronic lymphedema.

19. Ans. (a) **Single file pattern** (Ref: R 9th/pg 1065; 8th/pg 1085)

**Lobular carcinoma (invasive) of breast-Histologic hallmark: presence of discohesive infiltrating tumor**

**cells as single cells in single file pattern<sup>Q</sup>, including signet-ring cells<sup>Q</sup> containing intracytoplasmic mucin droplets<sup>Q</sup>**

20. Ans. (c, d, e); c. **Mitotic rate**; d. **Tubule formation**; e. **Nuclear polymorphism** (Ref: Robbins 9th/pg 1064)

- Invasive carcinoma are graded using **Modified Bloom Richardson Score<sup>Q</sup>**
- Carcinomas are scored for **tubule formation, nuclear pleomorphism, and mitotic rate<sup>Q</sup>**
- The points added to divide carcinomas into grade I (well differentiated), grade II (moderately differentiated), and grade III (poorly differentiated) types.

21. Ans. (b, d); b. **Apocrine metaplasia**; d. **Fibroadenoma**

(Ref: Robbins 9th/pg 1051)

Apocrine metaplasia and fibroadenoma do not increase the risk of breast cancer.

All other 3 increase the risk of breast carcinoma.

Pathologic Lesion	Relative Risk (Absolute Life time Risk)
<b>Nonproliferative Breast Changes</b> (Fibrocystic changes)	<b>1 (3%)<sup>Q</sup></b>
<ul style="list-style-type: none"> <li>• Duct ectasia</li> <li>• Cysts, Adenosis, mild hyperplasia</li> <li>• Apocrine change</li> <li>• Fibroadenoma <b>without<sup>Q</sup></b> complex features</li> </ul>	
<b>Proliferative Disease Without Atypia</b>	<b>1.5 to 2 (5% - 7%)<sup>Q</sup></b>
<ul style="list-style-type: none"> <li>• Moderate or florid hyperplasia</li> <li>• Sclerosing adenosis</li> <li>• Papilloma</li> <li>• Complex sclerosing lesion (radial scar)</li> <li>• Fibroadenoma <b>with<sup>Q</sup></b> complex features</li> </ul>	
<b>Proliferative Disease with Atypia</b>	<b>4 to 5 (13% - 17%)<sup>Q</sup></b>
<ul style="list-style-type: none"> <li>• Atypical ductal hyperplasia (ADH)</li> <li>• Atypical lobular hyperplasia (ALH)</li> </ul>	
<b>Carcinoma in Situ</b>	<b>8 to 10 (20% - 30%)<sup>Q</sup></b>
<ul style="list-style-type: none"> <li>• Lobular carcinoma in situ (LCIS)</li> <li>• Ductal carcinoma in situ (DCIS)</li> </ul>	

22. Ans. (a) **Ductal carcinoma** (Ref: Robbins 9th/pg 1057)

- Almost all (>95%) of breast malignancies are adenocarcinomas that first arise in the duct/lobular system as carcinoma in situ
- Distribution of invasive carcinoma, breast
- **Invasive Ductal Carcinoma-70 to 80%**
- **Invasive lobular Carcinoma-10%**
- **Mucinous- 2%**

23. Ans. (b) **Atypical epithelial hyperplasia**

(Ref: Robbins 9th/pg 1051)



[illegible]This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Endocrine System and its Disorders

## Key Points

- » Anterior lobe (adenohypophysis) consists of 80% of pituitary gland
- » MC cause of hyperpituitarism is anterior lobe adenoma
- » Graves disease is the most common cause of endogenous hyperthyroidism & thyrotoxicosis
- » Hürthle cell metaplasia or Oxyphil change are a feature of Hashimoto's Thyroiditis
- » Most common thyroid Cancer is Papillary carcinoma
- » Orphan Annie eye nuclei is a characteristic feature of papillary cell Ca thyroid
- » Integrity of capsule is most important in distinguishing follicular adenomas from follicular Ca
- » Most consistent feature of Diabetic nephropathy is diffuse thickening of basement membrane
- » Earliest manifestation of Diabetic Nephropathy is Microalbuminuria
- » Most common cause of primary adrenal insufficiency in developed countries is Autoimmune adrenalitis while in India, it is Tubercular adrenalitis in India
- » Most common subtype of congenital adrenal hyperplasia is caused by deficiency of the enzyme 21-hydroxylase
- » Primary hyperparathyroidism is the most common manifestation of Multiple Endocrine Neoplasia (MEN)-1
- » Duodenum is the most common site of gastrinomas in individuals with MEN-1

## Key Recent Updates


- » Most common mutation in papillary carcinoma, thyroid is BRAF
- » Wet keratin is characteristic of craniopharyngioma.



## PITUITARY GLAND

Composed of two **morphologically** and **functionally** distinct components:

- **Anterior lobe (adenohypophysis): 80% of gland<sup>Q</sup>**
- **Posterior lobe (neurohypophysis): 20% of the gland;**



### Pituitary Adenomas and Hyperpituitarism

- **Most Common cause of hyperpituitarism: Anterior lobe adenoma<sup>Q</sup> > Pituitary Carcinoma<sup>Q</sup>**
- GH and prolactin are the most common hormones secreted from Adenoma


### Genetic Alterations in Pituitary Tumors

- Gain of function of: GNAS, PRKAR 1A, Cyclin D1, HRAS genes
- Loss of function of: MEN1, CDKN1B (p27), AIP, RB genes

### Classification of Pituitary Adenomas

- Functional (i.e., associated with **hormone excess** and its clinical manifestations)
- **Non-functioning** (i.e., No hormone excess, **only mass effect**)
- **Microadenomas (<1 cm) (most common)<sup>Q</sup>**
- Macroadenomas (>1 cm)
- **Most common pituitary tumor is Adenoma<sup>Q</sup>**
- Most common cause of **hyperpituitarism** is **anterior lobe Microadenoma<sup>Q</sup>**
- Most common type of **microadenoma is Prolactinoma<sup>Q</sup> > GH-adenoma**

- **Distinctive morphologic features of pituitary Adenoma are: Cellular monomorphism & absence of a reticulin network staining<sup>Q</sup>**



Two distinctive morphologic features **cellular monomorphism** and **absence of a reticulin network**



### High Yield Facts

- Hormones secreted by posterior pituitary are: ADH (Vasopressin) & Oxytocin
- Crooke's hyaline change is seen in pituitary in Cushing syndrome. It is due to accumulation of cytokeratin intermediate filament



## Posterior Pituitary Syndromes

- **Diabetes insipidus: ADH deficiency** → Excessive urination (polyuria) due to inability of kidney to reabsorb water
- **Syndrome of inappropriate ADH (SIADH)<sup>Q</sup>: ADH excess** → resorption of excessive amounts of **free water**, resulting in hyponatremia.

## HYPOTHALAMIC SUPRASELLAR TUMORS

- Result in **hyperfunction/ hypofunction** of the pituitary
- Most commonly **gliomas** and **craniopharyngiomas**




### High Yield Facts

#### Craniopharyngiomas

3-4 cm, cause **mass effect<sup>Q</sup>** on optic chiasma or cranial nerves  
**Two distinct histologic variants: (Dystrophic calcification<sup>Q</sup> frequent in both)**

- **Adamantinomatous** type: (**children<sup>Q</sup>**): compact lamellar keratin formation
- **Papillary** type (**adults<sup>Q</sup>**)



Craniopharyngioma

## THYROID GLAND

### Hyperthyroidism and Thyrotoxicosis

- **Thyrotoxicosis: A hypermetabolic state** caused by elevated free T3 & T4<sup>Q</sup>
- **Hyperthyroidism** is hyperfunction of the thyroid gland<sup>Q</sup>

### High Yield Facts

- Thyroid **follicles** are lined by a **cuboidal to low columnar epithelium** filled with **PAS +ve thyroglobulin**
- **Hypothalamus** → **TRH** → **Thyrotrophs** (anterior pituitary)<sup>Q</sup> → **TSH (thyrotropin)**
- **Thyroid follicular epithelial cells** convert **thyroglobulin** into **thyroxine (T4)<sup>Q</sup>** & **triiodothyronine (T3)<sup>Q</sup>**
- **Parafollicular cells**, or **C cells**- synthesize and secrete the hormone **calcitonin<sup>Q</sup>**

### Disorders Associated with Thyrotoxicosis

With Hyperthyroidism	Without Hyperthyroidism
<b>Primary</b> <ul style="list-style-type: none"> <li>• <b>Graves disease<sup>Q</sup></b></li> <li>• <b>Toxic multinodular goiter<sup>Q</sup></b></li> <li>• <b>Toxic adenoma<sup>Q</sup></b></li> <li>• Iodine-induced hyperthyroidism</li> <li>• Neonatal thyrotoxicosis associated with maternal Graves disease</li> </ul> <b>Secondary:</b> TSH-secreting pituitary adenoma	<ul style="list-style-type: none"> <li>• <b>Granulomatous (de Quervain) thyroiditis<sup>Q</sup></b></li> <li>• <b>Subacute lymphocytic thyroiditis (painless)<sup>Q</sup></b></li> <li>• <b>Struma ovarii:</b> Ovarian teratoma with ectopic thyroid<sup>Q</sup></li> <li>• <b>Factitious thyrotoxicosis</b> (exogenous thyroxine intake)</li> </ul>

### Grave's Disease

- **Most common:**
  - **Most common cause of endogenous hyperthyroidism<sup>Q</sup>**
  - **Most common cause of thyrotoxicosis<sup>Q</sup>**
- **Epidemiology:**
  - **Mean age: 20-40 yrs of age<sup>Q</sup>, Female: Male = 10:1<sup>Q</sup>**
- **Genetic basis:**
  - Polymorphisms in immune-function genes like **CTLA4** and **PTPN22** and the **HLA-DR3 allele<sup>Q</sup>**
- **Pathogenesis:**
  - **Autoimmune disorder** characterized by **autoantibodies against TSH receptor:<sup>Q</sup>**
  - **Thyroid-stimulating immunoglobulin (TSI)<sup>Q</sup>** (90% cases)
- **Clinical features (Triad):**
  - **Hyperthyroidism<sup>Q</sup>** with diffuse enlargement of the gland
  - **Infiltrative ophthalmopathy<sup>Q</sup>** with resultant exophthalmos
  - **Localized, infiltrative dermopathy<sup>Q</sup>** (pretibial myxedema)
- **Morphology:**
  - **Symmetrically enlarged<sup>Q</sup>** gland due to **diffuse hypertrophy and hyperplasia<sup>Q</sup>** of **thyroid follicular epithelial cells<sup>Q</sup>**





- **Laboratory findings:**
  - Elevated free T4 and T3 levels and depressed TSH levels<sup>Q</sup>

## Hypothyroidism

Condition caused by a **structural or functional derangement<sup>Q</sup>** that **interferes** with **production of thyroid hormone<sup>Q</sup>**

- **Primary:** Intrinsic abnormality<sup>Q</sup> in the thyroid
- **Secondary:** Pituitary or Hypothalamic disease<sup>Q</sup>



## High Yield Facts

- Serum TSH<sup>Q</sup> level is the **most useful single screening<sup>Q</sup>** test for hyperthyroidism
- Low TSH is confirmed with **measurement of free T4** (increased)
- **Radioactive iodine uptake:**
  - Diffusely increased uptake by the whole gland (Grave's disease)<sup>Q</sup>
  - Increased uptake in a solitary nodule: Toxic adenoma<sup>Q</sup>
  - Decreased uptake: Thyroiditis<sup>Q</sup>

## Thyroiditis

**Inflammation** of the thyroid gland<sup>Q</sup>


### Hashimoto's Thyroiditis

- **Definition:**
  - Autoimmune destruction of thyroid gland → **gradually progressive thyroid failure.**<sup>Q</sup>
- **Most common:**
  - Cause of hypothyroidism in Iodine-sufficient areas of the world<sup>Q</sup>
  - Clinically apparent cause of chronic thyroiditis
- **Epidemiology:**
  - Mean age: 45 - 65 years of age; **Female: Male = 10-20:1<sup>Q</sup>** (Most Common in middle aged female)<sup>Q</sup>
- **Pathogenesis:**
  - Breakdown in self tolerance to thyroid autoantigens
- **Etiology:**
  - Autoimmune: Defect in **regulatory T-cells (Tregs)**
  - Genetic: polymorphisms in **immune regulation-associated genes**
    - Cytotoxic T lymphocyte-associated antigen-4 (CTLA4)<sup>Q</sup>
    - Protein tyrosine phosphatase-22 (PTPN22)<sup>Q</sup>
- **Clinical Features:**
  - Painless **enlargement** of the thyroid, usually associated with hypothyroidism
- **Increased risk for developing:**
  - B-cell Non Hodgkins lymphoma<sup>Q</sup>
  - Autoimmune diseases:
    - Endocrine (type 1 diabetes, autoimmune adrenalitis)<sup>Q</sup>
    - Non-endocrine (SLE, myasthenia gravis, and Sjögren's syndrome)<sup>Q</sup>

## RECENT EXAM<sup>Q</sup>

### Morphological changes in Thyroiditis

- Infiltration of the gland parenchyma by **small lymphocytes and plasma cells<sup>Q</sup>**
- Well-developed **germinal centers**, thyroid follicles are **atrophic<sup>Q</sup>**
- **Hallmark:** Hürthle cell metaplasia or Oxyphil change (cells with abundant eosinophilic, granular cytoplasm)
- **Intact capsule & fibrosis** confined to gland<sup>Q</sup>



Hashimoto's thyroiditis

### Granulomatous Thyroiditis/De Quervain Thyroiditis


- **Epidemiology:**
  - Most common age group: 40 - 50 yrs<sup>Q</sup>; **Female: Male = 4:1<sup>Q</sup>**
- **Etiology:**
  - Viral Infections: **Cox sackie virus<sup>Q</sup>**, mumps<sup>Q</sup>, measles<sup>Q</sup>, Adenovirus
- **Clinical feature:**
  - **Thyroid pain<sup>Q</sup>** and goiter
- **Morphology:**
  - **Scattered follicles** disrupted & replaced by **neutrophils forming microabscesses<sup>Q</sup>**
  - **Multinucleate giant cells** enclose fragments of colloid<sup>Q</sup>

### Subacute Lymphocytic (Painless) Thyroiditis/Painless Thyroiditis<sup>Q</sup>

- **Epidemiology**
  - Most common middle-aged adults ; F>M; **Self limiting<sup>Q</sup>**;
- **Clinical feature**
  - Mild **hyperthyroidism** with **goitrous enlargement<sup>Q</sup>** of the gland
- **Morphology**
  - **Lymphocytic infiltration** with **large germinal centers<sup>Q</sup>**
  - **Patchy disruption** and **collapse<sup>Q</sup>** of thyroid follicles
  - Fibrosis and Hürthle cell metaplasia are **not prominent<sup>Q</sup>**

### Riedel Thyroiditis

- **Characteristic:**
  - **Extensive fibrosis** involving thyroid & contiguous neck structures<sup>Q</sup>



■ **Associated with:**

- Fibrosis in retroperitoneum<sup>Q</sup>
- Systemic autoimmune **IgG4-related disease** (fibrosis & tissue infiltration by plasma cells producing IgG4)<sup>Q</sup>



**High Yield Facts**

- In **hyperthyroidism**, there is **proximal muscle weakness**<sup>Q</sup>
- **Generalized lymphadenopathy** can occur in **Grave's disease**<sup>Q</sup>
- **Most common** type of thyroiditis is **Hashimoto's thyroiditis**<sup>Q</sup>
- **Hyperthyroidism** may be seen initially in **Hashimoto's thyroiditis**<sup>Q</sup>
- Most common type of thyroiditis **postpartum** is **subacute painless thyroiditis**<sup>Q</sup>
- **HLA DR3/5** are associated with **Hashimoto's thyroiditis**<sup>Q</sup>
- **HLA B5** is associated with **De Quervain thyroiditis**<sup>Q</sup>


**Thyroid Carcinoma**

Thyroid Ca	%	Genes Mc mutated	Route of Metastasis
Papillary	MC 85%	BRAF (MC), RET	Lymphatic
Follicular	5–15%	RAS/P13K 10% PAX/PPARG	Hematogenous
Medullary	<5%	RET	Regional : Lymphatic Distant : Hematogenous
Anaplastic	<5%	RAS, P53	Direct & hematogenous


Mc: Most common

**Papillary Carcinoma**

- **Epidemiology:**
  - **Most common**<sup>Q</sup> form of thyroid cancer; most commonly seen at **20–40 yrs** of age
- **Morphology:**
  - **Solitary or multifocal**<sup>Q</sup>, Well **circumscribed** and **encapsulated**<sup>Q</sup>
  - **Infiltrate adjacent parenchyma**<sup>Q</sup> along with areas of fibrosis and calcification
- **Variants of Papillary Ca:**
  - **Follicular variant, Tall-cell variant, Diffuse sclerosing, Papillary Microcarcinoma**



Papillary carcinoma, thyroid. A classical papillary carcinoma shows multiple finger like branching



Orphan Annie Nuclei pseudoinclusions S/o papillary Ca

■ **Microscopic hallmarks:**

- Branching papillae with a fibrovascular stalk covered by cuboidal epithelial cells
- **Nuclear features: Hallmark of diagnosis**
  - **Orphan Annie eye nuclei**<sup>Q</sup> (clear or empty appearing ground-glass appearance),
  - **"Pseudo-inclusions"**<sup>Q</sup> – invaginations in cytoplasm-appearance of **intranuclear**<sup>Q</sup> inclusions or **intranuclear grooves**.<sup>Q</sup>
- **Psammoma bodies**<sup>Q</sup> (Absent in follicular and medullary Ca)
- **Lymphatic invasion** is common<sup>Q</sup>

**Mnemonic**

Clinical criteria that point towards **neoplastic nature** of thyroid nodule ("**SoMe CRY**")


- **Solitary** nodule<sup>Q</sup>
- **Male**<sup>Q</sup>
- **Cold** nodule on radioiodine uptake
- **Radiation**<sup>Q</sup> exposure history
- **Young** age<sup>Q</sup>





## Follicular Carcinoma

- **Epidemiology:**
  - 2<sup>nd</sup> Most common<sup>Q</sup> primary thyroid cancer; MC in mid to older age, Female > Males<sup>Q</sup>
  - More frequent in areas with dietary iodine deficiency<sup>Q</sup>
- **Morphology:**
  - **Gross:** Single well circumscribed or widely infiltrative nodule
- **Clinical Course:**
  - Slowly enlarging painless nodules
  - Most frequently they are cold nodule
  - Vascular (hematogenous) dissemination to bone, lungs, liver<sup>Q</sup>



Follicular carcinoma showing follicles

- Microscopically:
  - Small follicles containing colloid, Hurthle cell or oncocytic variant<sup>Q</sup>: cells with abundant eosinophilic cytoplasm, Capsular &/or vascular invasion is the sign of carcinoma & differentiates follicular adenoma<sup>Q</sup>, Lymphatic spread is uncommon<sup>Q</sup>


## Anaplastic (Undifferentiated) Carcinoma

- **Epidemiology**
  - Most common in older persons, (mean age of 65 years)<sup>Q</sup>
- **Microscopy**
  - Large, pleomorphic giant cells & spindle cells
- **Prognosis**
  - Most aggressive with a mortality rate approaching 100%<sup>Q</sup>

## Medullary Carcinoma

- **Definition:**
  - A Neuroendocrine neoplasm derived from the parafollicular cells,<sup>Q</sup> or C cells, of thyroid
- **Inheritance:**
  - 70% sporadic (old age); 30% familial (1<sup>st</sup> decade of life): Autosomal dominant
  - As a part of MEN syndrome 2A or 2B<sup>Q</sup> or familial medullary thyroid carcinoma, or FMTC<sup>Q</sup>
- **Clinical features:**
  - Sporadic cases: Mass in neck, dysphagia or hoarseness,
  - Paraneoplastic syndromes: diarrhea<sup>Q</sup> (due to VIP), Cushing syndrome<sup>Q</sup> (due to ACTH)

- **Familial cases:** endocrine neoplasms in other organs (e.g., adrenal or parathyroid in MEN)
- **Gross Morphology:**
  - Sporadic tumor present as Solitary nodule ("S for S")
  - Familial cases- are bilateral<sup>Q</sup> and multicentric<sup>Q</sup>
- **Tumor Markers:**
  - Calcitonin, serotonin, ACTH, and vasoactive intestinal peptide (VIP)



Medullary carcinoma thyroid

- **Microscopy:**
  - Polygonal spindle-shaped cells form nests, trabeculae and follicles
  - Acellular amyloid deposits (A cal)<sup>Q</sup> are characteristic feature
  - Electron microscopy: membrane-bound electron-dense granules<sup>Q</sup>
  - Multicentric C-cell hyperplasia in familial cases

## High Yield Facts

- Integrity of capsule is critical in distinguishing follicular adenomas from follicular Ca<sup>Q</sup>
- Follicular Carcinoma of Thyroid cannot be diagnosed by FNAC<sup>Q</sup>
- Most common thyroid Cancer is Papillary carcinoma
- Most common Thyroid Ca post radiation exposure is Papillary CA
- Thyroid Carcinoma with best (excellent) prognosis: Papillary Ca thyroid<sup>Q</sup>
- Thyroid Carcinoma with Worst prognosis: Anaplastic Carcinoma<sup>Q</sup>
- Hypocalcemia<sup>Q</sup> is not a prominent feature of Medullary Ca thyroid, despite presence of raised calcitonin levels
- Carcinoembryonic antigen (CEA) is a useful biomarker in calcitonin-negative Medullary Ca thyroid



## PARATHYROID GLANDS

The four parathyroid glands are composed of two cell types:

- **Chief cells:** Large amounts of **cytoplasmic glycogen** (water clear appearance)<sup>Q</sup>
- **Oxyphil cells:** **Acidophilic cytoplasm**, and are tightly packed with **mitochondria**

### Hyperparathyroidism

#### Primary Hyperparathyroidism

Underlying Parathyroid lesions can be Adenoma (90%) > Primary hyperplasia > Parathyroid carcinoma

Lesion	Frequency	Morphology
Adenoma	85% – 95% <sup>Q</sup> (most common)	<b>Solitary lesions</b> with uniform, polygonal chief cells along with nests of oxyphil cells ( <b>oxyphil adenomas</b> )
<b>Primary hyperplasia</b>	5%–10%	<b>Chief cell hyperplasia</b> with <b>water-clear cells</b> involving all 4 glands
Parathyroid carcinoma	~1%	Circumscribed lesion in a single gland <b>diagnosed by invasion of surrounding tissues</b> and <b>metastasis</b> as the only reliable criteria.

#### Molecular Defects

<b>Sporadic adenomas:</b> <ul style="list-style-type: none"> <li>• <b>Cyclin D1</b> overexpression</li> <li>• <b>MEN1 mutations</b> on chr 11q</li> </ul>	<b>Familial parathyroid adenomas:</b> <ul style="list-style-type: none"> <li>• Multiple Endocrine Neoplasia (MEN syndrome-1 &amp; 2)</li> <li>• <b>Familial hypocalciuric hypercalcemia:</b> (autosomal-dominant)</li> <li>• Loss-of-function mutations in <b>calcium-sensing receptor (CASR)</b> gene</li> </ul>
---	---

#### Clinical Features

Skeletal abnormalities:

- **Osteoporosis:** Cortical bone more severely involved than medullary bone
- **Brown tumors:** Brown color-vascularity, hemorrhage & hemosiderin deposition
- **Osteitis fibrosa cystica** (von Recklinghausen disease of bone)

#### Metastatic Calcification

Nephrolithiasis, nephrocalcinosis, calcific deposits in stomach, lungs, myocardium, blood vessels.

#### Secondary Hyperparathyroidism

- **Definition:**
  - **Chronic hypocalcemia** leading to **compensatory over-activity** of the parathyroid glands.
- **Etiology:**
  - **Renal Failure** (Most Common Cause)<sup>Q</sup>
  - **Inadequate Dietary Intake of Calcium**
  - **Steatorrhea**<sup>Q</sup>
  - **Vitamin D Deficiency**<sup>Q</sup>
- **Clinical Features:**
  - Can cause **calciophylaxis**<sup>Q</sup> (ischemic damage to skin and other organs)
- **Morphology:**
  - Hyperplastic parathyroid glands with **chief cells** (water-clear cells)<sup>Q</sup>

#### Tertiary Hyperparathyroidism

Autonomous activity of gland with **hypercalcemia**.

#### Pseudohypoparathyroidism

**End-organ resistance** to the actions of PTH<sup>Q</sup>.

- Serum **PTH levels are normal or elevated**<sup>Q</sup>
- Some cases have end-organ **resistance to TSH and FSH/LH and PTH**<sup>Q</sup>

### Hypoparathyroidism

#### Etiology

- **Surgical removal of parathyroid:** Most common cause<sup>Q</sup>
- **Genetic causes of hypoparathyroidism**
  - **Autoimmune hypoparathyroidism:** mutations in AIRE gene (AIRE).
  - **Autosomal-dominant hypoparathyroidism:** mutations in CASR gene<sup>Q</sup>
  - **Familial isolated hypoparathyroidism (FIH)**
- **DiGeorge syndrome** : 22q11 deletion syndrome<sup>Q</sup>





## PANCREAS

### Diabetes Mellitus

A Group of metabolic disorders sharing the common features of **hyperglycemia**.

#### Classification of Diabetes Mellitus

**Type-1 diabetes** ( $\beta$ -cell destruction causing absolute insulin deficiency).

**Type-2 diabetes** (combination of  $\beta$ -cell dysfunction and insulin resistance)

#### Genetic defects

- Maturity on set diabetes of the young (MODY).
  - **MODY-1** Hepatocyte nuclear factor 4 $\alpha$  (HNF-4 $\alpha$ )
  - **MODY-2** Glucokinase (GCK)
  - **MODY-3** Hepatocyte nuclear factor 1 $\alpha$  (HNF-1 $\alpha$ )
  - **MODY-4** Pancreatic and duodenal homeobox-1 (PDX-1).
  - **MODY-5** Hepatocyte necrotic factor 1 $\beta$  (HNF-1 $\beta$ ).
  - **MODY-6** Neurogenic differentiation factor 1 $\beta$  (Neuro D<sub>1</sub>)
    - ♦ Neonatal diabetes: Activating mutations in KCNJ 11 and ABCC 8.
    - ♦ Maternally inherited diabetes and deafness (MIDD), Mitochondrial DNA mutations (m. 3243A  $\rightarrow$  G)<sup>Q</sup>

#### Genetic defects in insulin action

- Type A insulin resistance
- Lipotrophic diabetes (PPARG mutation)

#### Exocrine pancreatic defects

- Chronic pancreatitis
- Fibrocalculous pancreatopathy
- Neoplasia
- Cystic fibrosis, Hemochromatosis

#### Infections

- CMV
- Congenital rubella
- Coxsackie B

#### Endocrinopathies

- Acromegaly
- Cushing's syndrome
- Hyperthyroidism
- Pheochromocytoma
- Glucagonoma

#### Gestational diabetes mellitus

- **Drugs**
  - Glucocorticoids
  - Thyroid hormone
  - $\beta$ -adrenergic agonists
  - Thiazides
  - Phenytoin

#### Genetic syndromes associated with diabetes

- Down's syndrome
- Klinefelter's syndrome
- Turner's syndrome
- Prader Willi syndrome

#### Diagnostic Criteria According to ADA and WHO

##### Diabetes

Presence of any 1 or more of the following:

- **Fasting** plasma glucose  $\geq 126$  mg/dL
- **Random** plasma glucose  $\geq 200$  mg/dL
- **2-hour plasma glucose**  $\geq 200$  mg/dL during oral glucose tolerance test (OGTT) with 75 g
- Glycated hemoglobin (**HbA<sub>1c</sub>**) level  $\geq 6.5\%$

##### Impaired Glucose Tolerance (Pre-diabetes)

Presence of any 1 or more of the following:

- **Fasting** plasma glucose between **100 and 125 mg/dL** (impaired fasting glucose)
- **2-hour plasma glucose** between **140 and 199 mg/dL** following a **75-gm glucose OGTT**
- A glycated hemoglobin (**HbA<sub>1c</sub>**) level between **5.7% and 6.4%**.

### Differences between Type 1 DM-(Insulin-dependent Diabetes Mellitus) and Type 2 DM (Non Insulin dependent Diabetes Mellitus)

Features	Type 1 Diabetes Mellitus	Type 2 Diabetes Mellitus
Onset	Childhood <sup>Q</sup> or adolescent	Mostly adults <sup>Q</sup>
Antibodies	Circulating <b>islet auto-antibodies</b> <sup>Q</sup> (anti-insulin <sup>Q</sup> , anti-GAD, anti-ICA 512)	<b>No auto-antibodies</b> <sup>Q</sup>
Complications	<b>Diabetic ketoacidosis</b> <sup>Q</sup> (DKA) in absence of insulin therapy	<b>Non-ketotic hyperosmolar coma</b> > DKA.
Genetics	1. Mostly linked to <b>MHC class I and II genes</b> <sup>Q</sup> . 2. <b>Polymorphism in CTLA4</b> <sup>Q</sup> and <b>PTPN 22</b> <sup>Q</sup> 3. Linked to insulin gene VNTR (variable number of tandem repeats)	1. No HLA-linkage <sup>Q</sup> 2. Related to diabetogenic and obesity related <b>candidates genes</b> (TCF7L2 <sup>Q</sup> , PPARG, FTO)
Pathogenesis	<b>Regulatory T-cell dysfunction</b> <sup>Q</sup> causing loss of self-tolerance to islet cell autoantigens.	1. <b>Insulin resistance</b> <sup>Q</sup> in peripheral tissue. 2. Failure of compensation by beta-cells. 3. Multiple obesity associated factors (non-esterified fatty acid, inflammatory cytokines)

Contd...



Features	Type 1 Diabetes Mellitus	Type 2 Diabetes Mellitus
Pathology	<ol style="list-style-type: none"> <li>1. Insulinitis<sup>Q</sup> (inflammatory infiltrates of T-cells and macrophages)</li> <li>2. <b>Beta-cell depletion</b><sup>Q</sup></li> <li>3. Islet atrophy</li> </ol>	<ol style="list-style-type: none"> <li>1. <b>No insulinitis</b><sup>Q</sup></li> <li>2. <b>Amyloid deposition</b><sup>Q</sup> in islets</li> <li>3. Islet cell hyperplasia (especially in non-diabetic newborns of diabetic mothers)</li> </ol>
Diagnostic methods	Reduced insulin, C peptide ↓	↑ Insulin level, C peptide ↑

## Pathological Changes in Diabetes

### Changes in Pancreas

More common in type 1 than type 2 diabetes.

- **Reduction in the number and size of islets**
- **Leukocytic infiltrates** in the islets (insulinitis) by T lymphocytes
- In type 2 diabetes → **amyloid deposition** within islets
- In **infants of diabetic mothers** → **Increase** in the number and size of islets

### Diabetic Macrovascular Disease

- **Endothelial dysfunction** which predisposes to **atherosclerosis**
- **Hallmark: Accelerated atherosclerosis** of **aorta & large-medium-sized arteries**.<sup>Q</sup>
- **Hyaline arteriosclerosis**: Amorphous hyaline thickening of arteriolar wall → Narrowing of lumen

### Diabetic Microangiopathy

Nephropathy, Retinopathy, Neuropathy

- **Most consistent**<sup>Q</sup> morphologic feature of **Diabetes**: **Diffuse thickening of basement membrane in capillaries of skin, skeletal muscles, Retina, Glomeruli & renal medulla**

### Diabetic Ocular Complications

- Diabetes-induced hyperglycemia can lead to **cataract**<sup>Q</sup>
- **Glaucoma** → **damage to optic nerve**<sup>Q</sup>
- **Retinal vasculopathy**:<sup>Q</sup> Can be non-proliferative or proliferative

### Diabetic Neuropathy

- Can affect **central and peripheral nerves**<sup>Q</sup>
- **Distal symmetrical polyneuropathy**<sup>Q</sup> – **most common pattern**
- **Autonomic nervous system involvement**<sup>Q</sup>

### Diabetic Nephropathy

Explained in chapter 16.



### High Yield Facts

- **Most common type of MODY is MODY3**<sup>Q</sup>
- Myocardial infarction due to atherosclerosis is the most common cause of death in diabetics.<sup>Q</sup>
- Renal failure is second most common cause of death in DM<sup>Q</sup>
- **Dry Gangrene** of lower extremities is 100 times<sup>Q</sup> more common in diabetics
- Most profound histopathological changes of Diabetes are seen in Retina<sup>Q</sup>
- Fundamental lesion of Retinopathy is Neovascularization<sup>Q</sup>
- Earliest manifestation of Diabetic Nephropathy is **Micro-albuminuria**<sup>Q</sup>
- Risk of Nephropathy & ESRD is more in type 1 than type 2 Diabetes<sup>Q</sup>

## Complications of Diabetes

### Acute Metabolic Complications of Diabetes:


- Diabetic ketoacidosis (Type 1 > type 2 DM)
- Hyperosmolar nonketotic hyperglycemia (Type 2 > type 1 DM)
- **Hypoglycemia**<sup>Q</sup> (**most common acute complication**)

### Chronic Complications of Diabetes: According to type of vessels involved

- **Macrovascular disease**: Large and medium sized
  - Accelerated atherosclerosis among diabetics
  - Increased risk of myocardial infarction
  - Stroke
  - Lower extremity ischemia
- **Microvascular disease**: Small vessels
  - Retina- diabetic retinopathy, Macular edema
  - Kidneys- nephropathy
  - Peripheral nerves- neuropathy



## Mechanisms of Chronic Complications in Diabetes



### Pancreatic Neuroendocrine Tumors (Pannets)

- Tumors of the pancreatic islet cells ("islet cell tumors")
- 2% of all pancreatic neoplasms; may be single or multiple and benign or malignant
- May elaborate pancreatic hormones, or may be non-functional
- **Surest criteria for malignancy: metastases, vascular invasion & local infiltration.**<sup>o</sup>
- 60%–90% of pancreatic endocrine neoplasms are malignant.



### High Yield Facts

- **Most common pancreatic endocrine neoplasm is Insulinoma**
- **VIPoma causes WDHA syndrome** (watery diarrhoea, Hypokalemia and Achlorhydria)
- Deposition of amyloid in the extracellular tissue is a characteristic feature of many insulinomas
- **Diarrhea** is the most common presenting symptom of **gastrinoma**



### Latest Update

#### Mutations in PanNETs involve:

- MEN-1, PTEN, TSC2
- Alpha-Thalassemia/mental Retardation syndrome, X-linked (ATRX)
- Death-domain associated protein (DAXX), which helps in telomere maintenance.

### Functional Pancreatic Endocrine Neoplasms

- Insulinoma
- Zollinger-ellison's syndrome (Gastrinoma)
- Multiple endocrine neoplasia (MEN)

#### Insulinoma

Most common pancreatic endocrine neoplasm<sup>o</sup>, Deposition of **amyloid**<sup>o</sup> is a characteristic feature, Focal or diffuse hyperplasia of the islets (nesidioblastosis)<sup>o</sup>

### Mnemonic

#### Causes of Hypoglycaemia<sup>o</sup> (Blood Glucose <60mg/dL) "LIFE"

- **L**iver disease<sup>o</sup>, **I**nsulin administration, **I**nsulinomas<sup>o</sup>, **I**nherited glycogenosis, **F**ibromas/Fibrosarcomas<sup>o</sup>, **E**ctopic insulin production



### High Yield Facts

#### Nesidioblastosis

- Also called **congenital focal/diffuse beta-islet hyperplasia**.
- It is often associated with **hyperinsulinemic hypoglycemia**.
- It occurs due to increase in expression of growth factors **IGF2, IGF1Ra and TGFBR3** in islets.
- **It is associated with:**
  - Beckwith-Weidman syndrome
  - Gastric bypass patients
  - Zollinger-Ellison syndrome

## ADRENAL GLANDS

Paired organs lying above kidneys, which consists of: Cortex and Medulla

**Adrenal cortex** has 3 zones (from outermost to innermost—**G-F-R**)

Zones	Hormone Secreted
Zona Glomerulosa	Mineralocorticoids like Aldosterone
Zona Fasciculata	Glucocorticoids (principally cortisol)
Zona Reticularis	Sex steroids (estrogens and androgens)

**Adrenal medulla** is composed of **Chromaffin (Specialized neural crest/ neuroendocrine)**<sup>o</sup> cells supported by **sustentacular**<sup>o</sup> cells. **Chromaffin cells** → **Epinephrine & Norepinephrine**<sup>o</sup>



## Adrenocortical Insufficiency

Caused by:

- Primary adrenal disease (primary hypoadrenalism) or
- Decreased stimulation of the adrenals due to a ACTH deficiency (secondary hypoadrenalism)

## Adrenocortical Hyperfunction (Hyperadrenalism)

Includes:

Disease	Hormone over-produced
<i>Cushing's syndrome</i>	Cortisol
<i>Hyperaldosteronism</i>	Aldosterone <sup>Q</sup>
<i>Adrenogenital or virilizing syndromes</i>	Androgens <sup>Q</sup>

## Hypercortisolism (Cushing's Syndrome)

Elevated glucocorticoid levels due to exogenous or endogenous causes. Homogeneous and paler cytoplasm of ACTH-producing cells due to intermediate keratin filaments in Pituitary is called: "Crooke hyaline change":



### High Yield Facts

- Most common cause of primary adrenal insufficiency in developed countries: Autoimmune Adrenalitis<sup>Q</sup>
- Most common cause of primary adrenal insufficiency in India: Tubercular Adrenalitis<sup>Q</sup>
- Patients with APS 1 develop antibody against IL-17 & IL-22
- Hyperpigmentation of skin in Addison's disease is caused by POMC<sup>Q</sup> → precursor of ACTH & MSH
- Hypertension is the most common manifestation of primary hyperaldosteronism<sup>Q</sup>
- Plasma renin levels distinguish primary from secondary hyperaldosteronism<sup>Q</sup>
- Most common cause of Cushing's syndrome is exogenous glucocorticoids ("iatrogenic")<sup>Q</sup>
- 'Cushing disease'<sup>Q</sup> refers to ACTH producing pituitary Adenoma
- Characteristic of aldosterone-producing adenomas is "Spironolactone bodies"<sup>Q</sup> (eosinophilic, laminated cytoplasmic inclusions), found after treatment with Spironolactone.

## Adrenal Medulla Neoplasms

Let us first understand what is a Paraganglion System

Paraganglion System		
Definition: Neuroendocrine cells in adrenal medulla & extraadrenal system location <sup>Q</sup>		
Classification: (Based on their anatomic distribution)		
Parasympathetic	Branchiomeric	Close to the major arteries & cranial nerves; e.g. carotid bodies <sup>Q</sup>
	Intravagal	Distributed along the vagus nerve. <sup>Q</sup>
Sympathetic	Aorticosympathetic	Segmental ganglia distributed mainly along the abdominal aorta. e.g: Organs of Zuckerkandl <sup>Q</sup>

## Neoplasms of Adrenal Medulla are of 2 types

- Pheochromocytoma:** Neoplasms of chromaffin cells that release catecholamines
- Neuroblastic tumors:** Neuroblastoma (Refer chapter 8)


### Pheochromocytoma

Familial Syndromes Associated with Pheochromocytoma and extra-adrenal Paragangliomas

Syndrome	Gene	Associated Neoplasms
<i>MEN-2A<sup>Q</sup></i>	<i>RET<sup>Q</sup></i>	Medullary Thyroid Ca <sup>Q</sup> & Parathyroid hyperplasia <sup>Q</sup>
<i>MEN-2B<sup>Q</sup></i>	<i>RET<sup>Q</sup></i>	Medullary thyroid Ca <sup>Q</sup> , Mucocutaneous Ganglioneuromas <sup>Q</sup>
<i>NF1</i>	<i>NF1<sup>Q</sup></i>	Optic nerve glioma <sup>Q</sup>
<i>Von Hippel-Lindau (VHL)</i>	<i>VHL<sup>Q</sup></i>	Renal cell Ca, Hemangioblastoma <sup>Q</sup> , PanNET
<i>Familial paraganglioma</i>	<i>SDHD</i>	Paraganglioma

### Diagnosis of Pheochromocytoma

- Gross Morphology:**
  - Small, circumscribed lesions to large hemorrhagic masses
  - Richly vascularized fibrous trabeculae producing lobular pattern.
  - Incubation of fresh tissue with a potassium dichromate solution turns the tumor a dark brown color (hence the name "chromaffin")<sup>Q</sup>.
- Markers:**
  - Chromogranin<sup>Q</sup> and Synaptophysin<sup>Q</sup> in the chief cells
  - S-100 in peripheral sustentacular cells<sup>Q</sup>
- Definitive diagnosis of malignancy:**
  - Based on the presence of metastases (vascular invasion) & not on histology.
  - Metastasis: Involves regional lymph nodes<sup>Q</sup>, liver<sup>Q</sup>, lung<sup>Q</sup>, and bone<sup>Q</sup>



Zellballen pattern characterized by well defined nests of epithelioid cells & vascular fibrous stroma separating the nests of cells





## Microscopy

- Tumors made of **clusters of polygonal to spindle-shaped chromaffin cells** surrounded by **sustentacular cells**, creating **small nests (zellballen)<sup>Q</sup>**, with rich vascularity.
- "Salt and pepper"<sup>Q</sup>** nuclear chromatin: **characteristic** of neuroendocrine tumors.
- Electron microscopy**: Membrane-bound, **electron-dense secretory<sup>Q</sup>** granules.



## High Yield Facts

### Adrenogenital Syndromes

- Congenital adrenal hyperplasia (CAH)**: Autosomal recessive inheritance<sup>Q</sup>
- Most common subtype is caused by deficiency of the enzyme **21-hydroxylase<sup>Q</sup>**.
- Bilateral hyperplasia of adrenal cortex due to excess ACTH is characteristic<sup>Q</sup>

CAH with:	Female virilization	Male incomplete virilization
<i>Hypertension</i>	11 hydroxylase deficiency <sup>Q</sup>	17 hydroxylase deficiency <sup>Q</sup>
<i>Salt crisis</i>	21 hydroxylase deficiency <sup>Q</sup>	3 β hydroxy steroid dehydrogenase <sup>Q</sup>

## MULTIPLE ENDOCRINE NEOPLASIA (MEN) SYNDROMES

- Definition:**
  - Genetically inherited diseases** resulting in **proliferative lesions (hyperplasia, adenomas, and carcinomas)** of multiple endocrine organs.
- Characteristics:**
  - Younger age** than sporadic tumors.
  - Involve multiple organs **synchronously (at same time)** or **metachronously (at different times)**
  - Tumors are often **multifocal<sup>Q</sup>**, **more aggressive<sup>Q</sup>** and **recurrent<sup>Q</sup>**
  - Preceded** by an asymptomatic stage of **endocrine hyperplasia<sup>Q</sup>**.



## High Yield Facts

- Primary hyperparathyroidism** is the most common manifestation of **MEN-1<sup>Q</sup>**.
- Most frequent anterior pituitary tumor in **MEN-1** is a **prolactinoma<sup>Q</sup>**.
- Most common thyroid Ca in MEN syndrome is **Medullary Ca thyroid**.
- Duodenum<sup>Q</sup>** is the most common site of **gastrinomas** in individuals with **MEN-1**.
- Medullary Thyroid Ca** in **MEN 2B** is **more aggressive<sup>Q</sup>** than in **MEN 2A**.
- All individuals with **RET** mutation are advised **prophylactic thyroidectomy**.

## Multiple Endocrine Neoplasia (MEN) Syndromes

## Mnemonic

MEN 1 (Wermer syndrome)	MEN 2A (Sipple syndrome)	MEN 2B
<b>MEN1 (Chr 11q), encodes <i>Menin</i></b>	<b>RET on Chr 10q</b>	
<b>3 "P" s</b> <b>P</b> arathyroid Hyperplasia/Adenoma (MC) <b>P</b> ituitary : Hyperplasia/Adenoma <b>P</b> ancreas: Hyperplasia/Adenoma/NET Less common manifestations: <b>"CAP"</b> <b>C</b> arcinoid of Foregut <b>A</b> ngiofibroma/Lipoma <b>P</b> heochromocytoma	<b>"HAPPY"</b> <b>H</b> irschsprung disease <b>A</b> myloidosis Cutaneous lichen <b>P</b> heochromocytoma <b>P</b> arathyroid hyperplasia/Adenoma <b>M</b> TC (thYroid Ca) • <b>N</b> euroblastoma • <b>W</b> ilms Tumor • <b>R</b> etinoblastoma	<b>3 Ms</b> <b>P</b> heochromocytoma <b>M</b> TC <b>M</b> ucosal & gastrointestinal neuromas <b>M</b> arfanoid features



R10<sup>th</sup>

## Latest Update

- Mc mutation in papillary carcinoma now is BRAF
- Variants of papillary ca

<b>Follicular variant</b>	Nuclear features of papillary Ca, but follicular architecture More angioinvasive with less lymphatic metastasis
<b>Tall-cell variant</b>	More common in older individuals; Histopath: Tall columnar eosinophilic cells Higher vascular invasion, extra-thyroid extension & distant metastases is common
<b>Diffuse sclerosing</b>	More common on young and children; Extensive, diffuse fibrosis; abundant psammoma bodies, squamous metaplasia, extensive lymphocytes (resembles Hashimoto thyroiditis)-No BRAF mutation
<b>Papillary Microcarcinoma</b>	< 1 cm size; precursors of typical papillary carcinomas




## NEXT Pattern Questions



Q's

1. A 45-year-old patient presented with features of hypothyroidism from 2 to 3 years the histopathology is shown below. Based on histological features what is your diagnosis?



- a. Hashimoto's thyroiditis
- b. Granulomatous thyroiditis
- c. Papillary carcinoma
- d. Rieders thyroiditis


### Ans. (a) Hashimoto's thyroiditis

- Hypothyroidism history with lymphocytic infiltration in gland making germinal follicles. On the right side of the image you can see the oncocytes. This is suggestive of Hashimoto thyroiditis.



Q's

2. A 24-year-old patient presented with thyroid nodule, and ultrasound shows calcification. The histological features are shown in the diagram below, so what is your diagnosis?



- a. Follicular carcinoma
- b. Medullary carcinoma
- c. Papillary carcinoma
- d. Anaplastic carcinoma

### Ans. (c) Papillary carcinoma

- Thyroid nodule with calcification on a sonogram may be secondary to dystrophic calcification seen in papillary thyroid carcinoma. The histopath shows papillary pattern and annie-eye nuclei which is the hallmark of the same.




Q's

3. A patient presented with neck swelling. Cytology showed parafollicular cells along with clusters of plasmacytoid and scant amyloid. What investigation should be done to follow up the patient?

- a. Calcitonin
- b. TSH level
- c. Anti TPO antibody
- d. TRH


### Ans. (a) Calcitonin

- Parafollicular cells along with clusters of plasmacytoid and scant amyloid is suggestive of medullary carcinoma thyroid gland. Serum calcitonin levels is used as a tumor marker in this case.




## Image-Based Questions

1. A 25-year-old female presented with features of weight gain and loss of appetite and easy fatigue. On examination a swelling was noticed in anterior aspect of neck which moved with deglutination. Biopsy performed from neck revealed the following. What is your diagnosis?




- Reidel's thyroiditis
- Hashimoto thyroiditis
- Follicular carcinoma thyroid
- Graves disease

3. A 40-year-old male presented with a thyroid swelling and dysphagia. He gave history of on and off watery diarrhea. Biopsy of the lesion is shown. What is your diagnosis?




- Follicular Ca thyroid
- Papillary Ca thyroid
- Medullary Ca thyroid
- Anaplastic cell Ca Thyroid

2. A 24-year-old male presented with a swelling on anterior aspect of neck. On examination the swelling was firm and moved with deglutination. Biopsy from the lesion has been shown. What is the typical finding and diagnosis?




- Orphan Annie eye nuclei; Follicular Ca thyroid
- Orphan Annie eye nuclei; Papillary Ca thyroid
- Hurthle cell change; Follicular Ca thyroid
- Hurthle cell change; Papillary Ca thyroid

4. Biopsy from resected mass has been shown. What is the finding and diagnosis?




- Zellballen in RCC
- Zellballen in pheochromocytoma
- Follicular Ca in renal metastasis
- Homer Wright rosettes in neuroblastoma






5. Kidney biopsy findings and ophthalmoscopy findings from a patient of Type II diabetes of 25 years has been shown below. Identify the findings and pathology?




- Kimmelsteil Wilson lesion, diabetic non proliferative retinopathy due to macroangiopathy
- Diffuse glomerulosclerosis, diabetic proliferative retinopathy due to macroangiopathy
- Kimmelsteil Wilson lesion, diabetic non proliferative retinopathy due to microangiopathy
- Diffuse glomerulosclerosis, diabetic proliferative retinopathy due to microangiopathy

6. Biopsy from parathyroid gland from a 55-year old male who presented to nephrology department of AIIMS has been shown below. He is a known case of chronic kidney disease with hypertension and type II diabetes. He has recently developed bone pain, lesions in skin and recurrent stones in kidney




- Water clear cells in parathyroid hyperplasia
- Water clear cells in parathyroid Ca
- TB parathyroid
- Parathyroid Necrosis

7. A 23-year-old female of Asian Indian origin was admitted to the orthopedic emergency department with pain in her left shoulder region, right knee, and left thigh following a trivial trauma. On physical examination of the patient, tenderness was found to be present in the left shoulder, right knee, and left thigh. On laboratory analysis, serum calcium level was 11.4 mg/dl (normal 8.4-10.7 mg/dl), serum alkaline phosphatase level was 780 IU/l (normal 50-240 IU/l), serum parathyroid hormone level was 456 pg/ml (normal 7-53 pg/ml), vitamin D3 (1,25-dihydroxy cholecalciferol) was 32 pg/ml (normal 25-45 pg/ml). Biopsy done from the lesion has been shown below. What is your diagnosis?



- Bone necrosis in secondary metastasis
- Bone fibrosis in tuberculosis
- Brown tumor in primary hyperparathyroidism
- Brown tumor in osteosarcoma





## Answers of Image-Based Questions

1. Ans. (b) **Hashimoto's thyroiditis**

- The history given suggests a thyroid swelling. The given biopsy from the same shows thyroid parenchyma containing a dense lymphocytic infiltrate with **germinal centers**. Residual thyroid follicles are lined by deeply **eosinophilic Hürthle cells**. This suggests Hashimoto's thyroiditis.

2. Ans. (b) **Orphan Annie eye nuclei; papillary Ca thyroid**

- High power shows nuclei of papillary carcinoma cells contain finely dispersed chromatin, which imparts an **optically clear** or **empty** appearance, giving rise to the **ground glass** or **Orphan Annie eye nuclei**. In addition, invaginations of the cytoplasm may give the appearance of intranuclear inclusions ("pseudoinclusions") or intranuclear grooves. **The diagnosis of papillary carcinoma can be made based on these nuclear features**, even in the absence of papillary architecture.

3. Ans. (c) **Medullary Ca thyroid**

- Diarrhea in thyroid tumor can be a paraneoplastic syndromes due to VIP in a case of medullary Ca thyroid. Histology demonstrates abundant **deposition of amyloid**, visible here as homogeneous extracellular material, derived from calcitonin molecules secreted by the neoplastic cells.

4. Ans. (b) **Zellballen in pheochromocytoma**

- The tumor is enclosed within an attenuated cortex and demonstrates areas of hemorrhage. The histology of the same shows characteristic **nests of cells ("Zellballen")** with abundant cytoplasm

5. Ans. (c) **Kimmelsteil Wilson lesion, Diabetic non proliferative retinopathy due to microangiopathy.**

- Given kidney biopsy shows spherical, laminated, **nodules of matrix** at the **periphery of glomerulus** which are PAS-positive referred to as Nodular or Inter-capillary Glomerulosclerosis or Kimmelstiel-Wilson disease. Remember retinopathy is a microangiopathy and not macroangiopathy.

6. Ans. (a) **Water clear cells in parathyroid hyperplasia.**

- The given histology shows abundant **optically clear cells** of variable size (hyperplasia and hypertrophy), with spherical clear vacuoles surrounded by thin eosinophilic material; basal nuclei, compact or alveolar patterns.

7. Ans. (c) **Brown tumor in primary hyperparathyroidism**

- Given histology shows brown **colour-vascularity, haemorrhage** & hemosiderin deposition in a case of primary hyperparathyroidism (Increased PTH hormone levels, Calcium levels)




## Multiple Choice Questions

### PITUITARY

- GNAS mutation is associated with malignancy of which cells?** (Recent Question 2016)  
a. Lactotroph                      b. Somatotroph  
c. Thyrotroph                    d. None
- Human chorionic thyrotropin is secreted from:** (Recent Question 2015)  
a. Placenta                        b. Pituitary  
c. Hypothalamus                d. Thyroid
- Which of these organs are not affected in autoimmune polyglandular syndrome type 2?** (Recent Question 2015)  
a. Parathyroid                    b. Thyroid  
c. Adrenal                         d. Pancreas
- Proopiomelanocortin is not released from?** (Recent Question 2015)  
a. Hypothalamus                b. Liver  
c. Lung                             d. Adrenal gland
- Posterior pituitary secretes-** (Recent Question 2014)  
a. GH                                b. TSH  
c. ADH                              d. FSH
- Findings of SIADH includes:** (PGI May 12)  
a. ↑ Urine Na<sup>+</sup>                      b. ↑ S. Na<sup>+</sup>  
c. ↑ Urine osmolality            d. ↑ Serum osmolality  
e. Postural hypotension
- A patient presents with Endocrinopathy, fibrous dysplasia of bone and hyperpigmentation. Diagnosis is?** (JIPMER 2011)  
a. McCune Albright syndrome  
b. Addison's disease  
c. Alagille syndrome            d. Lynch syndrome

### THYROIDITIS

- 25-year-old female presented with swelling in front of neck. TSH levels were elevated. Biopsy showed lymphocytic infiltration and Hurthle cells. Which of the following is the possible diagnosis?** (Recent Pattern Question 2020)




- Graves' disease
  - Hashimoto's thyroiditis
  - Medullary carcinoma thyroid
  - Papillary carcinoma thyroid
- Autoimmune thyroiditis is associated with all except-** (Recent Question 2014)  
a. DM                                b. Myasthenia gravis  
c. SLE                                d. Psoriasis

- Patients with Hashimoto's thyroiditis are at increased risk of developing:** (MH 10)  
a. Papillary carcinoma  
b. Follicular carcinoma  
c. T-cell lymphoma  
d. B-cell Lymphoma

### THYROID CARCINOMA

- Patient came with swelling in midline of neck measuring 2 cm in size. Histopathological examination showed Orphan Annie-eye nuclei. What is the most likely diagnosis?** (Recent Pattern Question 2020)  
a. Medullary carcinoma  
b. Papillary carcinoma thyroid  
c. Toxic nodular goiter  
d. Follicular thyroid carcinoma
- Risk factors for solitary thyroid to be malignant?** (PGI Nov 2018)  
a. Male sex                        b. Middle age  
c. Elderly age                      d. Iodine sufficiency
- Papillary carcinoma of thyroid features?** (PGI Nov 2018)  
a. Haematological spread is early  
b. Most commonly occurring thyroid carcinoma  
c. Capsular invasion is characteristic sign  
d. Shows worse prognosis than follicular type
- 40-year-old female presented with neck swelling. Gross and histology is shown below. What is your diagnosis?** (Recent exam 2018)



- Medullary carcinoma thyroid
  - Hashimoto's thyroiditis
  - Anaplastic carcinoma
  - Follicular carcinoma
- A patient presented with neck swelling. Cytology showed showed parafollicular cells along with clusters of plasmacytoid and few spindle shaped cells. What investigation should be done to follow up the patient?**  
a. Calcitonin                        b. TSH level (JIPMER 2017)  
c. Anti TPO antibody            d. TRH
  - All the following condition can be diagnosed by FNAC except?** (AIIMS Nov 2016)  
a. Follicular Ca thyroid            b. Anaplastic Carcinoma  
c. Papillary carcinoma            d. Medullary carcinoma
  - Lymphatic spread most commonly seen in which type of thyroid carcinoma?** (AIIMS May 2015)  
a. Papillary                        b. Medullary  
c. Follicular                        d. Lymphoma



18. Which one of the following variants of papillary carcinoma thyroid occurs in younger individuals including children with lymphonodal metastases in almost all cases and morphologically simulates Hashimoto thyroiditis is: (AP 2012)
- Tall-cell variant
  - Follicular variant
  - Diffuse sclerosing variant
  - Oncocytic variant
19. Thyroglossal cyst is associated with which type of thyroid Ca? (Recent Question 2015)
- Papillary
  - Medullary
  - Anaplastic
  - Lymphoma
20. About Reidel thyroiditis, which is true? (PGI Nov 2015)
- Extensive fibrosis involving thyroid
  - Fibrosis in retroperitoneum
  - IgG4-related disease
  - Occurs in middle-aged women
  - Painful gland
21. Hematogenous route of metastasis is seen in which type of thyroid carcinoma? (Recent Question 2016)
- Papillary
  - Medullary
  - Anaplastic
  - Follicular
22. Hurthle cell carcinoma is a variant of - (Recent Question 2014)
- Medullary carcinoma
  - Papillary carcinoma
  - Follicular carcinoma
  - Anaplastic carcinoma
23. Medullary ca of thyroid is associated with increase in: (Recent Question 2014)
- Calcitonin
  - Thyroglobulin
  - T3
  - T4
24. Which thyroid carcinoma has amyloid deposition: (Recent Question 2014, MH 11)
- Anaplastic
  - Follicular
  - Medullary
  - Papillary
25. True about Psammoma bodies are all except: (Recent Question 2013)
- Seen in meningioma
  - Concentric whorled appearance
  - Dystrophic calcification
  - Seen in teratoma
26. Most common Thyroid ca post radiation exposure: (Recent Question 2014)
- Papillary CA
  - Medullary CA
  - Follicular CA
  - Anaplastic CA
27. Orphan annie-eye nuclei appearance is characteristic of (Recent Question 2014, DNB Aug 12) (WB PG 2016)
- Papillary carcinoma thyroid
  - Carcinoma pituitary
  - Follicular ca thyroid
  - Medullary ca thyroid
28. Psammoma bodies can be seen in the following except? (AI 11)
- Follicular carcinoma of thyroid
  - Papillary carcinoma of thyroid
  - Meningioma
  - Serous cyadenoma of ovary
29. Most common thyroid Cancer is - (AP PGME 11, AI 00)
- Papillary carcinoma
  - Follicular carcinoma
  - Medullary carcinoma
  - Anaplastic carcinoma

## PARATHYROID GLAND

30. Active form of Vit D? (Recent Question 2015)
- $1,25(\text{OH})_2\text{Vit D}_3$
  - $25\text{ OH Vit D}_3$
  - $\text{Vit D}_3$
  - $\text{Vit D}_2$
31. The cut off for serum Calcium level below which it is called Hypocalcemia is: (Recent Question 2015)
- 6 mg/dl
  - 7 mg/dl
  - 8 mg/dl
  - 9 mg/dl
32. Diagnostic feature of parathyroid carcinoma is- (Recent Question 2014)
- Cytology
  - Metastasis
  - Clinical features
  - All
33. Most common cause of primary hyperparathyroidism is: (Recent Question 2014)
- Adenoma
  - Hyperplasia
  - Hypertrophy
  - Carcinoma
34. Gs-alpha mutation may lead to? (Recent Question 2013)
- Mccune Albright syndrome
  - Pseudohypoparathyroidism
  - Pituitary adenomas
  - All of the above
35. Brown tumor of bone is seen in - (AI 11)
- Hyperparathyroidism
  - Hypoparathyroidism
  - Hypo-thyroidism
  - Hyperthyroidism

## PANCREAS

36. Whipple's triad is diagnostic of: (AP PGME 2015)
- Gastrinoma
  - Insulinoma
  - somatostatinoma
  - Glucagonoma
37. Which of these is the most common cause for insulin resistance? (Recent Question 2015)
- Obesity
  - Post receptor defects
  - Liver dysfunction
  - Pancreatic dysfunction
38. Insulin resistance in liver disease is due to: (AIIMS May 2012)
- Decreased insulin release
  - Steatosis
  - Hepatocyte dysfunction
  - Decreased 'C' peptide level
39. True about neuroendocrine tumors of pancreas is/are? (PGI May 2011)
- Insulinoma is most common neuroendocrine tumors of pancreas
  - VIPoma causes diarrhea
  - Diarrhea is most common symptom of gastrinoma
  - Somatostatinoma causes gall stone formation
  - Gastrinoma has high chance of malignancy
40. Nesidioblastoma is due to hyperplasia of- (PGI Dec 2011)
- Alpha cell
  - Beta cell
  - Acinus
  - D cells

## DIABETES

41. 70 M presented to AIIMS OPD with fatigue. Fasting sugar was 110 mg%, PP was 180 mg%, HbA1c was 6.1 %. What is your diagnosis? (Recent Question 2016)
- Prediabetes
  - Stress induced
  - Normal
  - Diabetes



- 42. For diagnosis of DM, fasting blood glucose level should be more than?** (Recent Question 2014)  
 a. 126 mg/dl                      b. 140 mg/dl  
 c. 100 mg/dl                     d. 200 mg/dl
- 43. Which of the following drugs does not give rise to hyperglycemia?** (Recent Question 2015)  
 a. Thiazides                      b. Phenytoin  
 c. Chloroquine                  d. Prednisolone
- 44. Which is the most common acute complication of diabetes?** (Recent Question 2015)  
 a. Diabetic ketoacidosis  
 b. Hyperosmolar nonketotic hyperglycemia  
 c. Hypoglycemia  
 d. Stroke
- 45. Mechanisms responsible for chronic complications of Diabetes include all of the following except:** (Recent Question 2015)  
 a. Non-enzymatic Glycosylation  
 b. Protein Kinase C activation  
 c. Disturbances in Polyol Pathways  
 d. Chronic Inflammation
- 46. Most commonly seen feature in kidney biopsy of a patient with Diabetic Nephropathy is:** (Recent Question 2014)  
 a. Diffuse Mesangial Sclerosis  
 b. Diffuse glomerulosclerosis  
 c. Nodular glomerulosclerosis  
 d. Fibrin caps
- 47. Which of the following is not an example of Diabetic Microangiopathy?** (Recent Question 2014)  
 a. Nephropathy                  b. Stroke  
 c. Retinopathy                  d. Neuropathy
- 48. All of the following are ocular complications of Diabetes except:** (Recent Question 2014)  
 a. Cataract                      b. Corneal opacity  
 c. Proliferative Retinopathy  
 d. Glaucoma
- 49. Most common type of MODY is:** (Recent Question 2013)  
 a. MODY 1                      b. MODY 2  
 c. MODY 3                      d. MODY 4
- 50. All of the following statement about Type 1 Diabetes are true except?** (PGI Dec 11)  
 a. Family history is present in 90% cases  
 b. Dependent on Insulin to prevent DKA  
 c. Type of onset is usually predictable  
 d. Autoimmune destruction of  $\beta$ -cells  
 e. Often occurs in children

### PHEOCHROMOCYTOMA

- 51. True about pheochromocytoma:** (PGI May 2019)  
 a. Before surgery alpha blockers are given  
 b. Always forms part of MEN1 syndrome  
 c. Mostly malignant  
 d. Mostly familial  
 e. Associated with von Hippel-Lindau disease
- 52. All of the following are PNET Tumors except?** (PGI Nov 2016)  
 a. Rhabdomyosarcoma          b. Osteosarcoma  
 c. Ewings Sa                      d. Medulloepithelioma  
 e. Retinoblastoma

- 53. Sustentacular cells of pheochromocytoma are positive for which of the following markers?** (Recent Question 2016-17)  
 a. S-100                          b. Cytokeratin  
 c. Vimentin                      d. Desmin
- 54. Neuroendocrine tumors are positive for?** (Recent Question 2016-17)  
 a. Synaptophysin                b. Langerin  
 c. CD1 a                          d. CD 5a
- 55. Zellballen pattern is found in histology of which of the following condition?** (Recent Question 2016-17)  
 a. Neuroblastoma                b. Paraganglioma  
 c. Ewings Sarcoma                d. RCC
- 56. Carcinoid tumours commonly arise from:** (Recent Question 2016-17)  
 a. G. cells in pancreas  
 b. Argentaffin cells of small intestine  
 c. Pancreatic endocrine tumour  
 d. Colon polyps
- 57. Which of the following is paraganglioma:** (Recent Question 2016-17)  
 a. Adrenal Pheochromocytoma  
 b. Extra-adrenal Pheochromocytoma  
 c. Carotid body tumour  
 d. Carcinoid tumour              e. Glomus tympanicum
- 58. Pheochromocytomas has been associated with 'the rule of 10's. The following statements are true about Pheochromocytoma except?** (Recent Question 2015)  
 a. 10% are bilateral              b. 10% extra adrenal  
 c. 10% inherited                  d. 10% malignant
- 59. Pheochromocytoma is a tumour of** (MH PG 2014)  
 a. Parathyroid                    b. Adrenal medulla  
 c. Adrenal cortex                  d. Pituitary
- 60. Which of the following are true about pheochromocytoma?** (PGI May 2014)  
 a. Synthesizes epinephrine & norepinephrine  
 b. Increased urinary 5HIAA  
 c. Diagnosed by urinary catecholamine metabolites  
 d. Tumor of adrenal cortex and sympathetic ganglion  
 e. Tumor of adrenal medulla and sympathetic ganglion
- 61. Tumor that follows rule of 10 is** (Recent Question 2013, 2014)  
 a. Pheochromocytoma          b. Oncocytoma  
 c. Lymphoma                      d. Renal cell carcinoma
- 62. Which of the following is most reliable feature of malignant transformation of pheochromocytoma -**  
 a. Involvement of lymph nodes (Recent Question 2013)  
 b. Capsular invasion  
 c. Vascular invasion  
 d. Pleomorphism
- 63. Glomus Cells are found in -** (Recent Question 2013)  
 a. Carotid body Tumour          b. Thyroid carcinoma  
 c. Livercarcinoma                  d. None
- 64. Zellballen pattern is found in -** (PGI Nov 11)  
 a. Pheochromocytoma  
 b. Paraganglioma  
 c. Acoustic neuroma  
 d. Transitional renal cell carcinoma  
 e. Schwannoma





## MEN

- 65. Most common site of gastrinoma in MEN 1 is:** (Recent Pattern Question 2020)
- Stomach
  - Jejunum
  - Duodenum
  - Appendix
- 66. Which is found more in MEN2B than MEN2A?** (PGI Nov 2018)
- Medullary carcinoma thyroid
  - Hyperparathyroidism
  - Pheochromocytoma
  - Marfanoid features
  - Mucosal neuromas
- 67. All are the features of MEN-1 except:?**
- Pancreatic neuroendocrine tumour (AIIMS Nov 2016)
  - Midgut carcinoid
  - Posterior pituitary tumour
  - Parathyroid adenoma
- 68. What is most probable diagnosis of a patient who has prolactinoma, parathyroid hyperplasia and family history of renal stones?** (Recent Question 2016-17)
- MEN-I
  - MEN-II
  - NF1
  - Li fraumeni syndrome
- 69. Which of the following is not associated with a mutation in RET gene?** (Recent Question 2015)
- Leukemia
  - MEN2A
  - Hirschsprung's disease
  - MEN1

- 70. Most common neuroendocrine tumor in MEN -1 is?** (Recent Question 2016)
- Insulinoma
  - Gastrinoma
  - Glucagonoma
  - VIPoma
- 71. Which neuroendocrine tumor causes biliary sclerosis?** (Recent Question 2016)
- Somatostatinoma
  - Gastrinoma
  - Insulinoma
  - Glucagonoma
- 72. Commonest thyroid tumor in MEN (multiple endocrine neoplasia) is:** (Recent Question 2014)
- Follicular
  - Papillary
  - Anaplastic
  - Medullary
- 73. Which of the following is not involved in MEN type IIA-** (Recent Question 2014)
- Parathyroid
  - Adrenal
  - Thyroid
  - Pituitary
- 74. Wermer syndrome is -** (Recent Question 2014)
- MEN I
  - MEN IIA
  - MEN-IIB
  - APS
- 75. MEN type I includes tumors of all except -** (Recent Question 2014)
- Parathyroid
  - Pituitary
  - Pancreas
  - Medullary carcinoma of thyroid
- 76. Which of the following is not true about medullary carcinoma of thyroid?** (MAHA 10)
- Origin is from C cells of thyroid
  - Component of MEN-1
  - Multicentric in origin
  - Amyloid deposition



## Answers with Explanations

- 1. Ans. (b) Somatotrophs** (Ref: Robbins 9th/pg)
- The mutation of guanine nucleotide-activating alpha subunit (GNAS) gene is the somatic mutation related to the McCune-Albright syndrome.
- GNAS mutation is also detected in about 30% to 40% of sporadic growth hormone (GH) secreting tumors.
- 2. Ans. (a) Placenta** (Ref: Harrison's 18th/chapter 341)
- TSH secreted by placenta is called Human chorionic thyrotropin.
- 3. Ans. (c) Adrenal**
- (Ref: Robbins 9th/pg 1130; 8th/pg 1155-1156, Harrison 18th/chapter 318)

### Autoimmune polyglandular syndrome

Disease	Gene	Characteristics
APS-1	AIRE (Chr 21q)	Autoimmune Poly Endocrinopathy, Candidiasis, and Ectodermal Dystrophy <sup>a</sup> (APECED), Hypoparathyroidism, rarely lymphomas
APS-2	HLA DR3, CTLA-4	Hypothyroidism, hyperthyroidism, premature ovarian failure, vitiligo, type 1 diabetes mellitus, pernicious anemia

- 4. Ans. (a) Hypothalamus**
- (Ref: Harrison 18th/chapter 339 Robbins 9th/pg1074; 8th/pg 1098)
- The main source of POMC is pituitary gland
  - Other sources of POMC are Adrenal, gut, reproductive tract, placenta, leukocytes, spleen, lung, liver, thyroid, heart, skin & brain
- 5. Ans. (c) ADH** (Ref: Robbins 9th/pg1074; 8th/pg 1098)
- Posterior pituitary
- Produces two hormones:
- Arginine vasopressin (AVP), also known as antidiuretic hormone (ADH)
  - Oxytocin
- 6. Ans. (a) ↑ Urine Na<sup>+</sup>; (c) ↑ Urine osmolality**
- (Ref: Harrison 18th/chapter 100 Robbins 9th/pg 1081-1082; 8th/pg 1106)
- In syndrome of inappropriate ADH (SIADH):<sup>o</sup>
- ADH excess → resorption of excessive amounts of free water
  - Excessive retention of water expands extracellular & intracellular volume, increases glomerular filtration



and atrial natriuretic hormone, suppresses plasma renin activity, and **increases urinary sodium excretion**.

- This **natriuresis reduces total body sodium**, resulting in **hyponatremia**.
- **Reduced serum osmolality** occur in the setting of an inappropriately **normal or increased urine osmolality**

#### 7. Ans. (a) **McCune Albright syndrome**

(Ref: Robbins 9th/pg 1206-1207)

##### McCune-Albright syndrome

<b>Genetics</b>	It is due to <b>GNAS mutation</b> →continuous activation of stimulatory <b>G protein</b>
<b>Patho-physiology</b>	<b>Increased production of hormones</b> by glands regulated by the G protein system
<b>Clinical features</b>	Precocious puberty, Polyostotic fibrous dysplasia, unilateral Café au lait spots

#### 8. Ans. (b) **Hashimoto's thyroiditis**

(Ref: Robbins 9th/pg 1086)

#### 9. Ans. (d) **Psoriasis** (Ref: Robbins 9th/pg 1086-1087)

**Patients with (autoimmune) Hashimoto thyroiditis are at increased risk for developing:**

- **B-cell Non Hodgkins lymphoma<sup>o</sup>**
- **Autoimmune diseases:**
  - Endocrine (**type 1 diabetes, autoimmune adrenalitis**)<sup>o</sup>
  - Non-endocrine (**SLE, myasthenia gravis, and Sjögren syndrome**)

#### 10. Ans. (d) **B-cell Lymphoma**

(Ref: Robbins 9th/pg 1086-1087)

#### 11. Ans. (b) **Papillary carcinoma thyroid** (Ref: R 9<sup>th</sup> pg 1086)

#### 12. Ans. (c, d); **c. elderly age; d. Iodine sufficiency**

#### 13. Ans. (b) **Most commonly occurring thyroid carcinoma**

#### 14. Ans. (a) **Medullary carcinoma thyroid**

(Ref: Robbins 9th ed p 1099)

Sporadic medullary thyroid carcinomas present as a solitary nodule. In contrast, bilaterality and multicentricity are common in familial cases. Larger lesions often contain areas of necrosis and hemorrhage and may extend through the capsule of the thyroid. The tumor tissue is firm, pale gray to tan, and infiltrative.

Microscopically, medullary carcinomas are composed of polygonal to spindle-shaped cells, which may form nests, trabeculae, and even follicles. Small, more anaplastic cells are present in some tumors and may be the predominant cell type. Acellular amyloid deposits derived from calcitonin polypeptides are present in the stroma in many cases

#### 15. Ans. (a) **Calcitonin**

Cytology showed showed parafollicular cells along with clusters of plasmacytoid and few spindle shaped cells are suggestive of medullary ca thyroid and so should be followed up by its tumor marker calcitonin.

#### 16. Ans. (c) **Papillary carcinoma** (Ref: R 9th ed. Pg. 631)

The hallmark of all follicular adenomas is the presence of an intact, well-formed capsule encircling the tumor. **Careful evaluation of the integrity of the capsule is therefore critical in distinguishing follicular adenomas from follicular carcinomas**, which demonstrate capsular and/or vascular invasion. This can be demonstrated by biopsy of the gland and not FNAC which will demonstrate only the morphology of the tumor cells and not the entire gland histology

#### 17. Ans. (a) **Papillary** (Ref: Robbins 9th/pg 1097-1098)

Thyroid carcinoma type	Route of spread
Papillary	Lymphatic
Follicular	Hematogenous
Medullary	Regional lymphatic spread and hematogeneous routes to distant sites
Anaplastic	Direct

#### 18. Ans. (c) **Diffuse sclerosing variant**

(Ref: Robbins 9th/pg 1097-1098)

##### Diffuse sclerosing variant of papillary carcinoma

- It occurs in younger individuals, including children.
- The tumor has a prominent papillary growth pattern intermixed with solid areas containing nests of squamous metaplasia.
- There is extensive, diffuse fibrosis throughout the thyroid gland, often associated with a prominent lymphocytic infiltrate, simulating Hashimoto thyroiditis.
- Lymph node metastases are present in almost all cases.
- Diffuse sclerosing variant carcinomas lack BRAF mutations, but RET/PTCtranslocations are found in approximately half the cases.

#### 19. Ans. (a) **Papillary** (Ref: Robbins 9th/pg 1097-1098)

**Papillary carcinoma accounts for 80% of cases of thyroglossal duct carcinomas**, with the rest being squamous cell carcinoma


#### 20. Ans. (a, b, c, d) **a. Extensive fibrosis involving thyroid; b. Fibrosis in retroperitoneum; c. IgG4-related disease; d. Occurs in middle-aged women**

(Ref: R 9th/pg 1097-1098)

##### Riedel Thyroiditis

It is a rare disorder that typically occurs in middle-aged women.

It presents with an insidious, painless goiter with local symptoms due to compression of the esophagus, trachea, neck veins, or recurrent laryngeal nerves.



Dense fibrosis disrupts normal gland architecture and can extend outside the thyroid capsule.

Despite these extensive histologic changes, thyroid dysfunction is uncommon.

**Fibrosis of retroperitoneum can also occur**

Associated with Systemic autoimmune **IgG4-related disease** (fibrosis & tissue infiltration by plasma cells producing IgG4)

21. Ans. (d) **Follicular** (Ref: Robbins 9th/pg 1097-1098)

22. Ans. (c) **Follicular carcinoma** (Ref: R 9th/pg 1097-1098)

**Hurthle cell or oncocytic variant of follicular carcinoma** has characteristic cells with abundant eosinophilic cytoplasm.

**Other variants of Papillary Ca are:**

<b>Follicular variant</b>	Nuclear features of papillary Ca, but follicular architecture More angioinvasive with less lymphatic metastasis
<b>Tall-cell variant</b>	More common in older individuals; Histopath: Tall columnar eosinophilic cells Higher vascular invasion, extra-thyroid extension and distant metastases
<b>Diffuse sclerosing</b>	More common on young and children; Extensive, diffuse fibrosis;
<b>Papillary Microcarcinoma</b>	< 1 cm size; precursors of typical papillary carcinomas

23. Ans. (a) **Calcitonin** (Ref: Robbins 9th/pg 1099-1100)

- In **Medullary Carcinoma Thyroid**, serum **Calcitonin** is elevated.
- Elevated serum calcitonin also provides a marker of residual or recurrent disease.
- All patients with **Medullary Carcinoma Thyroid** should also be tested for *RET* mutations, so that genetic counseling and testing of family members can be offered;

24. Ans. (c) **Medullary** (Ref: Robbins 9th/pg 1099-1100)

**Medullary carcinoma thyroid** presents with **amyloid deposition of ACal type**.

25. Ans. (d) **Seen in teratoma** (Ref: Robbins 9th/pg 1096)

**Psammoma Bodies** are

Lamellated, concentrically calcified structures formed by progressive acquisition of outer layers;

26. Ans. (a) **Papillary CA** (Ref: Robbins 9th/pg 1095-1097)

Most common Thyroid Carcinoma post radiation exposure is **Papillary CA**

27. Ans. (a) **Papillary carcinoma thyroid**

(Ref: Robbins 9th/pg 1095-1097)

28. Ans. (a) **Follicular carcinoma of thyroid**

(Ref: Robbins 9th/pg 1096; 8th/pg 38, 1122)

29. Ans. (a) **Papillary carcinoma** (Ref: R 9th/pg 1095-1097)

Most common thyroid Cancer is **Papillary carcinoma**

30. Ans. (a) **1,25 (OH)<sub>2</sub> Vit D<sub>3</sub>**

(Ref: Harrison 18th/chapter 352)

Active form of Vit D is 1,25 (OH)<sub>2</sub> Vit D<sub>3</sub>

31. Ans. (c) **8 mg/dl** (Ref: Harrison 18th/Appendix)

Normal Calcium level

<b>Calcium</b>	Serum	2.2–2.6 mmol/L	8.7–10.2 mg/dL
<b>Calcium, ionized</b>	Whole Blood	1.12–1.32 mmol/L	4.5–5.3 mg/dL

Hypocalcemia refers to total serum Ca < 8 mg/dl or ionized Ca < 1 mmol/L

32. Ans. (b) **Metastasis**

(Ref: Robbins 9th/pg 1104; 8th/pg 1129)

Morphology in hyperparathyroidism

<b>Parathyroid adenomas</b>	<b>Solitary lesions</b> with uniform, polygonal chief cells along with nests of oxyphil cells ( <b>oxyphil adenomas</b> )
<b>Primary hyperplasia</b>	<b>Chief cell hyperplasia</b> with abundant water-clear cells (" <b>water-clear cell hyperplasia</b> ") involving all 4 glands
<b>Parathyroid Carcinomas</b>	Circumscribed lesion in a single gland <b>diagnosed by invasion of surrounding tissues and metastasis as the only reliable criteria.</b>

33. Ans. (a) **Adenoma** (Ref: Robbins 9th/pg 1104)

Most common cause of primary hyperparathyroidism is **Adenoma**

34. Ans. (d) **All of the above**

(Ref: Robbins 9th/pg 1077, 1101; 8th/pg 1101, 1127)

<b>GSα</b>		
<b>Loss of function</b>	<b>Gain of function</b>	<b>Gain or loss of function</b>
<ul style="list-style-type: none"> <li>• Pseudohypoparathyroidism type Ia</li> <li>• Pseudohypoparathyroidism type Ib</li> </ul>	<ul style="list-style-type: none"> <li>• Pituitary or thyroid adenomas</li> <li>• Leydig cell tumors</li> <li>• McCune-Albright syndrome</li> </ul>	<ul style="list-style-type: none"> <li>• Testotoxicosis with pseudohypoparathyroidism type Ia</li> </ul>
<b>Gi2α: Gain of function</b>		
<ul style="list-style-type: none"> <li>• Pituitary adenomas</li> <li>• Adrenal cortex and ovary tumors</li> </ul>		

35. Ans. (a) **Hyperparathyroidism** (Ref: R 9th/pg 1102)

Skeletal Abnormalities in Hyperparathyroidism:

- Osteoporosis- cortical bone more severely involved than medullary bone
- Brown tumors: brown colour- due to increased vascularity, hemorrhage & hemosiderin deposition
- Osteitis fibrocystica (von Recklinghausen disease of bone)



36. Ans. (b) **Insulinoma** (Ref: Robbins 9th/pg 1121; 8th 1146)

37. Ans. (b) **Post receptor defects**

(Ref: Robbins 9th/pg 1111)

“Postreceptor” defects in insulin-regulated **phosphorylation/dephosphorylation** appear to play the predominant role in insulin resistance.

- PI-3-kinase signaling defect might reduce **translocation of GLUT4** to the plasma membrane.
- **Accumulation of lipid** within skeletal myocytes may **impair mitochondrial oxidative phosphorylation** and **reduce insulin-stimulated mitochondrial ATP production**.
- **Impaired fatty acid oxidation** and **lipid accumulation** within skeletal myocytes also may generate **reactive oxygen species** such as lipid peroxides.

38. Ans. (c) **Hepatocyte dysfunction**

(Ref: World J Hepatol. 2011 May 27; 3(5): 99–107)

**Factors responsible for development of hepatogenous insulin resistance/diabetes:**

- Hepatic parenchymal cell damage
- Portal-systemic shunting
- Hepatitis C virus

HCV genotype 3 can give rise to insulin resistance and non alcoholic fatty liver disease (NAFLD)

39. Ans. (a, b, d, e); a. **Insulinoma is most common neuroendocrine tumors of pancreas; b. VIPoma causes diarrhea; d. Somatostatinoma causes gall stone formation, e. Gastrinoma has high chance of malignancy** (Ref: Harrison 18th/chapter 350)

40. Ans. (b) **Beta cell**

(Ref: Robbins 9th/pg 1121; 8th/pg 1146)

- **Nesidioblastosis** is Focal or diffuse hyperplasia of the  $\beta$ -cells<sup>Q</sup> in pancreatic islets
- **Nesidioblastosis** can also be seen in:
  - Infant of diabetic mother<sup>Q</sup>
  - Beckwith-Wiedemann syndrome<sup>Q</sup>
  - Rare mutations in the  $\beta$ -cell K<sup>+</sup> channel protein or sulfonylurea receptor

41. Ans. (a) **Prediabetes** (Ref: Robbins 9th/pg 1106; 8th 1131)

Diagnostic criteria according to ADA and WHO: Presence of any 1 or more of the following:

Criteria	Diabetes	Impaired glucose tolerance (pre-diabetes)
• Fasting plasma glucose	$\geq 126$ mg/dL	between 100 and 125 mg/dL
• Random plasma glucose	$\geq 200$ mg/dL	–
• 2-hour plasma glucose during oral glucose tolerance test (OGTT) with 75 g	$\geq 200$ mg/dL	between 140 and 199 mg/dL
• Glycated hemoglobin (HbA1C) level	$\geq 6.5\%$	between 5.7% and 6.4%.

42. Ans. (a) **126 mg/dl**

(Ref: Robbins 9th/pg 1106; 8th/pg 1131)

43. Ans. (c) **Chloroquine**

(Ref: Robbins 9th/pg 1107)

**Important drugs that give rise to hyperglycemia are:**

- Glucocorticoids<sup>Q</sup>
- Thyroid hormone
- $\beta$ -adrenergic agonists<sup>Q</sup>
- Thiazides<sup>Q</sup>
- Phenytoin<sup>Q</sup>

44. Ans. (c) **Hypoglycemia**

(Ref: Robbins 9th/pg 1115–1116)

#### Acute Metabolic Complications of Diabetes:

- Diabetic ketoacidosis (Type 1 > type 2 DM)
- Hyperosmolar nonketotic hyperglycemia (Type 2 > type 1 DM)
- **Hypoglycemia<sup>Q</sup> (most common acute complication)**

45. Ans. (d) **Chronic Inflammation**

(Ref: Robbins 9th/pg 1115–1116)

**Mechanisms responsible for chronic complications of Diabetes are:**

- Non-enzymatic Glycosylation
- Protein Kinase C activation
- Disturbances in Polyol Pathways

46. Ans. (a) **Diffuse Mesangial Sclerosis**

(Ref: Robbins 9th/pg 1118–1119)

47. Ans. (b) **Stroke**

(Ref: Robbins 9th/pg 1115–1116)

48. Ans. (b) **Corneal opacity**

(Ref: Robbins 9th/pg 1115–1116)

**Diabetic Ocular Complications**

- Diabetes-induced hyperglycemia can lead to **cataract<sup>Q</sup>**
- **Glaucoma** → **damage to optic nerve<sup>Q</sup>**
- **Retinal vasculopathy<sup>Q</sup>** Can be non-proliferative or proliferative

49. Ans. (c) **MODY 3**

(Ref: Robbins 9th/pg 1107; 8th/pg 1132)


Most common type of MODY is MODY3

**Genetic defects of  $\beta$ -cell function:**

**Maturity-onset diabetes of the young (MODY)**

- **MODY1:** Hepatocyte nuclear factor 4 $\alpha$  (**HNF4A**)<sup>Q</sup>
- **MODY2:** Glucokinase (**GCK**)<sup>Q</sup>
- **MODY3:** Hepatocyte nuclear factor 1 $\alpha$  (**HNF1A**)<sup>Q</sup>
- **MODY4:** Insulin promoter factor-1 (**IPF-1**)
- **Neonatal diabetes** (Mutations in **KCNJ11&ABCC8**)<sup>Q</sup>





50. Ans. (a, c); **a. Family history is present in 90% cases; c. Type of onset is usually predictable**

(Ref: Robbins 9th/pg 1106-1110; 8th/pg 1131-1134 Harrison 18th/chapter 344)

Discussing options about **Type 1 Diabetes**, one by one

- |    |   |
|----|---|
| a. | False as <b>Family history is present 10-20%cases</b>   |
| b. | True as <b>insulin deficiency predisposes to DKA</b>  |
| c. | False as the <b>rate of decline in beta cell mass varies widely</b> among individuals, with some patients progressing rapidly to clinical diabetes and others evolving more slowly. <b>Features of diabetes do not become evident until a majority of beta cells are destroyed (70–80%)</b> . |
| d. | True as <b>circulating islet autoantibodies</b> (anti-insulin, anti-GAD, anti-ICA512) causes immune destruction of B cells  |
| e. | True as <b>it usually involves normal wt children &amp; young adults</b>  |

51. Ans. (a) **Before surgery alpha blockers are given; (c) Mostly malignant; (d) Mostly familial; (e) Associated with von Hippel-Lindau disease** (Ref: R 9<sup>th</sup> pg 1135)

Pheochromocytoma is often associated with MEN-2A, MEN-2B, NF1, Von Hippel-Lindau (VHL), Familial paraganglioma

52. Ans. (c, d) **c. Ewings Sa d. Medulloepithelioma**

(Ref: Robbins 9th/pg 1134-1136)

Primitive neuroectodermal tumors (PNETs) are a group of highly malignant tumors comThe following tumors are classified as peripheral primitive neuroectodermal tumors (pPNETs):

- Ewing sarcoma (osseus and extraosseous)
- Malignant peripheral primitive neuroectodermal tumors (pPNETs) or peripheral neuroepithelioma of bone and soft tissues
- Askin tumor (peripheral neuroepithelioma of the thoracopulmonary region)
- Other less common tumors (eg, neuroectodermal tumor, ectomesenchymoma, peripheral medulloepithelioma)

53. Ans. (a) **S-100** (Ref: Robbins 9/1135)

In pheochromocytoma: Immunoreactivity for neuroendocrine markers (chromogranin and synaptophysin) is seen in the chief cells, while the peripheral sustentacular cells stain with antibodies against S-100 which is a calcium-binding protein.

54. Ans. (a) **Synaptophysin**

55. Ans. (b) **Paraganglioma** (Ref: Robbins 9/1135)

The histologic pattern in pheochromocytoma shows tumor cells composed of clusters of polygonal to spindle shaped chromaffin cells or chief cells that are surrounded by supporting sustentacular cells, creating small nests or alveoli called zellballen. These are supplied by a rich vascular network

56. Ans. (b) **Argentaffin cells of small intestine**

Carcinoid tumors arise from argentaffin cells of the crypts of Lieberkühn and are found from the distal duodenum to the ascending colon, areas embryologically derived from the midgut.

57. Ans. (b, c, e) **b. Extra-adrenal Pheochromocytoma c. Carotid body tumour e. Glomus tympanicum**

(Ref: Robbins 9th/ 741-42; Harrison 19th/ 2329-35; Davidson 22nd/ 781; CMDT 2016/ 1158; Danhart Radiology Review Manual 7th/401)

**Pheochromocytomas and paragangliomas are catecholamine- producing tumors derived from the sympathetic or parasympathetic nervous system**

58. Ans. (c) **10% inherited**

(Ref: Robbins 9th/pg 1134-1135/8th 1159-1160)

59. Ans. (b) **Adrenal medulla** (Ref: Robbins 9th/pg 1135)

60. Ans. (a, b, c, e); **a. Synthesizes epinephrine & norepinephrine; b. Increased urinary 5HIAA; c. Diagnosed by urinary catecholamine metabolites; e. Tumor of adrenal medulla and sympathetic ganglion**

(Ref: Robbins 9th/pg 1134-1135; 8th/pg 1159-1160 Harrison 18th/chapter 343)

**Biochemical tests for Pheochromocytoma:**

- Pheochromocytomas synthesize & store catecholamines, which include **norepinephrine (noradrenaline), epinephrine (adrenaline), and dopamine**.
- Elevated plasma and urinary levels of **catecholamines** and the methylated metabolites, **metanephrines**, are the cornerstone for the diagnosis.
- Urinary tests for **vanillylmandelic acid (VMA), metanephrines (total or fractionated), and catecholamines** are widely available and are used commonly for initial testing.

61. Ans. (a) **Pheochromocytoma**

(Ref: R 9th/pg 1134-1135)

62. Ans. (a) **Involvement of lymph nodes**


(Ref: Robbins 9th/pg 1134-1135)

- **There is no histologic feature that reliably predicts clinical behavior.**
- **Therefore, the definitive diagnosis of malignancy in pheochromocytomas is based exclusively on the presence of metastases.**
- **Metastasis:** Involve regional **lymph nodes<sup>o</sup>, liver<sup>o</sup>, lung<sup>o</sup>, and bone<sup>o</sup>**

63. Ans. (a) **Carotid body Tumour** (Ref: Robbins 9th/pg 517)

**Glomus tumour (Carotid body tumor)**

- Distinctive neoplasm which arises from **modified smooth muscle cells of the normal glomus body**.



- Glomus body is a specialized form of **arteriovenous anastomosis**, involved in **temperature regulation**.
- There is a central coiled canal known as Suquet-Hoyer canal which is lined by plump endothelial cells.

**64. Ans. (a, b); a. Pheochromocytoma; b. Paraganglioma**

(Ref: Robbins 9th/pg 1135; 8th/pg 1160)

**Microscopic Feature of Pheochromocytoma**

- Tumors made of **clusters of polygonal to spindle-shaped chromaffin cells** surrounded by **sustentacular cells**, creating **small nests (zellballen)**<sup>Q</sup>, with rich vascularity.
- **"Salt and pepper"** nuclear chromatin: **characteristic** of neuroendocrine tumors.
- **Electron microscopy:** Membrane-bound, **electron-dense secretory**<sup>Q</sup> granules.

Zellballen pattern is also found in Paraganglioma

**65. Ans. (c) Duodenum** (Ref: Robbins 9th/pg 1136)

**66. Ans. (d, e) d. Marfanoid features; e. Mucosal neuromas**

Multiple endocrine neoplasia type 2 (MEN 2) is classified into three subtypes: MEN 2A, FMTC (familial medullary thyroid carcinoma), and MEN 2B. All three subtypes involve high risk for development of medullary carcinoma of the thyroid (MTC); MEN 2A and MEN 2B have an increased risk for pheochromocytoma; MEN 2A has an increased risk for parathyroid adenoma or hyperplasia. Additional features in MEN 2B include mucosal neuromas of the lips and tongue, distinctive facies with enlarged lips, ganglioneuromatosis of the gastrointestinal tract, and a "marfanoid" habitus. MTC typically occurs in early childhood in MEN 2B, early adulthood in MEN 2A, and middle age in FMTC.

**67. Ans. (b) Midgut carcinoid** (Ref: Robbins 9th ed. Pg. 1130)

**68. Ans. (a) MEN1** (Ref: Robbins 9/1136)

MEN-1, or Wermer syndrome, characterized by abnormalities involving the parathyroid (*Primary hyperparathyroidism resulting in renal stones*), pancreas, and pituitary glands (*prolactinoma*); thus the mnemonic device, the 3Ps.

**69. Ans. (d) MEN1** (Ref: Robbins 9th/pg 1136/8th 1162)

Multiple Endocrine Neoplasia (MEN) Syndromes; Refer to pretexts of this chapter

**70. Ans. (b) Gastrinoma** (Ref: Robbins 9th/pg 1136/8th 1162)

**71. Ans. (a) Somatostatinoma** (Ref: Robbins 9th/pg 1136)

**Pancreatic endocrine tumors;** Refer to pretexts

**72. Ans. (d) Medullary**

(Ref: Robbins 9th/pg 1136; 8th/pg 1162)

Commonest thyroid tumor in MEN is Medullary Carcinoma thyroid

**73. Ans. (d) Pituitary** (Ref: Robbins 9th/pg 1136; 8th/pg 1162)

**74. Ans. (a) MEN 1** (Ref: Robbins 9th/pg 1136; 8th/pg 1162)

Wermer syndrome is MEN 1

**75. Ans. (d) Medullary carcinoma of thyroid**

(Ref: Robbins 9th/pg 1136)

**76. Ans. (b) Component of MEN-1**

(Ref: Robbins 9th/pg 1099-1100, 1136; 8th/pg 1124-1125, 1162)

[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines, typical of notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Skin and its Disorders

## Key Points

- » **Erythema multiforme:** Characteristic clinical lesion-**target lesion**
- » **Auspitz sign:** Multiple, minute, bleeding points when the scale is lifted from the plaque
- » **Lichen planus:** **Hypergranulosis**
- » **Psoriasis:** **Stratum granulosum is thinned or absent**
- » **Pemphigus vulgaris:** **Suprabasal** acantholytic vesicle
- » **Verruca vulgaris** is the **most common type** of wart
- » Epithelioid granulomas are seen in **tuberculoid leprosy**
- » **Marjolin's ulcer:** SCC arising at site of chronic inflammation, presenting as **persistent ulceration**
- » **Mycosis fungoides:** Histologic hallmark - **Sézary-Lutzner cells**


## Key Recent Updates

- » IgG antibodies to **desmoglein 1 & 3** are seen in **pemphigus vulgaris**
- » IgG autoantibodies to **hemidesmosomes** is seen in **Bullous pemphigoid**.







## NORMAL SKIN HISTOLOGY



## IMPORTANT DEFINITIONS




- **Acanthosis:** Diffuse **epidermal hyperplasia**<sup>o</sup>
- **Hyperkeratosis:** Thickening of the **stratum corneum**<sup>o</sup>
- **Parakeratosis:** Keratinization with **retained nuclei** in the stratum corneum. **On mucous membranes, parakeratosis is normal.**<sup>o</sup>
- **Dyskeratosis:** Abnormal, **premature keratinization** within cells **below the stratum granulosum**<sup>o</sup>
- **Spongiosis:** **Intercellular edema** of the epidermis<sup>o</sup>
- **Acantholysis:** **Loss of intercellular connections**<sup>o</sup>



## ACUTE INFLAMMATORY DERMATOSES

### Urticaria

- Localized **immediate hypersensitivity (type I)** reaction<sup>o</sup>




### POINTS TO REMEMBER

- Hereditary Angioneurotic edema-deficiency of C1 inhibitor
- Angiedema-edema of both dermis and subcutaneous fat

## ACUTE ECZEMATOUS DERMATITIS

- Eczematous dermatitis can be subdivided into the following categories:
  - Allergic contact dermatitis, Atopic dermatitis, Drug-related eczematous dermatitis, Photoeczematous dermatitis, Primary irritant dermatitis.<sup>Q</sup>
- Hallmark of acute eczema: spongiosis



## ERYTHEMA MULTIFORME


- Characteristic clinical lesion-**target lesion**<sup>Q</sup>-shows central necrosis surrounded by a rim of perivenular inflammation<sup>Q</sup>



## CHRONIC INFLAMMATORY DERMATOSES

- **Psoriasis**
  - Psoriasis is a **chronic inflammatory dermatosis**<sup>Q</sup> usually with **autoimmune basis**<sup>Q</sup>
  - **Most frequently** - *skin of the elbows, knees, scalp, lumbosacral areas, intergluteal cleft, and glans penis.*<sup>Q</sup>
  - **Koebner's phenomenon**<sup>Q</sup> -lesions can be induced in susceptible individuals by local trauma.
  - Strong association with **HLA-C**, particularly with the **HLA-Cw\*0602 allele**<sup>Q</sup>
  - **Auspitz sign**- multiple, minute, bleeding points when the scale is lifted from the plaque<sup>Q</sup>
  - **Nail changes occur in 30% of cases of psoriasis**<sup>Q</sup>





- Hyperkeratosis, parakeratosis, acanthosis
- Regular downward elongation of the rete ridges (**test tubes in a rack**)
- **Stratum granulosum is thinned or absent**
- Thinning of **suprapapillary plates**
- Intraepidermal infiltrates of neutrophils in the **stratum corneum (Munro microabscesses)** & in **spinous layer (spongiform pustules of Kogoj)**

#### ■ Seborrheic Dermatitis (SD)

- Classically involves regions with a high density of sebaceous glands, - scalp, forehead (especially the glabella), external auditory canal.



- **Dandruff is the common clinical expression** of SD of the scalp
- In infants, SD presents as **cradle cap**
- **Follicular lipping** - mounds of **parakeratosis** containing neutrophils and serum at the ostia of hair follicles

#### ■ Lichen Planus

- **"Six Ps"** : **"Pruritic, purple, polygonal, planar, papules, and plaques"**
- **Violaceous**, flat-topped papules coalesce focally to form plaques
- Papules are highlighted by white lines called **Wickham striae** created by areas of **hypergranulosis**
- Koebner's phenomenon may be seen

#### Microscopy

- Hyperkeratosis
- **Hypergranulosis**
- **Band of lymphocytes** are intimately associated with basal keratinocytes
  - **Squamization**, degeneration, necrosis seen in basal keratinocytes
  - **Colloid or Civatte bodies** -necrotic basal cells incorporated into the inflamed papillary dermis.




Lichen planus

### Mnemonic

- Pemphigus **S**-Superficial separation<sup>Q</sup>- intraepidermal acantholysis<sup>Q</sup>-**flaccid bulla**<sup>Q</sup> which ruptures easily
- Pemphigoid **D**-Deep separation at dermo-epidermal junction<sup>Q</sup>- **tense bulla**<sup>Q</sup> that **does not** rupture easily







## BLISTERING (BULLOUS) DISEASES

Caused by **autoantibodies**<sup>Q</sup> specific for epithelial or basement membrane proteins that lead to separation of keratinocytes (acantholysis)





Skin biopsy and direct immunofluorescence (DIF) are crucial for diagnosis of immune-bullous diseases.<sup>9</sup>

	Bullous Pemphigoid	Pemphigus Vulgaris	Dermatitis Herpetiformis
<b>MIC:</b>	Subepidermal bulla 	Suprabasal bulla (Tombstone pattern)  Row of tomb stone	Subepidermal bulla Neutrophil and fibrin accumulate at tip of dermal papule  Sub-epidermal bulla
<b>DIF:</b>	Linear IgG, C <sub>3</sub> in basement membrane zone  Linear pattern	IgG deposits in intercellular spaces-fishnet pattern  Fish net pattern	Granular IgA in tips of papillae  Granular pattern

#### Other Bullous disorders

- **Pemphigoid gestations:** Spongiotic dermatitis and subepidermal bulla
- **Epidermolysis bullosa acquisita:** subepidermal bulla sparse inflammatory infiltrate
- **Pemphigus foliaceus:** subcorneal bulla



#### High Yield Facts

- **Pemphigus:** IgG autoantibodies to various **intercellular desmogleins**. (Both desmoglein 1 & 3)
- **Bullous Pemphigoid:** IgG autoAb to **hemidesmosome** proteins c. (Bullous Pemphigoid Antigen (BPAG) that attach epidermal cells to basement membrane)
- **Dermatitis Herpetiformis:** IgA auto-Ab to **fibrils** that bind epidermal BM & produces **subepidermal blisters**. Associated with **gluten-sensitive enteropathy** in most cases<sup>9</sup>
- **Epidermolysis Bullosa:** **inherited defects in structural proteins that lend mechanical stability to the skin**<sup>9</sup>
- **Dystrophic epidermolysis bullosa:** **inherited defects in collagen type 7**<sup>9</sup>
- **Porphyria:** Vesicles are **subepidermal** in location and the adjacent dermis contains vessels with walls that are thickened by **glassy deposits of serum proteins, including immunoglobulins**<sup>9</sup>
- **Seborrheic dermatitis is not a disease of the sebaceous glands**
- **Leiner disease**<sup>9</sup> generalized seborrheic dermatitis in infants associated with diarrhea & failure to thrive

## DISORDERS OF PIGMENTATION AND MELANOCYTES

- **Freckle (Ephelis):** **Most common**<sup>9</sup> pigmented lesions of childhood in lightly pigmented individuals
- **Lentigo:** Essential histologic feature is linear (non-nested) melanocytic hyperplasia restricted to basal cell layer<sup>9</sup>
- **Melanocytic Nevus (Pigmented Nevus, Mole)**
  - Common benign neoplasms, occur by acquired activating mutations RAS signaling pathway<sup>9</sup>
- Acquired **melanocytic nevi** are the **most common type** and found in **all individuals**.<sup>9</sup>
- **Dysplastic Nevi:** Best regarded as **markers of melanoma risk** rather than premalignant lesions
  - They are characterized by architectural and cytologic atypia
  - Associated with **germline mutations in genes encoding cell cycle regulators (p16/INK4a, CDK4) and telomerase**.<sup>9</sup>
- **Melanoma:** Highly aggressive malignancy linked to sun exposure;





- The two most important predisposing factors are inherited genes and sun exposure<sup>Q</sup>
- Shows striking **variations in color**, in contrast to benign nevus<sup>Q</sup>
- The borders of melanomas are **irregular and often notched**<sup>Q</sup>
- **Radial growth** describes the horizontal spread of melanoma within the epidermis and superficial dermis- **this is not associated with risk of metastasis.**<sup>Q</sup>
- **Vertical growth phase is signaled by the appearance of a nodule & correlates with metastatic potential.**<sup>Q</sup>

### POINTS TO REMEMBER

- Acne, Rosacea, Panniculitis (Erythema nodosum) are disorders of epidermal appendages

### Mnemonic

#### Mole: Signs of Trouble ABCDE

- A**symmetry
- B**order irregular
- C**olour irregular
- D**iameter usually > 0.5 cm
- E**levation irregular




### High Yield Facts

- **Superficial spreading**, - most common type of melanoma,<sup>Q</sup> usually involving sun-exposed skin
- **Acral/mucosal lentiginous melanoma** - is **unrelated** to sun exposure.<sup>Q</sup>
- Melanoma is associated **with mutations** in **cell cycle regulators** (p16/INK4a, CDK4), growth factor receptors [e.g., KIT], **RAS**, **BRAF**), and **telomerase**<sup>Q</sup>
- **Marker- HMB-45** is very specific marker for melanocytes<sup>Q</sup>
  - Few other familial syndromes, notably **CDKN2 A** is a **important risk factor of melanoma**

## INFECTIONS

### ■ Verrucae (Warts):


- Caused by **human papilloma viruses**.
- **Verruca vulgaris** is the **most common type**<sup>Q</sup> of wart.
- **Condyloma acuminatum (venereal wart)** occurs on the penis, female genitalia, urethra, perianal areas, and rectum.
- Diagnosis is based primarily on hyperplastic papillary architecture zones of **koilocytosis**<sup>Q</sup>.



Verruca

### ■ Molluscum Contagiosum:

- Caused by a **poxvirus**.<sup>Q</sup>
- **Diagnostic feature** is the **molluscum body**-large (up to 35 µm), homogeneous, cytoplasmic inclusion in cells of the **stratum granulosum** and the **stratum corneum**<sup>Q</sup>



Molluscum bodies

### ■ Impetigo:

- **Common superficial bacterial infection** of skin caused by **Staphylococcus aureus**<sup>Q</sup>


### ■ Superficial Fungal Infections:

- Tinea capitis, Tinea barbae, Tinea corporis, **Tinea pedis (athlete's foot)** & Tinea versicolor<sup>Q</sup>
- All of above are caused by **dermatophyte infections** except **Tinea versicolor** which is caused by **Malassezia furfur** (a yeast, not a dermatophyte).<sup>Q</sup>



## Clinical, Bacteriologic, Pathologic, and Immunologic Spectrum of Leprosy

Features	Tuberculoid (TT, BT) Leprosy	Lepromatous (LL) Leprosy
Skin lesions	<b>Sharply defined</b> macules or plaques with a tendency toward central clearing, <b>elevated borders</b>	<b>Symmetric, poorly marginated, multiple infiltrated nodules and plaques<sup>q</sup></b> or diffuse infiltration; <b>leonine facies and eyebrow alopecia<sup>q</sup></b>
Nerve lesions	Skin lesions <b>anesthetize early</b> ; nerve near lesions sometimes <b>enlarged</b> ; <b>nerve abscesses most common in BT<sup>q</sup></b>	<b>Hyperesthesia, a late sign</b> ; nerve palsies variable; acral, distal, symmetric anesthesia common
Acid-fast bacilli (BI)	<b>0–1+</b>	<b>4–6+ GLOBI-</b> Macrophage cells laden with acid fast bacill <sup>q</sup>
Lymphocytes	<b>2+</b>	0–1+
Macrophage differentiation	<b>Epithelioid</b>	<b>Foamy cells change the rule; Lepra cells<sup>q</sup></b> - large, mononuclear histiocytes (macrophages) with a foam like cytoplasm
<b>Langhans' giant cells Histopathology<sup>q</sup></b>	<b>1–3+<sup>q</sup></b> <b>Epithelioid granulomas</b> in the papillary dermis, specially <b>around neurovascular structures<sup>q</sup></b>	— <b>Diffuse infiltrate of foamy macrophages</b> is present in the dermis below a subepidermal zone of uninvolved papillary dermis (i.e., <b>grenz zone</b> ). <sup>q</sup>
Lepromin skin test	<b>+++<sup>q</sup></b>	—
Lymphocyte transformation test	<b>Generally positive<sup>q</sup></b>	1–2%
<b>M. leprae PGL-1 antibodies</b>	60%	<b>95%<sup>q</sup></b>



**Image shows granuloma S/O Tuberculoid leprosy    Foamy cells S/O lepromatous leprosy**  
Acid-fast bacilli (BI): 0–1+    Acid-fast bacilli (BI): 4–6+ GLOBI- Macrophage cells laden with acid fast bacill<sup>q</sup>



### High Yield Facts

- **Globi:** Macrophage cells laden with acid fast bacill
- **Lepra cells<sup>q</sup>:** Large, mononuclear histiocytes (macrophages) with a foam-like cytoplasm
- **Grenz zone<sup>q</sup>** is seen in **lepromatous leprosy**
- **Epithelioid granulomas** are seen in **tuberculoid leprosy**

## BENIGN EPITHELIAL TUMORS

- Derived from keratinizing stratified **squamous epithelium** of epidermis, hair follicles & **ductular epithelium** of cutaneous glands.
- Telltale sign of syndromes associated with **visceral malignancies**, such as **multiple trichilemmomas in Cowden's syndrome<sup>q</sup>** or **multiple sebaceous neoplasms in Muir-Torre syndrome<sup>q</sup>**



## SEBORRHEIC KERATOSES

- Middle-aged or older individuals
- Appear as part of a **paraneoplastic syndrome (Leser-Trélat sign)**<sup>Q</sup> possibly due to stimulation of keratinocytes by **TGF-alpha**<sup>Q</sup> produced by tumor cells, **most commonly carcinomas of the gastrointestinal tract**.<sup>Q</sup>
- **Sporadic** activating mutations in the **fibroblast growth factor receptor-3 (FGFR3)**<sup>Q</sup>
- Clinically: **coin-like, waxy plaques**.
- Inspection with a hand lens usually reveals small, **round, pore-like ostia impacted with keratin**<sup>Q</sup>
- Microscopy shows following **characteristic features**:
  - Hyperkeratosis
  - **Horn cysts**-keratin-filled cysts
  - **Invagination cysts**-invaginations of keratin into the main mass

## ACANTHOSIS NIGRICANS

Thickened, hyperpigmented skin with a “**velvet-like**”<sup>Q</sup> texture, mostly seen in flexural areas

It is of two types

- In 80% of cases, associated with benign conditions like **obesity and diabetes**.<sup>Q</sup>


- In 20% cases, AN arises in association with cancers, **most commonly gastrointestinal adenocarcinomas**<sup>Q</sup>

## FIBROEPITHELIAL POLYP, (ACROCHORDON, SQUAMOUS PAPILLOMA, SKIN TAG)


- **Most common cutaneous lesions**<sup>Q</sup>
- Occurs at **neck, trunk, face, and intertriginous areas**<sup>Q</sup>
- Become **more numerous or prominent during pregnancy**

## ADNEXAL (APPENDAGE) TUMORS

- **Eccrine poroma**: Occurs predominantly on the palms and soles where sweat glands are numerous
- **Cylindroma: Turban tumor** (Microscopy shows zigzag puzzle pattern) see image below.
- **Brooke-Spiegler syndrome**: (associated with both trichoepithelioma and cylindroma)
- **Syringomas**
- **Sebaceous adenomas**-Can be associated with internal malignancy in **Muir-Torre syndrome**, a subset of hereditary non-polyposis colorectal carcinoma syndrome associated with germline deficits in **DNA mismatch repair**<sup>Q</sup> proteins.
- **Pilomatricomas**-shows **ghost cells**



Cells arranged in zigzag puzzle separated by basement membrane like material diagnostic of cylindroma



PAS+ basement membrane like material diagnostic of cylindroma



### High Yield Facts

- Most appendage tumors are **benign**.<sup>Q</sup>
- Apocrine tumors are unusual in that **malignant forms seem to be more common**<sup>Q</sup> than benign.
- Sebaceous carcinoma arises from the **meibomian glands**<sup>Q</sup> of the eyelid and may follow an aggressive course.
- **Birt-Hogg-Dubé syndrome**-fibroepithelial polyps and tumors of **perifollicular mesenchyme**.<sup>Q</sup>

## PREMALIGNANT AND MALIGNANT EPIDERMAL TUMORS

- **Carcinoma in Situ (Bowen's Disease)**
  - Occurs equally in **men and women**
  - Associated with HPV, Arsenic, solar radiation
  - **Morphology**: Full thickness dysplasia of the squamous epithelium.





## High Yield Facts

Differential diagnosis of Bowen's disease include Bowenoid papulosis and Erythroplasia of Queyrat occurs primarily in men.

- **Bowenoid papulosis:**

Histologically similar to squamous cell carcinoma in-situ.

Features **favoring the diagnosis of bowenoid papulosis** are:

- Presence of numerous mitotic figures in metaphase,
- Basophilic inclusions in the granular layer
- Koilocytes

- **Erythroplasia of Queyrat: Squamous cell carcinoma in situ** presenting on the **mucous membranes of the glans penis, vulva and oral mucosa**<sup>Q</sup>

- **Actinic Keratosis**


- **Sandpaper-like**<sup>Q</sup> consistency, grossly.
- Characterised by **cutaneous horns and pseudohorn cysts**<sup>Q</sup>.
- **Sites:** Sun-exposed sites (face, arms, dorsum of hands) are most frequently affected.
- The lips may also develop similar lesions (termed **actinic cheilitis**).
- Morphology
  - Hyperkeratosis, **parakeratosis**<sup>Q</sup>
  - **Basal cell and squamous layer atypia** and **disorderly maturation**<sup>Q</sup>

- **Squamous Cell Carcinoma (SCC):**

- It is the **2<sup>nd</sup> most common tumor**<sup>Q</sup> arising on sun-exposed sites in older people, exceeded only by basal cell Ca.
- **Most important cause DNA damage induced by exposure to UV light**<sup>Q</sup>

### Risk factors


<ul style="list-style-type: none"> <li>• Usually UV light / ionizing radiation</li> <li>• Actinic keratosis (precursor lesion)<sup>Q</sup></li> <li>• Albinism (lack of pigmentation in skin)</li> <li>• <b>Arsenic</b><sup>Q</sup></li> <li>• Burn scars</li> </ul>	<ul style="list-style-type: none"> <li>• Chronic ulcers</li> <li>• <b>Epidermodysplasia verruciformis</b><sup>Q</sup></li> <li>• Tars/oils</li> <li>• <b>Hidradenitis suppurativa</b><sup>Q</sup></li> <li>• Immunosuppression (post-transplant or HIV)</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Necrobiosis lipoidica</b><sup>Q</sup></li> <li>• Osteomyelitis - draining sinuses</li> <li>• PUVA treatment for psoriasis</li> <li>• <b>Xeroderma pigmentosa</b><sup>Q</sup>: disorder with diminished capacity for DNA repair after UV light exposure</li> </ul>
--	--	---




Mic: Keratin pearl and malignant squamous cells S/o SCC

- **Basal Cell Carcinoma-BCC**

- Basal cell carcinoma is the **most common**<sup>Q</sup> invasive cancer in humans
- Locally invasive-**rarely metastasize**<sup>Q</sup>.
- **Most common mutations is—Sonic Hedgehog pathway**



Peripheral palisading of nuclei S/O basal cell Ca



## High Yield Facts

- **Cancers associated with sunlight exposure: BCC & SCC**
- HPVs cause autosomal recessive condition, **epidermodysplasia verruciformis**, which is marked by a **high susceptibility to cutaneous squamous cell carcinomas**<sup>Q</sup>
- **Marjolin's ulcer**<sup>Q</sup>: SCC arising at site of chronic inflammation, presenting as **persistent ulceration**<sup>Q</sup>
- **Rodent ulcer**<sup>Q</sup>- BCC (Due to local invasiveness of tumor)
- **Nevoid basal cell carcinoma syndrome (Gorlin syndrome)**—autosomal dominant disorder; multiple basal cell Ca,<sup>Q</sup> often before age 20, accompanied by other tumors (medulloblastomas<sup>Q</sup> and ovarian fibromas<sup>Q</sup>), odontogenic keratocysts<sup>Q</sup>, pits of palms & soles.
- **Urticaria pigmentosa** is cutaneous form of mastocytosis





## TUMORS OF THE DERMIS

### Benign Fibrous Histiocytoma

- The **most common** form of fibrous histiocytoma is referred to as a **dermatofibroma**<sup>Q</sup>

### Dermatofibrosarcoma Protuberans

- Well-differentiated, primary fibrosarcoma of the skin**<sup>Q</sup>
- Deep extension from the dermis into subcutaneous fat, producing a characteristic **"honeycomb"** pattern<sup>Q</sup>
- The **molecular hallmark** - **balanced translocation between genes encoding collagen 1A1 (COL1A1) and the platelet-derived growth factor- $\beta$  (PDGFB)**<sup>Q</sup>
- Characteristically has **storiform arrangement of fibroblasts** (reminiscent of blades of a pinwheel)<sup>Q</sup>

## TUMORS OF CELLULAR MIGRANTS TO THE SKIN

In these lesions, progenitors arise elsewhere and then specifically home to the cutaneous microenvironment.


## Mnemonic

### Tumors that metastasize to skin BLOCK:

- B**reast
- L**ung
- O**vary
- C**olon
- K**idney

### Mycosis Fungoides (Cutaneous T-Cell Lymphoma)

- T-cell lymphoma** that presents in the skin and may evolve into generalized lymphoma
- Lesions involve truncal areas and include scaly, red-brown **patches**; raised, scaling **plaques** and fungating **nodules**
- Histologic hallmark** - **Sézary-Lutzner cells**<sup>Q</sup>, which form band-like aggregates within the superficial dermis
- These cells are **T-helper cells (CD4+)** have **hyperconvoluted or cerebriform contour**<sup>Q</sup>
- These invade the epidermis as single cells and small clusters (**Pautrier microabscesses**).<sup>Q</sup>
- Prognosis depends on **percentage** of body surface involved & **progression** from patch to plaque to nodule




E.M Mycosis fungoides. Several T-lymphocytes with deeply indented and irregular ("cerebriform") nuclei

Cerebriform nucleus



Sézary-Lutzner cell




Pautrier microabscess

### Mastocytosis

- Characterized by increased numbers of mast cells in the skin and in other organs
- Darier sign**-localized area of dermal edema and erythema (wheal) when lesional skin is rubbed<sup>Q</sup>
- Dermatographism** - area of **dermal edema** due to **localized stroking** of normal skin with a pointed instrument<sup>Q</sup>
- Diagnosis of mast cells-metachromatic stains (**toluidine blue or Giemsa**)<sup>Q</sup>



Dermatographism




## Image-Based Questions

1. A 24-year-old male presents with following skin condition. Before doing biopsy, provisional diagnosis for the following condition is:




- a. Erythema multiforme
- b. Psoriasis
- c. Lichen planus
- d. Lichen striatus

3. A 50-year-old female presents with blisters and erosions on the skin and mucous membranes, most commonly inside the mouth. Direct immunofluorescence (DIF) on normal-appearing perilesional skin was done. It shows IgG within suprabasal intercellular spaces—fishnet pattern. Diagnosis:




- a. Pemphigus vulgaris
- b. Dermatitis herpetiformis
- c. Bullous pemphigoid
- d. None

2. A 50-year-old female presents with blisters and erosions on the skin and mucous membranes, most commonly inside the mouth. Diagnosis:




- a. Pemphigus vulgaris
- b. Dermatitis herpetiformis
- c. Bullous pemphigoid
- d. None


4. A 60-year-old female presents with tense bullae on the flexural areas of the skin. Oral mucosa not involved. DIF shows Linear IgG, C3 in basement membrane zone. Diagnosis:



- a. Pemphigus vulgaris
- b. Bullous pemphigoid
- c. Pemphigus vegetans
- d. None




5. A 5-year-old child presents with Raised, firm, flesh colored nodules. On histopathological examination, following is seen. Diagnosis:



- a. Myrmecia  
b. Molluscum contagiosum  
c. Herpes zoster  
d. Herpes simplex

6. A 43-year-old smoker male, presents with growth in oral cavity. On examination, patient was diagnosed with well differentiated squamous cell carcinoma. Identify the structure marked with arrow.




- a. Keratin pearl  
b. Koilocytes  
c. Molluscum  
d. None



## Answers of Image-Based Questions

- Ans. (a) **Erythema multiforme**
  - Targetoid lesions are features of **Erythema multiforme**
  - Clinical **targetoid (target-like) lesion** shows central necrosis surrounded by a rim of perivenular inflammation
- Ans. (a) **Pemphigus vulgaris**
  - Early lesions of pemphigus vulgaris show **suprabasal** epidermal acantholysis, clefting and blister formation
  - Basal cells are intact giving tombstone pattern
  - The blister cavity may contain inflammatory cells including eosinophils and rounded acantholytic cells with intensely eosinophilic cytoplasm and a perinuclear halo.
- Ans. (a) **Pemphigus vulgaris**
  - DIF usually shows immunoglobulin G (IgG) deposited on the surface of the keratinocytes in and around lesions giving a fish net pattern. IgG1 and IgG4 are the most common subclasses.
  - Direct immunofluorescence (DIF) is usually done on normal-appearing perilesional skin
- Ans. (b) **Bullous pemphigoid**
  - Linear IgG, C3 in basement membrane zone on DIF is a feature of Bullous pemphigoid
- Ans. (b) **Molluscum Contagiosum**
  - Molluscum bodies are present (large cells with cytoplasmic, faintly granular eosinophilic inclusions that displace nuclei and contain viral particles) marked with arrow.
- Ans. (a) **Keratin pearl**
  - Typical SCC has nests of squamous epithelial cells arising from the epidermis and extending into the dermis
  - We differentiated neoplastic cells in nests have pink cytoplasmic keratin.
  - Keratin pearl-a focus of central keratinization within concentric layers of abnormal squamous cells is a feature of well-differentiated SCC.





## Multiple Choice Questions

### OVERVIEW AND DEFINITIONS


- Features of café au lait spots are all except:** (Recent Question 2015)
  - Larger
  - Arise independent of sun exposure
  - Contain aggregated melanosomes
  - Most common pigmented lesions
- Acanthosis means:** (AI 2014)
  - Loss of intracellular connections
  - Abnormal premature keratinization
  - Diffuse epidermal hyperplasia
  - Thickening of stratum corneum
- Spongiosis is seen in:** (Recent Question 2013)
  - Acute eczema
  - Lichen Planus
  - Psoriasis
  - Pemphigus
- True about Dyskeratosis congenita:** (PGI May 10)
  - Pancytopenia
  - Nail dystrophy
  - Hyperkeratosis
  - X linked
  - Leukoplakia

### INFLAMMATORY DERMATOSES


- Munro microabscesses are seen in:** (WB PGME 16, MH 16)
  - Lichen planus
  - Mycosis fungoides
  - Psoriasis
  - Eczema

### BLISTERING DISORDERS


- A 20-year-old female presented with painful blister in skin and oral mucosa. Direct immunofluorescence picture of which is given below. Which of the following is not true regarding the same?** (AIIMS May 2017)
  - Antibodies against hemidesmosomes
  - Antibodies against Desmoglein 1
  - Antibodies against Desmoglein 3
  - Basemene membrane deposition of IgG is the most common DIF picture in bullous lesions



- What is the diagnosis of the below histopathology image?** (AIIMS May 2017)



- Pemphigus
  - Leishmaniasis
  - Mycosis fungoides
  - Psoriasis
- Fish net pattern seen in** (AIIMS Nov 16)



- Pemphigus Vulgaris
  - Bullous pemphigoid
  - Behcet syndrome
  - All of above
- Acantholysis means:** (Recent Question 2014)
    - Diffuse epidermal hyperplasia
    - Loss of intercellular connections
    - Intercellular edema of the epidermis.
    - Abnormal keratinization
  - In congenital dystrophic variety of epidermolysis bullosa, mutation is seen in the gene coding for?** (AIIMS May 11)
    - Laminin 4
    - Collagen type VII
    - Alpha 6 integrin
    - Keratin 14
  - 'Row of tombstones' appearance is seen in:** (JIPMER 78, PGI 11)
    - Irritant dermatitis
    - Pemphigus
    - Pemphigoid
    - Herpes zoster





## INFECTIONS

12. The "Lepra cells" are: (Recent Question 2014)  
 a. histologically b. Histiocytes  
 c. Lymphocytes d. Neutrophils Plasma cells
13. In lepromatous leprosy, globi consist of: (Recent Question 2014)  
 a. Macrophage cells laden with acid fast bacilli  
 b. Lipid laden macrophages and degenerated tissue  
 c. Activated lymphocytes  
 d. Immunoglobulin-containing plasma cells

## SKIN TUMORS

14. Which of the following is a precancerous condition of the skin? (MH PG 2014)  
 a. Bowen disease b. Seborrheic keratosis  
 c. Leprosy d. Psoriasis
15. Mutation in malignant melanoma: (Kerala 2016), (Recent Question 2015)  
 a. N-myc b. CDKN2A  
 c. RbT d. None



## Answers with Explanations

1. Ans. (d) Most common pigmented lesions

(Ref: Robbins 9th/pg 1143)

Most common<sup>o</sup> pigmented lesions- freckles  
 Other 3 options are features of café au lait spots

2. Ans. (c) Diffuse epidermal hyperplasia

(Ref: Robbin 9th/pg 1143)

Microscopic Lesions	Definition
Acanthosis	Diffuse epidermal hyperplasia <sup>o</sup>
Hyperkeratosis	Thickening of the stratum corneum <sup>o</sup>

3. Ans. (a) Acute eczema (Ref: Robbins 9th/pg 1163; 1143)

Microscopic Lesions	Definition
Spongiosis	Intercellular edema of the epidermis <sup>o</sup>

4. Ans. (a, b, c, d, e); a. Pancytopenia; b. Nail dystrophy; c. Hyperkeratosis; d. X linked; e. Leukoplakia

(Ref: Table 41.4 Wintrobe's Clinical Hematology, 12th Edition)

### Dyskeratosis Congenita

- Inherited Bone Marrow Failure Syndrome
- Majority of patients present with pancytopenia
- Abnormal nails, reticular rash, leukoplakia
- X-linked recessive, autosomal dominant, and autosomal recessive inheritance patterns have been reported
- Patients exhibit a predisposition to bone marrow failure, malignancy, and pulmonary dysfunction.

5. Ans. (c) Psoriasis (Ref: Robbins 9th/pg 1165; 8th/pg 1185)

### Psoriasis

- Intraepidermal infiltrates of neutrophils
  - In the stratum corneum (Munro microabscesses)<sup>o</sup>
  - The spinous layer (spongiform pustules of Kogoj)<sup>o</sup>

6. Ans. (a) Antibodies against hemidesmosomes

(Ref: Lever dermatopathology/ 10<sup>th</sup> 273)

Antibody targets for Pemphigus:

Diseases	Antigens
Pemphigus Vulgaris	
Mucosal	Desmoglein 3
Mucocutaneous	Desmoglein 3 & 1
Pemphigus foliaceus	

The most common bullous lesion is bullous pemphigoid.  
 – And the most common pattern of Immuno-fluorescence is Linear IgG deposit along basement membrane. So, the best possible answer is A.

7. Ans. (a) Pemphigus

(Ref: Robbins 9th/pg 1167)

### Pemphigus Vulgaris

- Suprabasal blister formation
- Row of tombstone appearance

8. Ans. (a) Pemphigus Vulgaris

9. Ans. (b) Loss of intercellular connections

(Ref: Robbins 9th/pg 1165; 8th/pg 1168)

Acantholysis- Separation of keratinocytes- Loss of intercellular connections

Acanthosis- Diffuse epidermal hyperplasia

Spongiosis- Intercellular edema of the epidermis.

10. Ans. (b) Collagen type VII

(Ref: <http://ghr.nlm.nih.gov/condition/dystrophic-epidermolysis-bullosa>, J Med Genet. 2007 Mar; 44(3): 181–192.)

- Mutations in the COL7A1 gene seen in dystrophic epidermolysis bullosa
- COL7A1 mutations alter the structure or disrupt the production of type VII collagen, which impairs its ability to help connect the epidermis to the dermis.



# 11. Ans. (b) Pemphigus

(Ref: Robbins 9th/pg 1167)

Disease	Biopsy features	DIF
<b>Pemphigus vulgaris</b>	<ul style="list-style-type: none"> <li>Suprabasal acantholytic vesicle (tombstone pattern)<sup>Q</sup></li> <li>Mixed perivascular infiltrate with eosinophils<sup>Q</sup></li> </ul>	IgG within suprabasal intercellular spaces –fishnet pattern <sup>Q</sup>

# 12. Ans. (b) Histiocytes

(Ref: Sternberg's Diagnostic Surgical Pathology, 5th Edition. 862)

**Leptra cells<sup>Q</sup>** - large, mononuclear histiocytes (macrophages) with a foam like cytoplasm  
Seen in lepromatous leprosy

# 13. Ans. (a) Macrophage cells laden with acid fast bacilli

(Ref: Sternberg's Diagnostic Surgical Pathology, 5th Edition. 862)

**GLOBI**-Macrophage cells laden with acid fast bacilli<sup>Q</sup>  
Seen in lepromatous leprosy

# 14. Ans. (a) Bowen disease

(Ref: Clin Plast Surg. 1980 Jul;7(3):289-300.)

The most common precancerous skin lesions are actinic keratoses, Bowen's disease, and keratoacanthoma. Actinic keratoses appear over the exposed areas of the body as the result of actinic radiation. Bowen's disease is most probably secondary to the effects of internal carcinogens. Keratoacanthomas are self-limited lesions that occasionally may transform into invasive squamous cell carcinoma.

# 15. Ans. (b) CDKN2A (Ref: R 9th/pg 1148-50; 8th/pg 1172)

Melanoma is associated with mutations in **cell cycle regulators** (p16/INK4a, CDK4), growth factor receptors [e.g., **KIT**], **RAS**, **BRAF**), and **telomerase<sup>Q</sup>**

[illegible]This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.






# Central Nervous System and its Disorders

## Key Points

- » Neural tube defects are most common CNS malformations
- » Most common type of intracranial aneurysm is Berry aneurysm
- » HIV Meningoencephalitis is characterized by microglial nodules
- » Most common intracranial tumors is metastasis
- » Most common primary tumor of CNS: Meningioma
- » Most common primary intracranial tumor of CNS: Glioma (astrocytoma)
- » Most common malignant tumor of CNS: Glioma (glioblastoma multiforme)
- » Most aggressive tumor in children: Medulloblastoma
- » Most common tumor in children: Pilocytic astrocytoma

## Key Recent Updates

- » IDH<sub>1</sub> mutant glioblastomas have better prognosis
- » MC mutation in early onset familial Alzheimer's disease is PS<sub>1</sub>.



## CELLULAR PATHOLOGY OF THE CENTRAL NERVOUS SYSTEM



**Neurons and glia** of the CNS undergo a range of functional and morphologic **changes in the setting of injury**.

### Reactions of Neurons to Injury

- The **characteristic** histologic feature is **cell loss**<sup>Q</sup> and **reactive gliosis**<sup>Q</sup>

**Neuronal inclusions can be seen in**

- Viral infections**
  - Intranuclear inclusions-Herpes infection (Cowdry body)<sup>Q</sup>
  - Cytoplasmic inclusions-Rabies (Negri body)<sup>Q</sup>
  - Both nucleus and cytoplasm-Cytomegalovirus infection.<sup>Q</sup>
- Intracytoplasmic inclusions**<sup>Q</sup> in neurons,
  - Neurofibrillary tangles<sup>Q</sup> of Alzheimer's disease
  - Lewy bodies<sup>Q</sup> of Parkinson's disease
  - Lafora bodies<sup>Q</sup> in myoclonic epilepsy









## Reactions of Astrocytes to Injury

- **Gliosis**<sup>Q</sup> is the **most important histopathologic**<sup>Q</sup> indicator of **CNS injury** and is characterized by both **hypertrophy** and **hyperplasia** of astrocytes.




- **The Alzheimer type II astrocyte**<sup>Q</sup>:
  - Unrelated to Alzheimer's disease.<sup>Q</sup>
  - It is a gray matter cell with large nucleus with **intranuclear glycogen droplet**<sup>Q</sup>



- Seen in **Long-standing hyperammonemia** due to **chronic liver disease**, **Wilson's disease**, or **hereditary metabolic disorders of the urea cycle**.<sup>Q</sup>
- **Rosenthal fibers**:

- Brightly **eosinophilic structures / inclusions** within **cytoplasm** of<sup>Q</sup> astrocytic processes
- Contain two **heat-shock proteins** ( $\alpha$ B-crystallin and HSP27) as well as **ubiquitin**<sup>Q</sup>




- Seen in **Alexander's disease**, **pilocytic astrocytoma**<sup>Q</sup>.
- **Corpora amylacea**:
  - **Periodic acid-Schiff (PAS)**-positive,<sup>Q</sup> concentrically **lamellated structures** and contains **glycosaminoglycan** polymers, as well as **heat-shock proteins** and **ubiquitin**.
  - Seen in **advancing age** and are thought to represent a **degenerative change** in the Astrocyte.<sup>Q</sup>

## Reactions of Microglia to Injury

Microglia-Resident **macrophages** of the CNS<sup>Q</sup>.

They respond to injury by<sup>Q</sup>

- Developing elongated nuclei (**rod cells**)<sup>Q</sup>, as in **neurosyphilis**
- Forming **aggregates** around small foci of tissue necrosis → **microglial nodules**
- Forming **aggregates** around cell bodies of dying neurons → **neuronophagia**<sup>Q</sup>.




## Reactions of Other Glial Cells to Injury

- **Oligodendroglial nuclei** show viral inclusions in **progressive multifocal leukoencephalopathy**<sup>Q</sup>
- Ependymal cells-ciliated **columnar**<sup>Q</sup> epithelial cells **lining the ventricles**
- **Viral inclusions** in ependymal cells-CMV<sup>Q</sup>

## CEREBRAL HERNIATION


- Most common cause of herniation is increased intracranial pressure.





## Types of Herniation

- **Transtentorial:**
  - Occurs when the **medial aspect of the temporal lobe is compressed against the free margin of the tentorium**<sup>Q</sup>
  - **Duret hemorrhages**<sup>Q</sup> -linear or flame-shaped lesions<sup>Q</sup> -due to distortion or tearing of **penetrating veins and arteries**<sup>Q</sup> supplying the upper brainstem
- **Transfalcine:**
  - Caused by **asymmetric expansion of a cerebral hemisphere**<sup>Q</sup>
  - Displaces the cingulate gyrus-compresses the **anterior cerebral artery**<sup>Q</sup> and its branches.
- **Tonsillar:**
  - These are caused by displacement of the **cerebellar tonsils**<sup>Q</sup> through the **foramen magnum**<sup>Q</sup>.
  - Also occur if a **lumbar puncture (LP)** is performed in a **patient with increased intracranial pressure**<sup>Q</sup>.
  - May **compress the medulla and respiratory centers, causing death**<sup>Q</sup>.



## MALFORMATIONS AND DEVELOPMENTAL DISORDERS

### NEURAL TUBE DEFECTS


- Neural tube defects are most common CNS malformations
- **Most common site** of neural tube defects is the **spinal cord**<sup>Q</sup>

### Types of Neural Tube Defects

**Spinal dysraphism or spina bifida**-can be of two types:

- **Spina bifida occulta**<sup>Q</sup> -can be **asymptomatic bony defect**

- **Spina bifida aperta**<sup>Q</sup> disorganized segment of spinal cord, associated with an overlying meningeal outpouching
- **Myelomeningocele**-extension of **CNS tissue and meninges** through a defect in the vertebral column
- **Meningocele**-applies when there is only a **meningeal extrusion**.




### Encephalocele

- Diverticulum of **malformed brain tissue** extending through a **defect in the cranium**.
- Most common in **posterior fossa**<sup>Q</sup> **cribriform plate in the anterior fossa**




### Anencephaly

- Malformation of the anterior end of the neural tube, with absence of most of the brain and calvarium.
- Instead, there is a mass of disorganized glial tissue with vessels in this area called a **cerebrovasculosa**<sup>Q</sup>
- Neural tube defects are associated with **↑Alpha fetoprotein**.
- Mutation in **DOUBLECORTIN (Dcx)** results in Lissencephaly in males and subcortical band heterotopia in females






## FOREBRAIN ANOMALIES

Lissencephaly	Polygyria	Holoprosencephaly	Arrhen- cephaly	Agenesis of the corpus callosum	Neuronal Heterotopia
↓ gyri	Small ↑ gyri	Incomplete separation of cerebral hemispheres across midline	No olfactory nerves	<ul style="list-style-type: none"> <li>No white matter projections from one hemisphere to the other</li> <li><b>Bat wing anomaly</b></li> </ul>	Neurons in inappropriate locations along lines of migration
					
Lissencephaly	Polymicrogyria	Holoprosencephaly			


## POSTERIOR FOSSA ANOMALIES

Developmental abnormalities of the brain include the **Arnold-Chiari malformation**, the **Dandy-Walker malformation**<sup>Q</sup>

- **Arnold Chiari:**
  - Small posterior fossa
  - Herniation of cerebellum & 4th ventricle into foramen magnum
  - Associated with hydrocephalus and lumbar myelomeningocele



- **Dandy Walker:**
  - Enlarged posterior fossa
  - Severe hypoplasia/Absence of cerebellar vermis
  - Cystic dilation of 4th ventricle hydrocephalus and agenesis of corpus callosum



## Mnemonic

**Dandy-Walker syndrome: Components "Dandy Walker Syndrome":**

- Dilated 4th ventricle
- Water on the brain
- Small vermis



## High Yield Facts

- **Chiari type I malformation** is a less severe disorder in which low lying cerebellar tonsils extend down into the vertebral canal.<sup>Q</sup>
- Lucid interval is seen in epidural hemorrhage
- Worst headache is seen in SAH
- Self limited bleeding is SDH
- **Hypertension** MC causes **deep brain parenchymal hemorrhages**<sup>Q</sup>
- **Chronic hypertension** leads to development of **Charcot-Bouchard microaneurysms**<sup>Q</sup>
- **Cerebral amyloid angiopathy (CAA)**<sup>Q</sup> MC causes of **lobar hemorrhages**<sup>Q</sup>.
- Cerebral amyloid angiopathy (AB<sub>40</sub>)-MC cause of total hemorrhage



## Latest Update

- **Joubert syndrome:** Hypoplasia of the cerebellar vermis with elongation of the superior cerebellar peduncles and an altered shape of the brainstem; together these changes give rise to the '**molar tooth sign**'<sup>Q</sup> on imaging.
- Mutations affect genes that encode components of the **primary (non-motile) cilium**.<sup>Q</sup>

## CEREBRAL HEMORRHAGE

It can be in epidural, subdural, subarachnoid, and intraparenchymal compartments



Site of Hemorrhage	Cause
• Epidural	• Rupture of dural arteries MC-middle meningeal artery
• Subdural	• Rupture of bridging veins
• Subarachnoid	• Rupture of Berry aneurysm in cerebral artery
• Ultra parenchymal	• Due to hypertension and cerebral amyloid angiopathy

### Subarachnoid Hemorrhage (SAH)


- MC cause is rupture of a saccular ("Berry") aneurysm in a cerebral artery.<sup>Q</sup>
- "Worst headache ever"<sup>Q</sup>



### High Yield Facts

#### Berry Aneurysm

- Developmental abnormalities<sup>Q</sup>
- Due to **the structural abnormality of the involved vessel (absence of smooth muscle and intimal elastic lamina.)**<sup>Q</sup>
- Called **congenital**<sup>Q</sup>, although the aneurysm itself is **not present at birth**<sup>Q</sup> but develop over time
- Majority occur sporadically<sup>Q</sup>,
- Increased incidence in autosomal dominant polycystic kidney disease, Ehlers-Danlos syndrome type IV, neurofibromatosis type 1 [NF1], and Marfan's syndrome, fibromuscular dysplasia of extracranial arteries,<sup>Q</sup> and coarctation of the aorta.<sup>Q</sup>
- 90% found **near major arterial branch**<sup>Q</sup> points in the **anterior circulation**<sup>Q</sup>
- The chance of rupture increases with age (**rupture is rare in childhood**<sup>Q</sup>).



### Intraparenchymal Hemorrhage

- **Ganglionic hemorrhages**<sup>Q</sup>: Hemorrhage in the **basal ganglia and thalamus**
- **Lobar hemorrhages**<sup>Q</sup>: Hemorrhage is in the lobes of the **cerebral hemispheres**
- The two major causes: **Hypertension and cerebral amyloid angiopathy**.<sup>Q</sup>
- This deposition of **A $\beta$ 40** weaken the vessel wall and lead to hemorrhage.
- The presence of either  **$\epsilon$ 2 or  $\epsilon$ 4 allele**<sup>Q</sup> increases the risk of repeat bleeding.



### Latest Update

- **Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL)**-autosomal dominant disorder caused by mutations in the **NOTCH3 gene**.<sup>Q</sup>
- The disease is characterized clinically by recurrent strokes (usually infarcts, less often hemorrhages) and dementia.<sup>Q</sup>

### INTRACRANIAL ANEURYSMS

- **Charcot-Bouchard aneurysms:**
  - Results from weakening of the wall of cerebral artery by **lipohyalinosis**<sup>Q</sup> (deposition of lipids and hyaline material) caused by **hypertension**<sup>Q</sup>.
- **Saccular aneurysms (Berry aneurysms):**
  - **Most common type of intracranial aneurysm**.<sup>Q</sup>
  - Result of **congenital defects in the media of blood vessels**
  - Located at the **bifurcations of arteries**.
- **Atherosclerotic aneurysms:**
  - **Fusiform (spindle-shaped)**<sup>Q</sup> aneurysms
  - Located in **the major cerebral vessels**, most often found in the **anterior circulation**<sup>Q</sup>
  - They **rarely rupture**<sup>Q</sup>, but may become **thrombosed**.<sup>Q</sup>
- **Mycotic (septic) aneurysms:**
  - Result from **septic emboli**<sup>Q</sup>, most commonly from **subacute bacterial endocarditis**.<sup>Q</sup>
  - Causes **cerebral infarction**<sup>Q</sup>, rather than **subarachnoid hemorrhage**<sup>Q</sup>

### POINTS TO REMEMBER

- Hypertensive hemorrhage shows a **predilection for the distribution of the lenticulostriate arteries (branch of middle cerebral artery)** with small (lacunar) hemorrhages, or large hemorrhages **obliterating the corpus striatum**, including the **putamen and internal capsule**.<sup>Q</sup>

### CNS INFECTIONS

#### Meningitis

Inflammatory process of the leptomeninges and CSF within the subarachnoid space.





## Causative Organisms of Meningitis<sup>Q</sup>


Type	Age group	Etiology
Acute Pyogenic Meningitis	Neonates	<i>Escherichia coli</i> (India), group B streptococci (World)
	Infants	<i>S.pneumoniae</i> , <i>Haemophilus influenzae</i>
	Adolescents & young adults	<i>S.pneumoniae</i> , <i>Neisseria meningitidis</i>
	Elderly	<i>Streptococcus pneumoniae</i> <i>Listeria monocytogenes</i>
	Immunosuppressed individual	<i>Klebsiella</i> or anaerobic organisms
Chronic Meningitis		Tuberculous, spirochetal or cryptococcal
Aseptic meningitis		Viruses like enteroviruses

## CSF Findings in CNS Infections

Parameters	Normal values	Bacterial Meningitis	Tuberculous Meningitis	Viral Meningitis
Pressure	50–180 mm water	Raised	Raised	Raised
Gross appearance	Clear and colorless	Turbid <sup>Q</sup>	Clear (may clot) <sup>Q</sup>	Clear <sup>Q</sup>
Protein	20–50 mg/dL	High	Very High <sup>Q</sup>	Slightly high
Glucose	40–70 mg/dL	Very low <sup>Q</sup>	Low	Normal
Chloride	110–125 mEq/L	Low	Very low <sup>Q</sup>	Normal
Cells	< 5/microlitre	Neutrophils 1,000–1,00,000 neutrophils/uL	Pleocytosis 100-1000 mononuclear/uL	Lymphocytosis <sup>Q</sup> 10-100 mononuclear/uL

## Human Immunodeficiency Virus (HIV)

- Directly cause **meningoencephalitis**,<sup>Q</sup>
- Indirectly-increases the risk of **opportunistic infections**<sup>Q</sup> (toxoplasmosis, CMV) or EBV-positive CNS **lymphoma**.<sup>Q</sup>
- Meningoencephalitis-characterized by **microglial nodules**<sup>Q</sup> composed of **mononuclear cells**, **microglia**, and **scattered multinucleated giant cells**, usually found **near small blood vessels**<sup>Q</sup>, which shows **prominent endothelial cells** and **perivascular foamy or pigment-laden macrophages**<sup>Q</sup>
- HIV-associated central nervous system (CNS) lymphoma is a **diffuse, large-cell non-Hodgkin's lymphoma**<sup>Q</sup> of **B-cell origin**<sup>Q</sup> that usually occurs in the brain (rarely in the spinal cord).
- It is a **late complication**<sup>Q</sup> of HIV infection.
- **Epstein-Barr virus (EBV)**<sup>Q</sup> is identified in almost **all cases**<sup>Q</sup>.




Microglial nodules in HIV


## Rabies Virus

- Rabies is **only communicable disease of man** that is **always fatal**.<sup>Q</sup>
- Rabies is **dead end infection**<sup>Q</sup> caused by **enveloped, RNA (negative sense single stranded) virus**<sup>Q</sup>.
- Most characteristic pathologic finding in CNS is the formation of **cytoplasmic inclusion bodies** called **Negri bodies**<sup>Q</sup>.
- The **prominence of early brain stem dysfunction**<sup>Q</sup> distinguish it from other viral encephalitis.
- Diagnosis is made by **detection of rabies virus antigen by immunofluorescence**<sup>Q</sup>.
- **Antemortem specimen: corneal smear**<sup>Q</sup>, skin biopsy from neck or saliva.
- **Postmortem: Brain biopsy**<sup>Q</sup>
- **A definitive pathologic diagnosis of rabies can be based on the finding of negri bodies in the brain or spinal cord**<sup>Q</sup>.

Dense eosinophilic inclusion in Purkinje cells



Negri body S/o Rabies



## High Yield Facts

### Negri bodies

- Composed of finely fibrillar matrix (ribonuclear proteins produced by the virus) and rabies virus particles
- Found **most abundantly in cerebellum (purkinje cells) & pyramidal neurons of hippocampus**. They are also seen in neurons of Ammon's horn, cerebral cortex, brainstem, hypothalamus, and dorsal spinal ganglia.


## Neurosyphilis

- Tertiary stage of syphilis<sup>Q</sup>**, includes **syphilitic meningitis, paretic neurosyphilis, and tabes dorsalis<sup>Q</sup>**

Meningovascular Neurosyphilis	Paretic Neurosyphilis	Tabes Dorsalis
<ul style="list-style-type: none"> <li><b>Obliterative endarteritis (Heubner arteritis)<sup>Q</sup></b></li> <li>Perivascular infiltrates of lymphocytes and plasma cells<sup>Q</sup></li> <li><b>Cerebral gummas (plasma cell-rich mass lesions)<sup>Q</sup></b> can also occur.</li> </ul>	<ul style="list-style-type: none"> <li>Occurs due to <b>invasion of the brain by T. pallidum.<sup>Q</sup></b></li> <li>Characterized by <b>delusions of grandeur<sup>Q</sup></b> that terminate in <b>severe dementia</b> (general paresis of the insane).</li> <li>The lesions are characterized by <b>loss of neurons, proliferation of microglia (rod cells)<sup>Q</sup>, gliosis, and iron deposits.<sup>Q</sup></b></li> </ul>	<ul style="list-style-type: none"> <li>Result of <b>degeneration of the posterior columns of the spinal cord.<sup>Q</sup></b></li> <li>Impaired joint position sensation, ataxia<sup>Q</sup></li> <li>Loss of pain sensation (leading to joint damage, i.e. Charcot joints)<sup>Q</sup></li> <li><b>Argyll Robertson pupils</b> (pupils that react to accommodation but not to light).<sup>Q</sup></li> </ul>

## CMV

- Enlarged cells (cytomegaly) with intranuclear and intracytoplasmic inclusions are seen with cytomegalovirus infection.<sup>Q</sup>
- Owl eyed inclusions seen in CMV<sup>Q</sup>



### Features

- Cytomegaly
- Intranuclear inclusions (cowdry type)
- Cytoplasmic inclusions
- S/o CMV**

## PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY (PML)

- Demyelinating disease caused by the **JC polyomavirus<sup>Q</sup>**, preferentially infects **oligodendrocytes<sup>Q</sup>**.
- Occurs in **immunosuppressed individuals<sup>Q</sup>**.
- The **pathognomonic feature** of PML is **"ground-glass" appearance of oligodendrocyte nuclei<sup>Q</sup>**.

## PRION DISEASES


- Prions are **abnormal forms of a cellular protein<sup>Q</sup>**
- Can be **sporadic, familial or transmitted.<sup>Q</sup>**

Humans <sup>Q</sup>	Animals <sup>Q</sup>
<ul style="list-style-type: none"> <li>Creutzfeldt-Jakob disease</li> <li>Gerstmann-Sträussler-Scheinker syndrome</li> <li>Fatal familial insomnia</li> <li>Kuru</li> </ul>	<ul style="list-style-type: none"> <li>Scrapie in sheep</li> <li>Mink-transmissible encephalopathy</li> <li>Chronic wasting</li> <li>Disease of deer and elk</li> <li>Bovine spongiform encephalopathy</li> </ul>

- Clinically:** Rapidly progressive dementia
- Morphologically:** **Spongiform change<sup>Q</sup> without inflammation** hallmark of all prion diseases except **fatal familial insomnia,<sup>Q</sup> which is characterized by neuronal loss and reactive gliosis.<sup>Q</sup>**
- Pathology:**
  - Normal PrP is a 30-kD cytoplasmic protein present in neurons.<sup>Q</sup>
  - Disease occurs when PrP undergoes a conformational change from its normal alpha-helix-containing isoform (PrPc) to an abnormal B-pleated sheet isoform, usually termed PrPsc.<sup>Q</sup>
  - PrP acquires resistance to digestion with proteases<sup>Q</sup>

## High Yield Facts

- Vasculitis is characteristically absent in CNS involvement in AIDS
- Spinal cord involvement in AIDS leads to vacuolar myelopathy
- Herpes simplex virus produces Cowdry type A intranuclear inclusions<sup>Q</sup> in neurons and glial cells.<sup>Q</sup>
- Subacute sclerosing panencephalitis (SSPE)
  - Progressive clinical syndrome
  - Characterized by cognitive decline, spasticity of limbs, and seizures.
  - Occurs in children after an initial, early-age acute infection with measles.<sup>Q</sup>
- Babes nodules<sup>Q</sup> are seen in rabies encephalomyelitis



- **Prion Diseases:** PrP<sup>Sc</sup>, independent of the means from which it originates<sup>Q</sup>, then facilitates, in a **cooperative fashion**<sup>Q</sup>, the conversion of other PrP<sup>C</sup> molecules to PrP<sup>Sc</sup> molecules.
- This transformation can occur sporadically or in familial fashion.
- **Genetics:** Mutations in the gene encoding PrP<sup>C</sup> (PRNP).<sup>Q</sup>




Image shows spongiform change s/o prion disease

- **Diagnosis of Choice:** Western blotting of tissue extracts after partial digestion with proteinase K, **diagnostic**.<sup>Q</sup>

**Latest Update**

**Prion Diseases**  
A polymorphic locus in PRNP (codon 129 may be either Met or Val)<sup>Q</sup> Homozygosity increases risk of sporadic disease.<sup>Q</sup>

### Creutzfeldt-Jakob Disease (CJD)

- **Most common** prion disease<sup>Q</sup>
- Peak incidence in the **seventh decade**.<sup>Q</sup>
- **Familial forms**-mutations in PRNP<sup>Q</sup>
- Iatrogenic transmission-<sup>Q</sup>
  - **Corneal, transplantation<sup>Q</sup>, Deep implantation of electrodes in the brain<sup>Q</sup>, Administration of contaminated preparations of naturally derived human growth hormone<sup>Q</sup>**
- **Clinically-** dementia, startle myoclonus<sup>Q</sup>

### Variant Creutzfeldt-Jakob Disease

- **Young adults**<sup>Q</sup>
- **No alterations in the PRNP gene** are present<sup>Q</sup>
- Clinically-**behavioral disorders** prominently in the early stages<sup>Q</sup>
- **Extensive cortical plaques** surrounded by a "halo" of spongiform change<sup>Q</sup>

## NEURODEGENERATIVE DISEASES

- Diseases that affect the gray matter of brain with **damage to the neurons**.<sup>Q</sup>
- The **pathologic hallmark**<sup>Q</sup>-accumulation of **protein aggregates**<sup>Q</sup>, hence the use of the term "**proteinopathy**"<sup>Q</sup>.
- The protein aggregates are recognized histologically as **inclusions-diagnostic hallmark**<sup>Q</sup> of the disease.
- The protein aggregates typically are **resistant to degradation and are directly toxic to neuron**<sup>Q</sup>

Disease	Clinical pattern	Inclusions	Genetic causes
<b>Alzheimer's Disease</b>	Dementia	A $\beta$ (plaques) <sup>Q</sup> Tau (tangles) <sup>Q</sup>	APP, PS1, PS2 <sup>Q</sup>
<b>Parkinson's disease (PD)</b>	Hypokinetic movement disorder	<b><math>\alpha</math>-synuclein</b> <sup>Q</sup> Tau	<b><math>\alpha</math>-synuclein</b> <sup>Q</sup> LRRK2
<b>Huntington's disease (HD)</b>	Hyperkinetic movement disorder	Huntington <sup>Q</sup> (polyglutamine)	Htt
<b>Amyotrophic lateral sclerosis (ALS)</b>	Weakness with upper and lower motor neurons signs	SOD1 <sup>Q</sup> TDP-43 FUS	SOD1 TDP-43, C9orf72 FUS

### Alzheimer's Disease


- **Most common cause of dementia in elderly**<sup>Q</sup> (followed by **vascular multi-infarct dementia**)<sup>Q</sup>
- The **hallmark abnormality**<sup>Q</sup>-accumulation of two proteins (A $\beta$  and tau) in specific brain regions
- The etiology of AD is not well understood, Main risk factor- **age**<sup>Q</sup>
- Grossly, **cortical atrophy (narrowed gyri and widened sulci)**<sup>Q</sup> is predominant in **frontal, temporal, and parietal lobes**
- The **pathologic hallmark**<sup>Q</sup> are plaques and tangles.



### High Yield Facts

- Tangles are **not**<sup>Q</sup> specific to AD, being found in other diseases as well.
- AD is associated with decrease in cerebral cortical level of acetylcholine. This degeneration occurs in **nucleus basalis of Meyernet**<sup>Q</sup>
- Vascular amyloid in **cerebral amyloid angiopathy** is **A $\beta$ 40**.<sup>Q</sup>
- **Medial temporal lobe, including hippocampus, entorhinal cortex and amygdala**<sup>Q</sup>, are involved early in the course.
- **Mutations in the gene for tau cause frontotemporal lobar degenerations**<sup>Q</sup> rather than AD
- Beta-amyloid deposition is **necessary but not sufficient**<sup>Q</sup> for the development of Alzheimer's disease
- **Hirano bodies**<sup>Q</sup> and **Granulovacuolar degeneration**<sup>Q</sup> are other features seen in AD





**Latest Update**

1. Out of **APP**, **PSEN1** and **PSEN2** mutations in early-onset Alzheimer's disease occur in the order **PS1 > APP > PS2**.
2. Mutations in **PS-1** are the most common cause of early-age-of-onset **FAD**, representing perhaps 40–70% of all cases.
3. **ApoE4** remains the single most important biological marker associated with **AD** risk.


**Latest Update**

**Diagnosis:** Demonstrate **Aβ deposition** in the brain through **18F-labeled amyloid binding compounds on PET scan**.<sup>Q</sup>

**Biomarkers:** Increased phosphorylated tau and reduced **Aβ** in the **CSF**<sup>Q</sup> provide evidence of neuronal degeneration associated with AD.

**Plaques**


- Deposits of aggregated **Aβ** peptides in the neuropil.
- Plaques are of two types diffuse and neuritic (senile).
- **Neuritic plaques** contain both **Aβ 40** and **Aβ 42**,<sup>Q</sup> **Diffuse plaques** are predominantly made up of **Aβ 42**.<sup>Q</sup>
- **Diffuse plaques** –seen in brain of **trisomy 21**<sup>Q</sup> patients



Plaques are aggregated Ab

**Tangles**

- Aggregates of the microtubule binding protein **tau**.<sup>Q</sup>
- They are demonstrated by **silver (Bielschowsky) staining**<sup>Q</sup>
- They are commonly found in **cortical neurons**, especially in the **entorhinal cortex**, as well as in other sites such as **pyramidal cells of the hippocampus**<sup>Q</sup>, the amygdala, the basal forebrain, and the raphe nuclei<sup>Q</sup>



Neurofibrillary tangles are aggregates of tau protein


## Lewy Body Disorders

- Lewy bodies are intracytoplasmic eosinophilic inclusions.
- Major component of the Lewy body is **alpha-synuclein**<sup>Q</sup>, others being neurofilament antigens, parkin, and ubiquitin.
- The histologic features of Lewy body disorders depends on location of lewy bodies

Disease	Location	Symptoms
Classic Parkinson's disease	Nigrostriatal system <sup>Q</sup>	Extrapyramidal movement disorder
Lewy body dementia	Cerebral cortex <sup>Q</sup>	<b>Third most common cause of dementia</b> <sup>Q</sup>
Shy-Drager syndrome	Sympathetic neurons in the spinal cord <sup>Q</sup>	Autonomic dysfunction

**TUMORS**

- **Most common tumor** of CNS- **metastasis**<sup>Q</sup>
- Most common **primary** tumor of CNS- **meningioma**<sup>Q</sup>
- Most common **primary intracranial tumor** of CNS- **glioma (astrocytoma)**<sup>Q</sup>
- Most common **malignant** tumor of CNS- glioma (**glioblastoma**)<sup>Q</sup>
- Most common **malignant** tumor in **children**- **medulloblastoma** closely followed by **pilocytic astrocytoma**<sup>Q</sup>
- Most common primary CNS tumor in children: **Pilocytic astrocytoma**.



## GLIOMAS


- These are **most common group of primary brain tumors**.
- These include **astrocytoma, oligodendroglioma and ependymoma**.

### Astrocytoma

- Most common primary malignant brain tumors in adults**.
- The two major categories of astrocytic tumors are
  - Localized astrocytomas**\*: Most common are the **pilocytic astrocytomas**.<sup>Q</sup> (WHO grade I).
  - Diffusely Infiltrating Astrocytomas**: Now classified as IDH wildtype and IDH mutant Glioblastoma

#### Localized Astrocytomas, Pilocytic Astrocytoma

- WHO grade I**
- Occurs in **children and young**
- Located in the **cerebellum** but may also appear in the **floor and walls of the third ventricle, the optic nerves**.
- Microscopically: Tumors are often biphasic, with both loose "microcystic" and fibrillary areas.
- Rosenthal fibers and eosinophilic granular bodies, are characteristic findings.




Rosenthal fibers are thick elongated worm-like or "corkscrew" eosinophilic (pink) bundle that is found on H&E staining of the brain

### Diffusely Infiltrating Astrocytomas

- Range from diffuse astrocytoma (grade II) to anaplastic astrocytoma (grade III) to glioblastoma (grade IV).
- There are **no WHO grade I** infiltrating astrocytomas.

### Glioblastoma (WHO Grade IV Astrocytoma)


- A **highly malignant** tumor characterized histologically by **serpentine areas of necrosis**<sup>Q</sup> surrounded by **peripheral palisading of tumor cells and vascular/endothelial cell proliferation**.<sup>Q</sup>
- The minimal criterion for **vascular/endothelial cell proliferation**<sup>Q</sup> is a double layer of endothelial cells.
- Massive endothelial proliferation can lead to formation of **glomeruloid body**.<sup>Q</sup>
- It frequently crosses midline ("**butterfly tumor**").<sup>Q</sup>



### 2016 WHO CNS Tumor Classification

	IDH-wildtype glioblastoma	IDH-mutant glioblastoma
<b>Synonym</b>	Primary glioblastoma	Secondary glioblastoma
<b>Proportion of glioblastomas</b>	~90%	~10%
<b>Median age at diagnosis</b>	~62 years	~44 years
<b>Location</b>	Supratentorial	Preferentially frontal
<b>Necrosis</b>	Extensive	Limited
<b>Mutations</b>	TERT (72%) EGFR (35%) PTEN (25%)	P53 (80%) ATRX (70%)
<b>Prognosis</b>	Bad	Good





R10<sup>th</sup>

## Latest Update

### Molecular Genetics of malignant gliomas:


Genomically malignant gliomas are divided into four molecular subtypes: Classic, proneural, neural, and mesenchymal.<sup>Q</sup>

- **Classic subtype**-majority of primary glioblastoma mutations of the PTEN, deletions of chromosome 10, amplification of the EGFR oncogene<sup>Q</sup>
- **Proneural**-most common type associated with secondary glioblastoma<sup>Q</sup>; P53, and IDH, and IDH<sub>2</sub> mutations.
- **Neural type**-neuronal markers, including NEFL, GABRA1, SYT1, and SLC12A5.<sup>Q</sup>
- **Mesenchymal type** is characterized by deletions of the NF1 gene on chromosome 17 and TNF pathway genes, are highly expressed<sup>Q</sup>

## Oligodendroglioma

- **WHO grade II lesions**<sup>Q</sup>
- Most common in the **fourth and fifth decades**.<sup>Q</sup>
- **Mostly in the cerebral hemispheres, with a predilection for white matter**<sup>Q</sup>
- **The most common genetic alterations are mutations of the isocitrate dehydrogenase genes (IDH1 and IDH2)**-present in up to **90% of oligodendrogliomas**<sup>Q</sup>
- **Deletions of portions of chromosomes 1p and 19q, typically occurring together as a co-deletion, are seen in up to 80% of cases.**<sup>Q</sup>
- Cytogenetic abnormalities have therapeutic significance for this type of tumor, as only tumors involving 19q or 1p respond to chemotherapy
- **Microscopy:** Sheets of cells with clear halos ("fried-egg" appearance)<sup>Q</sup>
- **Calcification, present in up to 90% of these tumors.**<sup>Q</sup>
- **Perineuronal satellitosis.**<sup>Q</sup>

Fried egg appearance




Oligodendroglioma

## Ependymoma

- **WHO grade II lesions**<sup>Q</sup>
- These **most often** arise next to the ependyma-lined ventricular system, including central canal of spinal cord.
- **In first two decades**-, these typically occur *near fourth ventricle*<sup>Q</sup>
- **In adults, spinal cord**<sup>Q</sup> is the most common site.
- Spinal cord ependymoma frequently occur in setting of **neurofibromatosis type 2**.<sup>Q</sup>
- **Microscopy: Perivascular pseudorosettes**<sup>Q</sup>
- **CSF dissemination is a common occurrence**<sup>Q</sup>
- **Myxopapillary ependymomas-occur in the filum terminale of the spinal cord and contain papillary elements in a myxoid background, admixed with ependymoma-like cells.**<sup>Q</sup>

Perivascular pseudorosette



Ependymoma

## NEURONAL TUMORS

- **Gangliogliomas:**
  - **Most common** of the neuronal tumors of the CNS
  - **Most commonly found in the temporal lobe.**<sup>Q</sup>
- **Dysembryoplastic neuroepithelial tumor:**
  - Low-grade (WHO Grade I) tumor of childhood that often presents as a seizure disorder
  - Well-differentiated "**floating neurons**"<sup>Q</sup> that sit in the pools of mucopolysaccharide-rich fluid of the myxoid background
- **Central neurocytoma:**
  - **WHO Grade II**, most commonly the lateral or third ventricle
  - **Neurocytic rosette**<sup>Q</sup> are characteristic feature

## POORLY DIFFERENTIATED NEOPLASMS

- Medulloblastoma
- Atypical teratoid/rhabdoid tumor





## High Yield Facts

### Glial fibrillary acidic protein (GFAP):

- Intermediate filament expressed in astrocytes, ependymal cells and oligodendrocytes.<sup>Q</sup>

### Few GFAP positive tumors<sup>Q</sup>

- Astrocytomas
- Oligodendroglioma
- Ependymoma
- Medulloblastoma
- Pituitary adenoma
- Carcinoma of choroid plexuses
- Hemangioblastoma

## Medulloblastoma

- WHO grade IV.<sup>Q</sup>**
- Occurs predominantly in children and exclusively in the cerebellum<sup>Q</sup>**
- Largely undifferentiated tumor, appears as **Small round blue cell tumor<sup>Q</sup>**
- Mitoses are abundant, and markers of cellular proliferation, such as Ki-67, are detected in a high percentage of the cells<sup>Q</sup>**
- Homer-Wright rosettes<sup>Q</sup>**
- Dissemination through the CSF** is a common complication-giving rise to **nodular masses** at some distance from the primary tumor called as "drop metastases."<sup>Q</sup>
- Exquisitely radiosensitive<sup>Q</sup>**




Image shows Homer-Wright rosettes

## 2016 CNS TUMOR CLASSIFICATION-MEDULLOBLASTOMA-Genetic profile predicts prognosis

Genetic profile	Prognosis
WNT-Activated	Best
SHH-activated TP53-mutant	High-risk tumor of infancy
SHH-activated, TP53-wildtype	Low-risk tumor of infancy
Non-WNT/non-SHH, group 3	<ul style="list-style-type: none"> <li>myc amplification</li> <li><b>WORST prognosis</b></li> </ul>
Non-WNT/non-SHH, group 4	Intermediate prognosis (17q alteration)

## Atypical Teratoid/Rhabdoid Tumor


- WHO grade IV**
- Young children**
- Arise in **posterior fossa and supratentorial compartments<sup>Q</sup>**
- Highly malignant tumor**
- Genetic alterations in **chromosome 22** (>90% of cases) are a **hallmark** of rhabdoid tumor<sup>Q</sup>
- MC mutation INI-1

## PRIMARY CNS LYMPHOMA

- Primary CNS lymphoma (PCNSL)** are uncommon tumors, accounting for only 1% of malignant CNS tumors
- Most common** CNS neoplasm in HIV patient<sup>Q</sup>
- Predisposing factors include:
  - HIV/AIDS: approximately 2-6% of patients with HIV will develop PCNSL<sup>Q</sup>, Prior EBV infection, Post transplantation, IgA deficiency, Wiskott-Aldrich syndrome
- The vast majority (>90%) of PCNSL are **B cell in origin<sup>Q</sup>**: **diffuse large B-cell lymphoma** is most common followed by high-grade, Burkitt-like B-cell lymphoma
- Malignant cells tend to accumulate around blood vessels.
- Chemotherapy is highly effective<sup>Q</sup>**


## MENINGIOMA

- Benign tumors of Adults; Most common primary brain tumor
- Attached to the dura,<sup>Q</sup> they commonly arise along the venous sinuses (parasagittal, sphenoid wings, and olfactory groove).
- Arise from meningotheial cells of the arachnoid<sup>Q</sup>
- Risk factors: Prior radiation therapy to the head and neck, typically decades earlier<sup>Q</sup>
- Meningiomas often express progesterone receptors and may grow more rapidly during pregnancy<sup>Q</sup>
- Most common cytogenetic abnormality is loss of region 12 on chromosome 22q<sup>Q</sup>
- Common lesion in the setting of NF2<sup>Q</sup>
- They may also grow en plaque, in which the tumor spreads in a sheet-like fashion along the surface of the dura-This form is commonly associated with hyperostotic reactive changes in the adjacent bone.<sup>Q</sup>



Gross: meningioma


Psammoma bodies



## Mnemonic

### Tumors with Calcifications-COM

- **C**raniopharyngioma-suprasellar calcification.<sup>Q</sup>
- **O**ligodendroglioma<sup>Q</sup>
- **M**eningioma



Wet  
keratin

Craniopharyngioma shows wet keratin

## Metastatic Tumors

These are **most common intracranial tumors**. Five most common sites of metastasis are, Lung, Breast, Skin (melanoma), Kidney, GIT

Site of Primary Tumor	Brain Metastases, %	Leptomeningeal Metastases, %	Spinal Cord Compression, %
<b>Lung</b>	40	24	18
<b>Breast</b>	19	41	24
<b>Melanoma</b>	10	12	4
<b>Gastrointestinal tract</b>	7	13	6
<b>Genitourinary tract</b>	10	12	4

## NERVE SHEATH TUMORS

### Peripheral Nerve Sheath Tumor


These include the three common types- schwannoma, neurofibroma, and malignant peripheral nerve sheath tumor (MPNST).<sup>Q</sup>

Features	Schwannomas	Neurofibromas			
<b>General characteristics</b>	<ul style="list-style-type: none"> <li>• Benign tumors</li> <li>• Arise directly from peripheral nerves.<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Benign tumors</li> <li>• Heterogeneous in composition.</li> <li>• The neoplastic Schwann cells admixed with perineurial like cells, fibroblasts, mast cells, and CD34+ spindle cells<sup>Q</sup></li> </ul>			
<b>Gross</b>	Well-circumscribed, encapsulated	Non-encapsulated mass <sup>Q</sup>			
<b>Morphology</b>	<ul style="list-style-type: none"> <li>• Antoni A- cellular areas</li> <li>• Antoni B- loose edematous areas</li> <li>• Verocay bodies<sup>Q</sup> foci of palisaded nuclei in cellular areas</li> </ul>	<p>3 types of neurofibroma are there:</p> <table> <tr> <td> <b>Localized</b> <ul style="list-style-type: none"> <li>• Low cellularity<sup>Q</sup></li> </ul> </td><td> <b>Diffuse</b> <ul style="list-style-type: none"> <li>• Diffusely infiltrates the dermis giving <b>plaque-like appearance</b><sup>Q</sup></li> <li>• Pseudo-Meissner corpuscles or tactile-like bodies</li> </ul> </td><td> <b>Plexiform</b> <ul style="list-style-type: none"> <li>• These tumors grow within and expand nerve fascicles.</li> <li>• Bag of worms<sup>Q</sup> appearance</li> <li>• CARROT SHAVINGS</li> </ul> </td></tr> </table>	<b>Localized</b> <ul style="list-style-type: none"> <li>• Low cellularity<sup>Q</sup></li> </ul>	<b>Diffuse</b> <ul style="list-style-type: none"> <li>• Diffusely infiltrates the dermis giving <b>plaque-like appearance</b><sup>Q</sup></li> <li>• Pseudo-Meissner corpuscles or tactile-like bodies</li> </ul>	<b>Plexiform</b> <ul style="list-style-type: none"> <li>• These tumors grow within and expand nerve fascicles.</li> <li>• Bag of worms<sup>Q</sup> appearance</li> <li>• CARROT SHAVINGS</li> </ul>
<b>Localized</b> <ul style="list-style-type: none"> <li>• Low cellularity<sup>Q</sup></li> </ul>	<b>Diffuse</b> <ul style="list-style-type: none"> <li>• Diffusely infiltrates the dermis giving <b>plaque-like appearance</b><sup>Q</sup></li> <li>• Pseudo-Meissner corpuscles or tactile-like bodies</li> </ul>	<b>Plexiform</b> <ul style="list-style-type: none"> <li>• These tumors grow within and expand nerve fascicles.</li> <li>• Bag of worms<sup>Q</sup> appearance</li> <li>• CARROT SHAVINGS</li> </ul>			
<b>Genetics</b>	Loss of NF2 gene product, merlin	Loss of the NF1 gene product, neurofibromin <sup>Q</sup>			
<b>Malignant transformation</b>	Extremely rare	Occur especially in plexiform neurofibroma <sup>Q</sup>			


Antoni B

Verocay  
bodies

Antoni A



Schwannoma




Expanded nerve  
fascicles

Bland spindle cells  
with wavy collagen  
(CARROT SHAVINGS)

Plexiform neurofibroma





## High Yield Facts

- **Most common tumor of CNS- metastasis<sup>Q</sup>**
- **Most common tumor associated with Brain Metastases- lung ca<sup>Q</sup> > breast > melanoma**
- **Most common tumor associated with Leptomeningeal Metastases<sup>Q</sup>** (also called carcinomatous meningitis, meningeal carcinomatosis) - **breast cancer<sup>Q</sup> > lung**
- **Choriocarcinoma** has high likelihood of metastasizing to brain whereas **prostatic carcinoma almost never grow in the brain.<sup>Q</sup>**
- Acoustic neuroma- Schwannoma that arises from the vestibulocochlear nerve (CN VIII). Located at the **cerebello-pontine angle** or in the internal acoustic meatus

## Malignant Peripheral Nerve Sheath Tumors (MPNST)

- **Origin:**
  - Sporadic- **de novo<sup>Q</sup>**
  - **NF1-associated tumors<sup>Q</sup>**- malignant transformation of a (plexiform) neurofibroma.<sup>Q</sup>

## Morphology:

- Mitoses, necrosis, and nuclear anaplasia are common



## High Yield Facts

- **Triton tumor<sup>Q</sup> -MPNST with focal areas that exhibit other lines of differentiation**, including glandular, cartilaginous, osseous, or rhabdomyoblastic morphology

## Phakomatoses

Include **tuberous sclerosis, neurofibromatosis, von Hippel-Lindau disease, and Sturge-Weber syndrome.**

- **Tuberous sclerosis:** Explained in detail later
- **Von Hippel-Lindau disease:** Explained in detail later
- **Sturge-Weber syndrome** is a **non-familial<sup>Q</sup> congenital<sup>Q</sup>** disorder, display angiomas of the brain, leptomeninges, and ipsilateral face, which are called **port-wine stains (nevus flammeus)<sup>Q</sup>**.



Adenoma sebaceum



Subungual fibromas

## FAMILIAL TUMOR SYNDROMES

### Tuberous Sclerosis

- Autosomal dominant
- Mutations in **TSC I locus, which codes for hamartin<sup>Q</sup>**, and the **TSC2 locus, which codes for tuberlin.<sup>Q</sup>**
- These two proteins inhibit **mTOR<sup>Q</sup>**-plays a central role in the **regulation of cell growth<sup>Q</sup>**.
- Clinical triad of angiofibromas ("adenoma sebaceum"), seizures, and mental retardation.<sup>Q</sup>
- Syndrome is associated with the development of **tumors<sup>Q</sup>** over **childhood and adolescence<sup>Q</sup>**

### Systemic Changes

- **CNS:**
  - **Hamartomas<sup>Q</sup>** -
    - **Cortical tubers<sup>Q</sup>**—haphazardly arranged neurons- epileptogenic
    - **Subependymal nodules<sup>Q</sup>**—can protrude into ventricles (candle drippings)

- **Neoplasm<sup>Q</sup>**
- **Subependymal giant-cell astrocytomas<sup>Q</sup>**—low<sup>Q</sup> grade neoplasms, **develop from the hamartomatous (subependymal<sup>Q</sup>) nodules in the same location**
- **Kidney: Angiomyolipomas<sup>Q</sup>**
- **Retina: Glial hamartomas<sup>Q</sup>**
- **Lungs: Lymphangiomyomatosis<sup>Q</sup>**
- **Heart: Rhabdomyomas<sup>Q</sup>**
- **Skin:**
  - **Angiofibromas<sup>Q</sup>**,
  - **Shagreen patches<sup>Q</sup>**- localized leathery thickenings
  - **Ash-leaf patches<sup>Q</sup>**- hypopigmented areas
  - **Subungual fibromas.<sup>Q</sup>**

### Von Hippel-Lindau Disease

Autosomal dominant<sup>Q</sup>, **gene on chromosome 3p25.3<sup>Q</sup> (tumor suppressor gene)**

- **CNS: Hemangioblastomas<sup>Q</sup>** (in 60–80% patients): **most common in the cerebellum and retina<sup>Q</sup>**
- **Kidney: Clear cell<sup>Q</sup> Renal cell carcinomas** (in 75% cases): **multifocal and bilateral.<sup>Q</sup>** Multiple benign cysts





- **Adrenal:** Pheochromocytomas (often **bilateral**)
- **Pancreas:** Cysts and serous cystadenomas<sup>Q</sup> (most frequent lesions in pancreas)- seen in 77% cases. Neuroendocrine tumors occur in about 10-15 % of cases.<sup>Q</sup>
- **Miscellaneous:**
  - Extra adrenal paragangliomas
  - **Endolymphatic sac tumors**<sup>Q</sup> (11% of cases)
  - **Epididymal cysts**<sup>Q</sup> (often bilateral)- in 54% of men
  - Cystadenomas of the broad ligament ("adnexal papillary tumor of probable mesonephric origin")<sup>Q</sup> are highly specific

## Mnemonic

### Von Hippel-lindau: signs and symptoms (HIPPEL):

Hemanigoblastomas	Port-wine stains
Increased renal cancer	Eye dysfunction
Pheochromocytoma	Liver, pancreas, kidney cysts

## Neurofibromatosis (NF: Type 1 and 2)

Autosomal Dominant<sup>Q</sup> Inheritance

## Mnemonic

NF1 (von Recklinghausen disease)	NF2
Mutations in Neurofibromin gene (chr17q) <sup>Q</sup>	Mutations in NF2/Merlin gene (chr22q) <sup>Q</sup>
<b>"N-O S-P-A-C-E"</b> <ul style="list-style-type: none"> <li>• <b>N</b>-Neurofibroma<sup>Q</sup></li> <li>• <b>O</b>-Optic glioma<sup>Q</sup></li> <li>• <b>S</b>-Scoliosis</li> <li>• <b>P</b>-Positive family history</li> <li>• <b>A</b>-Axillary freckling</li> <li>• <b>C</b>-Café au lait spots<sup>Q</sup></li> <li>• <b>E</b>-Eye (Lisch nodules)<sup>Q</sup></li> </ul>	<b>"M-I-S-S M-E"</b> <ul style="list-style-type: none"> <li>• <b>M</b>: Multiple</li> <li>• <b>I</b>: Inherited</li> <li>• <b>S</b>: Schwannomas<sup>Q</sup></li> <li>• <b>M</b>: Meningiomas</li> <li>• <b>E</b>: Ependymomas<sup>Q</sup></li> </ul>

- NF1 patients have **increased risk** of **Meningioma**<sup>Q</sup>, **Pheochromocytoma**<sup>Q</sup> and **Wilm's Tumor**<sup>Q</sup>
- Most common **tumor** in NF1 is **optic nerve glioma**<sup>Q</sup>
- **Most common Leukemia** in NF1 is **JMML**<sup>Q</sup> (Juvenile Myelomonocytic Leukemia)

## Hereditary Syndromes Associated with Brain Tumors

Syndrome	Clinical Features	Mutation
Cowden syndrome	Dysplastic ganglioglioma of the cerebellum (Lhermitte-Duclos disease)	PTEN
Li-Fraumeni syndrome	Medulloblastomas	TP53
Turcot syndrome	Medulloblastoma or glioblastoma	APC or mismatch repair genes
Gorlin syndrome	Medulloblastoma	PTCH gene
Multiple endocrine neoplasia 1 (Werner's syndrome)	Pituitary adenoma, malignant schwannoma	<b>Menin</b> gene <i>MEN1</i> (11q13)

## Mnemonic

### Brain tumor spreading by CSF:

#### PGMEAL CSF

- Pineoblastoma
- Germinoma, Glioblastoma
- Medulloblastoma
- Ependymoma
- CNS Lymphoma
- Choroid plexuses carcinoma

R10<sup>th</sup>

### Latest Update

- Mutation of Rhabdoid tumor: **INI-1** mutation
- CNS tumors with mural nodules
  - Pilocytic astrocytoma
  - Pleomorphic xanthoastrocytoma
- CNS tumor with calcifications: Meningioma, oligodendroglioma, craniopharyngioma
- Genes in familial Alzheimer's disease-**PS<sub>1</sub> > APP > PS<sub>2</sub>**.




## NEXT Pattern Questions



Q's

1. A 5-year-old patient presented with unconsciousness and projectile vomiting. MRI brain shows lesion in cerebellum and characteristic histological diagram is shown below. All of the following are true except:



- a. It can be solid and cystic
- b. It is biphasic tumor shows microcystic and fibrillary area
- c. Rosenthal fiber and eosinophilic granular bodies
- d. Pseudopalisading necrosis is common


**Ans. (d) Pseudopalisading necrosis is common**

- Unconsciousness and projectile vomiting are suggestive of raised intracranial tension. Since the mass is in cerebellum, and biopsy shows the eosinophilic granular body and Rosenthal fibers, it is suggestive of Pilocytic astrocytoma. Pseudopalisading is a feature of Glioblastoma multiforme.



Q's


2. A patient presented with painless proptosis. Biopsy from the orbital mass showed the following image?



- a. Neurofibroma
- b. Rhabdomyoma
- c. Leiomyoma
- d. Schwannoma


**Ans. (d) Schwannoma**

- The image shows tumor cells arrange in palisading (Verocay body) and you can see the cellular areas Antoni A and acellular areas Antoni B. This is seen in Schwannoma.




## Image-Based Questions

1. A 10-year-old child with posterior fossa mass. On biopsy following features were seen. What are these dense eosinophilic fibers marked with arrow?




- a. Rosenthal fibers
- b. The Alzheimer type II astrocyte
- c. Corpora amylacea
- d. None

3. A 40-year-old male who died due to high fever. Autopsy finding?




- a. Cerebral abscess
- b. Cerebral infarction
- c. Cerebral tumor
- d. None

2. New born baby with lumbar mass. Diagnosis.




- a. Myelocoele
- b. Lumbar meningocele
- c. Spina bifida
- d. None

4. Identify the lesion. Also mention the special stain and disease associated.




- a. Tangles
- b. Plaques
- c. Hirano bodies
- d. None

5. Identify the Rosette and Diagnosis.




- a. Perivascular pseudorosette
- b. Homer-Wright rosette
- c. True rosette
- d. Flexner-Wintersteiner rosette

6. A 5-year-old child with posterior fossa mass. Biopsy from the mass shows the following rosettes.



- a. Homer-Wright rosette
- b. Perivascular pseudorosette
- c. Flexner-Wintersteiner rosette
- d. None





## Answers of Image-Based Questions

**1. Ans. (a) Rosenthal fibers**

- Rosenthal fibers are eosinophilic, corkscrew fibers found in pilocytic astrocytoma, the most common primary brain tumor in children. They contain two heat-shock proteins ( $\alpha$ B-crystallin and hsp27) as well as ubiquitin.
- They are seen in Alexander disease and, pilocytic astrocytoma.

**2. Ans. (b) Lumbar meningomyelocele**

- The lumbar region of a newborn baby with myelomeningocele. The skin is intact, and the placode-containing remnants of nervous tissue can be observed in the center of the lesion, which is filled with cerebrospinal fluid.

**3. Ans. (a) Cerebral Abscess**

- Note well defined outline of cerebral abscess

**4. Ans. (b) Plaques**

- Plaques (arrow) contain a central core of amyloid and a surrounding region of dystrophic neurites (Bielschowsky stain). These are seen in Alzheimer's disease.

**5. Ans. (a) Perivascular Pseudorosette**

- In this pattern, a spoke-wheel arrangement of cells with tapered cellular processes radiates around a wall of a centrally placed vessel
- Seen in ependymomas

**6. Ans. (a) Homer-Wright rosettes**

- Type of pseudo rosette in which differentiated tumor cells surround the neuropil seen in neuroblastoma, medulloblastoma, pinealoblastoma



## Multiple Choice Questions

### CELLS OF CNS AND FUNCTIONS OF BRAIN CELLS


- Most sensitive to hypoxia is:** (Recent Question 2016)
  - Neuron
  - Liver cell
  - Stem cell
  - Muscle
- Rosenthal fibres are:** (Recent Question 2015)
  - Intraclear inclusions
  - Intracytoplasmic inclusions
  - Present extracellularly
  - Part of cell membrane
- Rosenthal fibres in astrocytoma are composed of:** (Recent Question 2015)
  - Heat shock proteins
  - Fibrillar proteins
  - GFAP
  - Globulins
- Phagocytosis in brain is caused by:** (Recent Question 2014, 2013, AIIMS Dec 94)
  - Astrocytes
  - Microglia
  - Oligodendrocytes
  - Ependymal cells
- Which is a mesenchymal cell?** (Recent Question 2014)
  - Microglia
  - Astrocytoma
  - Oligodendrocyte
  - Ependymal cells
- The following cell types does not participate in repair after brain infarction:** (Recent Question 2014)
  - Microglia
  - Astrocytes
  - Fibroblasts
  - Endothelium
- Disease or infarction of neurological tissue causes it to be repaired by:** (AI 12)
  - Fluid
  - Neuroglia
  - Proliferation of adjacent nerve cells
  - Blood vessel
- Retraction ball is seen after injury to:** (PGI May 2011)
  - Liver
  - Spleen
  - Brain
  - Kidney
  - Lungs

### MALFORMATIONS AND DEVELOPMENTAL DISORDERS


- Foix-Alajouanine disease of the spinal cord is a:** (Recent Question 2016)
  - Arteriovenous malformation
  - Cavernous malformation
  - Capillary telangiectasia
  - Venous angioma
- All of the following are the classical presentation of Craniovertebral junction anomalies except:** (AIIMS Nov 13)
  - Pyramidal signs
  - Low hairline
  - Short neck
  - Pupillary asymmetry

### CNS INFECTIONS


- A HIV pt presented with fever and neck rigidity and few days later the patient died even before any investigation could be performed. After autopsy the gross and histopathological image is given below. What is your diagnosis?** (AIIMS Nov 16)



- Cryptococcus
  - Toxoplasmosis
  - Herpes
  - Echinococcus multilocularis
- In a patient suspected to be diagnosed with Rabies, a sample of corneal smear was taken. Which of the following investigations can be done from the specimen?** (JIPMER 16)
    - RT PCR
    - Immunofluorescence test
    - Negri body visualization
    - Virus isolation
  - True about image below all except** (Recent Question 2016-17)



- They are eosinophilic, sharply outlined bodies in nerve cells
  - These are Negri bodies seen in rabies
  - Consist of ribonuclear proteins produced by the virus
  - Consist of DNA
- Hutchinson triad includes all except?** (Recent Question 2015)
    - Sabre skin
    - Eight nerve deafness
    - Upper incisor notch
    - Interstitial keratitis



- 15. Finding in histopathology of brain in rabies includes:** (PGI May 2015)  
 a. Negri body  
 b. Nodule  
 c. Neuronophagia  
 d. Vacuolar degenerative changes  
 e. Inflammatory cell
- 16. The appearance of cobweb formation in CSF indicates** (WB PGME 2016), (MH PG 2014)  
 a. Pyogenic meningitis  
 b. Viral meningitis  
 c. Tuberculous meningitis  
 d. Fungal meningitis
- 17. Owl eye inclusion bodies are seen in:** (Recent Question 2015)  
 a. HSV  
 b. CMV  
 c. EBV  
 d. Hepatitis B
- 18. Progressive multifocal leucoencephalopathy spares:** (Recent Question 2014)  
 a. White matter of cerebrum  
 b. White matter of parietal lobe  
 c. White matter of periventricular area  
 d. Spinal cord and optic nerve
- 19. Perivascular lymphocytes & microglial nodules are seen in:** (Recent Question 2014)  
 a. Multiple sclerosis  
 b. HIV encephalitis  
 c. CMV meningitis  
 d. Bacterial meningitis
- 20. Which part of brain is not affected in HIV infection?** (JIPMER 2014)  
 a. Hippocampus  
 b. Subcortical white matter  
 c. Diencephalon  
 d. Brain stem
- 21. Negri bodies are abundant in the following cells except?** (JIPMER 2014)  
 a. Subcortical white matter  
 b. Purkinje cells  
 c. Hippocampus  
 d. Basal ganglia
- 22. Negri bodies are seen in:** (WB PGME 2016, Recent Question 2013)  
 a. Oligodendroglia  
 b. Neuron  
 c. Microglia  
 d. Astrocytes
- 23. In which stage of neurocysticercosis, there is no edema?** (AIIMS May 2012)  
 a. Vesicular  
 b. Vesicular colloidal  
 c. Granular nodular  
 d. Nodular calcified
- 24. Brain infarcts is/are seen in infection with:** (PGI May 2012, 2011)  
 a. Toxoplasmosis  
 b. Rabies  
 c. Cryptococcus  
 d. Aspergillosis  
 e. TB

### ANEURYSM

- 25. Binswanger disease is a form of?** (Recent Question 2016)  
 a. Hypertensive retinopathy  
 b. Hypertensive nephropathy  
 c. Hypertensive encephalopathy  
 d. subcortical leukoencephalopathy
- 26. Hypertensive hemorrhage is most commonly seen in?** (AIIMS May 2015)  
 a. Basal ganglia  
 b. Thalamus  
 c. Brain stem  
 d. Cerebrum

- 27. Berry aneurysm-Defect lies in:** (AIIMS May 10)  
 a. Degeneration of internal elastic lamina  
 b. Degeneration of medial muscle cell layer  
 c. Deposition of mucoid material in media  
 d. Low grade inflammation of vessel wall

### PRION DISEASE

- 28. What is the histological appearance of brain in Creutzfeldt-Jakob disease:** (Recent Question 2015)  
 a. Neuronophagia  
 b. Spongiform change in brain  
 c. Microabscesses  
 d. Demyelination
- 29. Spongiform degeneration of cerebral cortex occurs in** (Recent Question 2015)  
 a. Creutzfeldt-Jakob disease  
 b. Subacute sclerosing panencephalitis  
 c. Fatal familial insomnia  
 d. Cerebral toxoplasmosis
- 30. Infectious protein with correct primary structure and wrong tertiary structure is?** (DNB Nov 12 Pattern)  
 a. Prion  
 b. Tau  
 c. Huntingtin  
 d. Synuclein

### NEURODEGENERATIVE DISEASE

- 31. Which is the pathognomic feature of Alzheimer's disease?** (Recent exam 2018)  
 a. Lewy bodies  
 b. Ballooned neurons  
 c. Plaques and tangles  
 d. Pick bodies
- 32. Which of the following is/are features(s) of lewy body dementia:** (PGI May 16)  
 a. Plaque containing beta-amyloid peptide  
 b. Deposition of  $\alpha$ -synuclein protein  
 c. Often resistant to standard treatment  
 d. Common in elderly  
 e. Risk of falling may present
- 33. Features of Alzheimer's disease are all except?** (Recent Question 2016)  
 a. Narrowing of ventricles  
 b. Hirano bodies  
 c. amyloid  
 d. neuritic plaques
- 34. Knife-edge pattern of lobar atrophy of the brain is associated with which degenerative disease of the cerebral cortex:** (Recent Question 2015)  
 a. Alziemer's disease  
 b. Frontotemporal dementias  
 c. Pick's disease  
 d. Progressive supranuclear palsy
- 35. The nucleus involved in Alzheimer's disease is:** (AIIMS 14)  
 a. Basal nucleus of Meyernet  
 b. Raphe nucleus  
 c. Superior salivary nucleus  
 d. Basal lobe of cerebellum






36. In Alzheimer disease, pathology seen in brain is:
- Atrophy of parietal and temporal lobe (AIIMS May 13)
  - Atrophy of frontal and temporal lobe
  - Atrophy of occipital and temporal lobe
  - Atrophy of parietal and occipital lobe

### TUMORS

37. A 25-year-old male presented with swelling in the wrist joint. Histopathological examination showed spindle cells and Verocay bodies. What is the most likely diagnosis? (Recent Pattern Question 2020)




- Neurofibroma
  - Schwannoma
  - Lipoma
  - Squamous cell carcinoma
38. Which of the following condition is shown below? (AIIMS Nov 2019)




- Plexiform neurofibroma
  - Von Hippel-Lindau syndrome
  - Marfan syndrome
  - Tuberous sclerosis
39. A 15-year-old presents with a history of pain and swelling in the right thigh. Biopsy of the mass demonstrates osteosarcoma. His mother was diagnosed with breast cancer 1 year ago and his maternal grandmother died of breast cancer 10 years ago. The patient has three younger siblings. The siblings have an increased risk of developing which of the following cancers? (JIPMER Nov 2019)
- Wilms'
  - Neuroblastoma
  - Hepatoblastoma
  - Glioma


40. A patient presented with painless proptosis. Biopsy from the orbital mass showed the following image? (AIIMS Nov 2017)



- Neuofibroma
  - Rhabdomyoma
  - Leiomyoma
  - Schwannoma
41. A 40-year-old lady is diagnosed to have brain tumor in frontal lobe. The lesion is characterized by focal necrosis surrounded by ring like enhancement (Recent Question 2016-17, JIPMER\_May 2015)
- Glioblastoma multiforme
  - Oligodendroglioma
  - Ependymoma
  - Astrocytoma
42. Homer Wright rosette is seen in: (Recent Question 2016)
- Neuroblastoma
  - Nephroblastoma
  - Ependymoma
  - Rhabdomyosarcoma
43. Multiple schwannomas are seen with: (Recent Question 2015)
- NF1
  - NF2
  - Noonan's syndrome
  - Tuberous sclerosis
44. Perivascular pseudorosettes are classically seen in: (Recent Question 2015)
- Ependymoma
  - Oligodendroglioma
  - Astrocytoma
  - Medulloblastoma
45. Most common cause of leptomeningeal metastasis is adenocarcinoma arising from: (Recent Question 2015)
- Breast
  - Thyroid
  - Bone
  - Liver
46. In the following histopathology of schwannoma, the arrow marked lesion shows? (AIIMS Nov 2015)



- Antony A with verocay body
- Rosettes
- Antony B with verocay body
- Pallisading



47. Flexner wintersteiner's rosettes are seen in: (AP 2014)  
a. Hepatoblastoma b. Neuroblastoma  
c. Nephroblastoma d. Retinoblastoma
48. True statement about metastases of malignant tumors of brain is: (AP 2013)  
a. Drop metastases can occur in the spinal cord  
b. Lymph node metastases in patients who have had brain surgery  
c. Metastases through man made shunts  
d. All of the above
49. Most common CNS tumor in NF1? (Recent Question 2014)  
a. Optic nerve glioma b. Meningioma  
c. Astrocytoma d. Schamomma
50. Which of the following tumors is not derived from meninges: (Recent Question 2014)  
a. Hemangioblastoma b. Meningioma  
c. Fibrous tumor d. Hemangiopericytoma
51. Medulloblastoma most common metastasis is to: (Recent Question 2014)  
a. Lung b. CNS  
c. Liver d. Bone
52. Most common CNS neoplasm in HIV patient: (Recent Question 2014)  
a. Medulloblastoma  
b. Astrocytoma  
c. Primary CNS lymphoma  
d. Ependymoma
53. The commonest intracranial tumor is: (AP PGME 14)  
a. Glioma b. Pituitary tumor  
c. Meningioma d. Metastasis
54. Rosenthal fibres are seen in: (Recent Question 2013)  
a. Pilocytic astrocytoma b. Glioblastoma  
c. Medulloblastoma d. Ependymoma
55. Most common cerebellar tumor in children? (MH 2016, Recent Question 2013)  
a. Astrocytoma b. Medulloblastoma  
c. Ependymoma d. PNET
56. Following is true about medulloblastoma: (Recent Question 2013)  
a. It is seen mainly in over 50 age group  
b. It is radiosensitive tumour  
c. Only treatment is surgery  
d. Seen in anterior cranial fossa
57. Receptor on neuronal membrane that induces development of glioma: (AIIMS May 2013)  
a. CD-117 b. CD-133  
c. CD-33 d. CD-45
58. Which of the following statement(s) is/are true about pilocytic astrocytoma: (WB PGME 2016, PGI May 2013)  
a. Slow growing  
b. Eosinophilic granular bodies  
c. Most commonly involve cerebellum  
d. Mostly cystic in nature  
e. Mostly malignant  
f. Negative GFAP

59. Most common site for medulloblastoma is: (MH 2016, DNB 2012)  
a. Cerebellum b. Pituitary  
c. Cerebrum d. Basal ganglia
60. All of the following are neuronal tumors, except: (AI 11)  
a. Gangliocytoma b. Ganglioglioma  
c. Neurocytoma d. Ependymoma
61. Most common site of glioblastoma multiforme is: (DNB Pattern 11)  
a. CP angle b. Temporal lobe  
c. Brain stem d. Occipital lobe
62. True about meningioma: (JIPMER 11)  
a. More common in men  
b. 50% are malignant  
c. 95 % cure rate following treatment  
d. Arise from arachnoid layer
63. Which of the following brain tumors does not spread via CSF? (DPG 11)  
a. Germ cell tumors  
b. Medulloblastoma  
c. CNS lymphoma  
d. Craniopharyngioma

#### MISCELLANEOUS

64. Pseudolaminar necrosis is a feature of: (Recent Question 2014-15)  
a. Cerebral infarct  
b. Renal infarct  
c. Hepatic infarct  
d. Cardiac infarct
65. Onion bulb appearance on nerve biopsy is seen in: (AIIMS Nov 11)  
a. Amyloid neuropathy  
b. Diabetic  
c. CIDP  
d. Leprous neuritis

#### FAMILIAL SYNDROMES

66. Koener's tumor are seen in? (Recent Question 2015)  
a. Tuberous sclerosis b. Neurofibromatosis  
c. VHL d. NF
67. True about turcot syndrome (Recent Question 2015)  
a. Mutations in PTEN gene  
b. CNS tumors  
c. Non neoplastic polyps  
d. Congenital hypertrophy of retinal pigment epithelium
68. About Neurofibromatosis are true, except: (AIIMS May 14)  
a. Autosomal recessive b. Associated with cataract  
c. Scoliosis d. Multiple fibroma
69. Sturge-Weber syndrome is not associated with: (AIIMS 14)  
a. Seizures  
b. Hemiatrophy of cerebral cortex  
c. Gyriform calcification in brain  
d. Empty sella



## Answers with Explanations

1. **Ans. (a) Neurons** (Ref: Robbins 9th/pg 130; 8th/pg 129)
  - Neurons undergo irreversible damage after **3 to 4 minutes**<sup>Q</sup> of ischemia
2. **Ans. (b) Intracytoplasmic inclusions** (Ref: Robbins 9th/pg 1253)

**Rosenthal fibers-**

  - Brightly **eosinophilic structures/inclusions** within **cytoplasm of astrocytic processes**
  - Contain two **heat-shock proteins** ( $\alpha$ B-crystallin and hsp27) as well as **ubiquitin**<sup>Q</sup>
  - Seen in **Alexander disease, pilocytic astrocytoma**<sup>Q</sup>.
3. **Ans. (a) Heat shock proteins** (Ref: Robbins 9th/pg 1253)
4. **Ans. (b) Microglia** (Ref: Robbins 9th/pg 1253; 8th/pg 1282)
5. **Ans. (a) Microglia** (Ref: Robbins 9th/pg 1253; 8th/pg 1282)
  - Microglial cells are mesodermal in origin.
  - When the debris filled in the microglial cytoplasm is lipid- gitter cells
  - **Option B, C and D**-All the other cells are derived from neuroectoderm
6. **Ans. (c) Fibroblasts** (Ref: Robbins 9th/pg 1253; 8th/pg 1282)
7. **Ans. (b) Neuroglia** (Ref: Robbins 9th/pg 1253; 8th/pg 1282)
8. **Ans. (c) Brain**

(Ref: Refer The Journal of Neuroscience. 24 (19): 4605—4613).

Axons are normally elastic, but when rapidly stretched as in diffuse axonal injury they become brittle, and the axonal cytoskeleton can be broken

Axonal transport continues up to the point of the break in the cytoskeleton, leading to a **buildup of transport products and local swelling at that point**

When it becomes large enough, swelling can tear the axon at the site of the break in the cytoskeleton, causing it to **draw back toward the cell body and form a bulb**

This bulb is called a **retraction ball**, the **hallmark of diffuse axonal injury**
9. **Ans. (a) Arteriovenous malformation**

(Ref: Toole's Cerebrovascular Disorders pg 351)

Foix-Alajouanine syndrome is an arteriovenous (AV) malformation of the spinal cord predominantly affecting the lower thoracic and/or lumbosacral levels

Foix-Alajouanine syndrome usually occurs in older patients (>50 years).
10. **Ans. (d) Pupillary asymmetry**

(Ref: William wilkins neurosurgery: 2732-35)

  - Congenital anomalies included in craniovertebral anomalies-klippel fiell syndrome and down syndrome

In **Klippel-Feil syndrome**, head is cocked to one side with, **Low hairline posteriorly, Short neck**, Limited neck movement

11. **Ans. (a) Cryptococcus** (Ref: Harrisons 19th/pg 2010)

### In Cryptococcal Meningitis:

- Hematoxylin and eosin stain shows **lightly basophilic cell wall surrounded by a clear zone**.
- Cryptococcus neoformans will stain with **Periodic acid-Schiff or silver methenamine**.
- **Mucicarmine stains the capsule**-shows clear zone containing car-minophilic material. Alcian blue also stains the capsule
- **Negative staining can be done by India ink stain**

12. **Ans. (b) Immunofluorescence test**

(Ref: <http://www.cdc.gov/rabies/diagnosis/index.html>)

### Lab diagnosis of rabies

- RT-PCR: detecting the viral nucleic acid especially on **biological fluids** (saliva, cerebrospinal fluid, tears) and skin biopsy (ante mortem) and brain samples (post mortem)
- Serum and spinal fluid are **tested for antibodies to rabies virus**
- Skin biopsy specimens are examined for **rabies antigen in the cutaneous nerves at the base of hair follicles**
- Immunofluorescent antibody staining of the epithelial cells on the **corneal impression test (FAT)**

13. **Ans. (a, b, c, d) a. They are eosinophilic, sharply outlined bodies in nerve cells; b. These are Negri bodies seen in rabies; c. Consist of ribonuclear proteins produced by the virus d. Consist of DNA**

14. **Ans. (a) Sabre skin**

(Ref: Clinical Microbiology Reviews 12 (2): 187–209)

**Hutchinson's triad**-interstitial keratitis, Hutchinson incisors, and eighth nerve deafness

15. **Ans. (a, b, c, e); a. Negri body; b. Nodule; c. Neuronophagia; e. Inflammatory cell**

(Ref: Greenfield's Neuropathology, 8th ed/pg 1323)


Histopathologic evidence of rabies encephalomyelitis (inflammation) in brain tissue includes:

- Mononuclear infiltration
- Perivascular cuffing of lymphocytes or polymorphonuclear cells
- Neuronophagia
- Lymphocytic foci
- Babes nodules consisting of glial cells

16. **Ans. (c) Tuberculous meningitis**


(Ref: Robbins 9th/pg 1273 harshmohan 4th ed:857)





17. **Ans. (b) CMV** (Ref: Robbins 9th/pg 1266)  
Enlarged cells (**cytomegaly**) with **intranuclear and intracytoplasmic inclusions** are seen with **cytomegalovirus infection**.<sup>Q</sup>  
The intra nuclear inclusions appear like owl's eye inclusions in stained tissue sections
18. **Ans. (d) Spinal cord and optic nerve** (Ref: R 9th/pg 1278)  
**Progressive multifocal leukoencephalopathy (PML)** typically spares the optic nerve and the spinal cord
19. **Ans. (b) HIV encephalitis** (Ref: Robbins 9th/pg 1278)  
  - HIV Meningoencephalitis-characterized by **microglial nodules**<sup>Q</sup> composed of **mononuclear cells, microglia, and scattered multinucleated giant cells**.
20. **Ans. (a) Hippocampus** (Ref: R 9th/pg 1278; 8th/pg 1035)  
HIV encephalitis occurs especially in subcortical white matter, diencephalon and brainstem.
21. **Ans. (a) Subcortical white matter** (Ref: R 9th/pg 1277)  
  - Most characteristic pathologic finding in CNS in Rabies is the formation of **cytoplasmic inclusion bodies** called **Negri bodies**<sup>Q</sup> (composed of finely fibrillar matrix and rabies virus particles)
  - They are most commonly seen in **cerebellum (purkinje cells) & pyramidal neurons of hippocampus**
  - Basal ganglia involvement is common
  - The **prominence of early brain stem dysfunction**<sup>Q</sup> distinguish it from other viral encephalitis.
22. **Ans. (b) Neuron**  
(Ref: Robbins 9th/pg 1252)  
**Neuronal inclusions** can be seen in  
  - Viral infections**
  - Intranuclear inclusions**-herpes infection (Cowdry body)
  - Cytoplasmic inclusions**-rabies (Negri body)<sup>Q</sup>
  - Both nucleus & cytoplasm**-CMV.
23. **Ans. (d) Nodular calcified**  
(Ref: AJNR 2001 22: 677-680)  
4 main stages of NCC are:  
  - Vesicular:** viable parasite with intact membrane and therefore **no host reaction**.
  - Colloidal vesicular:** As the membrane becomes leaky oedema surrounds the cyst. This is the **most symptomatic stage**.<sup>Q</sup>
  - Granular nodular:** oedema decreases as the cyst retracts further; **enhancement persists**.
  - Nodular calcified:** end-stage quiescent calcified cyst remnant; **no oedema**.<sup>Q</sup>
24. **Ans. (d, e); d. Aspergillosis; e. TB**  
(Ref: Robbins 9th/pg 1280; The Brazilian Journal of Infectious Diseases 2004;8(2):175-17)  
Brain infarcts is/are seen in infection with:  
  - Aspergillus and mucor** have the tendency to invade blood vessels and cause thrombosis with cerebral infarction or vascular rupture with cerebral hemorrhage.

- Some infections, such as **cytomegalovirus, herpes zoster and tuberculosis** are known to produce vasculitis and infarction in the CNS.
  - HIV**
25. **Ans. (d) Subcortical leukoencephalopathy**  
(Ref: International Neurology textbook pg 12)  
Binswanger's disease (also known as **subcortical leukoencephalopathy**), also called *subcortical vascular dementia*, is a type of dementia caused by widespread, microscopic areas of damage to the deep layers of white matter in the brain  
Hypertension and old age are risk factors.
26. **Ans. (a) Basal ganglia** (Ref: Harrison 18th ed, chapter 370)  
Hypertensive hemorrhage is parenchymal and its most frequent sites of are the **basal ganglia, thalamus**, the cerebellum, the pons, and occasionally the subcortical white matter.
27. **Ans. (a, b); a. Degeneration of internal elastic lamina; b. Degeneration of medial muscle cell layer**  
(Ref: Robbins 9th/pg 1270; 8th/pg 1297)  
**Berry aneurysm**  
  - Developmental abnormalities**<sup>Q</sup>
  - Due to **the structural abnormality of the involved vessel (absence of smooth muscle and intimal elastic lamina.)**
  - Please note : new edition of robbins has specifically mentioned degeneration of internal elastic lamina also.
28. **Ans. (b) Spongiform change in brain**  
(Ref: Robbins 9th/pg 1281)
29. **Ans. (a) Creutzfeldt-Jakob disease** (Ref: R 9th/pg 1281)
30. **Ans. (a) Prion**  
(Ref: <http://www.rsc.org/chemistryworld/Issues/2005/October/prions>, J G Safar, Proc. Nat. Acad. Sci. US, 2005, 102, 3501)  
**The primary structure of the human prion protein, both in its benign and pathogenic forms, consists of a linear chain of 253 amino acids.**  
The pathogenic form, designated PrP<sup>sc</sup>, where sc stands for scrapie (the earliest known prion disease which occurs in sheep), is **folded quite differently, even though its primary structure is essentially identical**.
31. **Ans. (c) Plaques and tangles** (Ref: Robbins 9th ed p 1290)  
The major microscopic abnormalities of Alzheimer disease are neuritic (senile) plaques and neurofibrillary tangles.
32. **Ans. (b, c, d, e, f); b. Deposition of  $\alpha$ -synuclein protein; c. Often resistant to standard treatment; d. Common in elderly; e. Risk of falling may present**
33. **Ans. (a) Narrowing of ventricles** (Ref: Anderson 10th ed:2740; Robbins 9th/pg 1287-88; 8th/pg 1313)  
**Compensatory ventricular enlargement due to extensive cortical atrophy is called hydrocephalus ex vacuo.**



**This is seen in Alzheimer's and pick's disease.**

- The **pathologic hallmark**<sup>Q</sup> are plaques (both neuritic and senile) and tangles\_ Option d is true
- Vascular amyloid in **cerebral amyloid angiopathy** is A $\beta$ 40–Option c is true
- **Hirano bodies**<sup>Q</sup> and **Granulovacuolar degeneration**<sup>Q</sup> are other features seen in AD.–Option b is true

**34. Ans. (c) Pick's disease (Ref: R 9th/pg 1292; 8th/pg 1318)**

**PICK'S disease**

Asymmetric, atrophy of the frontal and temporal lobes  
Sparing of the posterior two thirds of the superior temporal gyrus, parietal or occipital lobe

**"Knife-edge" appearance-** atrophy can be severe, reducing the gyri to a wafer-thin appearance

**Pick cells-** characteristic swelling of surviving neurons (ballooning)

**Pick bodies-** cytoplasmic, 3R tau containing bodies, stain strongly with silver methods

**35. Ans. (a) Basal nucleus of Meyernet**

(Ref: Harrison 18th ed: 3306, 17th pg 2541)

AD is associated with decrease in cerebral cortical level of acetylcholine. This degeneration occurs in **nucleus basalis of Meyernet**<sup>Q</sup>

**36. Ans. (a) Atrophy of parietal and temporal lobe**

(Ref: Robbins 9th/pg 1290; 8th/pg 1313)

Grossly, brain shows a variable degree of cortical atrophy marked by widening of the cerebral sulci that is most pronounced in the frontal, temporal, and parietal lobes.

**37. Ans. (b) Schwannoma (Ref: R 9th pg 1317)**

**38. Ans. (a) Plexiform neurofibroma (Ref: R 9th pg/1317)**

Plexiform neurofibromas represent an uncommon variant of NF-1 in which neurofibromas arise from multiple nerves as bulging and deforming masses involving also connective tissue and skin folds—hence the clinical description of lesions as "bags of worms".

**39. Ans. (d) Glioma (Ref: Robbins 9th/pg 1317)**

**40. Ans. (d) Schwannoma**

Schwannomas often contain dense eosinophilic Antoni A areas(left) and loose, pale Antoni B areas (right), as well as hyalinized blood vessels (right). B, Antoni A area with the tumor cell nuclei aligned in palisading rows leaving anuclear zones and resulting in the formation of structures termed Verocay bodies

**41. Ans. (a) Glioblastoma multiforme (Ref: Robbins 9th/pg 1307, Journal of Neuro-Oncology 108 (1): 11–27)**

When viewed with MRI, glioblastomas often appear as ring-enhancing lesions. The appearance is not specific, however, as other lesions such as abscess, metastasis, tumefactive multiple sclerosis, and other entities may have a similar appearance. But necrosis with ring enhancement Is a f/o Glioblastoma multiforme.

Oligodendroglioma usually shows calcifications.

**42. Ans. (a) Neuroblastoma (Ref: Robbins 9th/pg 1306)**

Types of rosettes are given below:

<b>Homer Wright rosette:</b> Neuroblastoma, medulloblastoma	<b>Perivascular pseudorosette:</b> Ependymoma
<b>Flexner wintersteiner rosette:</b> Retinoblastoma	<b>True rosette:</b> Ependymoma

**43. Ans. (b) NF2**

(Ref: Robbins 9th/pg 1317)

NF2 is **most commonly** characterized by **bilateral schwannomas of the vestibulocochlear nerves** (cranial nerve VIII) and **multiple meningiomas**.

**44. Ans. (a) Ependymoma**

(Ref: Robbins 9th/pg 1306)

**45. Ans. (a) Breast (Ref: Harrison 18th ed 3390, 17th ed: 2608)**

Site of Primary Tumor	Leptomeningeal Metastases, %
Lung	24
Breast	41 <sup>Q</sup>
Melanoma	12
Gastrointestinal tract	13

**46. Ans. (a) Antony A with verocay body (Ref: R 9th/ 1314)**

<b>Morphology of schwannoma</b>	<b>Consists of:</b> <ul style="list-style-type: none"> <li>• Antoni A: cellular areas</li> <li>• Antoni B: loose edematous areas<sup>Q</sup></li> </ul> <b>Verocay bodies</b> <sup>Q</sup> (foci of palisaded nuclei) may be found in the more cellular areas.
---------------------------------	--

**47. Ans. (d) Retinoblastoma (Ref: Robbins 9th/pg 1306)**

**48. Ans. (d) All of the above**

(Ref: Robbins 9th/pg 1312; 8th/pg 1336)


**Dissemination through the CSF** is a common complication-giving rise to **nodular masses** at some distance from the primary tumor called as **"drop metastases" is a feature of medulloblastoma**

**Man made shunts provide a route for metastasis of malignant tumors**

Lymphomatous metastases, on the other hand, tend to produce a more diffuse pial enhancement, as breast and prostate metastases.

**49. Ans. (a) Optic nerve glioma (Ref: Robbins 9th/pg 1317)**

- NF1-autosomal dominant disorders
- Characterized by **neurofibromas of peripheral nerve, gliomas of the optic nerve, (Lisch nodules), and cutaneous hyperpigmented macules (café au lait spots).**
- NF2 is **most commonly** characterized by **bilateral schwannomas of the vestibulocochlear nerves** (cranial nerve VIII) and **multiple meningiomas**.



50. Ans. (d) **Hemangiopericytoma**

(Ref: Robbins 9th/pg 1317, textbook of neuro-oncology :512)

**Meningeal tumors**

Meningothelial tumor	Meningioma
Mesenchymal non	Hemangiopericytoma, meningeal
Meningothelial tumor	solitary fibrous tumor

Hemangioblastomas are highly vascular neoplasms that occur as a mural nodule associated with a large fluid-filled cyst. It is associated with Von Hippel-Lindau Disease

51. Ans. (b) **CNS** (Ref: Robbins 9th/pg 1312; 8th/pg 1336)

52. Ans. (c) **Primary CNS lymphoma** (Ref: R 9th/pg 1313)

- **MC neoplasm in HIV**-central nervous system (CNS) lymphoma is a **diffuse, large-cell non-Hodgkin lymphoma**<sup>Q</sup> of **B-cell origin**<sup>Q</sup> that usually occurs in the brain (rarely in the spinal cord).
- It is a **late complication**<sup>Q</sup> of HIV infection.
  - **Epstein-Barr virus (EBV)**<sup>Q</sup> is identified in almost all cases<sup>Q</sup>.

53. Ans. (d) **Metastasis** (Ref: Robbins 9th/pg 1306)

54. Ans. (a) **Pilocytic astrocytoma** (Ref: Robbins 9th/pg 1307)

55. Ans. (a) **Astrocytoma**

(Ref: Robbins 9th/pg 1306: See Ans 69)

- Most common pediatric brain tumor-Pilocytic astrocytoma
- Most aggressive pediatric brain tumor-Medulloblastoma

56. Ans. (b) **It is radiosensitive tumour** (Ref: R 9th/pg 1312)

**Medulloblastoma is exquisitely radiosensitive<sup>Q</sup>; Predominantly seen in children & exclusively in cerebellum<sup>Q</sup>**

57. Ans. (b) **CD-133**

(Ref: <http://cdn.intechopen.com/pdfs/14462/InTech-h->)

The most commonly used cell surface markers for glioma stem cells are CD133, CD15, and A2B5.

CD133 - on neuronal membrane that induces development of glioma.

58. Ans. (a, b, c, d); a. **Slow growing; b. Eosinophilic granular bodies; c. Most commonly involve cerebellum; d. Mostly cystic in nature** (Ref: Robbins 9th/pg 1309)

59. Ans. (a) **Cerebellum**

(Ref: Robbins 9th/pg 1312; 8th/pg 1336)

60. Ans. (d) **Ependymoma**

(Ref: Robbins 9th/pg 1306; 8th/pg 1330)

<b>Gliomas</b>	<b>Neuronal tumors</b>
Astrocytoma, oligodendroglioma and ependymoma	<b>Gangliogliomas</b> <b>Dysembryoplastic neuroepithelial tumor</b>

61. Ans. (b) **Temporal lobe** (Ref: BRS neuroanatomy 4th ed: 87)

Most common site of glioblastoma multiforme is temporal lobe, frontal lobe and basal ganglia

62. Ans. (d) **Arise from arachnoid layer** (Ref: R 9th/pg 1314)

**Meningioma**

- **Benign tumors of Adults**
- **Attached to the dura**<sup>Q</sup>, they commonly arise along the venous sinuses (parasagittal, sphenoid wings, and olfactory groove).
- **Arise from meningotheial cells of the arachnoid**<sup>Q</sup>

63. Ans. (d) **Craniopharyngioma** (Ref: Robbins 9th/pg 1314)

64. Ans. (a) **Cerebral infarct**

(Ref: AJNR Am J Neuroradiol. 2003 Apr;24(4):680-7)

**Cortical pseudolaminar necrosis**, also known as **laminar necrosis**, is the death of cells in the (cerebral) cortex of the brain in a band-like pattern, with a relative preservation of cells immediately adjacent to the meninges. It is a feature of subacute cerebral infarction.

65. Ans. (c) **CIDP**

(Ref: Current therapy in neurological disease vol 1: 447, Fundamentals of Neurology: An Illustrated Guide :175, Harrison 18th ed:3475-3478)

**Chronic inflammatory demyelinating polyneuropathy (CIDP)** - immune-mediated inflammatory disorder of the peripheral nervous system.

Biopsy typically **reveals little inflammation and onion-bulb changes** (imbricated layers of **attenuated Schwann cell** processes surrounding an axon) that result from recurrent demyelination and remyelination

25% of patients with clinical features of **CIDP neuropathy** also have a monoclonal gammopathy of undetermined significance (MGUS).

66. Ans. (a) **Tuberous sclerosis** (Ref: Robbins 9th/pg 516)

Ungual fibromas or Koenen's tumors are angiofibromas which occur in the lateral nail groove, along the proximal nail fold or under the nail. They are seen in **Tuberous Sclerosis**

67. Ans. (b) **CNS tumors**

(Ref: Russell & Rubinstein's Pathology of Tumors of the Nervous System 7Ed, pg 997)

**Turcot syndrome is characterised by:**

- Intestinal polyposis
- CNS tumours: glioblastoma or medulloblastoma

68. Ans. (a) **Autosomal recessive** (Ref: Robbins 9th/pg 1247)

69. Ans. (d) **Empty sella** (Ref: Robbins 9th/pg 516; 8th/pg 522)

- **Sturge-Weber syndrome** is a **non-familial**<sup>Q</sup> **congenital**<sup>Q</sup> disorder
- There is intra cranial calcification in occipito parietal region- **serpentine/ rail road track**<sup>Q</sup> **appearance**
- **Associated with facial port wine nevi**<sup>Q</sup>, ipsilateral venous angiomas in the cortical leptomeninges,
- **Mental retardation**<sup>Q</sup>, **seizures**<sup>Q</sup>, **hemiplegia**, **focal or diffuse atrophy**<sup>Q</sup> and skull radio-opacities.



[illegible]This image shows a single sheet of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

# Blood Banking and Transfusion Medicine

## Key Points

- » **Major** blood group systems (of clinical importance): **ABO and Rh** while **minor** blood groups: E.g. **MNS, Duffy, Kell, Kidd**
- » Genetic loci of **ABO is on chromosome 9** while **Rh on chromosome 1**
- » Bombay phenotype is absence of A, B, and H antigens on RBCs, and presence of **anti-A, anti-B, and anti-H** antibodies
- » **Rh D antigen** is the **most immunogenic<sup>o</sup>** RBC antigen **after A and B**
- » **Sodium Adenine Glucose Mannitol (SAGM)** is **blood bag anticoagulant with shelf life of 42 days**
- » Fresh frozen plasma contains all coagulation factors
- » **Massive transfusion is replacement of  $\geq 1$  time the total blood volume, within 24 hours or  $>50\%$  in 3 hours**
- » Most frequent transfusion reaction is FNHTR
- » **TRALI** is caused by **Ab against patient's HLA type II and HNA (human neutrophilic antigen)**

## Key Recent Updates

- » Leukoreduce blood products achieve 3 log reduction of WBC.



## RED CELL ANTIGENS/BLOOD GROUPS

- A total of **30 blood group systems** have been described
- Each blood group system is a series of **red cell antigens**, determined by **genetic loci**
- Major** blood group systems (of clinical importance): **ABO and Rh**
- Minor** blood groups: e.g. **MNS, Duffy, Kell, Kidd**

### The ABO and Rh Blood Group Systems

Properties	ABO system			Rh system
Blood groups	4 main blood groups: A, B, AB and O <sup>q</sup>			Rh positive and negative
Genetic loci	On chromosome 9 <sup>q</sup>			On chromosome 1 <sup>q</sup>
Antigens and Antibodies	Group	Antigen	Antibody	<ul style="list-style-type: none"><li>• C, c, D, E, e Antigens<sup>q</sup></li><li>• ‘d’ indicates the absence of D<sup>q</sup></li><li>• Anti D Antibody: Most important<sup>q</sup></li></ul>
	O	H	Anti-A and Anti-B <sup>q</sup>	
	A	A	Anti-B <sup>q</sup>	
	B	B	Anti-A <sup>q</sup>	
	AB	A and B	None <sup>q</sup>	
Antigens also seen on/in	Endothelial and epithelial cells <sup>q</sup> Plasma, saliva, semen (not in CSF) <sup>q</sup>			No other cells <sup>q</sup>
Type of Ab	IgM <sup>q</sup> ; naturally occurring <sup>q</sup> antibodies			IgG <sup>q</sup> ; Do not occur naturally
Clinical importance	<ul style="list-style-type: none"><li>• Anti-A and anti-B Ab can cause <b>severe intravascular hemolysis after incompatible transfusion.</b><sup>q</sup></li><li>• ABO-matching is required before transplantation<sup>q</sup> of solid organs</li></ul>			Rh -ve individuals <b>make anti-D Ab if</b> : <ul style="list-style-type: none"><li>• <b>Transfused with Rh +ve blood<sup>q</sup> or,</b></li><li>• <b>Rh –ve pregnant women, is exposed to Rh +ve fetal RBCs<sup>q</sup> that have crossed the placenta.</b></li></ul>

### Bombay Blood Group/Bombay Phenotype

- Absence of A, B, and H antigens on RBCs, and presence of **anti-A, anti-B, and anti-H** Antibodies
- Genetically: homozygous hh → cannot form the H precursor of A and B
- Clinical significance:
  - Their RBCs **type as group O**, but they **cannot receive blood group O** blood donation
  - Can only be safely transfused with other Bombay blood group which is rarely available



### High Yield Facts

- The **Rh D antigen** is the **most immunogenic<sup>a</sup>** RBC antigen **after A and B**.
- Immune **anti-K Ab (IgG)** is the **most common antibody<sup>a</sup>** found outside the ABO and Rh systems.
- Antibodies to K1 (Kell group) can cause **severe hemolytic disease of the newborn<sup>a</sup>**.
- Individuals **lacking Duffy antigens** are **immune to malaria<sup>a</sup>** caused by *P. vivax* and *P. knowlesi*
- Erythroid-specific RBC antigens include: **Rh, Kell and MNS, not expressed in other tissues<sup>a</sup>**.
- The **A, B and H antigens** are fully developed and reach **adult levels** by the age of **1 year<sup>a</sup>**.
- Most red cell genes are expressed as **co-dominant<sup>a</sup>** antigens (i.e. both genes expressed in heterozygote).
- Plasma **VWF and factor VIII levels** are **25% lower in Gr. O<sup>a</sup>** healthy individuals than other ABO groups.
- Auto-anti-P Ab ('Donath-Landsteiner antibody')<sup>a</sup>** is a potent **biphasic haemolysin<sup>a</sup> (IgG)<sup>a</sup>**, responsible for **paroxysmal cold hemoglobinuria<sup>a</sup>**.





## BLOOD AND ITS COMPONENTS IN CLINICAL USE


Name	Description	Volume/ Unit	Storage	Shelf Life	Indications	Special Remarks	Compatibility
<b>Whole Blood</b>	Donor blood plus anticoagulants	350 mL	2–6°C	35 days (CPDA) 42 days (SAGM)	<ul style="list-style-type: none"> <li>Acute blood loss with hypovolemia</li> <li>Exchange transfusion</li> </ul>	Per unit ↑ Hb by 1 gm/dL and Hct by 3–5%	ABO and Rh compatible with recipient
<b>Packed RBCs</b>	RBC concentrate (Hct 65–75%) <sup>a</sup>	350 mL	2–6°C	35 days (CPDA) 42 days (SAGM)	<ul style="list-style-type: none"> <li>Severe Anemia</li> <li>Exchange transfusion</li> </ul>	Same as above	Same as above
<b>Random Donor Platelets (RDP)</b>	Platelet Concentrate	50–70 mL	20–24°C in Platelet Agitator	5 days	Bleeding due to thrombocytopenia	Per unit ↑ platelet count by 10,000/uL	ABO compatible preferable
<b>Single Donor Platelet (SDP)</b>	Platelet concentrate from 1 donor by apheresis	200–300 mL	20–24°C in Platelet Agitator	5 days	Refractory/ severe thrombocytopenia	Per unit ↑ platelet by 30,000–50,000/uL	ABO and Rh compatible
<b>Fresh Frozen Plasma (FFP)</b>	Plasma from single donor frozen within 6 hrs of collection	200 mL	–30°C; thawed just before use	1 year	Hemophilia A/B, Liver disease, Warfarin overdose, DIC; TTP	Contains all coagulation factors and fibrinogen	ABO and Rh compatible
<b>Cryoprecipitate (CP)</b>	Precipitated proteins from FFP; Rich in Fibrinogen; f-VIII, XIII and vWF	10–20 mL	–30°C	1 year	vWD Hemophilia A Factor XIII def Fibrinogen def.	Each unit of factor VIII/kg plasma ↑ factor VIII by 2%	No compatibility testing required

## DURATION TIMES FOR TRANSFUSION

Blood products	Start transfusion	Complete transfusion
Whole blood/PRBC	Within 30 minutes of removing from refrigerator	≤4 hours <i>Discard unit this period is exceeded</i>
Platelet concentrate	Immediately	Within 30 minutes
FFP	As soon as possible	Within 30 minutes
Cryoprecipitate	As soon as possible	Within 30 minutes

## ADDITIVE SOLUTIONS USED FOR STORAGE OF BLOOD PRODUCTS


Name	Full form	Shelf life
<b>ACD</b>	Acid Citrate Dextrose	21 days
<b>CPD</b>	Citrate Phosphate Dextrose	21 days
<b>CPD-A</b>	Citrate Phosphate Dextrose with Adenine <sup>a</sup>	35 days
<b>SAGM</b>	Sodium Adenine Glucose Mannitol <sup>a</sup> (also contains CPD as anticoagulant)	42 days



## Action of Ingredients of Additive/Anticoagulant Solution

Ingredient	Action
Glucose	ATP generation by glycolysis <sup>a</sup>
Adenine	Synthesis of ATP; <sup>a</sup> Increases shelf life of RBCs to 42 days
Citrate	Prevents coagulation <sup>a</sup> by chelating Calcium
Sodium di phosphate	Prevents fall in pH <sup>a</sup>

## ADVERSE EFFECTS OF BLOOD TRANSFUSION




## High Yield Facts

- Cryoprecipitate is **not** useful in Hemophilia B.<sup>a</sup>
- FFP is **relatively** deficient in factor V and VIII.<sup>a</sup>
- To prevent hyperkalemia due to blood transfusion, it is **preferable** to use blood <7 days old.<sup>a</sup>
- TRALI is caused by **Ab against patient's HLA type II and HNA (Human Neutrophilic Antigen)**,<sup>a</sup> in donor plasma; usually 1-4 hrs after starting transfusion
- **Hepatitis C** is the most common cause of **transfusion associated viral hepatitis**
- **Acute hemolytic transfusion reactions** are **Type II Hypersensitivity** reactions caused by **complement mediated hemolysis**
- **Most frequent transfusion reaction** is FNHTR
- FNHTR is caused by **antibodies against donor lymphocytes and HLA antigens**

## Massive Transfusion

Definition	Replacement of $\geq 1$ time the total blood volume, within 24 hours <sup>a</sup> or Replacement of <b>more than 50% of the blood volume in 3 hours<sup>a</sup></b> in an adult
Complications	<ul style="list-style-type: none"> <li>• <b>Metabolic Alkalosis &gt; Acidosis<sup>a</sup></b></li> <li>• <b>Hyperkalemia<sup>a</sup></b> → Ventricular arrhythmia or Cardiac Arrest; Hypokalemia (rare)</li> <li>• <b>Hypocalcemia<sup>a</sup></b> and/ or Citrate toxicity and Hypomagnesemia (rare)</li> <li>• Depletion of coagulation factors → Increased risk of <b>DIC</b></li> <li>• Dilutional <b>thrombocytopenia</b></li> <li>• Hypothermia</li> </ul>



## Transfusion Related Acute Lung Injury (TRALI)


<b>Incidence</b>	TRALI is reported by FDA as the most common cause of transfusion-associated fatality. Mortality rate, of TRALI at 5 to 10%.
<b>Mechanism</b>	<ul style="list-style-type: none"> <li>Antileukocyte antibodies in donor or patient plasma react with WBC complement system triggered to produce C3a and C5a tissue basophils and platelets release histamine and serotonin, lung capillary bed <ul style="list-style-type: none"> <li>Interstitial edema and fluid in alveolar air spaces, injury decreases gas exchange and hypoxia.</li> </ul> </li> </ul>
<b>Clinical features</b>	<ul style="list-style-type: none"> <li>Severe respiratory distress of sudden onset, caused by a syndrome of noncardiogenic pulmonary edema resembling the adult respiratory distress syndrome (ARDS).</li> <li>Lung injury is generally transient with <math>PO_2</math> levels returning to pretransfusion levels within 48–96 hours and CXR returning to normal within 96 hours.</li> </ul>
<b>Diagnosis</b>	<ul style="list-style-type: none"> <li>Diagnosis of a TRALI reaction is based on the onset of acute lung injury (ALI) <b>within 6 hours of transfusion</b>.</li> <li>Characterized by an acute onset of <b>hypoxemia (oxygen saturation &lt;90% by pulse oximetry for a patient breathing room air or a <math>PaO_2/FiO_2 \leq 300</math> mm Hg)</b>, bilateral infiltrates on frontal chest radiograph, and no evidence of circulatory overload.</li> </ul>
<b>Management</b>	Management <b>involves supportive measures</b> for the pulmonary edema and hypoxia, including ventilatory support if required.

## Transfusion Associated Graft versus Host Disease (TA-GVHD)

- Unlike transplant associated GVHD, TA-GVHD it is usually a fatal condition.
- Occurs in patients such as: – Immuno-deficient recipients of bone marrow transplants. – Immuno-competent patients transfused with blood from individuals with whom they have a compatible HLA tissue type, usually blood relatives particularly 1st degree. Signs and symptoms typically occur 10-12 days after transfusion and are characterized by: – Fever. – Skin rash and desquamation. – Diarrhoea. – Hepatitis. – Pancytopenia.
- Management:** Treatment is supportive; there is no specific therapy.
- Prevention:** Do not use 1st degree relatives as donors, unless gamma irradiation of cellular blood components is carried out to prevent the proliferation of transfused lymphocytes.


<b>R10<sup>th</sup> Latest Update</b> <b>Quality Control of Cryoprecipitate</b> <i>(Ref: DGHS technical manual)</i>	
Parameter	Quality requirement
Volume	10–20 mL
Factor VIII	80-120 units
*von-Willebrand factor	40–70% of the original
*Factor XIII	20–30% of the original
Fibrinogen	150–250 mg
*Fibronectin	55 mg





## Image-Based Questions

1. Given below is blood group determination by slide method. What blood group does the results suggest?



- a. A<sup>+</sup>  
b. B<sup>+</sup>  
c. AB<sup>+</sup>  
d. B<sup>-</sup>

2. Identify the instrument used in blood banking?




- a. Platelet agitator  
b. Centrifuge  
c. Cold storage  
d. Apheresis

3. The given product is used in the following indications?



- a. Factor V deficiency  
b. Thrombocytopenia  
c. DIC  
d. vWD



## Answers of Image-Based Questions

### 1. Ans. (a) A<sup>+</sup>

To determine the ABO type, red cells must be tested with anti-A and Anti-B and the serum/plasma tested with A and B red cells

- **Forward grouping**-identifies the antigens on the red cells
- **Reverse grouping**-identifies the presence of antibodies in the serum/plasma


ABO group	ABO Antisera (Forward)			ABO Cells (Reverse)			Rh D Antisera
	Anti-A	Anti-B	Anti-AB	A Cells	B Cells	O cells	Anti-D
A	+	0	+	0	+	0	
B	0	+	+	+	0	0	
AB	+	+	+	0	0	0	
O	0	0	0	+	+	+	

### 2. Ans. (d) Apheresis

- The machine is used to separate out one particular component from donor's blood while returning the remainder back to the circulation. Mostly used to make single donor platelets (SDP)

### 3. Ans. (c) DIC


- This is a bag of fresh frozen plasma (FFP) used to replace all coagulation factors in blood.



## Multiple Choice Questions

1. **Why is CPDA better than ACD for storage of blood?**  
(Recent Pattern Question 2020)
  - a. Improves oxygen transport
  - b. More citrate ions
  - c. It is less acidic
  - d. Hypertonicity of blood
2. **Which of the following may show ABO incompatibility?**  
(JIPMER Nov 2019)
  - a. Cryoprecipitate
  - b. Single donor platelets
  - c. Platelets rich platelets
  - d. FFP
3. **A patient comes with severe bleeding, 2 units PRBCs and 4 units of platelet concentrates have been collected from the blood bank. With only one IV access what will you do?**  
(AIIMS Nov 18)
  - a. Transfuse PRBC first and store Platelets at room temperature
  - b. Transfuse PRBC first and store Platelets at 2-6°C
  - c. Transfuse platelets first and store PRBC at 2-6°C
  - d. Transfuse platelets first and store PRBC at room temp
4. **A trauma patient presents at emergency department. There is no time for cross matching. FFP of which blood group can be transfused safely?**  
(AIIMS Nov 18)
  - a. O RH D POSITIVE
  - b. O RH D NEGATIVE
  - c. AB RH D POSITIVE
  - d. AB RH D NEGATIVE
5. **Which component is not stored in cold temperature?**  
(PGI Nov 2018)
  - a. RBC
  - b. Whole blood
  - c. Leucocyte removed RBC
  - d. Platelet concentrate
  - e. FFP
6. **Which of the following is true regarding blood transfusion of packed RBC?**  
(AIIMS May 18)
  - a. Started within 4 hours of receiving it from blood bank
  - b. Completed within 4 hours of receiving from blood bank
  - c. Wait till the patient is stable then transfuse, irrespective of any timing.
  - d. Completed within 6 hours of receiving from blood bank.
7. **Storage temperature of RBC, Platelet, and Fresh Frozen Plasma (FFP) are?**  
(AIIMS May 18)
  - a. RBC 2-6 C, Platelet 20-22C, FFP -30 C
  - b. RBC -30C, FFP 2-6 C, Platelet 20-22 C
  - c. RBC 20-22 C, Platelet 2-6 C, FFP -30C
  - d. RBC 20-22 C, FFP -30C, PLATELET 2-6 C
8. **Most immunogenic RBC blood group system?**  
(JIPMER 18)
  - a. Kell
  - b. Duffy
  - c. Kidd
  - d. K antigen
9. **Contraindication for platelet transfusion are all except?**  
(JIPMER 18)
  - a. Flavivirus infection
  - b. Thrombotic thrombocytopenic purpura (TTP)
  - c. Immune thrombocytopenic purpura (ITP)
  - d. Heparin induced thrombocytopenia
10. **True about single donor platelet transfusion?**  
(JIPMER 18)
  - a. Equal to 6-8 RDP
  - b. Stored in 2-6 degree
  - c. 10 days shelf life
  - d. Bedside leukodepletion done
11. **Which of the following is false regarding TRALI?**  
(Recent exam 2018)
  - a. Due to release of mediators from the neutrophils in the lungs
  - b. Occurs within 6 hours of transfusion
  - c. Occurs more common when the donor is a multiparous lady
  - d. None
12. **A voluntary donor, underwent apheresis for platelet donation for the first time at a platelet count of  $1.9 \times 10^3$ /mL. He started having tingling sensation (perioral) and numbness because?**  
(AIIMS Nov 2017)
  - a. His platelet count was low for donation
  - b. It was his first donation
  - c. Due to fluid depletion
  - d. Due to citrate based anticoagulant
13. **Which of the following anticoagulant preservative can be used to store blood, so that it can be kept for 35 days?**  
(AIIMS Nov 2017)
  - a. Acid citrate dextrose (ACD)
  - b. Citrate phosphate dextrose adenine (CADDP-1)
  - c. CPD Citrate phosphate dextrose
  - d. CP2D citrate phosphate double dextrose
14. **What is the optimal use of ratio of RBC: FFP: PLATELETS in massive transfusion blood Products in severely Injured Trauma Patients?**  
(JIPMER 2016)
  - a. 1:1:1
  - b. 1:2:3
  - c. 1:3:5
  - d. 2:1:4
15. **Which of the following blood components has the shortest shelf life?**  
(Recent Question 2016-17)
  - a. Red Blood Cells
  - b. Platelets
  - c. Fresh frozen plasma
  - d. Cryoprecipitate
16. **Blood transfusion reaction can lead to:**  
(PGI June 2016)
  - a. Acute glomerulonephritis
  - b. Myoglobinuria
  - c. Hemoglobinuria
  - d. Transfusion-related acute lung injury
  - e. Acute renal tubular necrosis
17. **Packed RBC is stored at?**  
(AIIMS May 2016)
  - a. 2-6°C
  - b. -2 to -8°C
  - c. -20°C
  - d. 20-24°C
18. **About Transfusion related lung injury what is not true?**  
(AIIMS Nov 2016)
  - a. Mortality < 10 % of cases
  - b. Subsides within 2-3 weeks
  - c. Supportive care is mainstay of treatment
  - d. Steroids use is not well established.
19. **Which one of the following tests has the highest chance of detecting HIV infection in a blood donor during the window period?**  
(Recent Question 2016-17)
  - a. Demonstration of antibody to HIV by ELISA
  - b. CD4 count
  - c. P24 antigen detection
  - d. Western blot test





20. **Best blood product to be given in a patient of multiple clotting factor deficiency with active bleeding is?** (AIIMS May 2015)
  - a. Fresh frozen Plasma
  - b. Whole blood
  - c. Packed RBCs
  - d. Cryoprecipitate
21. **Which of the following is false about TRALI?** (AIIMS May 2015)
  - a. Develops within 24 hours
  - b. Mostly seen after sepsis and cardiac surgeries
  - c. It's a cause of non-cardiogenic pulmonary edema
  - d. Plasma is more likely to cause it than whole blood
22. **What is used in irradiation of blood products before transplant surgery?** (Recent Question 2016)
  - a.  $\alpha$  Rays
  - b.  $\beta$  Rays
  - c.  $\gamma$  Rays
  - d. X-Rays
23. **Shelf life of  $\gamma$  irradiated packed RBCs is?** (Recent Question 2016)
  - a. 21 d
  - b. 28 d
  - c. 35 d
  - d. 42 d
24. **Chromosome of Rh gene is located on which chromosome?** (Recent Question 2016)
  - a. Chr 1
  - b. Chr 3
  - c. Chr 9
  - d. Chr 19
25. **ABO is located on which chromosome?** (Recent Question 2016)
  - a. Chr 1
  - b. Chr 3
  - c. Chr 6
  - d. Chr 9
26. **Bombay blood group contains?** (Recent Question 2016)
  - a. Anti H
  - b. Anti A, Anti B, Anti H
  - c. Anti A, Anti B
  - d. H antibody
27. **The major compatibility test before blood transfusion of cross matching of** (MH PG 2014)
  - a. Donor's red cells and Recipients serum
  - b. Donor's serum and Recipients red cells
  - c. Donor's serum and Recipients serum
  - d. Donor's red cells and Recipients red cells
28. **Spontaneous bleeding occurs when platelet count falls** (MH PG 2014)
  - a. 50,000/uL
  - b. 40,000/uL
  - c. 30,000/uL
  - d. 20,000/uL
29. **Life span of transfused platelets is?** (Recent Question 2016)
  - a. <24 hrs
  - b. 1-3 days
  - c. 3-5 days
  - d. 7-14 days
30. **Platelets in Single donor platelets is?** (Recent Question 2016)
  - a.  $2 \times 10^{11}$
  - b.  $1 \times 10^{11}$
  - c.  $4 \times 10^{11}$
  - d.  $1 \times 10^{12}$
31. **Shelf life is maximum for?** (Recent Question 2015)
  - a. Whole blood
  - b. FFP
  - c. Platelet concentrate
  - d. PRBC
32. **Indication of Cryoprecipitate?** (Recent Question 2015)
  - a. DIC
  - b. vWD
  - c. Hemophilia B
  - d. Severe plasma loss
33. **Which disease is transmitted by all the components of blood-** (Recent Question 2014)
  - a. Malaria
  - b. Syphilis
  - c. Toxoplasma
  - d. H. pylori
34. **Leuko-reduced blood products have lower WBCs than normal by what fold?** (Recent Question 2014)
  - a. 1 log reduction
  - b. 2 log reduction
  - c. 3 log reduction
  - d. 4 log reduction
35. **Cryoprecipitate is useful in-** (Recent Question 2014)
  - a. Hemophilia B
  - b. Thrombasthenia
  - c. Afibrinogenemia
  - d. Warfarin reversal
36. **Not true regarding fresh frozen plasma** (Recent Question 2013)
  - a. Supplies major coagulation factors
  - b. ABO match required
  - c. Should be used in replacement of factors in DIC
  - d. To be used within 30 minutes of having trauma
37. **Which of the following regarding Bombay blood group is false?** (AIIMS May 12)
  - a. Lack of H, A and B antigen on RBCs
  - b. Lack of H, A and B substance in saliva
  - c. Lack of antigens of several blood group systems
  - d. H, A and B antibodies will always be present in serum
38. **True about Blood transfusions?** (PGI June 12)
  - a. Antigen D determines Rh positivity
  - b. Febrile reactions is due to HLA antigens
  - c. Anti D is naturally occurring Ab
  - d. Cryoprecipitate contains all coagulation factors
  - e. FFP is rich in factor V and VIII
39. **Rise in hemoglobin levels after one unit of whole blood transfusion is?** (DNB Aug 12 Pattern)
  - a. 0.55 g%
  - b. 1 g%
  - c. 1.5 g%
  - d. 2 g%
40. **In massive transfusion of blood, citrate toxicity is primarily due to?** (DNB Aug 12 Pattern)
  - a. Hemolysis
  - b. Coagulopathy
  - c. DIC
  - d. Direct binding to calcium
41. **Granulocyte transfusion is recommended when WBC count is below?** (DNB Aug 12 Pattern)
  - a. 2000/uL
  - b. 1000/uL
  - c. 500/uL
  - d. 150/uL
42. **Indication for fresh frozen plasma is/are:** (PGI Nov 2011)
  - a. Hypovolemia
  - b. Nutritional supplement
  - c. Coagulation factor deficiency
  - d. Warfarin toxicity
  - e. Hypoalbuminemia
43. **ABO antigens are not found in -** (DNB June 10)
  - a. CSF
  - b. Plasma
  - c. Saliva
  - d. Semen



## Answers with Explanations

1. **Ans. (a) Improves oxygen transport**  
(Ref: Wintrobe 12th ed/pg 677)
2. **Ans. (d) FFP** (Ref: Wintrobe 12th ed/pg 660)
3. **Ans. (d) Transfuse platelets first and store PRBC at room temp**  
The transfusion time of the PRBC is PRBC 100-150 mL/hour and platelets / plasma 150-300 mL/hour. So while the platelets are transfused, the PRBC are kept at room temperature as it can't be returned back to blood bank nor can be stored in ward refrigerator.
4. **Ans. (d) AB RH D NEGATIVE**  
This question is based on antibodies in the plasma. As you know, AB blood group has both A & B antigen on RBC, but no antibodies. So, when plasma of AB blood group is given, there will be no antibodies to react with recipient's antigen, so is most preferred in this case without any cross-match done. The reason of choosing AB negative group is because small amount of RBC which might get contaminated while preparing FFP will not induce any alloimmunization in the recipient.
5. **Ans. (d) Platelet concentrate**
6. **Ans. (d) Completed within 4 hours of receiving from blood bank**
7. **Ans. (a) RBC 2-6 C, Platelet 20-22C, FFP -30 C**
8. **Ans. (a) Kell**
9. **Ans. (a) Flavivirus infection**
10. **Ans. (a) Equal to 6-8 RDP**
11. **Ans. (d) None**
12. **Ans. (d) Due to citrate based anticoagulant**  
Citrate based anticoagulant may chelate the calcium causing hypocalcemia which may cause the tingling sensation (perioral) and numbness
13. **Ans. (b) Citrate phosphate dextrose adenine (CADDP-1)**
14. **Ans. (a) 1:1:1**

(Ref: ASH Education Book, December 4, 2010 vol. 2010 no. 1 465-469)

Current data indicate that the early identification of coagulopathy and its treatment with RBCs, plasma, and platelets in a 1:1:1 unit ratio achieved with the use of fresh RBCs, thawed plasma, and platelets; limited use of cryoprecipitate; and accompanied with rapid hemorrhage

control may improve survival in the uncommon patient who presents with severe traumatic injury and life-threatening bleeding.

15. **Ans. (b) Platelets** (Ref: Wintrobe 12th ed/pg 677)  
**Shelf life for various components:**
  - Red Blood Cells- 35 to 42 days
  - Platelets- 5-7 days (shortest)
  - Fresh frozen plasma- 1 year
  - Cryoprecipitate- 1 year
16. **Ans. (c, d, e) c. Hemoglobinuria d. Transfusion-related acute lung injury e. Acute renal tubular necrosis**  
(Ref: Robbins(SEA) 9th/665-66; Harrison 19th/138e5-6)
17. **Ans. (a) 2-60C** (Ref: Wintrobe 12th ed/pg 677)  
**Storage temperature for various components:**
  - Red Blood Cells- 2-6 c
  - Platelets- 22-24 c
  - Fresh frozen plasma- -30c
  - Cryoprecipitate- -30c
18. **Ans. (b) Subsides within 2-3 weeks**  
(Ref: Wintrobe 14th ed. pg. 575)  
**Transfusion-related Acute Lung Injury (TRALI)**
19. **Ans. (c) P24 antigen detection**  
(Ref: Centers for Disease Control and Prevention (CDC))  
Based on the assumption that HIV RNA is first detected approximately ten days after exposure, window periods are as follows.
  - First detection of HIV RNA: approximately ten days after exposure (7 to 21 days).
  - First detection of p24: approximately 17 days after exposure (13 to 28 days).
  - First detection of antibodies: approximately 22 days after exposure (18 to 34 days).
  - Nucleic acid test (NAT) looks for HIV virus in the blood 7 to 28 days after infection.


### Also Note

- Note that p24 is not used for HIV diagnosis, It is a test of immune status of a patient.
- Western blot is a specific test and not sensitive and so is not used for early diagnosis

20. **Ans. (a) Fresh Frozen Plasma**

(Ref: Wintrobe 12th ed/pg 677)

FFP contains all coagulation factors and fibrinogen, so is the choice when there is either multiple coagulation factor deficiency or unknown factor deficiency.



**21. Ans. (a) Develops within 24 hours**

(Ref: Wintrob's 12th ed/pg 699)

- **Transfusion related acute lung injury (TRALI)** is seen usually 1-4 hrs after starting transfusion.
- In here, all the options look correct, but to choose the best option A is the best choice as the diagnosis of a TRALI reaction is based on the onset of acute lung injury (ALI) within 6 hours of transfusion.

**22. Ans. (c)  $\gamma$  Rays** (Ref: Wintrob's 12th ed/pg 699)

**Irradiation of Blood Products**

**Done to** prevent Transfusion Associated Graft-Versus Host Disease (TA-GVHD) in immunocompromised host

**Uses:** The recommended dose for the irradiation is 2,500 cGy at the center of the irradiation field, with a minimum dose of 1,500 cGy at any point in the field.

**Cellular blood products:** whole blood, red blood cells, platelets, granulocytes

**Who is at risk?** Patients who are immunocompromised and patients receiving transfusion from a relative (directed donation) are at increased risk of TA-GVHD. TA-GVHD has also been reported rarely in patients with a 'normal' immune system.

**Consequences:** **Irradiation of red blood cells and whole blood** results in reduced post-transfusion red cell recovery and increases the rate of efflux of intracellular potassium. It has no clinically significant effect on red cell pH, glucose, 2,3 DPG levels or ATP. Packs irradiated within 14 days of collection **expire 28 days after collection.**

**23. Ans. (b) 28 d** (Ref: Wintrob's 12th ed/pg 677)

**24. Ans. (a) **Chr1**** (Ref: Dacie/pg 486, 487)

Genetic loci	ABO	Rh
Chromosome	9	1

**25. Ans. (d) **Chr 9**** (Ref: Dacie/pg 486, 487)

**26. Ans. (b) **Anti A, Anti B, Anti H**** (Ref: Dacie/pg 487)

Bombay phenotype is characterized by absence of A, B, and H antigens on RBCs, and presence of **anti-A, anti-B, and anti-H** Antibodies

**27. Ans. (a) **Donor's red cells and Recipients serum****

(Ref: Dacie/pg 489)

Major compatibility test before blood transfusion requires cross-matching of donor's red cells which have the red cell Antigens with recipients serum having antibodies against those antigens

**28. Ans. (d) **20,000/ $\mu$ L**** (Ref: Wintrob's 12th ed/pg 660)

In general, the risk of significant spontaneous hemorrhage increases gradually as the platelet count drops to  $<50 \times 10^9/L$  and is high at counts  $<5 \times 10^9/L$ . So among the options the best answer is 20,000/ $\mu$ L.

**29. Ans. (c) **3-5 days**** (Ref: Wintrob's 12th ed/pg 687)

In healthy adults, the half-life of transfused platelets is 3 to 5 days. In thrombocytopenic patients, however, platelet survival is reduced.

**30. Ans. (d)  **$1 \times 10^{12}$****  (Ref: Wintrob's 12th ed/pg 677)

- Single donor platelets contain at least  $3 \times 10^{11}$  platelets in approximately 300 mL of plasma
- Otherwise in platelets concentrates, there are  $5.5 \times 10^{10}$  platelets/unit

**31. Ans. (b) **FFP**** (Ref: Wintrob's 12th ed/pg 677)

**32. Ans. (b) **vWD**** (Ref: Wintrobe's 12th ed/pg 695)

**Indications of cryoprecipitate are:**

- vWD
- Hemophilia A (not Hemophilia B)
- Factor XIII deficiency
- Fibrinogen deficiency

**33. Ans. (a) **Malaria****

(Ref: Wintrobe's 12th ed/pg 702-706)

\*Singh et al. Asian J Transfus Sci. 2010 Jul; 4(2): 73-77,

\*\* McCutcheon et al. PLoS One. 2011;6(8):e23169.

**Transfusion transmitted infections are:**

Viral	Hepatitis: A, B, C, G, HIV, CMV, EBV, HTLV I/II, Parvo B19
Bacterial	Syphilis, Sepsis
Parasites	Malaria, Babesia
Others	Creutzfeldt-Jakob ds

- Most recognized infectious organisms, with the notable exception of non-lipid-enveloped viruses and prions, have been shown to be inactivated easily by plasma processing methods.
- All clinically-relevant blood components transmit prion disease following a single blood transfusion\*\*

**Transmission of malaria by transfusion\***

- Reported to occur mainly from **single-donor products: red cells, platelets or white cell concentrates** (because of contamination with residual red cells), **cryoprecipitate & frozen red cells** after thawing and washing.
- **Transmission from single-donor fresh-frozen plasma has not been reported.**
- Transmission from cryoprecipitate is rare and likely to reflect the preparation method and the degree to which the starting plasma is cell free.

So answer for this question should have been prion, but as **this is not given in options, Malaria is the best possible answer among options provided**

**34. Ans. (c) **3 log reduction****

(Ref: Wintrobe's 12th ed/pg 665-666)

- Leukoreduced blood products achieve 3 log reduction ( $10^{-3}$ ) of WBC, thereby reducing the risk of FNHTR and CMV transmission.
- Risk of TRALI is not reduced by leukoreduction





35. Ans. (c) **Afibrinogenemia**

(Ref: Wintrobe's 12th ed/pg695)

Cryoprecipitate lacks Factor IX so is not useful in Hemophilia B

36. Ans. (d) **To be used within 30 minutes of having trauma**

(Ref: Wintrobe's 12th ed/pg 677)

37. Ans. (c) **Lack of antigens of several blood group systems**

(Ref: Dacie pg 487)

38. Ans. (a) **Antigen D determines Rh positivity**

(Ref: Wintrobe's 12th ed/pg 677-680, Rossi Principles of Transfusion Medicine 4th ed, pg 833-834)

### Discussing options one by one

a.	True as presence of antigen D is Rh +ve while d (absence of D) denoted Rh -ve
b.	False as FNHTR is caused by cytokines & pyrogens released from lymphocytes
c.	False as Anti D is IgG <sup>a</sup> and do not occur naturally
d.	False as it is rich in Fibrinogen; f-VIII, XIII & vWF only
e.	False as FFP is relatively deficient in factor V & VIII. <sup>a</sup>

39. Ans. (b) **1 g%** (Ref: Wintrobe's 12th ed/pg 677)

Hb increases by 1 gm/dl & Hct by 3-5% on transfusion of per unit PRBCs

40. Ans. (d) **Direct binding to calcium**

(Ref: W 12th ed/pg 701)

In **massive transfusion**, **large amount of citrate** present in blood bag **can chelate the calcium** causing hypocalcemia

41. Ans. (c) **< 500/uL** (Ref: Wintrobe's 12th ed/pg 677)

Indications for using granulocyte transfusion include:

- **Severe neutropenia**, defined as an absolute neutrophil count < **0.5 x 10<sup>9</sup> /L**
- A documented or presumed **severe bacterial or fungal infection**
- **No response of the infection** after 48 hours of appropriate antibiotic treatment
- Expected prolonged neutropenia
- Neutrophil recovery is expected and/or there is **anticipated therapy of curative potential planned**.

42. Ans. (c, d, e) **c. Coagulation factor deficiency; d. Warfarin toxicity; e. Hypoalbuminemia**

(Ref: Wintrobe's 12th ed/pg 677)

Indications of FFP are: Hemophilia A/B, Liver ds, Hypoalbuminemia, Warfarin overdose, DIC, TTP, Refractory vitamin K deficiency.

43. Ans. (a) **CSF** (Ref: Wintrobe's 12th ed/pg 677)

**Antigens also seen on/in:** Endothelial & epithelial cells<sup>o</sup>  
**Plasma, saliva, semen (not in CSF)<sup>o</sup>**

# Tumors of Soft Tissue & Head & Neck

## Key Points



- » **Lipoma is the most common soft tissue tumor**
- » **Most common sarcomas of adulthood is liposarcoma**
- » Most common site of Rhabdomyoma and Rhabdomyosarcoma is **head and neck**
- » **Prognosis of Rhabdomyosarcoma depends on histologic type & location of tumor influence survival**
- » **Embryonal variety is the most common Rhabdomyosarcoma**
- » **Most important indicator of malignancy in soft tissue tumor is mitotic index**
- » **t (X;18) is the cytogenetics seen in synovial cell sarcoma**
- » **Most common** salivary gland tumor (overall): Pleomorphic adenoma
- » Ameloblastoma is the **most common** odontogenic tumor
- » Retinoblastoma is the most common **primary malignant tumor of the eye** in children

## Key Recent Updates

- » GIST & PNST are now included in soft tissue sarcoma.



## SOFT TISSUE TUMORS

Fatty tumors	
<b>Lipoma</b>	<ul style="list-style-type: none"> <li>• <b>Most common soft tissue tumor;</b><sup>Q</sup> Painless except angiolipoma<sup>Q</sup></li> <li>• Cytogenetics: Chr 6p, 12q and 13q involved</li> </ul>
<b>Liposarcoma</b>	<ul style="list-style-type: none"> <li>• <b>Most common sarcomas of adulthood</b><sup>Q</sup></li> <li>• <b>Most common: Proximal extremities &amp; retroperitoneum</b><sup>Q</sup></li> <li>• <b>Morphology:</b> Lipocytes with supernumerary rings &amp; giant rod chromosomes - 12q (<i>MDM2</i> oncogene) &amp; Lipoblasts with scalloping of nucleus</li> </ul>
Tumors & Tumor-like lesions of fibrous origin	
<b>Fibromatoses</b>	<ul style="list-style-type: none"> <li>• <b>Superficial:</b> Palmar (<b>Dupuytren's contracture</b>), Plantar, Penile (<b>Peyronie disease</b>)<sup>Q</sup></li> <li>• Deep: Desmoid tumors-Locally aggressive, Associated with Gardner's syndrome (APC)</li> </ul>
<b>Fibrosarcoma</b>	<ul style="list-style-type: none"> <li>• <b>Most common in extremities;</b><sup>Q</sup> Aggressive, Recurs &amp; Metastasizes</li> <li>• Herringbone pattern on Histology</li> </ul>
Tumors of skeletal muscle	
<b>Rhabdomyoma</b>	<ul style="list-style-type: none"> <li>• <b>Benign;</b> Most common in- <b>Head, Neck &amp; heart</b><sup>Q</sup>; Associated with Tuberous Sclerosis</li> <li>• Histology- Polygonal Rhabdomyoblast, <b>Spider cells</b></li> </ul>
<b>RMS</b>	Discussed below in detail
Tumors of Smooth muscles	
<b>Leiomyoma</b>	<b>Uterine leiomyoma</b> are the <b>most common Neoplasms in females</b> <sup>Q</sup>
Leiomyosarcoma	<b>Indicator of malignancy:</b> Mitotic index <sup>Q</sup> >Atypia> Necrosis
Tumors of uncertain histogenesis	
<b>Synovial Cell Sarcoma</b>	<ul style="list-style-type: none"> <li>• <b>Only 10% intraarticular;</b><sup>Q</sup> MC around knee</li> <li>• H/E: <b>Biphasic</b><sup>Q</sup>- epithelial &amp; mesenchymal (spindle cells)</li> <li>• <b>Stains:</b> Keratin, vimentin, S-100 &amp; EMA; t(X;18) → <b>poor prognosis</b><sup>Q</sup></li> </ul>
<div style="display: flex; justify-content: space-around; align-items: flex-end;"> <div style="text-align: center;">  <p>Lipoma</p> </div> <div style="text-align: center;">  <p>Herring bone pattern</p> </div> </div>	

## RHABDOMYOSARCOMA (RMS)

- Malignant primitive mesenchymal tumor of skeletal muscle (**arises from pluripotent muscle cells**)
- **MC soft-tissue sarcoma of childhood & adolescence**<sup>Q</sup>
- **Most common—head/neck;**<sup>Q</sup> 2<sup>nd</sup> MC—genitourinary tract
- **Diagnostic cell in all types is:**
  - **Rhabdomyoblast-** contains **eccentric eosinophilic granular cytoplasm**, rich in thick and thin filaments
  - **Tadpole or strap cells**<sup>Q</sup> - **Elongated Rhabdomyoblasts** that may contain cross-striations visible by light microscopy
- **Stains used-** Desmin, MYOD1 & myogenin.





Classification: Histological sub-types include- Embryonal, Alveolar & Pleomorphic	
<b>Embryonal RMS (Most common)<sup>Q</sup></b>	<ul style="list-style-type: none"> <li>Occurs in <b>children &lt; 10 yrs</b> of age</li> <li><b>Chr 11p</b> involved</li> <li>Histology: Consists of <b>primitive round &amp; spindle cells</b></li> <li>A submucosal zone of hypercellularity → <b>cambium layer<sup>Q</sup></b></li> </ul>
<b>Sarcoma botryoides</b>	<ul style="list-style-type: none"> <li>A variant of <b>Embryonal RMS</b></li> <li>More common in <b>age &lt; 5 yrs</b></li> <li>Arises from walls of hollow mucosal lined structures like vagina, bladder</li> <li><b>Grape-like clusters<sup>Q</sup></b> seen</li> <li>H/E: <b>Tennis racket cells<sup>Q</sup></b> on light microscopy</li> </ul>
<b>Alveolar RMS (20%)</b>	<ul style="list-style-type: none"> <li>Commonly arises in the deep musculature of the extremities</li> <li>Cytogenetics: t(2;13) &amp; t(1;13)</li> </ul>
<b>Pleomorphic RMS</b>	<ul style="list-style-type: none"> <li>Rare; poor prognosis</li> <li>Arise in the deep soft tissue of adults</li> </ul>
Prognosis of RMS: Histologic type & location of tumor influence survival	
<ul style="list-style-type: none"> <li>Best prognosis: Botryoid subtype</li> <li>Intermediate prognosis: Embryonal NOS</li> <li>Poor prognosis: pleomorphic and alveolar</li> </ul>	



Botryoides tumor

Sarcomas in which Lymphatic Metastasis is seen -

## Mnemonic

### RACE For MS

- R** : Rhabdomyosarcoma
- A** : Angiosarcoma
- C** : Clear cell sarcoma
- E** : Epithelial cell sarcoma
- For** : Fibrosarcoma
- M** : Malignant fibrous histiocytoma
- S** : Synovial cell sarcoma



### High Yield Facts

- Most common** salivary gland tumor (overall): **Pleomorphic adenoma<sup>Q</sup>**
- Most common benign** salivary gland tumor: Pleomorphic adenoma<sup>Q</sup>
- Most common malignant** salivary gland tumor: Mucoepidermoid carcinoma<sup>Q</sup>
- Most common salivary gland tumor in **children**: Hemangioma<sup>Q</sup>
- Most common **malignant** salivary gland tumor in **children**: Mucoepidermoid carcinoma.<sup>Q</sup>
- Salivary gland tumor with worst prognosis is Adenoid cystic carcinoma**
- Salivary gland tumor being malignant is inversely proportional to size of the gland.<sup>Q</sup>**

## SALIVARY GLAND TUMORS



Histological classification of common benign and malignant salivary gland tumors

Benign Tumors <sup>Q</sup>	Malignant Tumors <sup>Q</sup>
<ul style="list-style-type: none"> <li>Pleomorphic adenoma</li> <li>Warthin tumor's</li> <li>Oncocytoma</li> <li>Other adenoma (Basal cell adenoma)</li> </ul>	<ul style="list-style-type: none"> <li>Mucoepidermoid carcinoma</li> <li>Adenocarcinoma NOS</li> <li>Acinic cell carcinoma</li> <li>Adenoid cystic carcinoma</li> </ul>

### Benign Neoplasms


#### Pleomorphic Adenoma

- Commonest neoplasm of salivary gland.
- Most common salivary gland involved: **Parotid<sup>Q</sup>** > Submandibular > Minor salivary gland.
- Shows both epithelial and mesenchymal differentiation. Thus known as **mixed tumor<sup>Q</sup>**
- Carcinoma arising in pleomorphic adenoma is called carcinoma, e.g. pleomorphic adenoma.
- The malignant component of Ca ex pleomorphic adenoma is most often **adenocarcinoma not otherwise specified**



### Warthin's Tumor

- Also known as papillary **cystadenoma lymphomatosum**<sup>Q</sup>
- Second most common** benign salivary gland tumor (after pleomorphic adenoma)<sup>Q</sup>
- Common tumor of parotid gland with double layer of epithelial cells resting on dense lymphoid stroma<sup>Q</sup>
- Surface palisading of oncocytic** columnar cells are seen<sup>Q</sup>



## Malignant Neoplasms

### Mucoepidermoid Carcinoma

- Most common** malignant salivary gland tumor<sup>Q</sup>
- Occur mainly in **parotids**
- Most common radiation induced** neoplasm in salivary glands
- Consist of variable number of **squamous cell**, **mucus secreting cells** and **intermediate cells**.<sup>Q</sup>


### Adenoid Cystic Carcinoma

- Malignant tumor** most commonly seen in **minor salivary glands**

- Among major salivary glands, **parotid gland** is the **most common site**


- Microscopically:** Cribriform arrangements of bland cells arranged around cystic spaces - these are not truly cystic and are thought to be filled with material produced by tumour cells.

- Unpredictable tumor with **perineural invasion**.



### Acinic Cell Carcinoma

- Relatively **uncommon malignant tumor**
- 2nd childhood salivary gland** malignancy after mucoepidermoid carcinoma
- Most commonly arise in **parotid** > submandibular > minor salivary gland.
- Composed of cells resembling **normal serous acinar cells**<sup>Q</sup>




## ORAL CAVITY

### Ameloblastoma

- It is the **most common** odontogenic tumor<sup>Q</sup>
- Most commonly occur in 3-5 decade
- Most common site- **Posterior mandible (80%)**<sup>Q</sup>
- Risk factors:** impacted teeth, dentigerous cyst
- It is benign but locally aggressive





### Morphology of Ameloblastoma

- Columnar basal cells in palisading arrangement with vacuolated cytoplasm.
- Hyperchromatic nuclei **polarized away** from basement membrane.
- Suprabasal cells** loosely textured and non-cohesive, resembling **stellate reticulum**.
- Treatment include wide surgical excision.
- Metastasizes rarely to lungs or CNS.
- Metastases associated with tumor of **long duration**, **multiple** surgical procedures, **radiation therapy**.




### High Yield Facts

- Precancerous Lesions of Oral Cavity** are: Leukoplakia (most common) & Erythroplakia
- Most Common Malignancy of Oral Cavity** is Squamous Cell Carcinoma (classically linked to tobacco, HPV infection)

## RETINOBLASTOMA

- Most common **primary malignant tumor of the eye** in children<sup>Q</sup>
- Cell of origin is a neuronal progenitor**.<sup>Q</sup>



Rosette in Retinoblastoma

### Genetics

- Caused by **loss of function mutation of the Retinoblastoma (Rb) gene**<sup>Q</sup>
- Familial (40% cases; autosomal dominant)**:
  - Carrying **germline mutation** in one copy of the gene (**first hit**).<sup>Q</sup>
  - Spontaneous somatic mutation** in the second normal allele (**second hit**) → retinoblastoma<sup>Q</sup>
  - Often bilateral** with increased risk of **developing osteosarcoma**<sup>Q</sup> and other **soft tissue sarcomas**.
- Sporadic cases (60%)**:
  - Somatic mutation in both the alleles**.
  - Involves **only one eye** and there is **no increased risk of other cancer**<sup>Q</sup>

### Morphology

- Small, round cells with **hyperchromatic nuclei**.<sup>Q</sup>
- Flexner Wintersteiner rosettes**<sup>Q</sup> (a single layer of tumor cells aggregated around a central lumen)
- Fleurettes**<sup>Q</sup> (a single layer of cells with **tapering cytoplasmic processes** that protrude into the center of the rosette)

### Adverse Prognostic Factors

- Extraocular extension**<sup>Q</sup>, **invasion along optic nerve**<sup>Q</sup> and **choroidal invasion**.<sup>Q</sup>



### High Yield Facts

- Rb gene was the first tumor suppressor gene discovered
- Trilateral retinoblastoma**: **Bilateral** retinoblastoma along with **pinealoblastoma**
- Fleurettes** are seen in well differentiated tumors and represent **photoreceptor differentiation**.<sup>Q</sup>

### R10<sup>th</sup> Latest Updates


- GIST & peripheral nerve sheath tumors (PNST) are included in soft tissue sarcomas
- Undifferentiated-unclassified tumors is added & include pleomorphic sarcoma
- Round cell & mixed types of liposarcoma categories are removed





## Multiple Choice Questions

1. Which of the following translocation is seen in Myxoid liposarcoma? (Recent Pattern Question 2020)
  - a. t(11:22)
  - b. t(14:18)
  - c. t(x:18)
  - d. t(12:16)
2. Which of the following is not small round cell tumor? (AIIMS Nov 2019)
  - a. Neuroblastoma
  - b. Retinoblastoma
  - c. Hemangioblastoma
  - d. Ewing sarcoma
3. In alveolar variant of rhabdomyosarcoma, the resultant fusion protein is believed to function as: (JIPMER Nov 2019)
  - a. Activated growth factor receptor
  - b. Chimeric transcription factor
  - c. Constitutively active kinase
  - d. Novel growth factor
4. Sarcoma spreading by lymphatic spread is: (Recent Question 2016-17)
  - a. Alveolar RMS
  - b. Embryonal RMS
  - c. Liposarcoma
  - d. Fibrosarcoma
5. Spindle shaped cells is/are seen in which sarcoma: (PGI May 2015)
  - a. Osteosarcoma
  - b. Chondromyosarcoma
  - c. Embryonal rhabdomyosarcoma
  - d. Leiomyosarcoma
  - e. Fibrosarcoma
6. Most common site of Rhabdomyosarcoma is: (Recent Question 2016)
  - a. Head and neck
  - b. Trunk
  - c. Urogenital
  - d. Lungs
7. Hyperglycemia is associated with: (Recent Question 2016)
  - a. Osteosarcoma
  - b. Chondroblastoma
  - c. Chondrosarcoma
  - d. Osteoma
8. Olliers disease also known as? (Recent Question 2016)
  - a. Osteosarcoma
  - b. Enchondromatosis
  - c. Multiple myeloma
  - d. Enchondrosis
9. Cambium layer is seen in: (Recent Question 2015)
  - a. Embryonal rhabdomyosarcoma
  - b. Pleomorphic rhabdomyosarcoma
  - c. Alveolar rhabdomyosarcoma
  - d. Undifferentiated rhabdomyosarcoma
10. Which of the following are primarily spindle cell tumor? (PGI Nov 2015)
  - a. Rhabdomyosarcoma
  - b. Fibrosarcoma
  - c. Leiomyosarcoma
  - d. Kaposi sarcoma
11. Most common soft tissue tumor is: (Recent Question 2015)
  - a. Fibroma
  - b. Lipoma
  - c. Leiomyoma
  - d. Rhabdomyoma
12. Most common Sarcoma of adulthood is: (Recent Question 2015)
  - a. Liposarcoma
  - b. Fibrosarcoma
  - c. Rhabdomyosarcoma
  - d. Leiomyosarcoma
13. Most common site of Fibrosarcoma is: (Recent Question 2014)
  - a. Head and Neck
  - b. Extremities
  - c. Abdominal wall
  - d. Joints
14. Which of the following is associated with Tuberous Sclerosis: (Recent Question 2014)
  - a. Fibroma
  - b. Lipoma
  - c. Leiomyoma
  - d. Rhabdomyoma
15. Tadpole cells or comma shaped cells on histopathology are seen in: (Recent Question 2014)
  - a. Trichoepithelioma
  - b. Spideroma
  - c. Rhabdomyosarcoma
  - d. Histiocytoma
16. Most important indicator of malignancy in smooth muscle tumors is: (Recent Question 2014)
  - a. Mitotic index
  - b. Atypia
  - c. Necrosis
  - d. Cellularity
17. True about rhabdomyosarcoma? (PGI May 2014)
  - a. Arise from pluripotent cells
  - b. Tennis racket cells on light microscopy
  - c. Sarcoma botryoides, is a variant of embryonal rhabdomyosarcoma
  - d. Most common type is Embryonal
  - e. Most common site is lower extremity
18. Retinoblastomas arising in the context of germ-line mutations not only may be bilateral, but also may be associated with \_\_\_\_\_ (so called "trilateral" retinoblastoma) (AP PGME 14)
  - a. Medulloblastoma
  - b. Pinealoblastoma
  - c. Neuroblastoma
  - d. Hemangioblastoma
19. Immunohistochemical stain marker of Rhabdomyosarcoma (JIPMER 2014)
  - a. Desmin
  - b. Vimentin
  - c. Cytokeratin
  - d. Neurofilament
20. Primitive Neuro Ectodermal Tumor of bones and soft tissues shows which of the following cytogenetic abnormalities? (PGI May 2013)
  - a. EWS-ETVI
  - b. EWS-ERG
  - c. EWS-FLI1
  - d. EWS-ATFI
  - e. EWS-WTI
21. Bible Bump is a: (Recent Question 2016, 2013)
  - a. Synovial cyst
  - b. Malformation
  - c. Neurofibroma
  - d. Myxomatous degeneration
22. "Biphasic pattern" on histology is seen in which tumor: (DPG 10, MH 02)
  - a. Rhabdomyosarcoma
  - b. Synovial cell sarcoma
  - c. Osteosarcoma
  - d. Neurofibroma



23. **Glomus tumor is seen in:** (AIIMS Nov 10)  
a. Retroperitoneum      b. Soft tissue  
c. Distal portion of digits      d. Proximal portion of digits

#### TUMORS OF ORAL CAVITY

24. **Most common cancer of oral cavity histologically is:** (Recent Question 2016-17)  
a. Squamous cell Ca      b. Adeno ca  
c. Adenocystic Ca      d. Odontogenic ca  
(Recent Question 2016-17)
25. **True about Dentigerous cyst:**  
a. Arises in relation to unerupted teeth  
b. It most commonly encroaches maxillary antrum  
c. Mandibular third molar is common site  
d. Common in mandible
26. **Most malignant salivary gland tumor is:** (Recent Question 2016)  
a. MucoepidermoidCa  
b. Acinic cell carcinoma  
c. Adenoid cystic carcinoma  
d. Pleomorphic adenoma
27. **Cystic spaces lined by double layer of neoplastic epithelial cells resting on dense lymphoid tissue is a feature of:** (AP PGME 2015)  
a. Aneurismal bone cyst      b. Dermoid cyst  
c. Warthin tumor      d. Hashimoto's thyroiditis
28. **Warthins tumor true statement is:** (PGI Nov 2015)  
a. Sinuses are present  
b. Clefts are present  
c. Pallisading cells are seen  
d. Papillary CystadenomaLymphomatosum  
e. Arises from sublingual gland commonly

29. **Most common site of Ameloblastoma is:** (Recent Question 2014)  
a. Anterior Mandible  
b. Posterior Mandible  
c. Maxilla  
d. Zygomatic bone
30. **The most common pre-malignant condition of oral carcinoma is:** (Recent Question 2014)  
a. Leukoplakia      b. Erythroplakia  
c. Lichen planus      d. Fibrosis
31. **Most common salivary gland tumor?** (Recent Question 2013)  
a. Mucoepidermoid carcinoma  
b. Pleomorphic adenoma  
c. Warthins tumor  
d. Oncocytoma
32. **Pleomorphic adenoma has:** (Recent Question 2013)  
a. Columnar cells enclosing lymphoid stroma  
b. Exclusively myoepithelial cells  
c. Epithelial cells in chondroid matrix  
d. Nests of squamous cells and vacuolated cells containing mucin
33. **Which of the following head and neck tumor has worst prognosis?** (MH 11, DNB June 08)  
a. Adenoid cystic carcinoma  
b. Acinic cell carcinoma  
c. Cystadenolymphoma  
d. Mucoepidermoid carcinoma
34. **Most common type of salivary neoplasm:** (PGI Nov 10)  
a. Adenocysticcacinorna  
b. Mixed cell parotid neoplasm  
c. Epidermoid carcinoma  
d. Adenocarcinoma  
e. Adenolymphoma



## Answers with Explanations

### 1. Ans. (d) **t (12;16)** (Ref: Robbins 9th/pg 1220)

- Malignant tumor composed of primitive nonlipogenic mesenchymal cells, signet ring lipoblasts and prominent myxoid stroma with a highly characteristic branching vascular pattern. It shows Recurrent molecular alteration with either t(12;16)(q13;p11.2) *FUS-DDIT3* or very rarely (~2%) t(12;22)(q13;q12) *EWSR1-DDIT3* rearrangements

### 2. Ans. (c) **Hemangioblastoma** (Ref: Robbins 9th/pg/1222)

Diffuse round cell pattern is seen in: Ewing's sarcoma, Primitive neuroectodermal tumor (PNET), Merkel cell carcinoma, Embryonal rhabdomyosarcoma (ERMS), Small cell carcinoma, Lymphoma

#### Round cell pattern with rosettes

- Flexner's (also called Flexner - Winterstein, true rosettes) - e.g., Retinoblastoma, PNET
- Homer Wright rosette-center has no lumen, but abundant fibrillary material e.g., neuroblastoma.

### 3. Ans. (b) **Chimeric transcription factor** (Ref: Robbins 9th/pg 1253)

### 4. Ans. (d) **Fibrosarcoma** (Ref: Robbins 9th/pg 730-740)

### 5. Ans. (a, c, d, e); **a. Osteosarcoma c. Embryonal rhabdomyosarcoma, d. Leiomyosarcoma, e. Fibrosarcoma** (Ref: Robbins 9th/pg 474)

#### Differential diagnoses of spindle cell sarcomas:

- Fibrosarcoma
- Benign fibrous histiocytoma
- Embryonal rhabdomyosarcoma
- Leiomyosarcoma
- Synovial sarcoma: t(X;18) (p 11;q11) fusion of SYT-SSX.
- Malignant Peripheral Nerve Sheath Tumours (MPNST)
- **Vascular tumors:** hemangio-endothelioma, hemangio-pericytoma, angiosarcoma, lymphangiosarcoma, and Kaposi's sarcoma.

Note that osteosarcoma has osteoblasts which are plump cells and not typical spindle cells.

### 6. Ans. (a) **Head and neck** (Ref: Robbins 9th/pg 1253)

- MC site of rhabdomyosarcoma is head/neck; genitourinary tract

### 7. Ans. (c) **Chondrosarcoma**

(Ref: Cancer 42:603-610, 1978; Hypoglycemia as Paraneoplastic syndrome in some tumors: Mesenchymal tumors, sarcomas, adrenal, hepatic, gastrointestinal, kidney, prostate, Cervix (small-cell carcinoma)

The first association of hyperglycemia was made by Glicksman and Rawson who noted a 25% incidence of "diabetes" in patients with sarcomas of bone. Marcove

and Francis found hyperglycemia in 85% of their chondrosarcoma patient. and Turner and Horne reported similar abnormalities associated with fibrosarcoma

### 8. Ans. (b) **Enchondromatosis** (Ref: Robbins 9th/pg)

Ollier disease is a rare, non inherited disease of unknown cause characterized by multiple enchondromas.

### 9. Ans. (a) **Embryonal rhabdomyosarcoma**

(Ref: Robbins 9th/pg 1253)

In **Sarcoma botryoides**, a variant of embryonal rhabdomyosarcoma, histology shows a **cambium layer** where the tumors abut the mucosa of an organ, they form a submucosal zone of hypercellularity.

### 10. Ans. (a, b, d) **a. Rhabdomyosarcoma; b. Fibrosarcoma; d. Kaposi sarcoma** (Ref: Robbins 9th/pg 474)

### 11. Ans. (b) **Lipoma** (Ref: Robbins 9th/pg 1220; 8th/pg 1249)

### 12. Ans. (a) **Liposarcoma** (Ref: Robbins 9th/pg 1220)

### 13. Ans. (b) **Extremities**

(Ref: Robbins 9th/pg 1221; 8th/pg 1250)

#### **Fibrosarcoma**

- It is **most common in extremities**
- It shows **Herringbone pattern** on histopathology
- It is **Aggressive, Recurs and Metastasizes**

### 14. Ans. (d) **Rhabdomyoma**

(Ref: Nelson Textbook of Pediatrics, 19th Edition, 2011; pg 2049)

#### **Tuberous Sclerosis**

It is inherited as an **autosomal dominant trait** with variable expression

Important **clinical manifestations of Tuberous Sclerosis** include:


Skin Lesions	CNS Lesions	Other Tumors
<ul style="list-style-type: none"> <li>• <b>Facial angiofibroma</b></li> <li>• Ungual/ periungual fibroma (non-traumatic)</li> <li>• <b>Ash-leaf macules</b></li> <li>• <b>Shagreen patch</b></li> <li>• Confetti skin lesions</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Cortical tubers</b></li> <li>• <b>Subependymal nodule</b></li> <li>• <b>Subependymal giant cell astrocytoma</b></li> </ul>	<ul style="list-style-type: none"> <li>• <b>Multiple retinal hamartomas</b></li> <li>• <b>Cardiac rhabdomyoma</b></li> <li>• <b>Renal angiomyolipoma</b></li> </ul>

### 15. Ans. (c) **Rhabdomyosarcoma** (Ref: Robbins 9th/pg 1222)

In Rhabdomyosarcoma, **diagnostic cell in all types is:**

- **Rhabdomyoblast** - contains **eccentric eosinophilic granular cytoplasm**, rich in thick and thin filaments





- Tadpole or strap cells<sup>Q</sup> - Elongated Rhabdomyoblasts that may contain cross-striations visible by light microscopy

16. Ans. (a) **Mitotic index** (Ref: Robbins 9th/pg 1223)

Indicator of malignancy in soft tissue tumors (Leiomyosarcoma): Mitotic index<sup>Q</sup> > Atypia > Necrosis.

17. Ans. (a,b,c,d) a. Arise from pluripotent cells; b. Tennis racket cell on light microscopy; c. Sarcoma botryoides, is a variant of embryonal rhabdomyosarcoma; d. Most common type is Embryonal. (Ref: Robbins 9th/pg 1222)

18. Ans. (b) **Pinealoblastoma** (Ref: Robbins 9th/pg 1339)

- “Trilateral” retinoblastoma refers to bilateral Retinoblastomas plus pinealoblastoma, occurring in individuals inheriting a germline mutation of one RB allele; it is associated with a dismal outcome
- Retinoblastoma is the most common primary intra-ocular malignancy of children.

19. Ans. (a) **Desmin** (Ref: Robbins 9th/pg 1222; 8th/pg 1253)

20. Ans. (a,b,c) a. EWS-ETV1; b. EWS-ERG; c. EWS-FLI1

(Ref: Robbins 9th/pg 1219; 8th/pg 1249)

### Cytogenetics in Ewing's Sarcoma

- t(11;22) → fusion of EWS gene with FLI1 gene → EWS-FLI1 (most common, 90%)

21. Ans. (a) **Synovial cyst**

(Ref: Atlas of Soft tissue & bone pathology, 2015; pg 54)

- It is also known as a ‘ganglion cyst’, ‘myxoid cyst’, ‘Gideon's Disease’, ‘Bible Cyst’ or ‘Bible bump’
- It is a non-neoplastic soft tissue lump that may occur in any joint, but most often occurs in the hands or feet.
- Caused by leakage of fluid from the joint into the surrounding tissue.

22. Ans. (b) **Synovial cell sarcoma** (Ref: R 9th/pg 1220-1224)

A biphasic tumor refers to neoplastic tissue which is characterized by two different cellular elements.

### Biphasic tumors include

- |   |  |
|---|--|
| <ul style="list-style-type: none"> <li>• Spindle Cell Carcinoma</li> <li>• Nasopharyngeal Carcinoma –</li> <li>• Synovial Sarcoma (SC)</li> </ul> | <ul style="list-style-type: none"> <li>• Mesothelioma</li> <li>• Pleomorphic adenoma</li> <li>• Melanoma</li> <li>• Malignant Mixed Mullerian tumor</li> </ul> |
|---|--|

An example of Triphasic tumor is Wilm's tumor

23. Ans. (c) **Distal portion of digits** (Ref: Robbins 9th/pg 517)

24. Ans. (a) **Squamous cell Ca** (Ref: R 9/Pg 732)

25. Ans. (a) Arises in relation to unerupted teeth, (c) Mandibular third molar is common site, (d) Common in mandible

(Ref: Robbins 9th/ 734; Manipal Surgery 4th/292-93; Harshman 7th/511-12)

- Dentigerous cyst is defined as a cyst that originates around the crown of an unerupted tooth and is thought to be the result of fluid accumulation between the developing tooth and the dental follicle.
- Radiographically, these are unilocular lesions most often associated with impacted third molar (wisdom) teeth. Histologically, they are lined by a thin layer of stratified squamous epithelium. Often, there is a dense chronic inflammatory cell infiltrate in the connective tissue stroma.
- Complete removal of the lesion is curative.

26. Ans. (a) **Mucoepidermoid Carcinoma**

(Ref: R 9th/pg 757-758)

27. Ans. (c) **Warthin's tumor** (Ref: Robbins 9th/pg 757-758)

### Histology of Warthin tumour:

- Cystic spaces are lined by a double layer of neoplastic epithelial cells resting on a dense lymphoid stroma sometimes bearing germinal centers.
- The double layer of lining cells is distinctive; the upper layer consists of palisading columnar cells while the lower layer is comprised of cuboidal to polygonal cells.

28. Ans. (b, c, d); b. Clefts are present; c. Palisading cells are seen; d. Papillary Cystadenoma Lymphomatosum

(Ref: Robbins 9th/pg 757- 758)

29. Ans. (b) **Posterior Mandible** (Ref: Robbins 9th/pg 735)

30. Ans. (a) **Leukoplakia** (Ref: Robbins 9th/pg 731)

### Precancerous lesions of oral cavity

- Leukoplakia ( most common)
- Erythroplakia

Most common malignancy of oral cavity: squamous cell carcinoma (classically linked to tobacco, HPV infection)

31. Ans. (b) **Pleomorphic adenoma**

(Ref: Robbins 9th/pg 744)

32. Ans. (c) **Epithelial cells in chondroid matrix**

(Ref: Robbins 9th/pg 744)

33. Ans. (a) **Adenoid cystic carcinoma**

(Ref: R 9th/pg 744)

34. Ans. (b) **Mixed cell parotid neoplasm**

(Ref: R 9th/pg 744)



---

---

---

---

---

---

[illegible]

# Diseases of Muscles

## Key Points

- » **Duchenne & Becker Muscular dystrophy** are X-linked recessive myotonic dystrophy
- » **Central-core disease, Nemaline and Myotubular** (centronuclear) are congenital myopathies

## Key Recent Updates

- » Duchenne muscular dystrophy (DMD) is caused by mutations in DMD gene, largest gene in human body.





## INHERITED DISORDERS OF SKELETAL MUSCLES


**Muscular Dystrophies:** Inherited disorders of skeletal muscle with **progressive muscle damage**<sup>Q</sup>

Disease	Genetics	Clinical Features	Pathologic Findings
<b>Duchenne &amp; Becker Muscular dystrophy</b> (Becker: milder disease)	XR; <sup>Q</sup> Dystrophin gene <sup>Q</sup> on Chr X; <sup>Q</sup>	<ul style="list-style-type: none"> <li>Onset: 2-5 yrs</li> <li>Gower sign</li> <li>Pseudohypertrophy of calf muscles</li> <li>CPK level ↑<sup>Q</sup></li> </ul>	<ul style="list-style-type: none"> <li>Prominent <b>variation in size</b><sup>Q</sup> of muscle fibers</li> <li>Mixture of <b>atrophic &amp; regenerating myofibers</b><sup>Q</sup> seen</li> <li><b>Fatty infiltration</b><sup>Q</sup> (See fig 1)</li> </ul>
<b>Myotonic dystrophy</b>	AD; <sup>Q</sup> CTG repeats in DMPK gene on Chr19	<ul style="list-style-type: none"> <li>Skeletal muscle weakness (myotonia)</li> <li>Cataract</li> <li>Endocrinopathy</li> <li>Cardiomyopathy</li> </ul>	<ul style="list-style-type: none"> <li><b>Intrafusal fibers</b><sup>Q</sup> of muscle spindle affected</li> </ul>
<b>Fascioscapulohumeral muscular dystrophy</b>	AD; DUX4 gene on Chr 4	<ul style="list-style-type: none"> <li><b>Facial muscle &amp; Shoulder</b> girdle muscle weakness</li> </ul>	<ul style="list-style-type: none"> <li><b>Dystrophic</b> myopathy with <b>inflammatory infiltrates</b> in muscle</li> </ul>


### Congenital Myopathies

Congenital structural abnormalities of skeletal muscles, that present in infancy and remains **static or improves over time**.<sup>Q</sup>

Disease	Genetics	Clinical Findings	Pathologic Findings
<b>Central-core disease</b>	AD; Ryanodine receptor-1 <sup>Q</sup> gene; Chr 19	<b>Early-onset</b> weakness; "floppy infant"; <b>Malignant hyperthermia</b> <sup>Q</sup> may be seen	<b>Cytoplasmic cores</b> are <b>eosinophilic</b> <sup>Q</sup> <b>Disrupted Sarcomeres</b> <sup>Q</sup> <b>Decreased mitochondria</b> <sup>Q</sup>
<b>Nemaline myopathy</b> (See fig 2)	AD or AR; <b>NEM gene</b>	Hypotonia at birth; "floppy infant"	Aggregates of <b>sub-sarcolemmal spindle-shaped particles</b> ('Nemaline rods') <sup>Q</sup>
<b>Myotubular (centronuclear) myopathy</b>	XL, AD or AR myotubularin ( <i>MTM1</i> ) gene	Severe congenital hypotonia, "floppy infant"; poor prognosis	Abundance of <b>centrally located nuclei</b> in <b>type I fibers</b> <sup>Q</sup>



**Fig. 1:** End stage muscle with atrophic fibres and fibrofatty replacement (H&E) in Duchenne muscular dystrophy



**Fig. 2:** Rod (nemaline) myopathy. A. Muscle fibers contain dark aggregates of rods (toluidine blue, 1000×)



## INFLAMMATORY MYOPATHIES

- Disorders include polymyositis (PM), dermatomyositis (DM)
- **Features**
  - Symmetric proximal muscle weakness
  - Increased serum levels of muscle derived enzymes
  - Nonsuppurative inflammation of skeletal muscle.

### Polymyositis (PM)

- Primarily occurs in persons aged 40 to 60 years.
- Increased risk of malignant neoplasms (15%–20% of cases), particularly lung and bladder cancer, and non-Hodgkin malignant lymphomas.
- **Laboratory findings**
- Antibody findings
  - Serum ANA increased in 30% to 60% of cases.
  - Anti-transfer RNA synthetase (Jo-1) antibodies increased in 25% of cases.
- Muscle biopsies show necrotic and regenerating muscle and a lymphocytic and macrophage infiltrate.
- Muscle atrophy is not a prominent feature

### Dermatomyositis (DM)

Cutaneous findings are key.

- Reddish-purple papules called Gottron patches are noted over the knuckles and proximal interphalangeal (PIP) joints in both hands




- Purple-red eyelid discoloration occurs (called heliotrope eyelids or “racoon eyes”.




### Laboratory Findings

- Similar to those described for Polymyositis (PM)
- Muscle biopsies show an inflammatory reaction (primarily lymphocytic).
  - Unlike PM, atrophy of muscle fibers is a **prominent feature**.
  - Damage to the capillaries in the muscle leads to ischemia and atrophy of the muscle fibers.




## Image-Based Questions

1. A 5-year-old male presented to AIIMS pediatrics OPD with a chief complaint of difficulty in climbing stairs and getting up from sitting position. There was history of maternal uncle having the same illness. On examination, there was pseudohypertrophy of calf muscle. Biopsy of the muscle was performed as shown below. What is your diagnosis?



- a. Duchenne muscular dystrophy
- b. Myotonic dystrophy
- c. Fascioscapulohumeral muscular dystrophy
- d. Nemaline myopathy

2. Electron microscopy of muscles was performed in infant who presented with hypotonia at birth. What is the most likely diagnosis?



- a. Duchenne muscular dystrophy
- b. Myotonic dystrophy
- c. Fascioscapulohumeral muscular dystrophy
- d. Nemaline myopathy



## Answers of Image-Based Questions

1. Ans. (a) **Duchenne muscular dystrophy**

- This is myopathy with X-linked inheritance suggested by same history in maternal uncle. On muscle biopsy prominent variation in size of muscle fibers can be seen.

2. Ans. (d) **Nemaline myopathy**

- This is a congenital myopathy suggested by clinical features since birth. Electron microscopy shows aggregates of **subsarcolemmal spindle-shaped particles** ('Nemaline rods')






## Multiple Choice Questions

### BONE LESIONS

1. X-ray foot shows lytic lesion in the calcaneus, biopsy done from the lesion shows the following. What is your diagnosis? (AIIMS Nov 18)



- Pigmented villonodular synovitis.
- Onchocosis
- Osteomyelitis
- Eumycosis

### MUSCULAR DYSTROPHY AND MYOSITIS

2. Absence of dystrophin and presence of small muscle fibers are seen in: (Recent Pattern Question 2020)
- Duchenne muscular dystrophy
  - Becker muscular dystrophy
  - Myotonic dystrophy
  - Dermatomyositis

3. Juvenile myoclonic epilepsy is due to mutation in: (JIPMER Nov 2019)

- GABRA 1
- CHRNA 2
- COL4A1
- FMRI

4. Parking lot inclusions are seen in:

- Mitochondrial myopathy (Recent Question 2015)
- Nemaline myopathy
- Central myopathy
- Lipid Myopathies

5. Perifascicular atrophy of muscle fibres is seen in-

(Recent Question 2014, 2013)

- Steroid myopathy
- Dermatomyositis
- Inclusion body myositis
- Nemaline myopathy

6. Dystrophin is lacking in:

(Recent Question 2013, AIIMS May 93, 03)

- Polio
- Duchenne's muscular dystrophy
- Peroneal muscular atrophy
- None of the above

7. Myasthenia gravis is associated with:

(Recent Question 2013)

- Thymoma
- Thymic carcinoma
- Thymic hyperplasia
- Lymphoma



## Answers with Explanations

1. Ans. (a) Pigmented villonodular synovitis


Definition: Rare neoplastic-like villonodular hyperplasia of synovium and tendon sheaths in young adults composed of mononuclear cells and multinuclear giant cells with hemosiderin deposition

- It Develops in synovial lining of joints, tendon sheaths and bursae, usually of knee (80%), ankle, hip, shoulder, elbow joint; nodular variant occurs in hands and wrists
- Almost always monoarticular
- Occasionally invades underlying bone - radiograph shows pressure erosions of the bones about the ankle, including the calcaneus, caused by large hyperdense, irregular soft-tissue mass. The bone erosions have well-defined, sclerotic margins.



### Microscopic (Histologic) Description

- Hyperplastic synovium with papillary projections composed of foamy cells and hemosiderin containing macrophages

- Also large clefts, pseudoglandular or alveolar spaces lined by synovial cells, multinucleated (10 - 70 nuclei) giant cells, epithelioid cells



The papillary and villous structures show proliferation of polygonal cells in a background of fibroconnective tissue which is covered by synovial lining



Hemosiderin  
laden  
macrophages

2. **Ans. (a) Duchenne muscular dystrophy**  
(Ref: Robbins 9th/pg 1274)

3. **Ans. (a) GABRA 1** (Ref: Robbins 9th/pg 1242)

4. **Ans. (a) Mitochondrial myopathy** (Ref: R 9th/pg 1274)

In mitochondrial diseases, electron micrograph showing morphologically abnormal mitochondria with concentric membranous rings ("phonograph records") & rhomboid paracrystalline inclusions ("parking lot" inclusions).

5. **Ans. (b) Dermatomyositis** (Ref: Robbins 9th/pg 1238)

6. **Ans. (b) Duchenne's muscular dystrophy**

(Ref: Robbins 9th/pg 1242-1243)

7. **Ans. (c) Thymic hyperplasia** (Ref: Robbins 9th/pg 1236)

- Thymic hyperplasia is found in 65% and thymoma in 15% of patients with myasthenia gravis.

# Tumors of Bone and Joints

## Key Points

- » Most common mutation seen in osteosarcoma is germ line mutation in **retinoblastoma gene**
- » **Ext** gene mutations are seen in osteochondroma
- » **Giant cells** are seen in osteoclastoma, osteosarcoma, chondroblastoma, fibrous dysplasia, non ossifying fibrous and chondromyxoid fibroma





## BONE TUMORS

### Bone-Forming Tumors

#### Osteoma

- **Most common** site: head and neck
- Multiple lesions are a feature of Gardner's syndrome
- Composed of a mixture of woven and lamellar bone.


#### Osteoid osteoma

- Nocturnal pain- relieved by aspirin
- Less than 2 cm in diameter

#### Osteoblastoma

- **Most common** site: vertebral column
- Pain- not responsive to aspirin
- More than 2 cm in diameter

#### Osteosarcoma



Lacy osteoid

- **Most common** site: metaphyseal region of the long bones of the extremities
- Shows osteoid (eosinophilic, glassy appearance) or bone produced directly by tumor cells without interposition of cartilage
- Metastatize via blood mainly to lungs
- **Most common** mutation: RB gene mutations (70%)

### Cartilage-Forming Tumors

#### Osteochondroma

- **Most common** site: metaphysis near the growth plate of long bones
- **Most common** solitary tumor
- Multiple autosomal dominant disorder due to inactivity of EXT1 or EXT2 genes

#### Chondroma

##### Within Medulla: Enchondromas

##### On Bone surface: juxtacortical chondromas

- Maffucci syndrome: multiple chondromas associated with soft tissue spindle cell, Hemangiomas


- Ollier disease: multiple chondromas involving one side of the body
- Both syndromes have point mutations in isocitrate dehydrogenase I (IDH1) or IDH2

#### Chondroblastoma


- Mic: mixture of mononuclear cells with oval nuclei and longitudinal groove.
- Chicken wire calcification

#### Chondrosarcoma


- **Most common** site: pelvis, shoulder and ribs
- Gross: glistening white tumor image shows glistening white tumor S/o chondrosarcoma
- Mic: cartilaginous matrix and lack of direct bone formation by tumor cells



Glistening white tumor



Chondrosarcoma showing malignant chondrocytes with moderate pleomorphism



## Fibro-osseous Tumors

### Fibrous Dysplasia





Image showing curved trabeculae

- **Most common** Mutation: GNAS gene
- **Mic:** curved trabeculae of woven bone (mimicking Chinese characters), surrounded by cellular fibroblastic proliferation

### Miscellaneous Bone Tumors


#### Osteoclastoma/ Giant cell tumor



Osteoclastoma/Giant cell tumor showing giant cells in background of oval to spindle shaped mononuclear cells

- **Most common** site: epiphyses of long bones
- Benign but locally aggressive bone tumor
- Neoplastic component: mononuclear cells
- Mononuclear cells → RANK ligand → proliferation of non-neoplastic osteoclast-like cells.

### Ewings tumor/ PNET



Ewings tumor/PNET showing small round cells with rosettes

- **Most common** Mutation:  $t(11;22)(q24;q12)$  in 95% cases or  $t(21;22)(q22;q12)$ .
- **Most common** site: diaphyses of long bones
- **Gross:** characteristic periosteal reaction with deposition of bone in onion-skin pattern.
- **Mic:** uniform small, round cells with scant glycogen-rich cytoplasm
- Homer-Wright rosettes suggests neural differentiation (PNET)

## JOINTS

### Joint Disorders

#### Classification of Joint Disorders

##### Group I

- Noninflammatory
- Examples—osteoarthritis (OA), neuropathic joint

##### Group II

- Inflammatory
- Examples—rheumatoid arthritis (RA), gout

##### Group III

- Septic
- Examples—Lyme disease, disseminated gonococcemia


##### Group IV

- Hemorrhagic
- Examples—trauma, hemophilia A and B

### Osteoarthritis (OA)

- Definition**—progressive degeneration of articular cartilage
- Targets weight-bearing joints





### Joint findings

- Erosion and clefts in articular cartilage
- Reactive bone formation occurs at the joint margins (osteophytes)
- Subchondral cysts
- Bone eventually rubs on bone - This produces dense, sclerotic bone.
- No ankylosis (fusion) of the joint
- Joint mice- Refers to fragments of articular cartilage that break free into the joint space

### Rheumatoid Arthritis (RA)

**Definition**—systemic disorder associated with chronic joint inflammation that most commonly affects peripheral joints

#### Clinical findings

- Symmetric involvement of second/third metacarpophalangeal (MCP) and PIP joints
  - Causes ulnar deviation, morning stiffness
- **Swan neck deformity**
  - Flexion of the DIP joint
  - Extension of the PIP joint
- **Boutonnière deformity**
  - Extension of the DIP joint
  - Flexion of the PIP joint

#### Laboratory findings

- Positive serum antinuclear antibody (ANA) test (30% of cases)
- Positive serum RF (70%–90% of cases)
- Normal to increased serum C3, decreased synovial C3
- Increased serum total protein
  - Due to increase in  $\gamma$ -globulins (IgG) in chronic inflammation
  - Polyclonal gammopathy on serum protein electrophoresis

### Gouty Arthritis


**Definition**—Tissue deposition of monosodium urate (MSU) due to prolonged hyperuricemia

- Most commonly involve the first metatarsophalangeal joint (MTP; called podagra; joint in the foot with the most trauma)

#### Acute gout

##### Laboratory findings

- Hyperuricemia
  - Increased serum uric acid >7 mg/dL in men
  - Increased serum uric acid >6 mg/dL in women
- Absolute neutrophilic leukocytosis
- Joint aspiration is confirmatory. Shows Negatively birefringent MSU crystals




Showing Negatively Birefringent Crystals

### Chronic gout

Distal joints are preferential sites.

UA crystals accumulate in the joint and produce a tophus.


- MSU crystals leak into the soft tissue around the joint
- MSU excites a brisk giant cell reaction in the periarticular tissue.-Microscopic sections reveal numerous multinucleated giant cells within which are MSU crystals that polarize.




Large aggregations of urate crystals surrounded by an intense foreign body giant cell reaction

- Tophi destroy subjacent bone, causing erosive arthritis that breaks down bone and leaves overhanging edges (sometimes called rat bites)

### Pseudogout -Calcium Pyrophosphate Dihydrate Deposition (CPPD) Disease






- Most common joint involved is the knee.
- Calcium pyrophosphate crystals deposit in articular cartilage
  - Crystals produce linear deposits in articular cartilage
  - It is called chondrocalcinosis when it deposits in articular cartilage.
- If the patient has acute pain, redness, swelling, and limitation of motion in the joint, the combination is called pseudogout.
- Crystals-rhomboid, positively birefringent
- The crystals form chalky, white friable deposits, which are seen histologically in hematoxylin and eosin stained preparations as oval blue-purple aggregates

## Joint Tumors

### Tenosynovial Giant Cell Tumor

- Develop in the synovial lining of joints, tendon sheaths, and bursae
- 2 types:
  - Diffuse tumors-Also called as pigmented villonodular synovitis (MC site-knee)
  - Localized type -also known as giant cell tumor of tendon (MC site-wrist) -it is the most common mesenchymal neoplasm of the hand
- Microscopically both show similar changes (Giant cells are more in local variant whereas hemosiderin laden macrophages are predominant in PVNS)

- Both variants show proliferating synoviocytes, macrophages, and may contain hemosiderin or foamy lipid. Scattered multinucleated giant cells and patchy fibrosis are commonly present.



Showing Pigmented Macrophage and Proliferating Synoviocytes suggestive of Pigmented villonodular synovitis




## NEXT Pattern Question



Q's

1. 30 Y/male presented painful enlarging mass in the wrist associated with tenderness. After CT amputation was done, cut section of femur is shown along with corresponding histopathological image. Identify type of tumor?




- a. Osteosarcoma      b. Aneurysmal bone cyst      c. Osteoclastoma      d. Pagets disease

Ans. (c) **Osteoclastoma**


- Painful enlarging mass with the lytic lesion in wrist joint. The histopath is suggestive of giant cells in with the mononuclear tumor cells, this is suggestive of Osteoclastoma.






## Image-Based Questions

1. Histology from a bone biopsy from proximal femur is given below. What is your diagnosis? (Recent exam 2018)



- a. Osteomalacia
- b. Paget's disease
- c. Osteoporosis
- d. Osteosclerosis

2. X-ray shows lytic lesion in calcaneus in young adult. Biopsy shows following diagnosis:



Hyperplasia of synovium and tendon sheaths  
Giant Cells  
Hemosiderin-laden macrophages.

- a. Pigmented villonodular synovitis
- b. Ochronosis
- c. Osteomyelitis
- d. Eumycosis



## Answers of Image-Based Questions

1. Ans. (b) **Paget's disease**

Robbins 9th/pg 1190

- Paget disease shows remarkable histologic variation over time and from site to site. The hallmark is a mosaic pattern of lamellar bone, seen in the sclerotic phase. This jigsaw puzzle-like appearance is produced by unusually

prominent cement lines, which join haphazardly oriented units of lamellar bone.

2. Ans. (a) **Pigmented villonodular synovitis**

Image shows hyperplasia of synovium along with hemosiderin laden macrophages and giant cells s/o Pigmented villonodular synovitis.



## Multiple Choice Questions

1. All are true about Paget's disease of bone except: (PGIMay2019)
- a. Initially woven pattern, but ultimately lamellar pattern
  - b. Mosaic pattern in final stage
  - c. Osteoclasts may have up to 100 nuclei
  - d. Fibrous connective tissue replaces bone marrow
  - e. Cortical thickening
2. A 25-year-old female presented with swelling around the knee joint. Biopsy showed giant cells interspersed with mononuclear cells. What is your diagnosis? (Recent Pattern Question 2020)
- a. Rheumatoid arthritis
  - b. Osteosarcoma
  - c. Aneurysmal bone cyst
  - d. Giant cell tumor



## Answers with Explanations

1. Ans. (a) **Initially woven pattern, but ultimately lamellar pattern** (Ref: Robbins 9th/pg 1190)
2. Ans. (d) **Giant cell tumor** (Ref: R9th pg 1187)

A composite image featuring a close-up of a microscope's objective lenses and a gloved hand holding a slide. The image is partially obscured by a large, stylized red and white geometric shape that resembles a folded ribbon or a stylized 'V' shape, which frames the top and left sides of the page.

26


# Recent Techniques in Pathology



## FLOW CYTOMETRY

- Flow cytometry provides rapid analysis of **multiple characteristics** of **single cells** made to **flow in a single line**.
- Expression of **several antigens** can be assessed simultaneously
- The usefulness of FCM immunophenotyping are:

Field	Clinical Application
<b>Immunology</b>	<ul style="list-style-type: none"> <li>Histocompatibility cross-matching</li> <li>Transplantation rejection</li> <li><b>HLA-B27 detection</b></li> <li>Immunodeficiency studies</li> </ul>
<b>Oncology</b>	<ul style="list-style-type: none"> <li><b>DNA content</b> and S phase of tumors</li> <li>Measurement of <b>proliferation markers</b></li> </ul>
<b>Hematology</b>	<ul style="list-style-type: none"> <li><b>Leukemia and lymphoma</b></li> <li>Anti-platelet antibodies</li> <li>Anti-neutrophil antibodies</li> <li><b>Fetomaternal hemorrhage</b> quantification</li> <li>PNH</li> </ul>
<b>Blood banking</b>	<ul style="list-style-type: none"> <li>Immunohematology</li> <li>Assessment of leukocyte contamination of blood products</li> </ul>
<b>Genetic disorders</b>	<ul style="list-style-type: none"> <li>Leukocyte adhesion deficiency</li> </ul>




## PANEL OF MARKERS FOR FLOW CYTOMETRY

Leukemia	ALL		AML
	B-Cell	T-Cell	
<b>First-line</b>	CD19, CD22, CD79a, CD20	CD7, CD2, cCD3	CD13, CD33, CD117, anti-MPO
	TdT, HLA-DR, CD34		
<b>Second-line</b>	cμ, Smlg	CD1a, CD5, CD4, CD8 anti-TCR	CD41, CD42, CD62p, antiglycophorin-A (CD235a), CD11c, CD64

## Immunological Classification of Acute Leukemias

Lymphoblastic leukaemia/lymphomas (ALL)- (TdT +)			
B-ALL	MARKERS	T-ALL	MARKERS
<b>B-cell precursor</b>	CD19+ and/or cCD79a+ and/or cCD22+	<b>T-cell precursor</b>	Cytoplasmic CD3+, CD7+
<b>Pro-B-ALL</b>	No expression of other B-cell markers	<b>Pro-T-ALL</b>	No expression of other T-cell markers
<b>Common-ALL</b>	CD10+, cytoplasmic μ-	<b>Pre-T-ALL</b>	CD2+ and/or CD5+
<b>Pre-B-ALL</b>	Cytoplasmic μ+	<b>Cortical T-ALL</b>	CD1a+
		<b>Mature T-ALL</b>	Membrane CD3+

Miscellaneous leukemia	Markers
<b>Mixed phenotype acute leukemias (MPAL)</b> (coexpression of myeloid and lymphoid markers)	<b>MYELOID COMPONENT:</b> anti-MPO/cytochemical MPO and/ or monocytic component <b>LYMPHOID COMPONENT:</b> <ul style="list-style-type: none"> <li><b>B-lymphoid Differentiation:</b> <ul style="list-style-type: none"> <li><b>Strong CD19</b> plus B-cell marker (CD10, CD22 or CD79)</li> <li><b>Weak/negative CD19</b> and strong expression of two of the specified B-cell markers.</li> </ul> </li> <li><b>T-lymphoid Component-</b> CD3 whether cytoplasmic or membrane.</li> </ul>
<b>Myeloid antigen positive ALL</b>	<ul style="list-style-type: none"> <li>Leukemia with aberrant markers</li> </ul>
<b>Lymphoid antigen positive AML</b>	





## Common Phenotypes of B-cell Lymphoproliferative Disorders

Diagnosis	CD5	CD10	CD19	CD20	CD23	CD79b	FMC-7	CD25	CD11c	CD103
SLL/CLL*	+	–	+	+(W)	++	–	–	–/+	+/-	–
Mantle cell lymphoma	+	–	+	+	–	+	+	–	–	–
Follicle center lymphoma	–	+	+	+	–/+	+/-	+/-	–	–	–
Marginal zone lymphoma	–	–	+	+	–	+/-	+/-	–/+	+	–
Hairy cell leukemia	–	–	+	+	–	+/-	+/-	+	++	++

\*SLL/CLL small lymphocytic lymphoma/chronic lymphocytic leukemia: +, positive; –, negative; +/-, occasionally positive; w. weak; Red color indicates most important marker for the entity

## HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)

- A technique for separating mixtures into their components in order to **analyze**, **identify**, and **purify** the mixture or components.



### Important Terminologies used in HPLC

- Retention time:** Time taken for analyte to pass through the system under set conditions.
- Stationary Phase:** Fixed in place either in a column or on a planar surface, acts as a adsorbent (atoms that accumulate on the surface of the material)
- Mobile Phase:** Carries the analyte through the stationary phase, acts as eluent (**solvent that carry the analyte in elution**)

- Eluent:** Substance used as a solvent in elution
- Eluate:** Solution of the solvent and the substance that was adsorbed to another

### Applications

- Detection of hemoglobin abnormalities
- Toxicology
- Therapeutic and overdose drug monitoring
- HbA<sub>1c</sub> levels in DM monitoring
- Protein analysis (Transferrin)
- Nucleic acid sequencing (as replacement to PCR in HLA typing)




### High Yield Facts

- Normal Adult Hb %:
  - HbA: 96.5-97.5%
  - HbA<sub>2</sub>: 2.5-3.5%
  - HbF: <1%
- HbA<sub>2</sub>>4% suggests β-thalassemia trait
- HbA<sub>2</sub>>8% suggests HbE
- HbA<sub>2</sub>, HbE and LEPORE **co elute**.

### Hb HPLC

**Principle:** Depends on the interchange of **charged groups** on resin with charged groups on the hemoglobin molecule. Each hemoglobin variant has a specific elution time, which should always be matched with controls







## IMMUNOFLUORESCENCE

A technique that utilizes **fluorescent-labeled antibodies** to detect specific target antigens.

### Types


- **Direct Immunofluorescence**
  - Uses **fluorescent-tagged antibodies** to bind directly to the target antigen.
  - DIF techniques can also be used to detect non-antibody targets in the skin, such as infectious organisms. In this case, a fluorophore-labeled primary antibody directed against the suspected antigen is used to detect the presence or absence of the organism
- **Indirect Immunofluorescence:** A primary, unlabeled antibody binds to the target, after which a fluorophore-labeled second antibody (directed against the Fc portion of the primary antibody) is used to detect the first antibody.

### Purpose

To detect circulating autoantibodies

### Fluorescence Detection

It is based on the use of fluorochromes that emit light when excited by light of a shorter wavelength.





## NUCLEIC ACID TEST/NUCLEIC ACID AMPLIFICATION TEST/"NAAT"

It is a molecular technique used to detect a **virus or a bacterium**. These tests were developed to **shorten the window period**. The term includes any test that directly detects the genetic material of the infecting organism or virus.

### Polymerase Chain Reaction

- PCR was invented by **Kary Mullis** in 1983.
- He shared the **Nobel Prize** in chemistry with Michael Smith in 1993.





## Principle

- **DNA polymerase** adds nucleotides to the 3' end designed **primer (oligonucleotide)** when it is **annealed** to target template DNA.
- Thus, if primer is annealed to a single-stranded template that contains a region complementary to the oligonucleotide, DNA polymerase can use the oligonucleotide as a primer and **elongate its 3' end** to generate an extended region of **double stranded DNA**.

## Types: 2 main Types

- **Real Time PCR:** Quantitative estimation of sample (viral load, DNA, RNA etc)
- **Reverse Transcriptase PCR:** Diagnosis of genetic disorders & semi-quantitatively calculation of specific expression level of particular RNA.

**Reaction Requires:** Template DNA, target sequence, specific primers, mixture of dNTPs (Deoxynucleotides), and heat-stable *Taq* DNA polymerase (&  $Mg^{2+}$  for its activity).

### Taq Polymerase:

- Thermostable DNA polymerase named after the **thermo-philic bacterium**
- DNA polymerase from *E. coli* originally used in PCR

## FLUORESCENCE IN SITU HYBRIDIZATION (FISH)


- Cytogenetic technique that uses fluorescent probes binding to only **complementary sequences** of parts of the chromosome.
- Used to **detect and localize** the presence or absence of specific DNA sequences on chromosomes.
- FISH can also be used to detect and localize specific RNA targets (**mRNA, lncRNA and miRNA**) in cells, circulating tumor cells, and tissue samples. In this context, it can help define the **spatial-temporal patterns of gene expression** within cells and tissues.


### 1. Flow cytometry is used to detect? (JIPMER 2017)

- Antibody response
- To get differential leukocyte count
- T lymphocytes types**
- To separate blood cells

## Uses of FISH

Diagnostic	Research
Identification of specific chromosome abnormalities	Identification of new non-random abnormalities
The characterization of marker chromosomes	Gene mapping
Interphase FISH for specific abnormalities in cases of failed cytogenetics	Identification of regions of amplification or deletion by CGH
Monitoring disease progression	Identification of translocation breakpoints
Monitoring the success of bone marrow transplantation	Study of 3D chromosome organization in interphase nuclei






## Image-Based Questions

### 1. Forward scatter in Flowcytometry indicates?


(AIIMS May 2015)



- a. Cell Size  
b. Nucleus  
c. Granularity  
d. DNA content

### 2. Identify the HPLC pattern:


(Recent Question 2016)



- a.  $\beta$ -thalassemia trait  
b.  $\beta$ -thalassemia major  
c. HbE disease  
d. None

### 3. The given technique is very important in the diagnostic field. Which of the following is an important component for viewing by this technique?

(AIIMS May 2015)



- a. Cathode Ray tube  
b. Dichroic tube  
c. Dark field/phase condenser  
d. Phase plate



## Answers of Image-Based Questions

#### 1. Ans. (a) Cell Size

On flow cytometry

- Forward scatter-cell size
- Side scatter-granularity

#### 2. Ans. (a) $\beta$ -thalassemia trait-HbA2 >4%


#### 3. Ans. (b) Dichroic tube



A close-up photograph of a laboratory setting. A gloved hand in a blue nitrile glove is holding a small, light blue rectangular object, possibly a slide or a piece of tissue, near a microscope. The microscope's objective lenses are visible, and the background is blurred, showing other laboratory equipment. A large, stylized red and white geometric shape is overlaid on the left side of the image.

27

# Stains and Fixatives



## Multiple Choice Questions

### STAINS & FIXATIVES

1. Match the following viral intracellular bodies with respect to the disease? (AIIMS Nov 2019)

#### Column A

- HPV
- CMV
- Molluscum contagiosum
- Polyomavirus


#### Column B

- Henderson-Paterson bodies
- Decoy cells
- Owl eye inclusion
- Koilocyte

2. Which of the following stain is used for Acidic mucin? (AIIMS May 18)

- Alcian blue
- PAS
- Masson's trichrome stain
- PTAH

3. Which of the following labels corresponds to the condenser of the microscope? (AIIMS May 18)



- C
- B
- D
- A

4. Which of the following are false about electron microscopy? (PGI May 18)


- High resolving power
- Can see both live and dead objects
- Vacuum is required
- Three-dimensional images can be obtained
- Coloured images are seen

5. Microscopy which can be performed with minimum optical illumination? (PGI May 18)

- Dark field
- Phase contrast
- Bright-field microscopy
- Fluorescent microscopy
- Inverted microscopy

6. Toluidine blue staining is used for identification of?

(AIIMS May 18)




- Mast cell
- Fibroblast
- Melanocyte
- Macrophages

7. Best method for HbA1c estimation is? (AIIMS May 18)


- Affinity chromatography
- Ion exchange chromatography
- Electrophoresis
- HPLC

8. Gomori methenamine silver stain for fungus is shown below. Most likely diagnosis is? (AIIMS Nov 2017)




- Acute angle branching with septate hyphae - Aspergillus
- Right angle branching and aseptate hyphae - Mucor
- Acute angle branching with septate hyphae - Mucor
- Right angle branching and aseptate hyphae - Aspergillus

9. Identify the stain done on a section of liver shown below? (AIIMS May 2017)




- Sweets reticulin stain
- Gremilus silver stain
- Warthin starry silver stain
- Steiner silver stain



10. For detection of carcinoma lip, stain used is? (AIIMS Nov 2016)
- Giemsa
  - Crystal violet
  - Toulidine blue
  - Hematoxylin and eosin

11. Given below is the histopathology of liver biopsy of hemochromatosis. Which of the following stain is used? (AIIMS Nov 2016)



- Von kossa
  - Alcian blue
  - Prussian blue
  - Crystal violet
12. Which of the following is a correct match? (PGI Nov 2016)
- Perl stain-Iron
  - Von Gieson-Collagen
  - Mason Trichrome-elastin
  - PAS-glycogen
  - PAS- Acidic and neutral mucin
13. Which of the following stain and the material stained by it is correct? (PGI Nov 2016)
- Perl stain-Iron
  - Collagen- Von kossa
  - Elastin-Von Geison
  - Copper-modified rhodamine
  - Oil red O - Glycogen
14. Toluidine blue has been established as a diagnostic adjunct in detecting oral lesions related to invasive carcinomas, carcinoma in situ or early asymptomatic oral carcinomas Stain used for hemachromatosis (Recent Question 2016-17)
- Prussian blue stain
  - Von Kossa
  - Sudan black
  - Methenamine silver
15. Oil red O stain is used for? (AIIMS May 2015)
- Frozen specimen
  - Glutaraldehyde fixed specimen
  - Alcohol fixed specimen
  - Formalin fixed specimen
16. Which of the following cellular component gives purplish blue color with H & E reagent: (PGI May 2015)
- Reticulum
  - Elastin
  - P-selectin
  - Collagen
  - Heterochromatin
17. Which of the following stain is used for staining of glycogen? (Recent Question 2016)
- PAS
  - Oli Red -O
  - Sudan black
  - Von kossa
18. Perl' stain is for: (Recent Question 2015)
- Iron
  - Copper
  - Melanin
  - Glycogen

19. Stain used for copper: (Recent Question 2015)
- Congo red
  - Prussian blue
  - PAS
  - Rubeanic acid

20. Tissues for electron microscopy are fixed in: (Recent Question 2015)

- Carnoy's fixative
- 10% buffered formalin
- 4% gluteraldehyde
- 50% glycerine

21. Frozen section was introduced by: (Recent Question 2015)

- Virchow
- Feulgen
- Morgagni
- Cohnheim

22. Stain for collagen: (Recent Question 2015)

- Von gieson
- Von kossa
- Alizarin red
- Rubeanic acid

23. Stain for axons: (Recent Question 2015)

- Phosphotungstic acid-Hematoxylin (PTAH)
- Luxol fast blue
- Bileschowsky's silver
- Masson fontana

24. Stain for hepatitis B surface antigen: (Recent Question 2015)

- Fite-wade
- Grocott's silver methanamine
- Shiata's orciein
- Grimelius

25. Histopathological specimens are commonly preserved in: (Recent Question 2015)

- 10%formalin
- 4% gluteraldehyde
- Rectified spirit
- Saturated saline solution

26. PTAH stain is used for staining: (Recent Question 2015)

- Myelin
- Axon
- Muscle and glial filamets
- Melanin

27. Heart failure cells can be stained by: (Recent Question 2015)

- PAS
- Prussian blue
- Congo red
- Gram stain

28. Gauge of commonly used FNAC needle is? (Recent Question 2015)

- 26-29
- 22-26
- 18-22
- 16-18

29. Ocular basement membrane is stained by: (Recent Question 2015)

- Alcian blue
- PAS
- Methylene blue
- Geimsa stain

30. Acid mucin is best demonstrated by the stain: (Recent Question 2015)

- Alcian blue
- Periodic Acid Schiff (PAS)
- Van Giesen
- Reticulin

31. The fixative used in histopathology: (Recent Question 2014; AIIMS May 12)


- 10% buffered neutral formalin
- Bouins fixative
- Glutaraldehyde
- Ethyl alcohol





- 32. Stain used for glycogen:** (Recent Question 2014)  
 a. PAS                                      b. Congo red  
 c. Prussian blue                      d. Alician blue
- 33. Which of the following statement is TRUE?**  
 a. Gomori Methamine silver stain stains fungi green  
 b. Gram +ve stains bacteria black (AIIMS May 2014)  
 c. Gram -ve stains red  
 d. Calcoflor stains red in colour
- 34. Most common fixative used in electron microscopy:**  
 (AIIMS May 2013, Nov 2012)  
 a. Glutaraldehyde                      b. Formalin  
 c. Picric acid                              d. Absolute alcohol
- 35. Staining done for sebaceous cell carcinoma:**  
 a. Oil Red O                              b. PAS (AIIMS May 2013)  
 c. Methamine silver                      d. KOH
- 36. Liver in hemochromatosis is stained by:**  
 (Recent Question 2013)  
 a. Masson Fontana                      b. Prussian blue  
 c. Masson trichrome                      d. Congo red
- 37. Stain used for melanin:** (Recent Question 2013)  
 a. Masson Fontana                      b. Prussian blue  
 c. Masson trichrome                      d. Congo red
- 38. Which of the following is a negative stain:**  
 (Recent Question 2013)  
 α. Negrosin                              β. Fonatana  
 χ. ZN stain                              δ. Albert stain
- 39. The most common fixative used in pathology is?**  
 a. Gluteraldehyde                      b. Alcohol (AI 11)  
 c. Formaldehyde                      d. Picric acid
- 40. Fixative used for histopathology is?** (DNB Dec 11)  
 a. 10%formalin                      b. Normal saline  
 c. Rectified spirit                      d. 100% alcohol

- 41. Resolving power of a light microscope does not depend on?** (AIIMS Nov 2016)  
 a. Power of eyepiece  
 b. Wavelength of light used  
 c. Power of the lens  
 d. Thickness of the specimen
- 42. The following image is taken from a Neubauer chamber after charging of fluid. If the dilution factor is 20, what is the total cell count per cu.mm?**  
 (Recent Question 2016-17; AIIMS Nov 2016)



- a. 6000  
 b. 3000  
 c. 7000  
 d. 1100



## Answers with Explanations


1. Ans. (a) 4, (b) 3, (c) 1, (d) 2

### Viral Inclusion Bodies

Intracytoplasmic	Henderson-Peterson bodies (Molluscum Contagiosum)		
	Negri bodies (Rabies)		
	Guarnieri bodies (Smallpox)		
	Paschen bodies (Smallpox)		
	Bollinger bodies (Fowl pox)		
Intranuclear	Borrel bodies (Fowl pox)		
	Acidophilic	Cowdry type A	Varicella zoster virus
			Herpes simplex virus
			Yellow fever virus
	Basophilic	Cowdry type B	Polio virus
			Adeno virus
			Cytomegalo virus

2. Ans. (a) Alcian blue

3. Ans. (a) C



4. Ans. (b, e); b. Can see both live and dead objects; e. Coloured images are seen

5. Ans. (a) Dark field

Darkfield microscopy reduces the amount of light entering the lens system of a microscope in two ways. First, it blocks the center of the beam of light that would otherwise fill the objective lens. Second, only the light which is scattered by the specimen and enters the objective lens is seen

6. Ans. (a) Mast cell

Toluidine blue (also known as tolonium chloride) is an acidophilic metachromatic dye that selectively stains acidic tissue components (sulfates, carboxylates, and phosphate radicals), has an affinity for nucleic acids, and therefore binds to nuclear material of tissues with a high DNA and RNA content. Mast cell granules stain purple in color due to the presence of heparin and histamine.

7. Ans. (d) HPLC

Four basic types of methods are used most commonly to measure HbA1c: immunoassay, ion-exchange high-performance liquid chromatography (HPLC), boronate affinity HPLC, and enzymatic assays. The gold standard is HPLC method

8. Ans. (a) Acute angle branching with septate hyphae - Aspergillus

Mucor- Broad base with obtuse branching  
Aspergillus- Acute branching, Septate hyphae with narrow base

9. Ans. (a) Sweets reticulin stain

(Ref: Bancroft histochemical techniques. p. 180)

**Reticulin stain:** Demonstrates reticular fibers and basement membrane material. Reticular fibers are thin, usually type III collagen, widespread in connective tissue throughout the body. Staining procedures are Gordon and Sweet's reticulin stain (mc) & Gomori reticulin stain

10. Ans. (c) Toulidine blue

(Ref: Early Diagnosis and Treatment of Cancer Series: Head and Neck Cancers, Wayne Koch Pg. 54)

- Toluidine blue stain is used as a marker to **differentiate lesions at high risk of progression** in order to improve early diagnosis of oropharyngeal carcinomas.
- Toluidine blue, an **acidophilic metachromatic dye** of **thiazine group** selectively stains acidic tissue components (sulfates, carboxylates and phosphate radicals), thus staining DNA and RNA.

11. Ans. (c) Prussian blue


(Ref: Bancroft's histological techniques, 7th ed)

12. Ans. (a) Perl stain -Iron, (d) PAS-glycogen

(Ref: Bancroft's histological techniques, 7th ed)

Note:

- Verhoff-Von gieson stain is for elastin fibres
- Alcian blue is for differentiating Acidic and neutral mucin
- Masson trichrome is for collagen



13. Ans. (a) **Perl stain-Iron**, (c) **Elastin-Von Geison**, (d) **Copper-modified rhodamine**, (e) **Oil red O - Glycogen**

(Ref: Bancroft's histological techniques, 7th ed)

14. Ans. (a) **Prussian blue stain**

(Ref: Bancroft's histological techniques, 7th ed)

15. Ans. (a) **Frozen specimen**

(Ref: Bancroft's histological techniques, 7th ed)

16. Ans. (a, e) **a. Reticulum; e. Heterochromatin**

(Ref: Bancroft's histological techniques, 7th ed)

#### H&E Stain (Hematoxylin and Eosin)

- Hematoxylin, a natural dye product, acts as a basic dye that stains blue or black.
- Nuclear heterochromatin stains blue and the cytoplasm of cells rich in ribonucleoprotein also stains blue.
- The cytoplasm of cells with minimal amounts of ribonucleoprotein tends to be lavender in color.
- The aniline dye, eosin, is an acid dye that stains cytoplasm, muscle, and connective tissues various shades of pink and orange.
- This difference in staining intensity is useful in differentiating one tissue from another.

17. Ans. (a) **PAS** (Ref: Bancroft's staining 7th ed)

18. Ans. (a) **Iron** (Ref: Bancroft's staining 7th ed)

19. Ans. (d) **Rubeanic acid** (Ref: Bancroft's staining 7th ed)

20. Ans. (c) **4% glutaraldehyde**

(Ref: Bancroft's staining 7th ed)

21. Ans. (d) **Cohnheim** (Ref: Bancroft's staining 7th ed)

22. Ans. (a) **Von gieson** (Ref: Bancroft's staining 7th ed)

23. Ans. (c) **Bileschowsky's silver**

(Ref: Bancroft's staining 7th ed)

24. Ans. (c) **Shiata'sorciein** (Ref: Bancroft's staining 7th ed)

25. Ans. (a) **10% formalin** (Ref: Bancroft's staining 7th ed)

26. Ans. (c) **Muscle and glial filaments**

(Ref: Bancroft's staining 7th ed)

27. Ans. (b) **Prussian blue** (Ref: Bancroft's staining 7th ed)

28. Ans. (b) **22-26**

(Ref: Bancroft's histological techniques, 7th edition)

- FNAC is **Fine-Needle Aspiration Cytology** or Needle aspiration biopsy.
- A needle attached to a syringe is used to **collect cells from lesions or masses** in various body organs by microcoring, often with the application of **negative pressure (suction)** to increase yield.

- FNAC can be performed **under palpation or imaging guidance** (USG or CT scan)
- For FNAC, **Fine needles of 23 to 27 gauge** are used; most commonly used is a **25-gauge needle**.
- Core needle biopsy (CNB) is increasingly replacing FNAC because of the **inability of FNAC to distinguish carcinoma in-situ from invasive carcinoma**

29. Ans. (b) **PAS** (Ref: Bancroft's staining 7th ed)

PAS is used to stain **carbohydrates (polysaccharides)**, **neutral mucus (glycoproteins and glycolipids)**, **tissue basement membrane**, **fungal cell wall**.

30. Ans. (a) **Alcian blue** (Ref: Bancroft's staining 7th ed)

31. Ans. (a) **10% buffered neutral formalin**

(Ref: Bancroft's histological techniques, 7th edition)

32. Ans. (a) **PAS**

(Ref: Bancroft's histological techniques, 7th edition)

33. Ans. (c) **Gram -ve stains red**

(Ref: Bancroft's histological techniques, 7th edition)

#### Discussing the options one by one:

- False, as it stains fungi black, against a green background
- False, as Gram +ve stains bacteria blue
- True
- False, as it stains Acanthamoeba white & not red

34. Ans. (a) **Glutaraldehyde**

(Ref: Bancroft's histological techniques, 7th edition)

35. Ans. (a) **Oil Red O**

(Ref: Bancroft's histological techniques, 7th edition)

Oil red O stain, Sudan III, Sudan IV & Sudan black are used to stain Fat, seen in sebaceous cell carcinoma

36. Ans. (b) **Prussian blue**

(Ref: Bancroft's histological techniques, 7th edition)

**Iron** accumulates in Liver in **hemochromatosis**

37. Ans. (a) **Masson Fontana**

(Ref: Bancroft's histological techniques, 7th edition)

38. Ans. (a) **Negrosin**

(Ref: Bancroft's histological techniques, 7th edition)


Examples of **negative stain** are:

- **Negrosin:** Stains bacteria, Cryptococcus, test for viability (sperms)
- **India Ink:** Stains Cryptococcus

39. Ans. (c) **Formaldehyde**

(Ref: Bancroft's histological techniques, 7th edition)





40. Ans. (a) **10% formalin**

(Ref: Bancroft's histological techniques, 7th edition)

41. Ans. (d) **Thickness of the specimen**

(Ref: Laboratory Diagnosis of Infectious Diseases: Essentials of Diagnostic Pg. 130)

- **Resolving power:** It is defined as the **inverse of the distance or angular separation between two objects** which can be just resolved when viewed through the optical instrument.
- Resolving power of a microscope:
- For microscopes, the resolving power is the **inverse of the distance between two objects that can be just resolved**. This is given by the famous Abbe's criterion given by Ernst Abbe in 1873 as

$$\Delta d = \frac{\lambda}{2n \sin \theta}$$

$$\text{Resolving power} = \frac{\lambda}{\Delta d} = \frac{2n \sin \theta}{\lambda}$$

- Where n is the refractive index of the medium separating object and aperture. Note that to achieve high resolution  $n \sin \theta$  must be large. **This is known as the Numerical aperture.**


Thus, for good resolution:

- **$\sin \theta$  must be large.**
- To achieve this, the objective lens is kept as close to the specimen as possible.
- A **higher refractive index (n)** medium must be used. Oil immersion microscopes use oil to increase the refractive index.
- **Decreasing the wavelength** by using X-rays and gamma rays. While these techniques are used to study inorganic crystals, biological samples are usually damaged by x-rays and hence are not used.

42. Ans. (a) **6000**

(Ref: Text book of practical physiology GK Pal. Pg. 63)

Note the squares:



- Each of the squares (a to i) has a dimension of 1mm x 1mm
- When you keep a cover slip over this chamber, the depth is 0.1mm
- So the total volume is  $1 \times 1 \times 0.1 \text{ mm}^3 = 0.1 \text{ mm}^3$  or  $\mu\text{L}$
- Calculation:
- Concentration of cells =  $n/v \times d$
- Where ( $n$  = number of cells,  $V$  = volume,  $d$  = dilution)
- Coming back to the question: total number of cells in the 4 squares (a, c, g, i) =  $35 + 25 + 40 + 20 = 120$
- Volume of 4 squares =  $4 \times 0.1 = 0.4 \text{ mm}^3$
- So concentration =  $120 / 0.4 \times 20 = 6000 / \text{mm}^3$  or  $\mu\text{L}$

[illegible]This image shows a single sheet of white paper with horizontal blue or grey ruling lines, typical of notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



# it's here!



## Log in Now

And get ready to experience a smart new way of buying medical Books.

Visit **[www.prepshop.in](http://www.prepshop.in)** to avail amazing discounts, special offers, deals and much more...



## From the Authors' Desk

*"Give them Quality. That's the best kind of advertising in the World."*

We firmly believe in the above-mentioned quote that Quality does not need constant advertisement; it will be acknowledged anyhow. We are extremely grateful to you all for an overwhelming response for the 5th edition of the book, Complete Review of Pathology.

Join Author's Facebook Discussion Group – [www.facebook.com/Pathology by Dr Praveen](http://www.facebook.com/Pathology by Dr Praveen)

Join Author's Facebook Discussion Group – [www.facebook.com/Dr Vandana Pathology study circle](http://www.facebook.com/Dr Vandana Pathology study circle)

## Students' Reviews

This book is a concise version of whole Robbin's book which makes the job easier for students not only preparing for PG Entrance exam but also for those who write university theory exams. Most of the important points are either tabulated or given in flow chart which makes easier for me to revise at the last moment. I wonder how ma'am managed to include every single point in Robbin's yet in a concise manner. I recommend every single 2nd year juniors to have this book in hand so that you can attend every single question in university exam and fetch honors in the exams.

Raghunandan R



Main factor that stands out in the book is the mention of key recent updates at the start of each chapter. Those are pointers that are often missed by students who end up missing out on those. Extra attention is drawn. Presence of image-based questions at the end of each chapter is extremely useful and helps students to be prepared of all kinds of new question pattern being asked recently. Mnemonics are mentioned wherever possible.

Janvi Javia



This book comprises well illustrated diagrams and microscopic slides as well. This book is well conceptualized and one of the few to-the-point oriented book which comprises boxes and charts for better understanding of the subject. The book has range of MCQs which is very helpful on the PG point of view. If you are preparing for your PG entrance exams this is the must have book.

Shubham



Any book of pathology can't be a substitute for Robbin's if you are in 2nd proff. But to revise Robbin's and understand histopathology you must need a book to guide you, that what is important and what is not. So this book just solves all the problems and also have contents taken from Harrison and Robbin's which are must for PG entrance. Tables and diagrams are very helpful. Annexures are also very important. Separate chapters on transfusion medicine and stains make this book unique. According to me, read Robbins in 2nd professional but read this book also side by side to make difficult concepts very easy. It is a must buy book for NEET and AIIMS (LOTS OF IMAGE-BASED QUESTIONS).

Aman Kashyap



Book is truly helpful for PG exams. I found it very helpful during my practice sessions. Please try to go through the book. I'm sure it will be very useful. Thanks to the Publisher and the Author

Arghdip Dey



Mam really liked your book and way of presentation... I got PGI Rank 96. I took anaesthesia... really liked your way of telling the concept. Fortunately I read your book. Thanks Mam.

Dinesh Prathick



Awesome book. The only downside if any is that the content is not exactly catered to just NEET PG, and tends to focus more on Central Institutes and is an excellent book for the same. There are loads of high-yield boxes and tables, appendices too. Not to mention the vibrant colors overall, which is great for visual learners. My only regret is not buying it earlier on, in prep as Patho being a very expansive subject, its MCQ section alone takes a huge chunk of time to complete satisfactorily.

Anonymous



The features that I liked in this book are:

1. Annexures—Perfect for last minute revision.
2. All recent exam questions included.
3. AIIMS new pattern model questions and sample video questions included.
4. Covers ample image-based questions.
5. Point wise concise theory with all recent updates provided.

Bhadra



**CBS Publishers & Distributors Pvt. Ltd.**

4819/XI, Prahlad Street, 24 Ansari Road, Daryaganj, New Delhi 110 002, India

E-mail: [delhi@cbspd.com](mailto:delhi@cbspd.com), [cbspubs@airtelmail.in](mailto:cbspubs@airtelmail.in); Website: [www.cbspd.com](http://www.cbspd.com)

New Delhi | Bengaluru | Chennai | Kochi | Kolkata | Mumbai | Pune

Hyderabad | Nagpur | Patna | Vijayawada

ISBN: 978-81-945783-3-8



9 788194 578338